

A REVERSE MULHOLLAND-TYPE INEQUALITY IN THE WHOLE PLANE WITH MULTI-PARAMETERS

Michael Th. Rassias and Bicheng Yang*

By introducing multi-parameters, and applying weight coefficients, we prove a reverse Mulholland-type inequality in the whole plane with a best possible constant factor. Moreover, the equivalent forms and a few particular cases are also considered.

1. INTRODUCTION

Let $p > 1$, $\frac{1}{p} + \frac{1}{q} = 1$, $a_m, b_n \geq 0$,

$$0 < \sum_{m=1}^{\infty} a_m^p < \infty \text{ and } 0 < \sum_{n=1}^{\infty} b_n^q < \infty.$$

We have the following Hardy-Hilbert inequality (cf. [1]):

$$(1.0.1) \quad \sum_{n=1}^{\infty} \sum_{m=1}^{\infty} \frac{a_m b_n}{m+n} < \frac{\pi}{\sin(\pi/p)} \left(\sum_{m=1}^{\infty} a_m^p \right)^{1/p} \left(\sum_{n=1}^{\infty} b_n^q \right)^{1/q},$$

with the best possible constant factor $\frac{\pi}{\sin(\pi/p)}$. The Mulholland inequality with the same best possible constant factor $\frac{\pi}{\sin(\pi/p)}$ was provided as follows (cf. Theorem

*Corresponding author. Michael Th. Rassias

2010 Mathematics Subject Classification. 26D15 47A07.

Keywords and Phrases. Mulholland-type inequality, parameter, weight coefficient, equivalent form, reverse.

343 of [1], replacing $\frac{a_m}{m}, \frac{b_n}{n}$ by a_m, b_n):

$$(1.0.2) \quad \sum_{n=2}^{\infty} \sum_{m=2}^{\infty} \frac{a_m b_n}{\ln mn} < \frac{\pi}{\sin(\pi/p)} \left(\sum_{m=2}^{\infty} \frac{a_m^p}{m^{1-p}} \right)^{1/p} \left(\sum_{n=2}^{\infty} \frac{b_n^q}{n^{1-q}} \right)^{1/q}.$$

The inequalities (1.0.1)-(1.0.2) are important in the field of analysis and its applications (cf. [1], [2]).

In 2007, the following Hilbert-type integral inequality in the whole plane was presented (cf. [3]):

$$(1.0.3) \quad \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} \frac{f(x)g(y)}{(1+e^{x+y})^{\lambda}} dx dy \\ < B\left(\frac{\lambda}{2}, \frac{\lambda}{2}\right) \left(\int_{-\infty}^{\infty} e^{-\lambda x} f^2(x) dx \int_{-\infty}^{\infty} e^{-\lambda y} g^2(y) dy \right)^{\frac{1}{2}},$$

where, the constant factor $B\left(\frac{\lambda}{2}, \frac{\lambda}{2}\right)$ ($\lambda > 0, B(u, v)$ is the Beta function) is the best possible. Some extensions of (1.0.1)-(1.0.3) as well as other types of inequalities were established in [4]-[29].

In 2016, Yang and Chen [30] obtained the following extension of (1.0.1) in the whole plane with multi-parameters:

$$(1.0.4) \quad \sum_{|n|=1}^{\infty} \sum_{|m|=1}^{\infty} \frac{a_m b_n}{(|m - \xi| + |n - \eta|)^{\lambda}} < 2B(\lambda_1, \lambda_2) \\ \times \left[\sum_{|m|=1}^{\infty} |m - \xi|^{p(1-\lambda_1)-1} a_m^p \right]^{\frac{1}{p}} \left[\sum_{|n|=1}^{\infty} |n - \eta|^{q(1-\lambda_2)-1} b_n^q \right]^{\frac{1}{q}},$$

where the constant factor $2B(\lambda_1, \lambda_2)$ ($0 < \lambda_1, \lambda_2 \leq 1, \lambda_1 + \lambda_2 = \lambda, \xi, \eta \in [0, \frac{1}{2}]$) is the best possible. Another result on this kind of inequalities was provided by Xin et al. [31].

In the present paper, by introducing multi-parameters and applying weight coefficients, we obtain a reverse Mulholland-type inequality in the whole plane with a best possible constant factor similar to the reverse of (1.0.4). Moreover, the equivalent forms and a few particular cases are also considered.

2. SOME LEMMAS

In the sequel, we shall assume that $0 < p < 1$ ($q < 0$), $\frac{1}{p} + \frac{1}{q} = 1$, $\sigma \in \mathbf{R} = (-\infty, \infty)$, $-\sigma < \lambda_1, \lambda_2 \leq 1 - \sigma$, $\lambda_1 + \lambda_2 = \lambda$, $\rho \geq 1$,

$$(2.0.5) \quad \arccos \sqrt{1 - \frac{1}{\rho}} \leq \gamma \leq \pi - \arccos \sqrt{1 - \frac{1}{\rho}} (\gamma = \alpha, \beta),$$

and $\zeta \in (-1, 1)$, satisfying

$$(2.0.6) \quad \frac{1}{\rho(1 - \cos \gamma)} - 1 \leq \zeta \leq 1 - \frac{1}{\rho(1 + \cos \gamma)}$$

$((\zeta, \gamma) = (\xi, \alpha) \text{ or } (\eta, \beta))$. In particular, for $\rho = 1$, it follows that $\alpha = \beta = \frac{\pi}{2}$ and $\xi = \eta = 0$.

Remark 1. *In view of (2.0.6) we have*

$$\rho(1 \pm \zeta)(1 \mp \cos \gamma) \geq 1, (\zeta, \gamma) = (\xi, \alpha) \text{ (or } (\eta, \beta)).$$

We define

$$(2.0.7) \quad h_\gamma(\lambda_1) := \frac{2(\lambda + 2\sigma) \csc^2 \gamma}{(\lambda_1 + \sigma)(\lambda_2 + \sigma)} \in \mathbf{R}_+ = (0, \infty) \text{ } (\gamma = \alpha, \beta).$$

For $|t| > 1$ ($t = x, y$), $(\zeta, \gamma, t) = (\xi, \alpha, x)$ (or (η, β, y)), we set the following functions

$$A_{\zeta, \gamma}(t) := |t - \zeta| + (t - \zeta) \cos \gamma,$$

and

$$H(x, y) := \frac{[\min\{\ln \rho A_{\xi, \alpha}(x), \ln \rho A_{\eta, \beta}(y)\}]^\sigma}{[\max\{\ln \rho A_{\xi, \alpha}(x), \ln \rho A_{\eta, \beta}(y)\}]^{\lambda+\sigma}}.$$

Lemma 2.1. *Define two weight coefficients as follows:*

$$(2.0.8) \quad \omega(\lambda_2, m) := \sum_{|n|=2}^{\infty} \frac{H(m, n)}{A_{\eta, \beta}(n)} \frac{\ln^{\lambda_1} \rho A_{\xi, \alpha}(m)}{\ln^{1-\lambda_2} \rho A_{\eta, \beta}(n)}, |m| \in \mathbf{N} \setminus \{1\},$$

$$(2.0.9) \quad \varpi(\lambda_1, n) := \sum_{|m|=2}^{\infty} \frac{H(m, n)}{A_{\xi, \alpha}(m)} \frac{\ln^{\lambda_2} \rho A_{\eta, \beta}(n)}{\ln^{1-\lambda_1} \rho A_{\xi, \alpha}(m)}, |n| \in \mathbf{N} \setminus \{1\},$$

where

$$\sum_{|j|=2}^{\infty} \dots = \sum_{j=-2}^{-\infty} + \dots + \sum_{j=2}^{\infty} \dots \quad (j = m, n).$$

Lemma 2.2. *(cf. [31]) Let us assume that $g(t) (> 0)$ is strictly decreasing in $(1, \infty)$, satisfying*

$$\int_1^\infty g(t) dt \in \mathbf{R}_+.$$

We have

$$(2.0.10) \quad \int_2^\infty g(t) dt < \sum_{n=2}^{\infty} g(n) < \int_1^\infty g(t) dt.$$

If $(-1)^i g(t) > 0$ ($i = 0, 1, 2; t \in (\frac{3}{2}, \infty)$),

$$\int_{\frac{3}{2}}^{\infty} g(t) dt \in \mathbf{R}_+,$$

then we have the following Hermite-Hadamard inequality (cf. [32]):

$$(2.0.11) \quad \sum_{n=2}^{\infty} g(n) < \int_{\frac{3}{2}}^{\infty} g(t) dt.$$

Lemma 2.3. For $\lambda_1 > -\sigma, -\sigma < \lambda_2 \leq 1 - \sigma$, the following inequalities are valid:

$$(2.0.12) \quad h_{\beta}(\lambda_1)(1 - \theta(\lambda_2, m)) < \omega(\lambda_2, m) < h_{\beta}(\lambda_1), |m| \in \mathbf{N} \setminus \{1\},$$

where

$$(2.0.13) \quad \begin{aligned} \theta(\lambda_2, m) &:= \frac{(\lambda_1 + \sigma)(\lambda_2 + \sigma)}{\lambda + 2\sigma} \int_0^{\frac{\ln \rho(2+\eta)(1+\cos \beta)}{\ln \rho A_{\xi,\alpha}(m)}} \frac{(\min\{1, u\})^{\sigma} u^{\lambda_2 - 1}}{(\max\{1, u\})^{\lambda + \sigma}} du \\ &= O\left(\frac{1}{\ln^{\lambda_2 + \sigma} \rho A_{\xi,\alpha}(m)}\right) \in (0, 1). \end{aligned}$$

Proof. For $|m| \in \mathbf{N} \setminus \{1\}$, we set

$$\begin{aligned} H^{(1)}(m, y) &:= \frac{[\min\{\ln \rho A_{\xi,\alpha}(m), \ln \rho(y - \eta)(\cos \beta - 1)\}]^{\sigma}}{[\max\{\ln \rho A_{\xi,\alpha}(m), \ln \rho(y - \eta)(\cos \beta - 1)\}]^{\lambda + \sigma}}, y < -1, \\ H^{(2)}(m, y) &:= \frac{[\min\{\ln \rho A_{\xi,\alpha}(m), \ln \rho(y - \eta)(\cos \beta + 1)\}]^{\sigma}}{[\max\{\ln \rho A_{\xi,\alpha}(m), \ln \rho(y - \eta)(\cos \beta + 1)\}]^{\lambda + \sigma}}, y > 1, \end{aligned}$$

wherefrom,

$$H^{(1)}(m, -y) = \frac{[\min\{\ln \rho A_{\xi,\alpha}(m), \ln \rho(y + \eta)(1 - \cos \beta)\}]^{\sigma}}{[\max\{\ln \rho A_{\xi,\alpha}(m), \ln \rho(y + \eta)(1 - \cos \beta)\}]^{\lambda + \sigma}}, y > 1.$$

Then we deduce that

$$(2.0.14) \quad \begin{aligned} \omega(\lambda_2, m) &= \sum_{n=-2}^{-\infty} \frac{H^{(1)}(m, n)}{(n - \eta)(\cos \beta - 1)} \frac{\ln^{\lambda_1} \rho A_{\xi,\alpha}(m)}{\ln^{1-\lambda_2} \rho(n - \eta)(\cos \beta - 1)} \\ &\quad + \sum_{n=2}^{\infty} \frac{H^{(2)}(m, n)}{(n - \eta)(\cos \beta + 1)} \frac{\ln^{\lambda_1} \rho A_{\xi,\alpha}(m)}{\ln^{1-\lambda_2} \rho(n - \eta)(\cos \beta + 1)} \\ &= \frac{\ln^{\lambda_1} \rho A_{\xi,\alpha}(m)}{1 - \cos \beta} \sum_{n=2}^{\infty} \frac{H^{(1)}(m, -n)}{(n + \eta) \ln^{1-\lambda_2} \rho(n + \eta)(1 - \cos \beta)} \\ &\quad + \frac{\ln^{\lambda_1} \rho A_{\xi,\alpha}(m)}{1 + \cos \beta} \sum_{n=2}^{\infty} \frac{H^{(2)}(m, n)}{(n - \eta) \ln^{1-\lambda_2} \rho(n - \eta)(1 + \cos \beta)}. \end{aligned}$$

By the fact that $\lambda_1 > -\sigma$, $-\sigma < \lambda_2 \leq 1 - \sigma$ ($\lambda_1 + \lambda_2 = \lambda$), we obtain that for $y > 1, i = 1, 2$,

$$\begin{aligned} & \frac{H^{(i)}(m, (-1)^i y)}{[y - (-1)^i \eta] \ln^{1-\lambda_2} \rho[y - (-1)^i \eta][1 + (-1)^i \cos \beta]} \\ &= \begin{cases} \frac{1}{[y - (-1)^i \eta](\ln \rho A_{\xi, \alpha}(m))^{\lambda+\sigma} \ln^{1-\lambda_2-\sigma} \rho[y - (-1)^i \eta][1 + (-1)^i \cos \beta]}, \\ \frac{1}{\rho} < [y - (-1)^i \eta][1 + (-1)^i \cos \beta] \leq A_{\xi, \alpha}(m) \\ \frac{(\ln \rho A_{\xi, \alpha}(m))^{\sigma}}{[y - (-1)^i \eta] \ln^{1+\lambda_1+\sigma} \rho[y - (-1)^i \eta][1 + (-1)^i \cos \beta]}, \\ [y - (-1)^i \eta][1 + (-1)^i \cos \beta] > A_{\xi, \alpha}(m) \end{cases} \end{aligned}$$

are strictly decreasing in $(1, \infty)$. By (2.0.14) and (2.0.10) we get

$$\begin{aligned} \omega(\lambda_2, m) &< \frac{\ln^{\lambda_1} \rho A_{\xi, \alpha}(m)}{1 - \cos \beta} \int_1^\infty \frac{H^{(1)}(m, -y) dy}{(y + \eta) \ln^{1-\lambda_2} \rho(y + \eta)(1 - \cos \beta)} \\ &+ \frac{\ln^{\lambda_1} \rho A_{\xi, \alpha}(m)}{1 + \cos \beta} \int_1^\infty \frac{H^{(2)}(m, y) dy}{(y - \eta) \ln^{1-\lambda_2} \rho(y - \eta)(1 + \cos \beta)}. \end{aligned}$$

Setting

$$u = \frac{\ln \rho(y + \eta)(1 - \cos \beta)}{\ln \rho A_{\xi, \alpha}(m)} \quad \left(u = \frac{\ln \rho(y - \eta)(1 + \cos \beta)}{\ln \rho A_{\xi, \alpha}(m)} \right)$$

in the above first (second) integral, in view of Remark 1, by simplifications, we obtain

$$\begin{aligned} \omega(\lambda_2, m) &< \left(\frac{1}{1 - \cos \beta} + \frac{1}{1 + \cos \beta} \right) \int_0^\infty \frac{(\min\{1, u\})^\sigma u^{\lambda_1-1}}{(\max\{1, u\})^{\lambda+\sigma}} du \\ &= 2 \csc^2 \beta \left(\int_0^1 u^{\lambda_1+\sigma-1} du + \int_1^\infty \frac{u^{\lambda_1-1}}{u^{\lambda+\sigma}} du \right) = h_\beta(\lambda_1). \end{aligned}$$

By (2.0.14) and (2.0.10), similarly, we have

$$\begin{aligned} \omega(\lambda_2, m) &> \frac{\ln^{\lambda_1} \rho A_{\xi, \alpha}(m)}{1 - \cos \beta} \int_2^\infty \frac{H^{(1)}(m, -y) dy}{(y + \eta) \ln^{1-\lambda_2} \rho(y + \eta)(1 - \cos \beta)} \\ &+ \frac{\ln^{\lambda_1} \rho A_{\xi, \alpha}(m)}{1 + \cos \beta} \int_2^\infty \frac{H^{(2)}(m, y) dy}{(y - \eta) \ln^{1-\lambda_2} \rho(y - \eta)(1 + \cos \beta)} \end{aligned}$$

$$\begin{aligned}
 &\geq 2 \csc^2 \beta \int_{\frac{\ln \rho(2+\eta)(1+\cos \beta)}{\ln \rho A_{\xi, \alpha}(m)}}^{\infty} \frac{(\min \{1, u\})^{\sigma} u^{\lambda_1-1}}{(\max \{1, u\})^{\lambda+\sigma}} du \\
 &= h_{\beta}(\lambda_1) - 2 \csc^2 \beta \int_0^{\frac{\ln \rho(2+\eta)(1+\cos \beta)}{\ln \rho A_{\xi, \alpha}(m)}} \frac{(\min \{1, u\})^{\sigma} u^{\lambda_1-1}}{(\max \{1, u\})^{\lambda+\sigma}} du \\
 &= h_{\beta}(\lambda_1)(1 - \theta(\lambda_2, m)) > 0,
 \end{aligned}$$

where $\theta(\lambda_2, m)$ is as in (2.0.13). It follows that $\theta(\lambda_2, m) < 1$ and for

$$A_{\xi, \alpha}(m) \geq (2 + \eta)(1 + \cos \beta),$$

we have

$$\begin{aligned}
 0 &< \theta(\lambda_2, m) = \frac{(\lambda_1 + \sigma)(\lambda_2 + \sigma)}{\lambda + 2\sigma} \int_0^{\frac{\ln \rho(2+\eta)(1+\cos \beta)}{\ln \rho A_{\xi, \alpha}(m)}} u^{\lambda_2+\sigma-1} du \\
 &= \frac{\lambda_1 + \sigma}{\lambda + 2\sigma} \left[\frac{\ln \rho(2+\eta)(1+\cos \beta)}{\ln \rho A_{\xi, \alpha}(m)} \right]^{\lambda_2+\sigma}.
 \end{aligned}$$

Hence, (2.0.12) and (2.0.13) are valid. This completes the proof of the lemma. \square

Similarly, we obtain the lemma below:

Lemma 2.4. *For $\lambda_2 > -\sigma, -\sigma < \lambda_1 \leq 1 - \sigma$, the following inequalities are valid:*

$$(2.0.15) \quad h_{\alpha}(\lambda_1)(1 - \tilde{\theta}(\lambda_1, n)) < \varpi(\lambda_1, n) < h_{\alpha}(\lambda_1), |n| \in \mathbf{N} \setminus \{1\},$$

where

$$\begin{aligned}
 \tilde{\theta}(\lambda_1, n) &:= \frac{(\lambda_1 + \sigma)(\lambda_2 + \sigma)}{\lambda + 2\sigma} \int_0^{\frac{\ln \rho(2+\xi)(1+\cos \alpha)}{\ln \rho A_{\eta, \beta}(n)}} \frac{(\min \{1, u\})^{\sigma} u^{\lambda_1-1}}{(\max \{1, u\})^{\lambda+\sigma}} du \\
 (2.0.16) \quad &= O\left(\frac{1}{\ln^{\lambda_1+\sigma} \rho A_{\eta, \beta}(n)}\right) \in (0, 1).
 \end{aligned}$$

Lemma 2.5. *If $(\zeta, \gamma, k) = (\xi, \alpha, m)$ (or (η, β, n)), then for $\varepsilon > 0$, we have*

$$(2.0.17) \quad H_{\varepsilon}(\zeta, \gamma) := \sum_{|k|=2}^{\infty} \frac{\ln^{-1-\varepsilon} \rho A_{\zeta, \gamma}(k)}{A_{\zeta, \gamma}(k)} = \frac{1}{\varepsilon} (2 \csc^2 \gamma + o(1)) \quad (\varepsilon \rightarrow 0^+).$$

Proof. By (2.0.11), we get that

$$\begin{aligned}
 H_{\varepsilon}(\zeta, \gamma) &= \sum_{k=-2}^{-\infty} \frac{\ln^{-1-\varepsilon} \rho(k - \zeta)(\cos \gamma - 1)}{(k - \zeta)(\cos \gamma - 1)} \\
 &\quad + \sum_{k=2}^{\infty} \frac{\ln^{-1-\varepsilon} \rho(k - \zeta)(\cos \gamma + 1)}{(k - \zeta)(\cos \gamma + 1)}
 \end{aligned}$$

$$\begin{aligned}
&= \sum_{k=2}^{\infty} \left[\frac{\ln^{-1-\varepsilon} \rho(k+\zeta)(1-\cos\gamma)}{(k+\zeta)(1-\cos\gamma)} + \frac{\ln^{-1-\varepsilon} \rho(k-\zeta)(\cos\gamma+1)}{(k-\zeta)(\cos\gamma+1)} \right] \\
&< \int_{\frac{3}{2}}^{\infty} \left[\frac{\ln^{-1-\varepsilon} \rho(y+\zeta)(1-\cos\gamma)}{(y+\zeta)(1-\cos\gamma)} + \frac{\ln^{-1-\varepsilon} \rho(y-\zeta)(\cos\gamma+1)}{(y-\zeta)(\cos\gamma+1)} \right] dy \\
&= \frac{1}{\varepsilon} \left[\frac{\ln^{-\varepsilon} \rho(\frac{3}{2}+\zeta)(1-\cos\gamma)}{1-\cos\gamma} + \frac{\ln^{-\varepsilon} \rho(\frac{3}{2}-\zeta)(1+\cos\gamma)}{1+\cos\gamma} \right] \\
&= \frac{1}{\varepsilon} \left(\frac{1}{1-\cos\gamma} + \frac{1}{1+\cos\gamma} + o_1(1) \right) (\varepsilon \rightarrow 0^+).
\end{aligned}$$

By (2.0.10), we derive that

$$\begin{aligned}
H_{\varepsilon}(\zeta, \gamma) &= \sum_{n=2}^{\infty} \left[\frac{\ln^{-1-\varepsilon} \rho(n+\zeta)(1-\cos\gamma)}{(n+\zeta)(1-\cos\gamma)} + \frac{\ln^{-1-\varepsilon} \rho(n-\zeta)(\cos\gamma+1)}{(n-\zeta)(\cos\gamma+1)} \right] \\
&> \int_2^{\infty} \left[\frac{\ln^{-1-\varepsilon} \rho(y+\zeta)(1-\cos\gamma)}{(y+\zeta)(1-\cos\gamma)} + \frac{\ln^{-1-\varepsilon} \rho(y-\zeta)(\cos\gamma+1)}{(y-\zeta)(\cos\gamma+1)} \right] dy \\
&= \frac{1}{\varepsilon} \left[\frac{\ln^{-\varepsilon} \rho(2+\zeta)(1-\cos\gamma)}{1-\cos\gamma} + \frac{\ln^{-\varepsilon} \rho(2-\zeta)(1+\cos\gamma)}{1+\cos\gamma} \right] \\
&= \frac{1}{\varepsilon} \left(\frac{1}{1-\cos\gamma} + \frac{1}{1+\cos\gamma} + o_2(1) \right) (\varepsilon \rightarrow 0^+).
\end{aligned}$$

Hence, (2.0.17) is valid. This completes the proof of the lemma. \square

3. MAIN RESULTS AND SOME PARTICULAR CASES

Setting the constant

$$(3.0.18) \quad k_{\alpha,\beta}(\lambda_1) := h_{\beta}^{1/p}(\lambda_1)h_{\alpha}^{1/q}(\lambda_1) = \frac{2(\lambda+2\sigma) \csc^{2/p} \beta \csc^{2/q} \alpha}{(\lambda_1+\sigma)(\lambda_2+\sigma)},$$

and using the kernel

$$H(m, n) = \frac{[\min\{\ln \rho A_{\xi,\alpha}(m), \ln \rho A_{\eta,\beta}(n)\}]^{\sigma}}{[\max\{\ln \rho A_{\xi,\alpha}(m), \ln \rho A_{\eta,\beta}(n)\}]^{\lambda+\sigma}},$$

it follows that

Theorem 3.6. *If $a_m, b_n \geq 0$ ($|m|, |n| \in \mathbb{N} \setminus \{1\}$), satisfying*

$$\begin{aligned}
0 &< \sum_{|m|=2}^{\infty} \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi,\alpha}(m)}{(A_{\xi,\alpha}(m))^{1-p}} a_m^p < \infty, \\
0 &< \sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho A_{\eta,\beta}(n)}{(A_{\eta,\beta}(n))^{1-q}} b_n^q < \infty,
\end{aligned}$$

then we have the following reverse equivalent inequalities:

$$\begin{aligned}
 I &:= \sum_{|n|=2}^{\infty} \sum_{|m|=2}^{\infty} H(m, n) a_m b_n \\
 &> k_{\alpha, \beta}(\lambda_1) \left[\sum_{|m|=2}^{\infty} (1 - \theta(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi, \alpha}(m)}{(A_{\xi, \alpha}(m))^{1-p}} a_m^p \right]^{\frac{1}{p}} \\
 (3.0.19) \quad &\times \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho A_{\eta, \beta}(n)}{(A_{\eta, \beta}(n))^{1-q}} b_n^q \right]^{\frac{1}{q}},
 \end{aligned}$$

$$\begin{aligned}
 J_1 &:= \left[\sum_{|n|=2}^{\infty} \frac{\ln^{p\lambda_2-1} \rho A_{\eta, \beta}(n)}{A_{\eta, \beta}(n)} \left(\sum_{|m|=2}^{\infty} H(m, n) a_m \right)^p \right]^{\frac{1}{p}} \\
 (3.0.20) \quad &> k_{\alpha, \beta}(\lambda_1) \left[\sum_{|m|=2}^{\infty} (1 - \theta(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi, \alpha}(m)}{(A_{\xi, \alpha}(m))^{1-p}} a_m^p \right]^{\frac{1}{p}},
 \end{aligned}$$

$$\begin{aligned}
 J_2 &:= \left[\sum_{|m|=2}^{\infty} \frac{\ln^{q\lambda_1-1} \rho A_{\xi, \alpha}(m)}{(1 - \theta(\lambda_2, m))^{q-1} A_{\xi, \alpha}(m)} \left(\sum_{|n|=2}^{\infty} H(m, n) b_n \right)^q \right]^{\frac{1}{q}} \\
 (3.0.21) \quad &> k_{\alpha, \beta}(\lambda_1) \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho A_{\eta, \beta}(n)}{(A_{\eta, \beta}(n))^{1-q}} b_n^q \right]^{1/q}.
 \end{aligned}$$

In particular, (i) for $\alpha = \beta = \frac{\pi}{2}$, $\frac{1}{\rho} - 1 \leq \xi, \eta \leq 1 - \frac{1}{\rho}$, setting

$$\begin{aligned}
 \theta_1(\lambda_2, m) &:= \frac{(\lambda_1 + \sigma)(\lambda_2 + \sigma)}{\lambda + 2\sigma} \int_0^{\frac{\ln \rho(2+\eta)}{\ln \rho|m-\xi|}} \frac{(\min\{1, u\})^\sigma u^{\lambda_2-1}}{(\max\{1, u\})^{\lambda+\sigma}} du \\
 &= O\left(\frac{1}{\ln^{\lambda_2+\sigma} \rho|m-\xi|}\right) \in (0, 1),
 \end{aligned}$$

we have the following reverse equivalent inequalities:

$$\begin{aligned}
& \sum_{|n|=2}^{\infty} \sum_{|m|=2}^{\infty} \frac{(\min\{\ln \rho|m-\xi|, \ln \rho|n-\eta|\})^{\sigma}}{(\max\{\ln \rho|m-\xi|, \ln \rho|n-\eta|\})^{\lambda+\sigma}} a_m b_n \\
& > \frac{2(\lambda+2\sigma)}{(\lambda_1+\sigma)(\lambda_2+\sigma)} \left[\sum_{|m|=2}^{\infty} (1-\theta_1(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho|m-\xi|}{|m-\xi|^{1-p}} a_m^p \right]^{\frac{1}{p}} \\
(3.0.22) \quad & \times \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho|n-\eta|}{|n-\eta|^{1-q}} b_n^q \right]^{\frac{1}{q}},
\end{aligned}$$

$$\begin{aligned}
& \left\{ \sum_{|n|=2}^{\infty} \frac{\ln^{p\lambda_2-1} \rho|n-\eta|}{|n-\eta|} \left[\sum_{|m|=2}^{\infty} \frac{(\min\{\ln \rho|m-\xi|, \ln \rho|n-\eta|\})^{\sigma} a_m}{(\max\{\ln \rho|m-\xi|, \ln \rho|n-\eta|\})^{\lambda+\sigma}} \right]^p \right\}^{\frac{1}{p}} \\
(3.0.23) \quad & \frac{2(\lambda+2\sigma)}{(\lambda_1+\sigma)(\lambda_2+\sigma)} \left[\sum_{|m|=2}^{\infty} (1-\theta_1(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho|m-\xi|}{|m-\xi|^{1-p}} a_m^p \right]^{\frac{1}{p}}.
\end{aligned}$$

$$\begin{aligned}
& \left\{ \sum_{|m|=2}^{\infty} \frac{\ln^{q\lambda_1-1} \rho|m-\xi|}{(1-\theta_1(\lambda_2, m))^{q-1}|m-\xi|} \right. \\
& \times \left. \left[\sum_{|n|=2}^{\infty} \frac{(\min\{\ln \rho|m-\xi|, \ln \rho|n-\eta|\})^{\sigma} b_n}{(\max\{\ln \rho|m-\xi|, \ln \rho|n-\eta|\})^{\lambda+\sigma}} \right]^q \right\}^{\frac{1}{q}} \\
(3.0.24) \quad & > k_{\alpha, \beta}(\lambda_1) \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho|n-\eta|}{|n-\eta|^{1-q}} b_n^q \right]^{\frac{1}{q}}.
\end{aligned}$$

(ii) For $\xi = \eta = 0$, $\arccos \sqrt{1 - \frac{1}{\rho}} \leq \alpha, \beta \leq \pi - \arccos \sqrt{1 - \frac{1}{\rho}}$, setting

$$\begin{aligned}
\theta_2(\lambda_2, m) &:= \frac{(\lambda_1+\sigma)(\lambda_2+\sigma)}{\lambda+2\sigma} \int_0^{\frac{\ln 2\rho(1+\cos \beta)}{\ln \rho(|m|+m \cos \alpha)}} \frac{(\min\{1, u\})^{\sigma} u^{\lambda_2-1}}{(\max\{1, u\})^{\lambda+\sigma}} du \\
&= O\left(\frac{1}{\ln^{\lambda_2+\sigma} \rho(|m|+m \cos \alpha)}\right) \in (0, 1),
\end{aligned}$$

we have the following reverse equivalent inequalities:

$$\begin{aligned}
 & \sum_{|n|=2}^{\infty} \sum_{|m|=2}^{\infty} \frac{[\min\{\ln \rho(|m| + m \cos \alpha), \ln \rho(|n| + n \cos \beta)\}]^{\sigma} a_m b_n}{[\max\{\ln \rho(|m| + m \cos \alpha), \ln \rho(|n| + n \cos \beta)\}]^{\lambda+\sigma}} \\
 & > k_{\alpha, \beta}(\lambda_1) \left[\sum_{|m|=2}^{\infty} (1 - \theta_2(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho(|m| + m \cos \alpha)}{(|m| + m \cos \alpha)^{1-p}} a_m^p \right]^{\frac{1}{p}} \\
 (3.0.25) \quad & \times \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho(|n| + n \cos \beta)}{(|n| + n \cos \beta)^{1-q}} b_n^q \right]^{\frac{1}{q}},
 \end{aligned}$$

$$\begin{aligned}
 & \left\{ \sum_{|n|=2}^{\infty} \frac{\ln^{p\lambda_2-1} \rho(|n| + n \cos \beta)}{|n| + n \cos \beta} \right. \\
 & \times \left. \left[\sum_{|m|=2}^{\infty} \frac{[\min\{\ln \rho(|m| + m \cos \alpha), \ln \rho(|n| + n \cos \beta)\}]^{\sigma} a_m}{[\max\{\ln \rho(|m| + m \cos \alpha), \ln \rho(|n| + n \cos \beta)\}]^{\lambda+\sigma}} \right]^p \right\}^{\frac{1}{p}} \\
 (3.0.26) \quad & > k_{\alpha, \beta}(\lambda_1) \left[\sum_{|m|=2}^{\infty} (1 - \theta_2(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho(|m| + m \cos \alpha)}{(|m| + m \cos \alpha)^{1-p}} a_m^p \right]^{\frac{1}{p}},
 \end{aligned}$$

and

$$\begin{aligned}
 & \left\{ \sum_{|m|=2}^{\infty} \frac{\ln^{q\lambda_1-1} \rho(|m| + m \cos \alpha)}{(1 - \theta_2(\lambda_2, m))^{\eta-1} (|m| + m \cos \alpha)} \right. \\
 & \times \left. \left[\sum_{|n|=2}^{\infty} \frac{[\min\{\ln \rho(|m| + m \cos \alpha), \ln \rho(|n| + n \cos \beta)\}]^{\sigma} b_n}{[\max\{\ln \rho(|m| + m \cos \alpha), \ln \rho(|n| + n \cos \beta)\}]^{\lambda+\sigma}} \right]^q \right\}^{\frac{1}{q}} \\
 (3.0.27) \quad & > k_{\alpha, \beta}(\lambda_1) \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho(|n| + n \cos \beta)}{(|n| + n \cos \beta)^{1-q}} b_n^q \right]^{\frac{1}{q}}.
 \end{aligned}$$

Proof. By the reverse Hölder inequality with weight (cf. [33]) and (2.0.9), we obtain

that

$$\begin{aligned}
& \left(\sum_{|m|=2}^{\infty} H(m, n) a_m \right)^p \\
= & \left\{ \sum_{|m|=2}^{\infty} H(m, n) \left[\frac{(A_{\xi, \alpha}(m))^{1/q} \ln^{(1-\lambda_1)/q} \rho A_{\xi, \alpha}(m)}{\ln^{(1-\lambda_2)/p} \rho A_{\eta, \beta}(n)} a_m \right] \right. \\
& \quad \times \left. \left[\frac{\ln^{(1-\lambda_2)/p} \rho A_{\eta, \beta}(n)}{(A_{\xi, \alpha}(m))^{1/q} \ln^{(1-\lambda_1)/q} \rho A_{\xi, \alpha}(m)} \right] \right\}^p \\
\geq & \sum_{|m|=2}^{\infty} H(m, n) \frac{(A_{\xi, \alpha}(m))^{p/q} \ln^{(1-\lambda_1)p/q} \rho A_{\xi, \alpha}(m)}{\ln^{1-\lambda_2} \rho A_{\eta, \beta}(n)} a_m^p \\
& \times \left[\sum_{|m|=2}^{\infty} H(m, n) \frac{\ln^{(1-\lambda_2)q/p} \rho A_{\eta, \beta}(n)}{A_{\xi, \alpha}(m) \ln^{1-\lambda_1} \rho A_{\xi, \alpha}(m)} \right]^{p-1} \\
= & \frac{(\varpi(\lambda_1, n))^{p-1} A_{\eta, \beta}(n)}{\ln^{p\lambda_2-1} \rho A_{\eta, \beta}(n)} \sum_{|m|=2}^{\infty} \frac{H(m, n) (A_{\xi, \alpha}(m))^{\frac{p}{q}} \ln^{(1-\lambda_1)\frac{p}{q}} \rho A_{\xi, \alpha}(m)}{A_{\eta, \beta}(n) \ln^{1-\lambda_2} \rho A_{\eta, \beta}(n)} a_m^p.
\end{aligned}$$

By (2.0.15), for $0 < p < 1$, it follows that $(\varpi(\lambda_1, n))^{p-1} > h_{\alpha}^{p-1}(\lambda_1)$ and

$$\begin{aligned}
J_1 & > h_{\alpha}^{1/q}(\lambda_1) \left[\sum_{|n|=2}^{\infty} \sum_{|m|=2}^{\infty} \frac{H(m, n) (A_{\xi, \alpha}(m))^{p/q} \ln^{(1-\lambda_1)p/q} \rho A_{\xi, \alpha}(m)}{A_{\eta, \beta}(n) \ln^{1-\lambda_2} \rho A_{\eta, \beta}(n)} a_m^p \right]^{\frac{1}{p}} \\
& = h_{\alpha}^{1/q}(\lambda_1) \left[\sum_{|m|=2}^{\infty} \sum_{|n|=2}^{\infty} \frac{H(m, n) (A_{\xi, \alpha}(m))^{p/q} \ln^{(1-\lambda_1)p/q} \rho A_{\xi, \alpha}(m)}{A_{\eta, \beta}(n) \ln^{1-\lambda_2} \rho A_{\eta, \beta}(n)} a_m^p \right]^{\frac{1}{p}} \\
(3.0.28) & = h_{\alpha}^{1/q}(\lambda_1) \left[\sum_{|m|=2}^{\infty} \omega(\lambda_2, m) \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi, \alpha}(m)}{(A_{\xi, \alpha}(m))^{1-p}} a_m^p \right]^{\frac{1}{p}}.
\end{aligned}$$

By (2.0.12) and (3.0.18) we have (3.0.20).

Using again the reverse Hölder inequality, we have

$$\begin{aligned}
I & = \sum_{|n|=2}^{\infty} \left[\frac{\ln^{\lambda_2-(1/p)} \rho A_{\eta, \beta}(n)}{(A_{\eta, \beta}(n))^{1/p}} \sum_{|m|=2}^{\infty} H(m, n) a_m \right] \left[\frac{\ln^{(1/p)-\lambda_2} \rho A_{\eta, \beta}(n)}{(A_{\eta, \beta}(n))^{-1/p}} b_n \right] \\
(3.0.29) & \geq J_1 \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho A_{\eta, \beta}(n)}{(A_{\eta, \beta}(n))^{1-q}} b_n^q \right]^{\frac{1}{q}},
\end{aligned}$$

and then by (3.0.20) we deduce (3.0.19).

On the other hand, assuming that (3.0.19) is valid, we set

$$b_n := \frac{\ln^{p\lambda_2-1} \rho A_{\eta,\beta}(n)}{A_{\eta,\beta}(n)} \left(\sum_{|m|=2}^{\infty} H(m, n) a_m \right)^{p-1}, \quad |n| \in \mathbf{N} \setminus \{1\},$$

and find

$$J_1 = \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho A_{\eta,\beta}(n)}{(A_{\eta,\beta}(n))^{1-q}} b_n^q \right]^{\frac{1}{p}}.$$

By (3.0.28), it follows that $J_1 > 0$. If $J_1 = \infty$, then (3.0.20) is trivially valid; if $J_1 < \infty$, then we have

$$\begin{aligned} & \sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho A_{\eta,\beta}(n)}{(A_{\eta,\beta}(n))^{1-q}} b_n^q = J_1^p = I > k_{\alpha,\beta}(\lambda_1) \\ & \times \left[\sum_{|m|=2}^{\infty} (1 - \theta(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi,\alpha}(m)}{(A_{\xi,\alpha}(m))^{1-p}} a_m^p \right]^{\frac{1}{p}} \\ & \times \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho A_{\eta,\beta}(n)}{(A_{\eta,\beta}(n))^{1-q}} b_n^q \right]^{\frac{1}{q}}, \\ J_1 &= \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho A_{\eta,\beta}(n)}{(A_{\eta,\beta}(n))^{1-q}} b_n^q \right]^{\frac{1}{p}} \\ &> k_{\alpha,\beta}(\lambda_1) \left[\sum_{|m|=2}^{\infty} (1 - \theta(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi,\alpha}(m)}{(A_{\xi,\alpha}(m))^{1-p}} a_m^p \right]^{\frac{1}{p}}. \end{aligned}$$

Hence, (3.0.20) is valid, and thus (3.0.19) and (3.0.20) are equivalent. Next we will show that inequalities (3.0.19) and (3.0.21) are equivalent, which will show that all the inequalities (3.0.19), (3.0.20) and (3.0.21) are equivalent.

Having proved that (3.0.19) is valid, we then set

$$a_m := \frac{\ln^{q\lambda_1-1} \rho A_{\xi,\alpha}(m)}{(1 - \theta(\lambda_2, m))^{q-1} A_{\xi,\alpha}(m)} \left(\sum_{|n|=2}^{\infty} H(m, n) b_n \right)^{q-1}, \quad |m| \in \mathbf{N} \setminus \{1\},$$

and find

$$J_2 = \left[\sum_{|m|=2}^{\infty} (1 - \theta(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi,\alpha}(m)}{(A_{\xi,\alpha}(m))^{1-p}} a_m^p \right]^{\frac{1}{q}}.$$

If $J_2 = 0$, then (3.0.21) is impossible, namely $J_2 > 0$. If $J_2 = \infty$, then (3.0.21) is trivially valid; if $J_2 < \infty$, then we have

$$\begin{aligned} & \sum_{|m|=2}^{\infty} (1 - \theta(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi,\alpha}(m)}{(A_{\xi,\alpha}(m))^{1-p}} a_m^p = J_2^q = I \\ & > k_{\alpha,\beta}(\lambda_1) \left[\sum_{|m|=2}^{\infty} (1 - \theta(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi,\alpha}(m)}{(A_{\xi,\alpha}(m))^{1-p}} a_m^p \right]^{\frac{1}{p}} \\ & \quad \times \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho A_{\eta,\beta}(n)}{(A_{\eta,\beta}(n))^{1-q}} b_n^q \right]^{\frac{1}{q}}, \\ J_2 &= \left[\sum_{|m|=2}^{\infty} (1 - \theta(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi,\alpha}(m)}{(A_{\xi,\alpha}(m))^{1-p}} a_m^p \right]^{\frac{1}{q}} \\ &> k_{\alpha,\beta}(\lambda_1) \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho A_{\eta,\beta}(n)}{(A_{\eta,\beta}(n))^{1-q}} b_n^q \right]^{\frac{1}{q}}. \end{aligned}$$

Hence, (3.0.21) holds true.

On the other-hand, assuming that (3.0.21) is valid, using the reverse Hölder inequality, we have

$$\begin{aligned} I &= \sum_{|m|=2}^{\infty} \left[\frac{\ln^{(1/q)-\lambda_1} \rho A_{\xi,\alpha}(m)}{(1 - \theta(\lambda_2, m))^{-1/p} (A_{\xi,\alpha}(m))^{-1/q}} a_m \right] \\ &\quad \times \left[\frac{\ln^{\lambda_1-(1/q)} \rho A_{\xi,\alpha}(m)}{(1 - \theta(\lambda_2, m))^{1/p} (A_{\xi,\alpha}(m))^{1/q}} \sum_{|n|=2}^{\infty} H(m, n) b_n \right] \\ (3.0.30) \quad &\geq \left[\sum_{|m|=2}^{\infty} (1 - \theta(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi,\alpha}(m)}{(A_{\xi,\alpha}(m))^{1-p}} a_m^p \right]^{\frac{1}{p}} J_2, \end{aligned}$$

and then by (3.0.21) we have (3.0.19), which is equivalent to (3.0.21).

Therefore, inequalities (3.0.19), (3.0.20) and (3.0.21) are equivalent. This completes the proof of the theorem. \square

Theorem 3.7. *With respect to the assumptions of Theorem 1, the constant factor $k_{\alpha,\beta}(\lambda_1)$ in (3.0.19), (3.0.20) and (3.0.21) is the best possible.*

Proof. For $0 < \varepsilon < p(\lambda_1 + \sigma)$, we set $\tilde{\lambda}_1 = \lambda_1 - \frac{\varepsilon}{p}$ ($\in (-\sigma, 1 - \sigma)$), $\tilde{\lambda}_2 = \lambda_2 + \frac{\varepsilon}{p}$ ($> -\sigma$), and

$$\begin{aligned}\tilde{a}_m &:= \frac{\ln^{\lambda_1 - (\varepsilon/p) - 1} \rho A_{\xi, \alpha}(m)}{A_{\xi, \alpha}(m)} = \frac{\ln^{\tilde{\lambda}_1 - 1} \rho A_{\xi, \alpha}(m)}{A_{\xi, \alpha}(m)} (|m| \in \mathbf{N} \setminus \{1\}), \\ \tilde{b}_n &:= \frac{\ln^{\lambda_2 - (\varepsilon/q) - 1} \rho A_{\eta, \beta}(n)}{A_{\eta, \beta}(n)} = \frac{\ln^{\tilde{\lambda}_2 - \varepsilon - 1} \rho A_{\eta, \beta}(n)}{A_{\eta, \beta}(n)} (|n| \in \mathbf{N} \setminus \{1\}).\end{aligned}$$

By (2.0.17) and (2.0.15) we find

$$\begin{aligned}\tilde{I}_1 &:= \left[\sum_{|m|=2}^{\infty} (1 - \theta(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho A_{\xi, \alpha}(m)}{(A_{\xi, \alpha}(m))^{1-p}} \tilde{a}_m^p \right]^{\frac{1}{p}} \\ &\quad \times \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} \rho A_{\eta, \beta}(n)}{(A_{\eta, \beta}(n))^{1-q}} \tilde{b}_n^q \right]^{\frac{1}{q}} \\ &= \left[\sum_{|m|=2}^{\infty} \frac{\ln^{-1-\varepsilon} \rho A_{\xi, \alpha}(m)}{A_{\xi, \alpha}(m)} - \sum_{|m|=2}^{\infty} \frac{O(\ln^{-1-(\lambda_2+\sigma+\varepsilon)} \rho A_{\xi, \alpha}(m))}{A_{\xi, \alpha}(m)} \right]^{\frac{1}{p}} \\ &\quad \times \left[\sum_{|n|=2}^{\infty} \frac{\ln^{-1-\varepsilon} \rho A_{\eta, \beta}(n)}{A_{\eta, \beta}(n)} \right]^{\frac{1}{q}} \\ &= \frac{1}{\varepsilon} (2 \csc^2 \alpha + o(1) - \varepsilon O(1))^{1/p} (2 \csc^2 \beta + \tilde{o}(1))^{1/q} (\varepsilon \rightarrow 0^+),\end{aligned}$$

$$\begin{aligned}\tilde{I} &:= \sum_{|m|=2}^{\infty} \sum_{|n|=2}^{\infty} H(m, n) \tilde{a}_m \tilde{b}_n \\ &= \sum_{|n|=2}^{\infty} \sum_{|m|=2}^{\infty} H(m, n) \frac{\ln^{\tilde{\lambda}_1 - 1} \rho A_{\xi, \alpha}(m) \ln^{\tilde{\lambda}_2 - \varepsilon - 1} \rho A_{\eta, \beta}(n)}{A_{\xi, \alpha}(m) A_{\eta, \beta}(n)} \\ &= \sum_{|n|=2}^{\infty} \varpi(\tilde{\lambda}_1, n) \frac{\ln^{-1-\varepsilon} \rho A_{\eta, \beta}(n)}{A_{\eta, \beta}(n)} < h_{\alpha}(\tilde{\lambda}_1) \sum_{|n|=2}^{\infty} \frac{\ln^{-1-\varepsilon} \rho A_{\eta, \beta}(n)}{A_{\eta, \beta}(n)} \\ &= \frac{1}{\varepsilon} h_{\alpha}(\lambda_1 - \frac{\varepsilon}{p}) (2 \csc^2 \beta + \tilde{o}(1)).\end{aligned}$$

If there exists a positive number $K \geq k_{\alpha, \beta}(\lambda_1)$, such that (3.0.19) is still satisfied when replacing $k_{\alpha, \beta}(\lambda_1)$ by K , then in particular, we have

$$\varepsilon \tilde{I} = \varepsilon \sum_{|m|=2}^{\infty} \sum_{|n|=2}^{\infty} H(m, n) \tilde{a}_m \tilde{b}_n > \varepsilon K \tilde{I}_1.$$

In view of the above results, it follows that

$$\begin{aligned} & h_\alpha(\lambda_1 - \frac{\varepsilon}{p})(2 \csc^2 \beta + \tilde{o}(1)) \\ & > K \cdot (2 \csc^2 \alpha + o(1) - \varepsilon O(1))^{1/p} (2 \csc^2 \beta + \tilde{o}(1))^{1/q}, \end{aligned}$$

and then

$$\frac{4(\lambda + 2\sigma) \csc^2 \alpha}{(\lambda_1 + \sigma)(\lambda_2 + \sigma)} \csc^2 \beta \geq 2K \csc^{2/p} \alpha \csc^{2/q} \beta \ (\varepsilon \rightarrow 0^+),$$

namely,

$$k_{\alpha, \beta}(\lambda_1) = \frac{2(\lambda + 2\sigma)}{(\lambda_1 + \sigma)(\lambda_2 + \sigma)} \csc^{2/p} \beta \csc^{2/q} \alpha \geq K.$$

Hence, $K = k_{\alpha, \beta}(\lambda_1)$ is the best possible constant factor in (3.0.19).

The constant factor $k_{\alpha, \beta}(\lambda_1)$ in (3.0.20) ((3.0.21)) is still the best possible. Otherwise we would reach a contradiction by (3.0.29) ((3.0.30)) that the constant factor in (3.0.19) is not the best possible. This completes the proof of the theorem. \square

Remark 2. (i) For $\rho = 1, \xi = \eta = 0$ in (3.0.22), setting

$$\begin{aligned} \tilde{\theta}_1(\lambda_2, m) &:= \frac{(\lambda_1 + \sigma)(\lambda_2 + \sigma)}{\lambda + 2\sigma} \int_0^{\ln |m|} \frac{(\min\{1, u\})^\sigma u^{\lambda_2 - 1}}{(\max\{1, u\})^{\lambda + \sigma}} du \\ &= O\left(\frac{1}{\ln^{\lambda_2 + \sigma} |m|}\right) \in (0, 1), \end{aligned}$$

we have the following inequality:

$$\sum_{|n|=2}^{\infty} \sum_{|m|=2}^{\infty} \frac{(\min\{\ln |m|, \ln |n|\})^\sigma a_m b_n}{(\max\{\ln |m|, \ln |n|\})^{\lambda + \sigma}} > \frac{2(\lambda + 2\sigma)}{(\lambda_1 + \sigma)(\lambda_2 + \sigma)}$$

$$(3.0.31) \quad \times \left[\sum_{|m|=2}^{\infty} (1 - \tilde{\theta}_1(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} |m|}{|m|^{1-p}} a_m^p \right]^{\frac{1}{p}} \left[\sum_{|n|=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} |n|}{|n|^{1-q}} b_n^q \right]^{\frac{1}{q}},$$

It follows that (3.0.21) is an extension of (3.0.31).

(ii) If $a_{-m} = a_m$ and $b_{-n} = b_n$ ($m, n \in \mathbb{N} \setminus \{1\}$), setting

$$\begin{aligned} \hat{\theta}_1(\lambda_2, m) &:= \frac{(\lambda_1 + \sigma)(\lambda_2 + \sigma)}{\lambda + 2\sigma} \int_0^{\ln \rho(2+\eta)} \frac{(\min\{1, u\})^\sigma u^{\lambda_2 - 1}}{(\max\{1, u\})^{\lambda + \sigma}} du \\ &= O\left(\frac{1}{\ln^{\lambda_2 + \sigma} \rho(m - \xi)}\right) \in (0, 1), \end{aligned}$$

$$\begin{aligned}\tilde{\theta}_1(\lambda_2, m) &:= \frac{(\lambda_1 + \sigma)(\lambda_2 + \sigma)}{\lambda + 2\sigma} \int_0^{\ln \rho(2+\eta)} \frac{(\min\{1, u\})^\sigma u^{\lambda_2-1}}{(\max\{1, u\})^{\lambda+\sigma}} du \\ &= O\left(\frac{1}{\ln^{\lambda_2+\sigma} \rho(m + \xi)}\right) \in (0, 1),\end{aligned}$$

then (3.0.22) reduces to

$$\begin{aligned}&\sum_{n=2}^{\infty} \sum_{m=2}^{\infty} \left\{ \frac{[\min\{\ln \rho(m - \xi), \ln \rho(n - \eta)\}]^\sigma}{[\max\{\ln \rho(m - \xi), \ln \rho(n - \eta)\}]^{\lambda+\sigma}} \right. \\ &\quad \left. + \frac{[\min\{\ln \rho(m - \xi), \ln \rho(n + \eta)\}]^\sigma}{[\max\{\ln \rho(m - \xi), \ln \rho(n + \eta)\}]^{\lambda+\sigma}} + \frac{[\min\{\ln \rho(m + \xi), \ln \rho(n - \eta)\}]^\sigma}{[\max\{\ln \rho(m + \xi), \ln \rho(n - \eta)\}]^{\lambda+\sigma}} \right. \\ &\quad \left. + \frac{[\min\{\ln \rho(m + \xi), \ln \rho(n + \eta)\}]^\sigma}{[\max\{\ln \rho(m + \xi), \ln \rho(n + \eta)\}]^{\lambda+\sigma}} \right\} a_m b_n \\ &> \frac{2(\lambda + 2\sigma)}{(\lambda_1 + \sigma)(\lambda_2 + \sigma)} \left\{ \sum_{m=2}^{\infty} \left[(1 - \tilde{\theta}_1(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho(m - \xi)}{(m - \xi)^{1-p}} \right. \right. \\ &\quad \left. \left. + (1 - \tilde{\theta}_1(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} \rho(m + \xi)}{(m + \xi)^{1-p}} \right] a_m^p \right\}^{\frac{1}{p}} \\ (3.0.32) \quad &\times \left\{ \sum_{n=2}^{\infty} \left[\frac{\ln^{q(1-\lambda_2)-1} \rho(n - \eta)}{(n - \eta)^{1-q}} + \frac{\ln^{q(1-\lambda_2)-1} \rho(n + \eta)}{(n + \eta)^{1-q}} \right] b_n^q \right\}^{\frac{1}{q}}.\end{aligned}$$

In particular, for $\rho = 1, \xi = \eta = 0$, setting

$$\begin{aligned}\bar{\theta}_1(\lambda_2, m) &:= \frac{(\lambda_1 + \sigma)(\lambda_2 + \sigma)}{\lambda + 2\sigma} \int_0^{\ln 2} \frac{(\min\{1, u\})^\sigma u^{\lambda_2-1}}{(\max\{1, u\})^{\lambda+\sigma}} du \\ &= O\left(\frac{1}{\ln^{\lambda_2+\sigma} m}\right) \in (0, 1),\end{aligned}$$

we have the following reverse Mulholland-type inequality:

$$\begin{aligned}&\sum_{n=2}^{\infty} \sum_{m=2}^{\infty} \frac{(\min\{\ln m, \ln n\})^\sigma}{(\max\{\ln m, \ln n\})^{\lambda+\sigma}} a_m b_n > \frac{\lambda + 2\sigma}{(\lambda_1 + \sigma)(\lambda_2 + \sigma)} \\ (3.0.33) \quad &\times \left[\sum_{m=2}^{\infty} (1 - \bar{\theta}_1(\lambda_2, m)) \frac{\ln^{p(1-\lambda_1)-1} m}{m^{1-p}} a_m^p \right]^{\frac{1}{p}} \left[\sum_{n=2}^{\infty} \frac{\ln^{q(1-\lambda_2)-1} n}{n^{1-q}} b_n^q \right]^{\frac{1}{q}}.\end{aligned}$$

Competing interests.

The authors declare that they have no competing interests.

Acknowledgments.

We would like to thank Professors N. K. Govil, M. Krnić and J. Kuang for their very useful comments and suggestions which helped improve the presentation of this paper.

M. Th. Rassias: I would like to express my gratitude to the John S. Latsis Public Benefit Foundation for their financial support provided under the auspices of my current “Latsis Foundation Senior Fellowship” position.

B. Yang: This work is supported by the National Natural Science Foundation (No. 61772140), and Science and Technology Planning Project of Guangdong Province (No.2017A010101021). We are grateful for this help.

REFERENCES

1. G. H. HARDY, J. E. LITTLEWOOD, AND G. PÓLYA: *Inequalities*, Cambridge University Press, 1934.
2. D. S. MITRINOVIĆ, J. E. PEČARIĆ, A. M. FINK: *Inequalities involving functions and their integrals and derivatives*, Kluwer Academic Publishers, Boston, Mass, USA, 1991.
3. B. YANG: *A new Hilbert's type integral inequality*, Soochow Journal of Mathematics, vol. 33, no. 4, pp. 849–859, 2007.
4. M. GAO AND B. YANG: *On the extended Hilbert's inequality*, Proc. Amer. Math. Soc., vol. 126, no. 3, pp. 751–759, 1998.
5. M. TH. RASSIAS, B. YANG AND A. RAIGORODSKII: *Two Kinds of the Reverse Hardy-Type Integral Inequalities with the Equivalent Forms Related to the Extended Riemann Zeta Function*, Applicable Analysis and Discrete Mathematics, 12(2018), 273–296.
6. Y. HONG: *All-sided generalization about Hardy-Hilbert integral inequalities*, Acta Mathematica Sinica, vol. 44, no. 4, pp. 619–626, 2001.
7. E. A. LAITH: *On some extensions of Hardy-Hilbert's inequality and applications*, Journal of Inequalities and Applications, Volume 2008, Article ID 546828, 14 pages.
8. M. KRNIĆ AND J. E. PEČARIĆ, "GENERAL HILBERT'S AND HARDY'S INEQUALITIES", Mathematical inequalities and Applications, vol. 8, no. 1, pp. 29–51, 2005.
9. I. PERIĆ, AND P. VUKOVIĆ: *Multiple Hilbert's type inequalities with a homogeneous kernel*, Banach Journal of Mathematical Analysis. vol. 5, no. 2, pp. 33–43, 2011.
10. G. V. MILOVANOVIĆ: *On some integral inequalities*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. No 498–No 541 (1975), 119–124.
11. P. M. VASIĆ AND G. V. MILOVANOVIĆ: *On an inequality of lyengar*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. No 544–No 576 (1976), 18–24.
12. G. V. MILOVANOVIĆ: *On some functional inequalities*, Univ. Beograd. Publ. Elektrotehn. Fak. Ser. Mat. Fiz. No 599 (1977), 1–59 (Serbian, English summary).
13. G. V. MILOVANOVIĆ (ED.): *Recent Progress in Inequalities*, Kluwer Academic Publishers, Dordrecht, 1998.
14. G. V. MILOVANOVIĆ AND M.TH. RASSIAS (EDS.): *Analytic Number Theory Approximation Theory and Special Functions*, Springer, 2014.

15. G. V. MILOVANOVIĆ, D. S. MITRINOVIC AND TH. M. RASSIAS: *Topics in Polynomials: Extremal Problems, Inequalities, Zeros*, World Scientific Publ. Co., Singapore, 1994.
16. B. HE: *A multiple Hilbert-type discrete inequality with a new kernel and best possible constant factor*, Journal of Mathematical Analysis and Applications, vol. 431, pp. 990 – 902, 2015.
17. V. ADIYASUREN, B. TSERENDORJ AND M. KRNIĆ: *Multiple Hilbert-type inequalities involving some differential operators*, Banach J. Math. Anal., vol. 10, no. 2, pp. 320-337, 2016.
18. Y. LI AND B. HE: *On inequalities of Hilbert's type*, Bulletin of the Australian Mathematical Society, vol. 76, no. 1, pp. 1–13, 2007.
19. W. ZHONG: *A Hilbert-type linear operator with the norm and its applications*, Journal of Inequalities and Applications, Vol. 2009, Art. ID 494257, 18 pages.
20. Q. HUANG: *On a multiple Hilbert's inequality with parameters*, Journal of Inequalities and Applications Volume 2010, Article ID 309319, 12 pages
21. M. KRNIĆ AND P. VUKOVIC: *On a multidimensional version of the Hilbert type inequality*, Analysis Mathematica, vol. 38, no. 4, pp. 291-303, 2012.
22. Q. HUANG: *A new extension of Hardy-Hilbert-type inequality*, Journal of Inequalities and Applications (2015), 2015: 397.
23. Z. T. XIE, Z. ZENG AND Y. F. SUN: *A new Hilbert-type inequality with the homogeneous kernel of degree -2*, Adv. Appl. Math. Sci., vol. 12, no. 7, pp. 391—401, 2013.
24. V. ADIYASUREN AND TS. BATBOLD: *On a generalization of a Hilbert-type integral inequality in the whole plane with a hypergeometric function*, J. Inequal. Appl. 2013, Article ID189, 8pp (2013).
25. Z. ZHENG, K. RAJA RAMA GANDHI AND Z. T. XIE: *A new Hilbert-type inequality with the homogeneous kernel of degree -2 and with the integral*, Bull. Math. Sci. Appl., vol. 3, no.1, 11—20, 2014.
26. G. V. MILOVANOVIĆ AND M. TH. RASSIAS: *Some properties of a hypergeometric function which appear in an approximation problem*, J. Glob. Optim. vol. 57, pp. 1173–1192, 2013.
27. Q. HUANG, A. WANG AND B. YANG: *A more accurate half-discrete Hilbert-type inequality with a genreal nonhomogeneous kernel and operator expressions*, Mathematical Inequalities and Applications, vol. 17, no. 1, pp. 367–388, 2014.
28. Q. HUANG, S. WU AND B. YANG: *Parameterized Hilbert-type integral inequalities in the whole plane*, The Scientific World Journal, 2014, Art. ID 169061:1-8.
29. B. HE AND Q. WANG: *A multiple Hilbert-type discrete inequality with a new kernel and best possible constant factor*, Journal of Mathematical Analysis and Applications, vol. 431, no. 2, pp. 889–902, 2015.
30. B. YANG AND Q. CHEN: *A New Extension of Hard-Hilbert's inequality in the whole plane*, Journal of Function Spaces, 2016, Art. ID 9197476: 1-8.
31. D. XIN, B. YANG AND Q. CHEN: *A discrete Hilbert-type inequality in the whole plane*, Journal of Inequalities and Applications (2016), 2016: 133.

32. Q. CHEN AND B. YANG: *On a more accurate half-discret Hardy-Hilbert-type inequality related to the kernel of arc tangent function*, SpringerPlus (2016)5: 1317.
33. J. KUANG: *Applied inequalities*, Shangdong Science Technic Press, Jinan, 2010 (in Chinese).

Michael Th. Rassias

Institute of Mathematics,
University of Zurich, CH-8057,
Zurich, Switzerland

& Moscow Institute of Physics and Technology,
141700 Dolgoprudny,
Institutskiy per, d. 9,
Russia

& Institute for Advanced Study,
Program in Interdisciplinary Studies,
1 Einstein Dr, Princeton,
NJ 08540, USA.

E-mail: *michail.rassias@math.uzh.ch*

(Received 24.12.2018)

(Revised 05.04.2019)

Bicheng Yang

Department of Mathematics,
Guangdong University of Education,
Guangzhou, Guangdong 510303,
P. R. China.

E-mail: *bctyang@gdei.edu.cn*
bctyang818@163.com