

Original Article

Asian Pacific Journal of Reproduction

Journal homepage: www.apjr.net



doi: 10.4103/2305-0500.326721

In-vitro effect of *Peganum harmala* total alkaloids on spermatozoa quality and oxidative stress of epididymal ram semenHanane Derbak, Mohamed Moussaoui, Amine Benberkane, Abdelhanine Ayad[✉]

Department of Biological Sciences of the Environment, Faculty of Nature and Life Sciences, Université de Bejaia, 06000 Bejaia, Algeria

ABSTRACT

Objective: To determine the *in-vitro* effect of the total alkaloid extract of *Peganum* (*P.*) *harmala* seeds on ram epididymal sperm.**Methods:** Semen was divided into six groups according to the following concentrations of the *P. harmala* total alkaloids: 1, 5, 10, 50, and 100 µg/mL, and the control group. The samples were incubated at ambient temperature (21 °C–24 °C) for 24 h, and analyzed in terms of motility, membrane integrity, and oxidative status.**Results:** The sperm kinematic parameters, *i.e.* straight-line velocity, curvilinear velocity, average path velocity, were significantly higher when treated with *P. harmala* at concentrations ranging from 1 to 10 µg/mL compared to the control group ($P < 0.05$). In addition, the highest amplitude of the lateral head displacement value was found in the groups treated with concentrations 1 and 5 µg/mL of *P. harmala* compared to the control group ($P < 0.05$). Total and progressive motilities showed that the extracts at 1, 5, and 10 µg/mL exhibited a high percentage after 24 h of incubation. The effect of *P. harmala* extracts on the membrane integrity of ram epididymal sperm was concentration-dependent and significantly different compared to the control group ($P < 0.05$). Non-significantly lower lipid peroxidation levels were observed after 24 h of incubation of ram epididymal sperm treated with concentrations 1, 5, and 10 µg/mL of *P. harmala* extracts compared to the control group ($P > 0.05$).**Conclusions:** Low concentrations (1–10 µg/mL) of *P. harmala* extracts stimulate sperm motility, preserve membrane integrity and protect ram spermatozoa from lipid peroxidation.**KEYWORDS:** Alkaloids; Epididymal sperm; Motility; *Peganum harmala*; Ram; Semen; Membrane integrity; Lipid peroxidation

1. Introduction

The addition of a substance as a seminal component that would stimulate motility and sperm transport in the uterine tract would

be desirable when artificial insemination is employed in domestic animals. It is known that plant extracts are excellent sources of natural antioxidants and could be used to prevent free radical-induced harmful effects. Several studies experimented with the antioxidant effects of plant extract were associated with an improvement in spermatozoa quality[1,2].

The use of medicinal plants is widespread throughout the world. The herbal plants are increasingly recognized as a valuable source of natural products and drugs against various disorders and diseases. Medicinal plants have been used to develop new molecules with potential effects that are considered a very interesting research approach. The beneficial uses of these plant species as an antioxidant to improve fertility were reported[3]. More recently, attention has focused on the alkaloid extracts from the plants, *e.g.* *Glycine soja*[4], *Epichloë coenophiala*[5], *Tinospora*

Significance

Several natural extracts are used to preserve animal sperm. The development of a new one in extenders with new natural molecules to improve sperm quality is therefore an interesting goal. This study shows that the addition of the total alkaloid extract of *Peganum harmala* seeds at low doses 1–10 µg/mL can improve the quality of epididymal ram sperm by stimulating velocity, maintaining membrane integrity and preserving sperm cells from lipid peroxidation.

[✉]To whom correspondence may be addressed. E-mail: abdelhanine.ayad@univ-bejaia.dz

This is an open access article distributed under the terms of the Creative Commons Attribution-Non Commercial-Share Alike 4.0 License, which allows others to remix, tweak and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

©2021 Asian Pacific Journal of Reproduction Produced by Wolters Kluwer- Medknow. All rights reserved.

How to cite this article: Derbak H, Moussaoui M, Benberkane A, Ayad A. *In-vitro* effect of *Peganum harmala* total alkaloids on spermatozoa quality and oxidative stress of epididymal ram semen. *Asian Pac J Reprod* 2021; 10(5): 232–238.

Article history: Received: 12 May 2021; Revision: 29 July 2021; Accepted: 8 August 2021; Available online: 5 October 2021

cordifolia[6]. Alkaloids are secondary metabolites that have attracted a great deal of research interest due to their different effects on human health[7]. They have multiple physiological properties, such as anti-proliferative[8], anti-inflammatory[9], antimicrobial[10], antioxidant[11], and antidepressant effects[12].

Peganum (P.) harmala L., called harmal, is a perennial plant rich in alkaloids of the Zygophyllaceae family distributed in a large part of the world. The seeds of *P. harmala* are the most important part of the plant. They contain a great structural diversity of alkaloids such as β -carbolines (harmine, harmaline, harmalol, and harman) and quinazoline derivatives (vasicin and vasicinone)[13]. There are several reports in the literature indicating a great variety of pharmacological activities from *P. harmala* L. It is usually used to treat lumbago, asthma, colic, jaundice, and as a stimulant emmenagogue[14]. Moreover, numerous studies have demonstrated that *P. harmala* alkaloids exhibit antitumor[15], anti-inflammatory[16], antiparasitic[17], and antimicrobial effects[18]. On the reproductive system, some studies have been carried out *in-vivo* using extracts of *P. harmala*. Nevertheless, their results have been divergent, with improved sperm quality sometimes[19,20] and adverse changes in their characteristics other times[21].

In the last time, a large body of research has employed a wide variety of antioxidant herbal extracts to enhance sperm performance *in-vitro* and potentially sperm-fertilizing capacity. Notably, polyphenols and essential oils are the most used[2,22]. Alkaloid extracts have been poorly studied despite their properties, as mentioned above. This work will shed light on the effects of alkaloid extract on spermatozoa, thus contributing to the veterinary and human reproduction field.

To our knowledge, the effect of *P. harmala* alkaloids on sperm cells *in-vitro* has never been reported, and their mode of action is still unknown. For this reason, we aimed in the present study to assess the effects of different concentrations of the *P. harmala* extract on motility, membrane integrity, and membrane lipid peroxidation level of fresh epididymal ram spermatozoa extended and incubated in Tris-medium for 24 h.

2. Materials and methods

2.1. Plant materials

P. harmala seeds were collected in June 2016 from areas far from any pollution in the fields of Batna provinces (35°32'N, 6°10'E) in Algeria. The scientific authentication of plants was carried out in the Ecology Laboratory, University of Bejaia, Algeria. A specimen was kept in the Herbarium of the Department of Pharmacy at Batna-2 (voucher number: PhB080). Seeds were put into drying at room temperature (25 °C–30 °C) for four weeks. Thereafter, plant material was pounded by using a coffee grinder resulting in a fine powder and kept in the dark.

2.2. Preparation of plant extract

The extraction of alkaloids was carried out according to the method described by Kartel *et al*[13] with some modifications. Briefly, dried plant materials (20 g) were crushed and extracted with 200 mL ethanol (96%, v/v) in a Soxhlet apparatus for 8 h. The ethanol extract was concentrated under rotavapor, acidified with HCl (2%, v/v), and extracted with petroleum ether (100 mL) to remove fatty materials. The aqueous layer was brought to pH 9 with ammoniac and extracted three times with dichloromethane (20 mL). The organic layer was dried in the open air to obtain a total alkaloid extract.

2.3. Epididymal semen collection

Six testes of mature and healthy rams were obtained from a local slaughterhouse and transported at a cooler temperature to the laboratory. The sperm collection was performed by the retrograde flushing method as previously described by Martinez-Pastor *et al*[23]. The epididymides were dissected free from the testicles and cleaned of connective tissue. Cauda epididymis blood vessels were cut, and the surface area of the cauda was rinsed and wiped. After isolating the cauda epididymis and the vas deferens, 1 mL of the sperm washing solution was injected into the lumen of the vas deferens making a small incision near the junction of the corpus and the proximal cauda with a surgical blade. Sperm were collected in conical tubes and checked, *i.e.* samples with volume ≥ 0.8 mL, concentration $> 2 \times 10^9$ spermatozoa/mL, mass motility $> 3+$ and individual motility $> 70\%$ were selected in this study.

2.4. Sperm dilution

Pooled semen was diluted with Tris buffer (3.028 g hydroxymethyl aminomethane, 1.70 g citric acid, 1.25 g fructose, 800 UI/mL penicillin G, 100 mL distilled water) in order to reach an approximate concentration of 50×10^6 sperm/mL. The sperm sample was divided into 6 groups according to the following concentrations of the total alkaloids of *P. harmala*: the control group (Tris), group 1 (1 μ g/mL), group 2 (5 μ g/mL), group 3 (10 μ g/mL), group 4 (50 μ g/mL) and group 5 (100 μ g/mL). All alkaloid solutions were prepared from Tris solution and were mixed with an equal volume of sperm suspensions. The samples were incubated at laboratory temperature (21 °C–24 °C) for 24 h and then were analyzed in terms of motility, hypoosmotic test, and oxidative status.

2.5. Computer-assisted sperm analysis (CASA)

Computer-assisted sperm analysis was performed by using the Sperm Class Analyzer (Version 3.2.0; SCA® 2014, Microptic SL, Barcelona, Spain); 10 μ L of sperm of each sample was placed in a pre-warmed Makler cell chamber (Makler Counting chambers, Sefi-Medical Instruments Ltd., Biosigma S.r.l., Italy). During the spermatozoa analysis, five fields were examined by using a phase-

contrast microscope at 10× magnification. Six sperm parameters were measured, namely: straight-line velocity (VSL, $\mu\text{m/s}$), curvilinear velocity (VCL, $\mu\text{m/s}$), average path velocity (VAP, $\mu\text{m/s}$), the amplitude of the lateral head displacement (ALH, μm), total motility (%) and progressive motility (%).

2.6. Membrane integrity

Sperm plasma membrane integrity was evaluated by the hypo-osmotic swelling test (HOST)[24]. Briefly, 30 μL of semen from each sample was incubated at 37 °C for 1 h with 300 μL of 100 mOsm hypo-osmotic (9.0 g fructose, 4.9 g sodium citrate, 1 L distilled water). A total of 200 cells per slide were assessed under phase-microscopy (20× magnification). The spermatozoa showing a coiled and swollen tail was considered to have functional membranes.

2.7. Sperm lipid peroxidation

Malondialdehyde (MDA) has been identified as the end product of lipid peroxidation. Measurement of MDA level was determined by quantifying the thiobarbituric acid reacted substances according to Buege & Aust[25], with some modifications. In order to precipitate protein, 0.5 mL sonicated sperm treated at different concentrations of *P. harmala* was added to 1 mL of TBA-TCA-HCl solution (trichloroacetic acid 15%, w/v, thiobarbituric acid 0.375%, w/v in hydrochloric acid 0.25N). The mixture in a glass tube was placed into a boiling water bath (95 °C) for 15 min and then cooled in ice bath to stop the chemical reaction. Subsequently, the suspension was separated (2 000 × *g* for 10 min) and kept in the ice bath. The thiobarbituric acid reactive substances (TBARS) were then quantified by using a spectrophotometer (Biotech Engineering Management Co. Ltd. UK VIS-7220G) at a wavelength of 532 nm. The molar extinction coefficient for MDA was $1.56 \times 10^5 \text{ M}^{-1}\text{cm}^{-1}$. The results were expressed as nmol MDA/ 10^8 spermatozoa.

2.8. Statistical analysis

Data were analyzed by using Statview 4.02 software (Abacus Concepts Inc., Berkeley, CA, USA). The results of experiments were expressed as mean±standard deviation of mean (mean±SD). The normality of the data was confirmed and the differences between concentrations of motility parameters, sperm membrane integrity, and lipid peroxidation were determined by using one-way analysis of variance, followed by *post hoc* Fisher's LSD test. Values were considered significant when $P < 0.05$.

2.9. Ethics statement

This research was approved by the Scientific Council of the Faculty of Natural and Life Sciences (Report of Faculty Science Council #05 dated December 14, 2016), University of Bejaia, Algeria. Concerning the ethical aspects, the experimental procedure was performed according to good veterinary practice under farm conditions.

3. Results

3.1. Kinematic parameters

The effects of the total alkaloid extract of *P. harmala* on the kinematic parameters of ram epididymal sperm are illustrated in Figure 1. Over the incubation time, a perfect similarity was observed when comparing sperm velocities (VSL, VCL, VAP) recorded by the CASA. The incubation of spermatozoa with the total alkaloid extract caused a significant increase in VSL in the groups treated with concentrations ranging from 1 to 10 $\mu\text{g/mL}$ compared to the control group ($P < 0.05$), except for the beginning of the incubation time (0 h). A significant decrease in VCL and VAP was observed in the group treated with 100 $\mu\text{g/mL}$ compared to the control group in all time points ($P < 0.05$), while a significant enhancement of VCL and VAP was observed with the concentrations ranging from 1 to 10 $\mu\text{g/mL}$. Also, the highest ALH values were found in the groups treated with concentrations 1 and 5 $\mu\text{g/mL}$ of *P. harmala* ranging from $(2.94 \pm 0.48) \mu\text{m}$ to $(3.94 \pm 0.62) \mu\text{m}$.

3.2. Sperm motility

Total motility and progressive motility showed that the extracts at 1, 5, and 10 $\mu\text{g/mL}$ exhibited a high percentage after 24 h of incubation of ram epididymal sperm. At a concentration of 50 $\mu\text{g/mL}$, the progressive motility percentage was slightly similar compared to the control group $[(11.87 \pm 8.20)\%$ and $(12.96 \pm 7.15)\%$, respectively]. On the other hand, the treated group with a concentration of 100 $\mu\text{g/mL}$ of *P. harmala* extract showed a significant low value $[(2.92 \pm 2.31)\%]$ compared to semen diluted in Tris at 24 h of incubation ($P < 0.05$) (Figure 2).

3.3. Membrane integrity

The variation of the membrane integrity percentage after 24 h of incubation of ram epididymal sperm, according to the concentration ranging from 1 to 100 $\mu\text{g/mL}$ of *P. harmala* extracts, is illustrated in Figure 3. The effect of *P. harmala* extracts on the membrane integrity of ram epididymal sperm was concentration-dependent and significantly different ($P < 0.05$) compared to the control group. The results indicated a significant alteration of the structural integrity of spermatozoa treated with 100 $\mu\text{g/mL}$ of *P. harmala* extract $[(62.86 \pm 4.33)\%]$. However, the concentrations 1, 5, and 10 $\mu\text{g/mL}$ of *P. harmala* extract seemed to improve the sperm membrane integrity $[(73.98 \pm 5.17)\%$, $(75.11 \pm 4.15)\%$, and $(77.52 \pm 1.83)\%$, respectively] compared to the control group $[(66.88 \pm 6.87)\%]$, ($P < 0.05$). On the other hand, there were no significant differences between the groups treated with 50 $\mu\text{g/mL}$ of *P. harmala* extract and the control group.




Figure 1. Values for straight linear velocity (A), curvilinear velocity (B), average path velocity (C), and the lateral movement amplitude (D) after 0, 1, 2, 4, 24 h of incubation of ram epididymal sperm in the control group and groups treated with different concentrations of *Peganum harmala* alkaloid extracts (1, 5, 10, 50, 100 µg/mL). Data are expressed as mean±SD. At the same time point, different superscripts (a, b, c, d, e, f) in the different concentration groups denote significant differences, $P < 0.05$. VSL: straight-line velocity; VCL: curvilinear velocity; VAP: average path velocity; ALH: the amplitude of the lateral head displacement.




Figure 2. Percentages of total motility (A) and progressive motility (B) after 24 h of incubation of ram epididymal sperm in the control group and groups treated with different concentrations of *Peganum harmala* alkaloid extracts. Data are expressed as mean±SD. The same superscript letter denotes no significant difference.




Figure 3. Percentages of membrane integrity after 24 h of incubation of ram epididymal sperm in the control group and groups treated with different concentrations of *Peganum harmala* alkaloid extracts. Data are expressed as mean±SD. The same superscript letter denotes no significant difference. HOST: hypo-osmotic swelling test.




Figure 4. Percentages of lipid peroxidation after 24 h of incubation of ram epididymal sperm in the control group and groups treated with different concentrations of *Peganum harmala* alkaloid extracts. Data are expressed as mean±SD. The same superscript letter denotes no significant difference. MDA: malondialdehyde. Spz: spermatozoa.

3.4. Lipid peroxidation level

The production of MDA reflects the level of lipid peroxidation of sperm. MDA levels of sperm treated with different concentrations of *P. harmala* extract are shown in Figure 4. After 24 h of incubation of ram epididymal sperm, MDA level non-significantly decreased in the groups treated with concentrations 1, 5, and 10 µg/mL of *P. harmala* extract [(1.66±0.22), (1.57±0.42) and (1.68±0.22) nmol/10⁸ spermatozoa] compared to the control group [(1.78±0.27) nmol/10⁸ spermatozoa] ($P>0.05$), while the highest MDA level was observed in the treated group with concentrations 50 and 100 µg/mL of *P. harmala* extract.

4. Discussion

Since the 20th century, the consumption of plant extracts known for their rich source of beneficial compounds has increased hugely. However, the interest in herbal products needs thorough research in order to ensure their safety, particularly on the reproductive system. Also, it would be important to establish experiments about plant extracts' direct effects on sperm performance *in-vitro*. According to the literature, numerous scientific papers showed significant biological activity of *P. harmala* extract[15,18]. To the author's knowledge, no study investigating the impact of plant extracts from *P. harmala* L. on ram epididymal sperm was reported. The present study explores the potential effects of total alkaloids from *P. harmala*, in the reproductive field, by analyzing motility, oxidative status, and membrane integrity of ram epididymal sperm.

The results of our study have revealed that *P. harmala* extracts have a positive *in vitro* effect on sperm motility in comparison with the control group. Low concentrations of the extract ranging from 1 µg/mL to 10 µg/mL stimulate sperm motility, preserve the membrane's structural integrity, and maintain low lipid peroxidation in ram spermatozoa. In contrast, the highest dose (100 µg/mL) had opposite effects. Our results are consistent with those reported by Hanon[24] in an *in vivo* study on male mice, who showed that the alkaloid compounds of *P. harmala* L. had a significant improvement in most parameters of epididymal sperm. A previous study conducted by Tvrdá *et al*[26] has suggested that supplementation of berberine to bovine spermatozoa, particularly at concentrations ranging between 1 µmol/L and 50 µmol/L, may protect sperm by enhancing their motility and viability. Ros-Santaella and Pintus[27] reported that *Aspalathus linearis* extract significantly affected boar sperm kinetics except for beat cross frequency, ALH, and VCL parameters, which suggests that the rooibos extract enhances sperm velocity. In addition, the lower concentrations (<2.5 µg/mL) of *Eurycoma longifolia* have no deleterious effects on sperm functions; however, at very high concentrations, chemical alkaloid compounds may have harmful effects *in-vitro*[28]. Recently, one study conducted by Page *et al*[29] reported that overall motility and quality of bovine sperm are decreased by ergonovine and ergotamine, alkaloid compounds

found in ergot. A similar observation was reported by Condorelli *et al*[30] using the flow cytometry, in which nicotine was suppressed in a concentration-dependent manner in the sperm progressive motility. Otherwise, Halo *et al*[31] suggested any dose and/or time dependent significant effect of *Viscum albumquercus* on rabbit spermatozoa motility and viability characteristics *in-vitro*.

Although some studies are sometimes contradictory in their results, our findings observed an improvement in the performance of epididymal sperm when treated at low doses with alkaloid extracts of *P. harmala*. Nevertheless, the mechanisms of action of alkaloid compounds on sperm motility are still unclear. In the literature, it is noted that mammalian sperm express neuronal receptors on their plasma membrane and are involved in the regulation of sperm function[32]. The results obtained concerning the effect of *P. harmala* alkaloid extracts on kinetic parameters, measured by CASA system, of ram epididymal sperm could be due to β-carboline alkaloids and their ability to bind with sperm membrane receptors. This difference can be explained by the concentration of the active ingredients in the plant extracts and the kind of alkaloid compounds[29]. It is important to underline that sperm motility is dependent on many cellular functions, including cyclic adenosine monophosphate and calcium concentrations[33]. Also, it should be noted that *P. harmala* alkaloids are lipid-soluble; therefore, they probably can penetrate across sperm membranes and interact with intracellular signaling molecules. However, the adverse effects observed at high concentrations are probably due to harmaline and harmine, which have been previously reported *in-vivo* and *in-vitro* on cancer cell lines[34]. Moreover, the harmine binds to DNA by intercalation, consequently inducing DNA fragmentation[35], affecting fertilization rates and sperm motility[36].

Interestingly, results of both sperm plasma membrane parameters were significantly concentration-dependent when compared to the control group, in agreement with the results reported by Shalaweh *et al*[37] and Tvrdá *et al*[26]. A similar observation was reported by Suresh *et al*[38], in which the effect of seed extracts of *Mucuna pruriens*, rich in alkaloids such as prurienine, prurieninine, and prurienidine, showed remarkable potential reducing sperm damages induced by oxidative stress, with a well preserved epididymal spermatozoa DNA. Our results are corroborated with those reported by Ros-Santaella & Pintus[27], whose alkaloid extracts of rooibos (*Aspalathus linearis*) seem to preserve membrane integrity during sperm storage. Likewise, Baghshahi *et al*[39] showed that plasma membrane integrity was higher in the semen exposed to a medium containing 75 µg/mL of clove buds extract (*Syzygium aromaticum*) extract after cryopreservation processes.

Several scientific papers reported that oxidative stress causes defective changes in cellular functions[40]. It is generally accepted that antioxidants have shown the ability to reduce free radical production, and their addition to spermatozoa could improve the sperm performances[1,2]; a number of studies have shown the biological effects of alkaloid extracts of *P. harmala*, including antioxidant power on oxidative stress in mammalian. This would be due to the fact that β-carboline alkaloids of *P. harmala* can act

as scavengers of free radicals[41]. One possible explanation is the prevention of the oxidation of biogenic amines by inhibiting the monoamine oxidase, which is responsible for the production of toxic substances such as hydrogen peroxide, inducing oxidative stress[42]. Indeed, in agreement with our observation, many investigations demonstrated that the essential oils of *Rosmarinus officinalis* displayed a high antioxidant activity on sperm quality[3,22]. Moreover, the treatment by cholesterol and vitamin E, known for their antioxidant property, preloaded in cyclodextrins, reduces the lipid peroxidation of spermatozoa and increases the membrane integrity[43]. It is reported that the antioxidant action at low doses protects spermatozoa from membrane damages[44], thus preserving their fluidity, integrity, flexibility and consequently the conservation of sperm quality.

This research represents a preliminary study of the impact of total alkaloids of *P. harmala* on ram semen. Some limitations of this study may include the source of the epididymal sperm collected from rams intended for meat consumption. Further, more intermediate concentrations and a longer incubation time are needed to yield more interesting results in the sperm conservation field.

In conclusion, this study reports the effect of *P. harmala* alkaloids on the performance of ram epididymal sperm. The current investigation demonstrated that sperm motility considerably increases at low concentrations (1-10 µg/mL) of alkaloid extracts of *P. harmala*. Also, based on our findings, *P. harmala* alkaloids could protect sperm against oxidative damages and improve the membrane integrity of ram epididymal sperm. This plant could eventually be beneficial from infertility therapy. However, future studies should be carried out to understand the biological and biochemical mechanisms underlying this beneficial effect.

Conflict of interest statement

The authors declare there is no conflict of interest.

Acknowledgements

The authors would like to thank the management of the Bejaia communal slaughterhouse for their help in sampling ram testicles and for their hospitality. Also, our sincere thanks goes to Dr. R. Kebbi for her the technical assistance in plant alkaloids extraction. This study was conducted in Associated Laboratory in Marine Ecosystems and Aquaculture, University of Bejaia, Algeria.

Authors' contributions

Hanane Derbak performed the experimental work, the statistical analysis, and wrote the manuscript. Mohamed Moussaoui conducted the testicular samples from the slaughterhouse and analyzed the

oxidative stress. Amine Benberkane contributed in the technical aspect of the CASA system and the statistical analysis. Abdelhanine Ayad designed and supervised the experimental study, wrote and revised the manuscript. All authors have read and approved the final version of the manuscript.

References

- [1] Khan H, Khan M, Qureshi MS, Shakoor A, Gohar A, Ullah H, et al. Effect of green tea extract (*Camellia sinensis*) on fertility indicators of post-thawed bull spermatozoa. *Pak J Zool* 2017; **49**(4): 1243-1249.
- [2] Sapanidou VG, Margaritis I, Siahos N, Arsenopoulos K, Dragatidou E, Taitzoglou IA, et al. Antioxidant effect of a polyphenol-rich grape pomace extract on motility, viability and lipid peroxidation of thawed bovine spermatozoa. *J Biol Res-Thessaloniki* 2014; **21**(1): 19.
- [3] Touazi L, Aberkane B, Bellik Y, Moula N, Iguer-Ouada M. Effect of the essential oil of *Rosmarinus officinalis* (L.) on rooster sperm motility during 4 °C short-term storage. *Vet World* 2018; **11**(5): 590-597.
- [4] Gofur A, Lestari SR. Effect of black soybean natto extract (*Glycine soja*) on reproduction system of hypercholesterolemia male mice. *Asian Pac J Reprod* 2016; **5**(5): 387-390.
- [5] Burnett CR, Bridges WC, Pratt SL. Effects of grazing tall fescue containing ergot alkaloids on bull sperm cryopreservation. *Anim Reprod Sci* 2017; **181**: 24-29.
- [6] Jayaganthan P, Perumal P, Balamurugan TC, Verma RP, Singh LP, Pattanaik AK, et al. Effects of *Tinosporacordifolia* supplementation on semen quality and hormonal profile in rams. *Anim Reprod Sci* 2013; **140**(1): 47-53.
- [7] Macáková K, Afonso R, Saso L, Mlad nka P. The influence of alkaloids on oxidative stress and related signaling pathways. *Free Radic Biol Med* 2019; **134**: 429-444.
- [8] Habli Z, Toumeh G, Fatfat M, Rahal ON, Gali-Muhtasib H. Emerging cytotoxic alkaloids in the battle against cancer: Overview of molecular mechanisms. *Molecules* 2017; **22**(2): 250.
- [9] Alves de Almeida AC, de-Faria FM, Dunder RJ, Manzo LPB, Souza-Brito ARM, Luiz-Ferreira A. Recent trends in pharmacological activity of alkaloids in animal colitis: Potential use for inflammatory bowel disease. *Evid Based Complement Alternat Med* 2017; **2017**: e8528210.
- [10] Gurrapu S, Mamidala E. *In vitro* antibacterial activity of alkaloids isolated from leaves of *Eclipta alba* against human pathogenic bacteria. *Pharmacogn J* 2017; **9**(4): 573-577.
- [11] Dalimunthe A, ZaitunHasibuan PA, Silalahi J, Satria D. Antioxidant activity of alkaloid fractions of *Litsea cubeba* Lour fruits. *Asian J Pharm Clin Res* 2018; **11**(13): 31.
- [12] Hamid H, Aizi RNM, Mashitah YM. Indole alkaloids from plants as potential leads for antidepressant drugs: A mini review. *Front Pharmacol* 2017; **8**: 96. doi: 10.3389/fphar.2017.00096.
- [13] Kartal M, Altun ML, Kurucu S. HPLC method for the analysis of harmol, harmalol, harmine and harmaline in the seeds of *Peganum harmala* L. *J Pharm Biomed Anal* 2003; **31**(2): 263-269.
- [14] Mazandarani M, Sepehr KS, Baradaran B, Khuri V. Autecology,

- phytochemical and antioxidant activity of *Peganum harmala* L. seed extract in north of Iran (Tash Mountains). *J Med Plants By-Prod* 2012; **2**: 151-156.
- [15] Bournine L, Bensalem S, Fatmi S, Bedjou F, Mathieu V, Iguer-Ouada M, et al. Evaluation of the cytotoxic and cytostatic activities of alkaloid extracts from different parts of *Peganum harmala* L. (Zygophyllaceae). *Eur J Integr Med* 2017; **9**: 91-96.
- [16] Pradeepkumar MR, Joshi SD, Kulkarni VH, Savant C. Phytochemical screening and evaluation of analgesic, anti-inflammatory activities of *Peganum harmala* Linn., seeds in rodents. *J Appl Pharm Sci* 2015; **5**(6): 066-069. doi: 10.7324/JAPS.2015.50610.
- [17] Kebbi R, Besseboua O, Nait-Mouloud M, Ayad A. *In-vitro* acaricidal activity of *Peganum harmala* and *Glaucium flavum* alkaloid against *Rhipicephalus* sp. of dog. *Eurasian J Vet Sci* 2020; **36**(1): 28-35.
- [18] Shaheen HA, Issa MY. *In vitro* and *in vivo* activity of *Peganum harmala* L. alkaloids against phytopathogenic bacteria. *Sci Hortic* 2020; **264**: 108940.
- [19] Subhan F, Sultan S, Alam W, Tahir F, Dil AS. Aphrodisiac potential of *Peganum harmala* seeds. *Hamdard Medicus* 1998; **41**: 69-72.
- [20] Hanon HA. Effects of aqueous and alcoholic extracts of *Peganum harmala* L. seeds on male fertility of white mice treated with olanzapin drug. *J Global Pharm Technol* 2009; **10**(6): 426-436.
- [21] EL-Dwairi QA, Banihani SM. Histo-functional effects of *Peganum harmala* on male rat's spermatogenesis and fertility. *Neuroendocrinol Lett* 2007; **28**(3): 305-310.
- [22] Benberkane A, Khellouf A, Benhenia K, Fatmi S, Iguer-Ouada M. *Rosmarinus officinalis* essential oil preloaded in β -cyclodextrin: Effect on ram spermatozoa motility, membrane integrity and oxidative status during 4 °C storage. *Cryoletters* 2019; **40**(4): 219-225.
- [23] Martinez-Pastor F, Garcia-Macias V, Alvarez M, Chamorro C, Herraez P, de Paz P, et al. Comparison of two methods for obtaining spermatozoa from the cauda epididymis of Iberian red deer. *Theriogenology* 2006; **65**(3): 471-485.
- [24] Revell SG, Mrode RA. An osmotic resistance test for bovine semen. *Anim Reprod Sci* 1994; **36**(1): 77-86.
- [25] Buege JA, Aust SD. Microsomal lipid peroxidation. In: Fleischer S, Packer L (eds.) *Methods in enzymology*. New York: Academic Press; 1978, p. 302-310.
- [26] Tvrdá E, Greifová H, Ivani P, Lukáč N. *In vitro* effects of berberine on the vitality and oxidative profile of bovine spermatozoa. *Int J Anim Vet Sci* 2019; **13**(10): 244-249.
- [27] Ros-Santaella JL, Pintus E. Rooibos (*Aspalathus linearis*) extract enhances boar sperm velocity up to 96 hours of semen storage. *PLOS One* 2017; **12**(8): e0183682.
- [28] Erasmus N, Solomon MC, Fortuin KA, Henkel RR. Effect of *Eurycoma longifolia* Jack (Tongkat ali) extract on human spermatozoa *in vitro*. *Andrologia* 2012; **44**(5): 308-314.
- [29] Page R, Lester T, Rorie R, Jr CR. Ergot alkaloid effects on bovine sperm motility *in vitro*. *Adv Reprod Sci* 2019; **7**: 7-15.
- [30] Condorelli RA, La Vignera S, Giaccone F, Iacoviello L, Vicari E, Mongioi L, et al. *In vitro* effects of nicotine on sperm motility and bio-functional flow cytometry sperm parameters. *Int J Immunopathol Pharmacol* 2013; **26**(3): 739-746.
- [31] Halo Jr M, Massanyi P, Gren A, Lasak A, Slanina T, Ondruska L, et al. Time and dose-dependent effects of viscum album quercus on rabbit spermatozoa motility and viability *in vitro*. *Physiol Res* 2019; **68**(6): 955-972. doi: 10.33549/physiolres.934223.
- [32] Ramírez AR, Castro MA, Angulo C, Ramió L, Rivera MM, Torres M, et al. The presence and function of dopamine type 2 receptors in boar sperm: A possible role for dopamine in viability, capacitation, and modulation of sperm motility. *Biol Reprod* 2009; **80**(4): 753-761.
- [33] Li X, Wang L, Li Y, Zhao N, Zhen L, Fu J, et al. Calcium regulates motility and protein phosphorylation by changing cAMP and ATP concentrations in boar sperm *in vitro*. *Anim Reprod Sci* 2016; **172**: 39-51.
- [34] Jiménez J, Riverón-Negrete L, Abdullaev F, Espinosa-Aguirre J, Rodríguez-Arnaiz R. Cytotoxicity of the β -carboline alkaloids harmine and harmaline in human cell assays *in vitro*. *Exp Toxicol Pathol* 2008; **60**(4): 381-389.
- [35] Mota NSRS, Kwiecinski MR, Felipe KB, Grinevicius VMAS, Siminski T, Almeida GM, et al. β -carboline alkaloid harmine induces DNA damage and triggers apoptosis by a mitochondrial pathway: Study *in silico*, *in vitro* and *in vivo*. *Int J Funct Nutr* 2020; **1**(1): 1-12. doi: 10.3892/ijfn.2020.1.
- [36] Huang CC, Lin DPC, Tsao HM, Cheng TC, Liu CH, Lee MS. Sperm DNA fragmentation negatively correlates with velocity and fertilization rates but might not affect pregnancy rates. *Fertil Steril* 2005; **84**(1): 130-140.
- [37] Shalaweh SM, Erasmus N, Weitz F, Henkel RR. Effect of *Cissampelos capensis* rhizome extract on human spermatozoa *in vitro*. *Andrologia* 2015; **47**(3): 318-327.
- [38] Suresh S, Prithiviraj E, Venkata Lakshmi N, Karthik Ganesh M, Ganesh L, Prakash S. Effect of *Mucuna pruriens* (Linn.) on mitochondrial dysfunction and DNA damage in epididymal sperm of streptozotocin induced diabetic rat. *J Ethnopharmacol* 2013; **145**(1): 32-41.
- [39] Baghshahi H, Riasi A, Mahdavi AH, Shirazi A. Antioxidant effects of clove bud (*Syzygium aromaticum*) extract used with different extenders on ram spermatozoa during cryopreservation. *Cryobiology* 2014; **69**(3): 482-487.
- [40] Rahman MS, Kang KH, Arifuzzaman S, Pang WK, Ryu DY, Song WH, et al. Effect of antioxidants on BPA-induced stress on sperm function in a mouse model. *Sci Rep* 2019; **9**(1): 10584.
- [41] Berrougui H, Isabelle M, Cloutier M, Hmamouchi M, Khalil A. Protective effects of *Peganum harmala* L. extract, harmine and harmaline against human low-density lipoprotein oxidation. *J Pharm Pharmacol* 2006; **58**(7): 967-974. doi: 10.1211/jpp.58.7.0012.
- [42] Herraiz T, González D, Ancín-Azpilicueta C, Arán VJ, Guillén H. β -carboline alkaloids in *Peganum harmala* and inhibition of human monoamine oxidase (MAO). *Food Chem Toxicol* 2010; **48**(3): 839-845.
- [43] Khellouf A, Benhenia K, Fatmi S, Iguer-Ouada M. The complementary effect of cholesterol and vitamin E preloaded in cyclodextrins on frozen bovine semen: Motility parameters, membrane integrity and lipid peroxidation. *Cryo Letters* 2018; **39**(2): 113-120.
- [44] Herraiz T, Guillén H. Monoamine oxidase-A inhibition and associated antioxidant activity in plant extracts with potential antidepressant actions. *Biomed Res Int* 2018; **2018**: 4810394. doi: 10.1155/2018/4810394.