

# Hypoxia and metabolic inhibitors alter the intracellular ATP:ADP ratio and membrane potential in human coronary artery smooth muscle cells

Mingming Yang<sup>1,2</sup>, Caroline Dart<sup>3</sup>, Tomoko Kamishima<sup>2</sup> and John M. Quayle<sup>2</sup>

<sup>1</sup> Department of Cardiology, Zhongda Hospital, Medical School of Southeast University, Nanjing, People's Republic of China

<sup>2</sup> Department of Cellular and Molecular Physiology, Institute of Translational Medicine, Liverpool, UK

<sup>3</sup> Department of Biochemistry, Institute of Integrative Biology, Liverpool, UK

## ABSTRACT

ATP-sensitive potassium ( $K_{ATP}$ ) channels couple cellular metabolism to excitability, making them ideal candidate sensors for hypoxic vasodilation. However, it is still unknown whether cellular nucleotide levels are affected sufficiently to activate vascular  $K_{ATP}$  channels during hypoxia. To address this fundamental issue, we measured changes in the intracellular ATP:ADP ratio using the biosensors Perceval/PercevalHR, and membrane potential using the fluorescent probe DiBAC<sub>4</sub>(3) in human coronary artery smooth muscle cells (HCASMCs). ATP:ADP ratio was significantly reduced by exposure to hypoxia. Application of metabolic inhibitors for oxidative phosphorylation also reduced ATP:ADP ratio. Hyperpolarization caused by inhibiting oxidative phosphorylation was blocked by either 10  $\mu$ M glibenclamide or 60 mM  $K^+$ . Hyperpolarization caused by hypoxia was abolished by 60 mM  $K^+$  but not by individual  $K^+$  channel inhibitors. Taken together, these results suggest hypoxia causes hyperpolarization in part by modulating  $K^+$  channels in SMCs.

**Subjects** Cell Biology, Molecular Biology, Anatomy and Physiology, Cardiology

**Keywords** Metabolic inhibitor, Hypoxia, ATP, Membrane potential, Potassium channels

## INTRODUCTION

When the oxygen supply cannot meet the demand of metabolically active tissues, the systemic arteries dilate, causing an increase in local blood flow. This phenomenon known as hypoxic vasodilation was first reported as far back as 1880 by *Roy & Brown (1880)*. Arteries supplying organs with changeable metabolic requirements need to produce fast and graded contractile responses. Coronary arteries are one such example since they supply blood to one of the most metabolically active organs. Thus, the coronary circulation is adapted to provide high basal rates of blood flow that can further increase more than ~5 fold in response to enhanced cardiac work, for example during exercise (*Duncker & Bache, 2008; Levick, 2010*). This direct and tight link between metabolic demand and blood supply is a hallmark of the coronary circulation. A number of parallel mechanisms to achieve this may exist, and interpretation of studies where one pathway is blocked has been problematic due to this redundancy (*Duncker & Bache, 2008*). Presumably, these multiple

Submitted 5 June 2020  
Accepted 20 October 2020  
Published 10 November 2020

Corresponding author  
Mingming Yang,  
mmyang@seu.edu.cn

Academic editor  
Jeffrey Stuart

Additional Information and  
Declarations can be found on  
page 18

DOI 10.7717/peerj.10344

© Copyright  
2020 Yang et al.

Distributed under  
Creative Commons CC-BY 4.0

**OPEN ACCESS**

mechanisms act as a safety net, ensuring supply of blood is guaranteed. This is a reflection of the absolute importance of coronary metabolic vasodilation, and also why it still remains poorly understood nearly 140 years after the original publication by *Roy & Brown (1880)*. With this caveat, two of the major hypotheses that have been advanced to account for coronary metabolic vasodilation are: (i) the release of vasodilator metabolites from cardiac myocytes (e.g., adenosine,  $K^+$  and  $H^+$ ) acting on vascular smooth muscle cells (VSMCs), and (ii) the direct vasodilatory effects of hypoxia on VSMCs. Of course, these possibilities are not mutually exclusive, but if the latter scenario plays an important role, then hypoxia and metabolic insult should affect VSMCs in vitro.

One possible pathway for the second hypothesis described above is that hypoxia causes the opening of ATP-sensitive potassium ( $K_{ATP}$ ) channels, leading to VSMC hyperpolarization, a closure of voltage-dependent  $Ca^{2+}$  channels (VDCCs) and relaxation (*Daut et al., 1990*).  $K_{ATP}$  channels are ideally placed to couple cellular metabolic state to membrane excitability.  $K_{ATP}$  channels are perhaps best known for glucose sensation in pancreatic  $\beta$ -cells where an increase in blood glucose level causes channel closure through cellular ATP production (or, more precisely, via a change in the ATP:ADP ratio) (*Misler et al., 1992*). The resulting  $\beta$ -cell membrane depolarization triggers  $Ca^{2+}$  entry through VDCCs and so insulin secretion. The anti-diabetic sulphonylurea drugs such as glibenclamide cause insulin secretion by directly inhibiting  $K_{ATP}$  channels in  $\beta$ -cells (*Sturgess et al., 1985*). In cardiac myocytes, the  $K_{ATP}$  channel opens during severe and prolonged anoxia due to reduced ATP and increased ADP levels. This reduces cellular excitability, so preventing action potential generation and cell contraction (*Flagg et al., 2010*). Ultimately this protects the myocardium from damage by preventing complete depletion of intracellular ATP, which would trigger cell death. In the vasculature,  $K_{ATP}$  channels have diverse functions (*Flagg et al., 2010; Quayle, Nelson & Standen, 1997*).  $K_{ATP}$  channels are activated by vasodilators coupled to cAMP-dependent protein kinase (PKA) and are inhibited by vasoconstrictors coupled to PKC (*Kubo, Quayle & Standen, 1997*). The resulting membrane potential changes are important in the actions of vasodilators and vasoconstrictors. Co-localization of  $K_{ATP}$  channels and protein kinases within specific plasma membrane microdomains, termed caveolae, appears important for modulation (*Davies et al., 2010; Sampson et al., 2007; Sampson et al., 2004*). Importantly, vascular  $K_{ATP}$  channels are activated by the anti-anginal compound nicorandil, and are inhibited by the sulphonylureas.

Many studies using intact coronary artery or whole heart preparation have provided evidence for a role of  $K_{ATP}$  channels in hypoxic vasodilation. The first report was made by *Daut et al. (1990)*, in which vasodilation induced by hypoxia was inhibited by glibenclamide using a Langendorf preparation in intact guinea-pig heart. It was suggested that hypoxic vasodilation occurred via activation of coronary artery  $K_{ATP}$  channels, probably as a consequence of the change in cellular nucleotide levels (*Von Beckerath et al., 1991*). The results of a large number of functional studies in animal models are summarized in recent reviews (*Duncker & Bache, 2008; Flagg et al., 2010; Shimoda & Polak, 2011*). However, we know of only two attempts to address this issue in human coronary arteries. In arteries isolated from the right atrial appendage, hypoxia caused dilation that was not dependent

on surrounding cardiac myocytes or the presence of the endothelium (Lynch *et al.*, 2006; Miura *et al.*, 2003). In one study pharmacological inhibition of  $K_{ATP}$  channels using a sulphonylurea (glibenclamide) blunted the hypoxic response (Miura *et al.*, 2003), whilst in the other study glibenclamide was without effect (Lynch *et al.*, 2006).  $K_{ATP}$  channels have been identified in cultured human coronary artery SMCs (HCASMCs) but there have been no studies looking at metabolic or other regulation of these channels in single cells (Yoshida *et al.*, 2004). Regarding the physiological role of the  $K_{ATP}$  channel, it is of interest that genetic knockout in mice results in hypertension, coronary artery vasospasm and sudden cardiac death (Chutkow *et al.*, 2002a; Miki *et al.*, 2002). The effects of knockout have been linked to Prinzmetal angina, although it is not known if  $K_{ATP}$  channel mutations cause this disease in humans.

Despite the functional studies cited above, the apparently simple yet crucial question whether hypoxia changes cellular nucleotide levels to a degree sufficient to activate vascular  $K_{ATP}$  channels has proved to be difficult to address (Flagg *et al.*, 2010).  $K_{ATP}$  channels appear to be mostly closed at levels of ATP normally expected in cells, and cytochrome oxidase, the key oxygen sensitive enzyme involved in mitochondrial ATP generation, is fully saturated with oxygen unless oxygen drops to very low levels ( $<1$  mmHg, essentially equivalent to anoxia). In contrast, hypoxic vasodilation can occur following relatively mild drops in oxygen levels ( $\sim 30$  mmHg) (Duncker & Bache, 2008). Indeed, a major recent review on the subject states that it remains unclear what the adenosine nucleotide levels are in the VSMCs and whether cellular metabolism changes sufficiently to dynamically regulate VSMC  $K_{ATP}$  channel activity under physiological conditions (Flagg *et al.*, 2010).  $K_{ATP}$  currents in pig coronary artery are activated by hypoxia ( $PO_2 \sim 35$  mmHg) (Dart & Standen, 1995). In contrast,  $K_{ATP}$  currents in rat femoral artery are not activated by hypoxia but are activated during anoxia ( $PO_2 < 1$  mmHg) (Quayle *et al.*, 2006). This difference may reflect origin of artery (i.e., species, vascular bed, small vs large artery).

In this study, we exploited fluorescent probes (Perceval/PercevalHR) to directly measure changes in cellular nucleotide levels caused by hypoxia and metabolic inhibitors in real-time in HCASMCs. Changes in membrane potential were also measured, and effects of various  $K^+$  channel inhibitors were investigated. We report that hypoxia and metabolic inhibition cause membrane hyperpolarization at least in part through the activation of  $K^+$  channels.

## MATERIAL AND METHODS

### Cell culture

HCASMCs (Cat no. C-12511) were obtained from Promocell and maintained in smooth muscle growth medium 2 supplemented with smooth muscle growth medium 2 supplement mix (Promocell, Heidelberg, Germany) with a final supplement concentration of (per ml): 5% v/v fetal calf serum, 0.5 ng epidermal growth factor (recombinant human), 2 ng basic fibroblast growth factor (recombinant human) and 5  $\mu$ g insulin (recombinant human). Penicillin/streptomycin (Thermo Fisher Scientific, Waltham, MA) was added to cell culture medium with a final concentration of 100 U/mL and 100  $\mu$ g/mL, respectively. Cells were used between P7-P13. The expression of signature smooth muscle proteins, SMC  $\alpha$ -actin,

calponin and myosin heavy chain (MHC) was maintained for the duration of culture as detected by Western blot (WB) (Fig. S1) and immunocytochemistry (ICC) (Fig. S2).

### Subcultivation of HCASMCs

HCASMCs were grown in T75 cm<sup>2</sup> flasks with 15 ml fully supplemented smooth muscle growth medium 2 in a 5% CO<sub>2</sub> humidified incubator at a temperature of 37 °C. Cells were routinely passaged every 48–72 h when reaching about 70–80% confluency. Cell culture medium was aspirated, and the cells were washed with 15 ml pre-warmed Dulbecco's Phosphate Buffered Saline (DPBS, no Ca<sup>2+</sup>/Mg<sup>2+</sup>, Cat no. 14190-144, Gibco) or HEPES Buffered Saline Solution (HepesBSS). Cells were then trypsinised with 1.5 ml versene (Gibco) containing 0.025% trypsin (Gibco) or 100 µl per cm<sup>2</sup> Trypsin/EDTA Solution (0.04%/0.03%) for approximately 1 min. Enzyme was neutralized by addition of 8 ml fully supplemented cell culture medium or 100 µl per cm<sup>2</sup> Trypsin Neutralizing Solution (TNS). HepesBSS, Trypsin/EDTA Solution and TNS were all contained in Promocell DetachKit (Cat no. C-41200). Cells were then re-suspended in fully supplemented fresh cell culture medium and cultured in T75 cm<sup>2</sup> flasks at a volume of 15 ml.

### Western blotting

Proteins were separated according to their size by 10% sodium dodecyl sulphate polyacrylamide gel electrophoresis (SDS-PAGE). Separated proteins were transferred from polyacrylamide gel onto nitrocellulose membrane (Amersham Hybond ECL, Cat no. RPN303D) for western blot analyses. Ponceau S solution (Sigma, Cat no. P7170) for 5 min to check the transfer efficiency of the protein. Membranes were blocked in 1% w/v non-fat powdered milk (Marvel) before incubation with the following primary and secondary antibodies: anti  $\alpha$ -actin (Sigma-Aldrich, Cat no. A2547) at 500 fold dilution, anti calponin (Sigma-Aldrich, Cat no. C2687) at 500 fold dilution, anti actin (Sigma-Aldrich, Cat no. A4700) at 500 fold dilution and anti-mouse horseradish peroxidase (HRP)-conjugated secondary antibody (Fitzgerald Industries International, Cat no. 43C-CB1569-FIT) at 10,000 fold dilution. Protein bands were detected with ECL western blotting detection reagents (Amersham, Cat no. RPN2109). Bands were sized by referencing to the rainbow protein mass marker (Amersham full-range rainbow MW marker, Cat no. RPN800E).

### Immunocytochemistry

HCASMCs were fixed with 2% w/v paraformaldehyde in PBS (in mM: 2.7 KCl, 1.5 KH<sub>2</sub>PO<sub>4</sub>, 137 NaCl, 8.0 Na<sub>2</sub>HPO<sub>4</sub>, pH 7.4) for 10 min and then incubated in permeabilisation solution (0.1% v/v Triton X-100 in PBS, pH adjusted to 7.4 using 1 M NaOH) for 10 min at room temperature. Next, the cells were incubated with 200 µl antibody diluting solution (in mM: SSC containing 250 NaCl and 15 Na<sub>3</sub>Citrate, 2% v/v goat serum, 0.05% v/v Triton X-100, 1% w/v bovine serum albumin, pH 7.2) for 30 min at room temperature to block non-specific binding. Cells were then incubated overnight at 4 °C with the following primary antibodies: anti  $\alpha$ -actin (Sigma-Aldrich, Cat no. A2547) at 300 fold dilution, anti calponin (Sigma-Aldrich, Cat no. C2687) at 300 fold dilution and anti MHC (Novocastra, Cat no. NCL-MHCs) at 300 fold dilution. On the following day, primary antibodies were indirectly stained using fluorescent Alexa Fluor 488-conjugated goat anti-mouse IgG

secondary antibody (Molecular probes, Cat no. A-11029) at the dilution of 1:500 for 2 h at room temperature. A LSM510 multiphoton laser-scanning confocal microscope was used to visualize the labelled cells. Alexa Fluor 488 was excited at 488 nm using an argon-ion laser, and emission light was collected at 500–550 nm.

### Cell transfection

Plasmids encoding Perceval, PercevalHR and pHRed were obtained from Addgene (Cambridge, MA). HCASMCs expressing PercevalHR or pHRed were transiently transfected using either FuGENE6 (Promega UK, Southampton, UK) or Promofectin (Promocell, Heidelberg, Germany) following manufacturer's instruction. HCASMCs were seeded 24 h prior to transfection at a density of  $1.25 \times 10^5$  per well in 35 mm glass-bottom dishes (Greiner Bio-One, Gloucestershire, UK). The cells expressing Perceval and PercevalHR were used after 24–48 h of transfection, and cells expressing pHRed were also used after 24–48 h of transfection. The cells expressing PercevalHR based on a conventional 3rd generation lentivirus packaging system which has been described previously were used in PercevalHR in vitro calibration experiments (Yang *et al.*, 2017).

### Confocal imaging

Experiments were carried out using either culture medium or physiological saline solution (PSS) containing [in mM]: 120 NaCl, 5 KCl, 1 MgCl<sub>2</sub>, 2 CaCl<sub>2</sub>, 0.42 Na<sub>2</sub>HPO<sub>4</sub>, 0.44 NaH<sub>2</sub>PO<sub>4</sub>, 24 NaHCO<sub>3</sub>, and 10 Glucose. Fluorescence signal of Perceval/PercevalHR and pHRed was measured with LSM510 (Carl Zeiss AG, Oberkochen, Germany), which allowed control of CO<sub>2</sub>, humidity and temperature while environmental O<sub>2</sub> can also be changed to create hypoxic conditions. Manipulating O<sub>2</sub> tension in the custom made micro-imaging chamber (atmospheric gas) was achieved by an O<sub>2</sub> controller (# 0508.000, PeCon GmbH, Germany). This system has been specifically designed to image cultured cells under conditions of varying O<sub>2</sub> tension, achieved by changing the O<sub>2</sub> level in the atmosphere of the micro-imaging chamber. Images in the experiments of PercevalHR calibration were taken using LSM510 high speed multiphoton confocal laser scanning microscope (Carl Zeiss AG, Oberkochen, Germany). Software analysis of the data from LSM510 and LSM510 Multiphoton were carried out by region of interest analysis using AIM software, version 3.2 SP2 (Carl Zeiss AG).

### Measurement of intracellular ATP:ADP ratio, pH and membrane potential

Cells expressing Perceval or PercevalHR were excited with 488 nm light from an argon laser, and emission was collected at 500–550 nm. Cells expressing pHRed were excited at 458 nm, and emission signal over 575 nm was collected. To examine change in membrane potential, HCASMCs were plated into 35 mm glass-bottom dishes a day before experiments. Next day, cells were washed twice with 2 ml PSS, then incubated with 1 μM DiBAC<sub>4</sub>(3) dissolved in PSS for 15 min in a 37 °C/5%CO<sub>2</sub> incubator. Cells were then used without washing the dye as the continued presence of the dye in the extracellular media was required for this experiment. Cells were excited with 488 nm and the emission signal was captured at 500–550 nm. The photon counts (F) from Perceval/PercevalHR, pHRed and DiBAC<sub>4</sub>(3)

were normalized against the initial reading ( $F_0$ ) at the start of experiments and expressed as a ratio (relative fluorescence ( $F/F_0$ )).

### Permeabilisation of HCASMCs with *Staphylococcus aureus* $\alpha$ -toxin

It is of high importance that the signals from biosensors are calibrated in vitro to validate that sensitivity and range are satisfactory for planned experiments. Furthermore, although Perceval and PercevalHR are supposed to detect ATP:ADP ratio, it has been also reported that their signals are independent of ADP (*Li et al., 2013*). Thus, attempts were made to calibrate the PercevalHR using solutions containing known concentration of ATP and ADP. To achieve this, HCASMCs have to be permeabilised in such a way that hydrophilic nucleotides can be introduced into the cells without loss of biosensor. Thus, permeabilisation with  $\alpha$ -toxin was carried out for PercevalHR in vitro calibration. Preliminary experiments resulted in, however, loss of HCASMCs during solution exchange due to cell detachment from the glass. This was due to the excessive HCASMC contraction in the solution once cells were permeabilised. The reason for this is unknown, but in order to minimize loss of cells, 35 mm glass-bottom dishes were coated with Poly D-Lysine and 10  $\mu$ M wortmannin was added to solutions.

HCASMCs were washed with calibration solution A containing (mM) 110 KCl, 30 NaOH, 10 KCl, 10 HEPES, 10 EGTA, and 0.05 EDTA (pH adjusted to 7.2) to remove  $Ca^{2+}$ . Cells were then further washed with calibration solution B containing (in mM) 140 KCl, 10 NaCl, 10 HEPES, 2 EGTA, and 0.05 EDTA. Next, Cells were permeabilised with calibration solution B containing 200  $\mu$ g/ml  $\alpha$ -toxin for about 1 h at room temperature. Finally, cells were washed with calibration solution B and in vitro calibration was carried out using calibration solution B containing required concentrations of nucleotides while keeping free  $Mg^{2+}$  constant at 0.5 mM (All calculations were done using MAXCHELATOR: <http://maxchelator.stanford.edu/>).

### Drugs

DiBAC<sub>4</sub>(3) was obtained from Thermo Fisher Scientific. Rotenone, antimycin, oligomycin B and carbonyl cyanide 3-chlorophenylhydrazone (CCCP) (all from Sigma-Aldrich, Dorset, UK) were dissolved in DMSO as 1, 1, 6 and 10 mM stock solutions, respectively.  $K^+$  channels inhibitors, glibenclamide (Sigma-Aldrich), penitrem A (Alomone labs, Jerusalem, Israel), tram34 (Sigma-Aldrich), apamin (Tocris Bioscience, Abingdon, UK) and XE991 (Tocris Bioscience) were also prepared in DMSO as 10, 1, 10, 1 and 10 mM stock solutions, respectively. Other chemicals were obtained from Sigma-Aldrich or VWR (Leicestershire, UK).

### Statistical analysis

When appropriate, values are expressed as a mean  $\pm$  standard error of mean (SEM). N indicates number of cells unless otherwise stated. Statistical significance was evaluated with a paired Student's *t*-test or with ANOVA with Tukey's test for post hoc analysis using SPSS software package (version 20.0, IBM, New York, NY).

## RESULTS

### **In vitro calibration using $\alpha$ -toxin permeabilized HCASMCs confirms PercevalHR detects ATP concentration over physiological range and is sensitive to intracellular ADP**

Perceval is a fluorescent biosensor of adenylate nucleotides (Berg, Hung & Yellen, 2009) with T loop of ATP binding bacterial protein GlnK1 containing a yellow cpFP and circularly permuted monomeric Venus (cpmVenus). The ATP/ADP binding site is always occupied due to very high affinity, and the competition between ATP and ADP makes the sensor a reporter of intracellular ATP:ADP ratio (Berg, Hung & Yellen, 2009). PercevalHR is an improved fluorescent biosensor and is better tuned to sense the ATP:ADP ratios expected in healthy mammalian cells (Tantama *et al.*, 2013; Tantama & Yellen, 2014).


First, the sensitivity of PercevalHR to ATP over a physiological range and the effect of ADP on signal were assessed using  $\alpha$ -toxin permeabilized HCASMCs. Figure 1A indicates that relative fluorescence ( $F/F_0$ ), determined in the presence of 10 mM ATP at the start of the experiment was reduced by a third with ATP-free solution. Upon the re-introduction of 10 mM ATP, relative fluorescence recovered to the previous level (Fig. 1A). The response of fluorescence probe to 10 mM and 0 mM ATP was stable and repeatable over 100 min, establishing that PercevalHR can report ATP level for a prolonged period.

Next, experiments were carried out to determine the  $K_D$  of PercevalHR in vitro using extracellular solutions containing 10, 3, 1 and 0 mM ATP. Figure 1B depicts a representative result showing confocal images of  $\alpha$ -toxin-permeabilized HCASMCs expressing PercevalHR (Figs. 1B–1F) and the time-course of relative fluorescence change with varying concentrations of extracellular ATP (Fig. 1G). The original data show that the signal strength of PercevalHR was proportional to the concentration of extracellular ATP. When concentration–response curve was constructed, curve fitting showed a half maximal effect occurring at  $\sim 3.3$  mM of ATP (Fig. 1H).

As described earlier, one publication reported that Perceval is insensitive to ADP (Li *et al.*, 2013). This is an important unresolved issue as some metabolically sensitive detectors such as  $K_{ATP}$  channels are regulated by not only absolute amount of ATP but also ADP (Quayle *et al.*, 1994). We therefore studied the effect of intracellular ADP on PercevalHR fluorescence measured using 1 mM ATP in the presence of either 0.1 or 1 mM ADP. As shown in Fig. 1I, increasing ADP concentration from 0.1 mM to 1 mM reduced relative fluorescence significantly ( $p < 0.01$ ) indicating that PercevalHR reports change in ATP:ADP ratio. Taken together, results obtained with permeabilised HCASMCs showed that PercevalHR can be used as a reporter of physiological ATP:ADP change.

### **Removal of glucose from extracellular solution reduces the ATP:ADP ratio in HCASMCs**


Figures 2A–2C shows that Perceval fluorescence remained stable over a period of 40 min when the cell was perfused continuously with PSS. However, upon the removal of 10 mM extracellular glucose, the fluorescent signal dimmed (Figs. 2D–2E) and relative fluorescence decreased by almost 50% over 30 min (Fig. 2F). Removal of glucose significantly reduced relative fluorescence by  $25.7 \pm 8.3\%$  ( $p < 0.05$ ,  $n = 6$ , Fig. 2G). This reduction was not due



**Figure 1** In vitro calibration of PercevalHR. (A) PercevalHR fluorescence from a single HCASMC permeabilized with  $\alpha$ -toxin repeatedly exposed to 10 and 0 mM ATP. (B–G) PercevalHR fluorescence determined over physiological range of ATP concentrations (G) with confocal images of HCASMC (B–F). Scale bar is 50  $\mu$ m. (H) Concentration-response relationship between ATP concentration and corrected fractional fluorescence ( $F/F_0$ ) with  $K_D$  of  $\sim 3.3$  mM ATP ( $n = 4$  from four independent experiments). (I) Mean  $\pm$  SEM of PercevalHR fluorescence determined with 0.1 mM ADP/1 mM ATP is significantly different from that with 1 mM ADP/1 mM ATP ( $p < 0.01$ , paired Student's  $t$ -test,  $n = 5$  from five independent experiments).


Full-size DOI: 10.7717/peerj.10344/fig-1

to the bleaching of biosensor or loss of focus as the re-introduction of 10 mM extracellular glucose restored the signal (original trace, Fig. 2H; summary of 3 cells, Fig. 2I). When the glucose analogue, 2-deoxyglucose (2-DG, 5mM) that cannot be metabolized by the cell, was used, fluorescent signal became weaker (Figs. 2J–2L). Mean results from 3 cells showed that Perceval fluorescence signal was reduced by  $54.0 \pm 15.3\%$  after the application of 2-DG ( $n = 3$ , Fig. 2M). These results indicate that removal of extracellular glucose affects ATP:ADP ratio to a degree that is detectable by Perceval.



**Figure 2** Effect of glucose removal on Perceval signal. (A–C) Perfusion with PSS for 40 min caused little change in fluorescence signal. (D–G) Images of HCASMC expressing Perceval before (D) and after glucose removal (E). The Perceval signal decreased after glucose removal (F). Fractional fluorescence just before (10 Glu) and after (0 Glu) glucose removal was significantly different (G,  $p < 0.05$ , paired Student's  $t$ -test,  $n = 6$  from 6 independent experiments). (H–I) Time course of fractional fluorescence caused by glucose removal and re-administration (H), and Mean  $\pm$  SEM of fractional fluorescence for control, 0 mM glucose and 10 mM glucose (I,  $n = 3$  from three independent experiments). (J–M) Images of HCASMC expressing Perceval at control condition (J) and after the application of 2-DG (K) with time course of fluorescence signal (L). Mean  $\pm$  SEM of fractional fluorescence of Perceval was measured just before (control) and after administration of 2-DG (M, right panel,  $P = 0.112$ , paired Student's  $t$ -test,  $n = 3$  from three independent experiments).

Full-size DOI: [10.7717/peerj.10344/fig-2](https://doi.org/10.7717/peerj.10344/fig-2)



**Figure 3** Effects of metabolic inhibitors on intracellular ATP:ADP ratio. (A–E): Time course of fractional fluorescence change from the cells treated with DMSO (A), 1  $\mu$ M rotenone (B), 1  $\mu$ M antimycin (C), 6  $\mu$ M oligomycin (D) and 1  $\mu$ M CCCP (E). (F): Mean  $\pm$  SEM of fractional fluorescence after application of DMSO ( $p = 0.074$ , paired Student's  $t$ -test,  $n = 4$  from four independent experiments), rotenone ( $p = 0.111$ , paired Student's  $t$ -test,  $n = 3$  from three independent experiments), antimycin ( $p < 0.05$ , paired Student's  $t$ -test,  $n = 4$  from four independent experiments), oligomycin ( $p = 0.085$ , paired Student's  $t$ -test,  $n = 3$  from three independent experiments) and CCCP, measured at the peak ( $p < 0.05$ , paired Student's  $t$ -test,  $n = 3$  from three independent experiments).

Full-size DOI: 10.7717/peerj.10344/fig-3

### Inhibition of oxidative phosphorylation causes a measurable decrease in cellular ATP:ADP ratio

ATP can be generated using extracellular glucose by oxidative phosphorylation and glycolysis pathways. Therefore, whether inhibition of oxidative phosphorylation affects PercevalHR signal was examined next in the cell culture medium. Oxidative phosphorylation was blocked by a number of approaches: inhibiting the electron transport chain with rotenone (1  $\mu$ M), an inhibitor of mitochondrial complex I, or antimycin (1  $\mu$ M), an inhibitor of mitochondrial complex III; inhibiting ATPase activity with oligomycin (6  $\mu$ M), or through use of the uncoupling agent CCCP (1  $\mu$ M). As shown in Figs. 3B–3E, application of metabolic inhibitors leads to a decrease in relative fluorescence. The effects of metabolic inhibitors, along with that of vehicle (DMSO), are summarized in Fig. 3F. When compared to the fluorescence ratio just before the administration of metabolic inhibitors, application of antimycin and CCCP induced significant reduction in ATP:ADP ratio signal (both  $p < 0.05$ ).

Like other biosensors, Perceval may respond to changes in intracellular pH, which may accompany hypoxia (Berg, Hung & Yellen, 2009). Thus, the effect of hypoxia on the fluorescence of a pH sensor, pHRed, was also examined (Tantama, Hung & Yellen, 2011).

pHRed responds robustly to intracellular alkalization and acidification induced by the addition and subsequent removal of extracellular  $\text{NH}_4\text{Cl}$  (20 mM) (Dart & Vaughan-Jones, 1992) (Fig. S3). There was little change in intracellular pH in response to metabolic interventions, as assessed by pHRed in cell culture medium (Fig. S4).


### Effect of hypoxia on cellular ATP:ADP ratio and pH

Figure 4A-C show confocal images of HCASMC transiently transfected with Perceval under normoxia (20% oxygen), hypoxia (1%  $\text{O}_2$ ), and following the restoration of oxygen (20%). Fractional ATP:ADP ratio was reduced by about 10% by hypoxia and recovered upon oxygen reintroduction (Fig. 4D). Exposure to hypoxia significantly reduced Perceval signal by  $10.3 \pm 2.1\%$ , and the change was reversible ( $p < 0.05$ ,  $n = 5$ , Fig. 4E). There was no change in Perceval signal when mild hypoxia (10% and 5%  $\text{O}_2$ ) was tested (Fig. S5). pHRed however detected no change in intracellular pH after exposure to hypoxia (Figs. 4F–4I), suggesting that the fluorescence change induced in Perceval was due to changes in the ATP:ADP ratio and not an artefact caused by shift in intracellular pH. All the experiments were conducted in cell culture medium.

### Identification of $\text{K}^+$ channels setting resting membrane potential of HCASMCs

DiBAC<sub>4</sub>(3) is a slow-response potential-sensitive fluorescent probe (Apell & Bersch, 1987; Mustafa et al., 2011), and thus a useful tool for HCASMCs where changes in membrane potential are generally slow and steady. It is permeable to the plasma membrane, accumulating in the cytoplasm following a Nernst equilibrium distribution (Apell & Bersch, 1987). The anionic dye in the cell binds to intracellular proteins or to membranes, resulting in an enhanced fluorescence and also a red spectral shift (Epps, Wolfe & Groppi, 1994; Klapperstuck et al., 2009). Depolarization of membrane potential causes entry of the anionic dye into the cell, increasing the fluorescence signal. Hyperpolarization, on the other hand, results in a decrease in the signal. Compared to cationic carbocyanines, DiBAC<sub>4</sub>(3) is largely excluded from mitochondria due to their overall negative charge, and so largely measures changes in the plasma membrane potential (Apell & Bersch, 1987; Baczkó, Giles & Light, 2004). To establish that DiBAC<sub>4</sub>(3) can detect moderate changes in membrane potential in HCASMCs, initial experiments were designed to characterize the changes in the relative fluorescence ( $F/F_0$ ) using the extracellular solutions containing various concentrations of  $[\text{K}^+]_o$  (5, 30, and 80 mM). Resting membrane potential, assumed to mainly reflect  $\text{K}^+$  equilibrium potential, can be calculated with the Nernst equation. A representative characterization experiment is depicted in supplementary data, showing that the relationship of relative fluorescence and predicted membrane potential, note that the cultured HCASMCs have a more depolarized membrane potential at rest (Fig. S6).


Like other excitable cells, HCASMCs express multiple types of  $\text{K}^+$  channels (Nelson & Quayle, 1995; Standen & Quayle, 1998). In order to investigate which  $\text{K}^+$  channel(s) are important in setting the resting membrane potential, a series of  $\text{K}^+$  channel inhibitors were applied. These included glibenclamide (10  $\mu\text{M}$ ), penitrem A (200 nM), tram34 (1  $\mu\text{M}$ ), apamin (200 nM),  $\text{BaCl}_2$  (25  $\mu\text{M}$ ) and XE991 (10  $\mu\text{M}$ ) which are inhibitors of



**Figure 4** Effect of hypoxia on intracellular ATP:ADP ratio. (A–C) Confocal images of HCASMC expressing Perceval under control condition (A), after 10 minutes exposure to hypoxia (B), and recovery after re-oxygenation (C). (D) Time course of Perceval fractional fluorescence. (E) Exposure to hypoxia (1%  $O_2$ ) significantly changed Perceval fluorescence compared to control (basal) and recovery (20%  $O_2$ ) ( $p < 0.05$ , ANOVA with Tukey's test for post hoc analysis,  $n = 5$  from five independent experiments). (F–G) Confocal images of HCASMC expressing pHRed under control condition and during exposure to 1%  $O_2$ . (H): Time course of fractional fluorescence. (I) Hypoxia caused little change in intracellular pH (bottom panel,  $n = 5$  from five independent experiments).

Full-size DOI: 10.7717/peerj.10344/fig-4

$K_{ATP}$  channels, calcium-activated large, intermediate and small conductance potassium ( $BK_{Ca}$ ,  $IK_{Ca}$ ,  $SK_{Ca}$ ) channels, inward rectifying potassium ( $K_{ir}$ ) channels and voltage-gated potassium ( $K_v7$ ) channels respectively (Asano *et al.*, 2012; Blatz & Magleby, 1987; Gluais *et al.*, 2005; Ko *et al.*, 2008; Mackie & Byron, 2008; Nelson & Quayle, 1995; Wulff & Castle, 2010). These inhibitors were used at concentrations thought to be selective for their respective class of channel (Alexander, Mathie & Peters, 2009). An increase in DiBAC<sub>4</sub>(3) signal (membrane depolarization) upon inhibition of a particular class of  $K^+$  channel would indicate that this channel is important in maintaining the resting membrane potential. Figs. 5A–5F show the representative time-course of relative fluorescence observed with different  $K^+$  channel inhibitors. The fractional fluorescence, determined as the difference between the relative fluorescence of DiBAC<sub>4</sub>(3) before and after drug application and expressed as percentage change is summarized in Fig. 5G. Application of glibenclamide and  $BaCl_2$  caused a significant change in fractional fluorescence ( $p < 0.001$ ), suggesting



**Figure 5** The identification of  $K^+$  channels in setting HCASMC resting membrane potential. (A–F) The change in the fractional fluorescence of DiBAC<sub>4</sub>(3) treated with 10  $\mu$ M glibenclamide (A), 200 nM penitrem A (B), 1  $\mu$ M tram34 (C), 200 nM apamin (D), 25  $\mu$ M BaCl<sub>2</sub> (E), and 10  $\mu$ M XE991 (F). (G) A summary of percent change in fractional fluorescence of DiBAC<sub>4</sub>(3) after application of  $K^+$  channel inhibitors: glibenclamide ( $p < 0.001$ , paired Student's  $t$ -test,  $n = 37$  from three independent experiments), penitrem A  $n = 18$  from two independent experiments), tram34 ( $n = 13$  from 1 experiment), apamin ( $n = 32$  from two independent experiments), BaCl<sub>2</sub> ( $p < 0.001$ , paired Student's  $t$ -test,  $n = 26$  from two independent experiments), and XE991 ( $n = 14$  from two independent experiments).


Full-size [DOI: 10.7717/peerj.10344/fig-5](https://doi.org/10.7717/peerj.10344/fig-5)

that  $K_{ATP}$  and  $K_{ir}$  channels play major roles in setting resting membrane potential of HCASMCs.

### Effects of metabolic inhibitors and hypoxia on membrane potential

Of the  $K^+$  channels involved in setting the resting membrane potential of HCASMCs,  $K_{ATP}$  channels seem a likely candidate responsible for hypoxic vasodilation. Initial experiments using the  $K_{ATP}$  channel opener, pinacidil (10  $\mu$ M), caused a significant decrease in DiBAC<sub>4</sub>(3) fluorescence (Fig. S7), suggesting that  $K_{ATP}$  channels have spare capacity to increase their open probability and/or active channel number from the resting condition. Therefore, the effect of metabolic insult and hypoxia on the DiBAC<sub>4</sub>(3) signal was examined.

Inhibiting mitochondrial complex III with antimycin (1  $\mu$ M) induced a significant decrease in DiBAC<sub>4</sub>(3) fluorescence (Figs. 6A–6B), which was blocked by glibenclamide (10  $\mu$ M) or 60  $K^+$  (Figs. 6C–6D), an indication that  $K_{ATP}$  channels may be involved. The



**Figure 6** Effects of antimycin and hypoxia on membrane potential. (A–B) Application of 1  $\mu\text{M}$  antimycin caused significant reduction in fractional fluorescence ( $p < 0.001$ , paired Student's  $t$ -test,  $n = 25$  from one experiment). (C–D) Pretreating the cells either with 10  $\mu\text{M}$  glibenclamide (C,  $n = 22$  from one experiment) or 60  $\text{K}^+$  (D,  $n = 6$  from one experiment) abolished the change induced by antimycin. (E–F) Hypoxia caused significant decrease in fractional fluorescence ( $p < 0.001$ , paired Student's  $t$ -test,  $n = 72$  from 11 independent experiments). (G–H): The effect of hypoxia was not blocked by 10  $\mu\text{M}$  glibenclamide (G  $p < 0.001$ , paired Student's  $t$ -test,  $n = 36$  from four independent experiments), but hypoxia was ineffective with extracellular 60  $\text{K}^+$  (H,  $n = 11$  from two independent experiments, NS).

Full-size [DOI: 10.7717/peerj.10344/fig-6](https://doi.org/10.7717/peerj.10344/fig-6)

effect induced by hypoxia on DiBAC<sub>4</sub>(3) fluorescence was highly significant (Figs. 6E–6F). The effect from hypoxia was blocked by 60  $\text{K}^+$  (Fig. 6H), suggesting the involvement of  $\text{K}^+$  channels. However, exposure to hypoxia resulted in significant decrease in DiBAC<sub>4</sub>(3) fluorescence when in the presence of 10  $\mu\text{M}$  pinacidil ( $n = 5$ ), 10  $\mu\text{M}$  glibenclamide ( $n = 36$ ), 25  $\mu\text{M}$  BaCl<sub>2</sub> ( $n = 14$ ), 100 nM IbTX ( $n = 17$ ), 200 nM penitrem A ( $n = 14$ ), and 10  $\mu\text{M}$  glibenclamide plus 25  $\mu\text{M}$  BaCl<sub>2</sub> ( $n = 25$ ) (Fig. S8). These results indicate that  $\text{K}^+$  channels are likely to be involved in hypoxia-induced membrane hyperpolarization reported by DiBAC<sub>4</sub>(3), but classic  $\text{K}^+$  channel inhibitors, when applied singly or in combination in the case of glibenclamide plus BaCl<sub>2</sub>, failed to block the effect of hypoxia.

## DISCUSSION

Many studies have reported that hypoxia activates various  $\text{K}^+$  channels in systemic arteries at different locations and of different species (Dart & Standen, 1995; Gebremedhin, Yamaura & Harder, 2008; Park et al., 2005; Park et al., 2007; Shimoda & Polak, 2011). Of these,  $\text{K}_{\text{ATP}}$  channels seem uniquely suited to act as the molecular targets, but to our knowledge, changes in nucleotide concentrations have not been measured in real time using VSMCs. The key issue addressed in the current paper was, therefore, whether the ATP:ADP ratio changes when HCASMCs are challenged by hypoxia and metabolic insults. This is the next step from our previous paper using the Seahorse technique showing that metabolic intervention resulted in a decrease, measured at the end point, in both cellular ATP and ATP:ADP ratio (Yang et al., 2017). The main finding of current paper is that

1% O<sub>2</sub> hypoxia caused a significant decrease in ATP:ADP ratio in HCASMCs. Although this is somewhat surprising as we found that approximately half of ATP in these cells was produced by glycolysis that does not require oxygen (Yang *et al.*, 2017), our present result shows that severe hypoxia can change the ATP:ADP ratio and that Perceval/PercevalHR can report such a change.

Compared to total ATP levels, the ATP:ADP ratio is arguably a more important parameter in cellular energy status. Perceval is originally constructed to monitor the changes in cellular ATP:ADP ratio (Berg, Hung & Yellen, 2009; Li *et al.*, 2013). While Perceval, an original nucleotide biosensor, saturates at low ATP:ADP ratio (ATP:ADP <5) and is therefore tuned to extreme metabolic inhibition (Berg, Hung & Yellen, 2009), mammalian ATP:ADP ratios are estimated to range from 1 to 100 in healthy conditions (Traut, 1994; Veech *et al.*, 1979). The improved sensor PercevalHR measures ATP:ADP ratio over the range expected in mammalian cells (Tantama *et al.*, 2013; Tantama & Yellen, 2014). We report here that, using  $\alpha$ -toxin permeabilized HCASMCs, K<sub>D</sub> of PercevalHR for ATP is ~3.3 mM (Fig. 1). It could be argued that, because the absolute amount of ATP, ADP and AMP can vary widely among and possibly within the cells, measurement of intracellular ATP:ADP and ATP:AMP ratios provide more reliable information of cellular metabolism (Berg, Hung & Yellen, 2009; Tantama & Yellen, 2014). When Perceval/PercevalHR were introduced to the cells to report intracellular ATP:ADP ratio, removal of glucose and inhibition of either glycolysis or oxidative phosphorylation with metabolic inhibitors all resulted in decreases in cellular ATP:ADP ratio (Figs. 2 and 3). Hypoxia induced a decrease in cellular ATP:ADP ratio (Fig. 4). The effect was not caused by a change in pH as parallel studies showed hypoxia exerted no change in pHRed signal (Fig. 4).

Although [ATP]<sub>i</sub> is usually high enough to inhibit K<sub>ATP</sub> channels, K<sub>ATP</sub> channels may not sense global cytosolic ATP ([ATP]<sub>c</sub>). Rather, local concentrations of nucleotides in the microdomains close to K<sub>ATP</sub> channels may be crucial, and it is conceivable that subcellular nucleotide concentrations dynamically change due to the activities of membrane ATPases and phosphotransfer enzymes. It has been reported that ATP binding site on K<sub>ir</sub>6.2 subunit is estimated to be ~2 nm below the membrane and at the interface between adjacent K<sub>ir</sub>6.2 (Antcliff *et al.*, 2005). Evidence has also suggested that local nucleotide level is modified when bulk ATP was transmitted over the diffusional barrier into the sub-membrane space due to the effect from metabolic enzymes (Flagg *et al.*, 2010; Foster & Coetzee, 2016; Selivanov *et al.*, 2004). AMP is suggested to regulate ATP:ADP ratio in the near membrane area by being involved in the phosphotransfer reactions mediated by Adenylate kinase (AK), resulting in a change sufficient to cause K<sub>ATP</sub> channel opening (Carrasco *et al.*, 2001). Taken together, ATP compartmentalization/intracellular concentration gradient may change sufficiently to regulate K<sub>ATP</sub> channels whilst [ATP]<sub>c</sub> is unchanged.

A biosensor to monitor pH was used in this study to ensure that the change in Perceval/PercevalHR signal was not an artifact due to acidification/alkalization of cells. A genetically encoded pH sensor, pHRed, made by mutagenesis of the red fluorescent protein mKeima, was used in this study (Tantama, Hung & Yellen, 2011). Metabolic inhibition of HCASMCs with 5 mM 2-DG when measured in bicarbonate buffered PSS or hypoxia when measured in cell culture medium induced no change in intracellular

pH, and this is consistent with the results of other researchers (Imamura *et al.*, 2009). Therefore, it seems safe to ascribe changes in Pervceval/PercevalHR signal to true changes in ATP:ADP. Indeed, intracellular pH of HCASMCs is rather well maintained as cells can handle pH changes via ion transporters on the cell membrane and by intracellular buffering (Boedtkjer, Praetorius & Aalkjaer, 2006; Madden, Keller & Kleinman, 2000). There is, however, reports showing that pH can change with metabolic inhibition (Imamura *et al.*, 2009). In response to 2-DG, Kiang, McKinney & Gallin (1990) reported a  $\sim 0.2$  pH change in A-431 cells (Kiang, McKinney & Gallin, 1990), and Brown *et al.* (1991) found the same degree of pH change in isolated epithelial cells (Brown *et al.*, 1991). It has been also reported that, in Neuro2A cells, metabolic inhibition by oligomycin and carbonyl cyanide 4-(trifluoromethoxy)phenylhydrazone (FCCP) showed no effect on cytosol pH whilst complete withdrawal of glucose acidified the cytosol (Tantama, Hung & Yellen, 2011). In the same study, it was shown that when pHRed was targeted to the mitochondrial matrix by COX8, it was reported an acidification of the mitochondrial matrix after the application of FCCP (Tantama, Hung & Yellen, 2011). Due to their crucial and multi-tasking function including ATP production cell signaling, cell differentiation, cell cycle and cell death, it would also be interesting to look at mitochondrial ATP specifically, especially when measured in combination with other substrates, such as  $\text{Ca}^{2+}$ . However, it is beyond the scope of this study.

Many studies suggested that hypoxia causes vasodilation by decreasing  $[\text{Ca}^{2+}]_i$ . Some investigations indicated that hypoxia has a direct effect on  $\text{Ca}^{2+}$  channels while others have shown that hypoxia causes membrane potential hyperpolarization and a closure of VDCCs (Dart & Standen, 1995; Flagg *et al.*, 2010; Francoobregon, Urena & Lopezbarneo, 1995; Welsh, Jackson & Segal, 1998). The observation in our lab that hypoxia inhibited  $\text{Ca}^{2+}$  oscillations in the presence of vasoconstrictors also indicates that  $\text{Ca}^{2+}$  handling itself is affected by hypoxia where the net effect would be reduction in  $[\text{Ca}^{2+}]_i$  and muscle relaxation (Fig. S9, Table S1). Thus, the activation of  $\text{K}^+$  channels during hypoxic coronary vasodilation is aided by other mechanisms such as suppression of  $[\text{Ca}^{2+}]_i$  transient during hypoxic vasodilation. Recent studies have also suggested that heme-protein myoglobin in VSMCs can produce nitric oxide (NO) through the reduction of endogenous nitrite under hypoxia and there leads to vasodilation, which regulates vasodilation via myoglobin-dependent NO generation and is independent of the well-known endothelial NO synthase (eNOS) (Hendgen-Cotta, Kelm & Rassaf, 2014; Totzeck *et al.*, 2012).

At least two types of  $\text{K}^+$  channels are involved in setting resting membrane potential in HCASMCs, as application of  $10 \mu\text{M}$  glibenclamide and  $25 \mu\text{M}$   $\text{BaCl}_2$  caused significant increase in DiBAC<sub>4</sub>(3) signal (Fig. 5).  $25 \mu\text{M}$   $\text{BaCl}_2$  also caused membrane potential depolarization in the presence of  $10 \mu\text{M}$  glibenclamide (data not shown). Although  $\text{Ba}^{2+}$  inhibits multiple  $\text{K}^+$  channels including  $\text{K}_{\text{ATP}}$  channels at a higher concentration, at the concentration used in this study,  $\text{BaCl}_2$  should act as a specific blocker of  $\text{K}_{\text{ir}}$  channels (Nelson & Quayle, 1995; Quayle *et al.*, 1993). The additive effect of glibenclamide and  $\text{BaCl}_2$  further supports the hypothesis that both  $\text{K}_{\text{ATP}}$  channels and  $\text{K}_{\text{ir}}$  channels regulate resting membrane potential in HCASMCs. Further characterization of  $\text{K}^+$  channels using selective activators and inhibitors showed that HCASMCs functionally express  $\text{K}_{\text{ATP}}$ ,  $\text{BK}_{\text{Ca}}$ , and

IK<sub>Ca</sub>/SK<sub>Ca</sub> channels. In HCASMCs, changes in membrane potential caused by 10 μM NS1619 and 10 μM NS11021, both activators of BK<sub>Ca</sub> channels, were not inhibited by 100 nM iberitoxin (IbTX) (data not shown), and this could be due to the presence of a specific β subunit in human coronary BK<sub>Ca</sub> channels. A total of four types of modulatory β subunits have been identified, and in HCASMCs the majority of BK<sub>Ca</sub> channels contain β1 subunit (KCNMB1) (Tanaka *et al.*, 1997), which has higher sensitivity to Ca<sup>2+</sup> and a lower sensitivity to IbTX (Dworetzky *et al.*, 1996).

The most puzzling result regarding hypoxia effects on membrane potential was that none of the traditional K<sup>+</sup> channel inhibitors eliminated the reduction in signal caused by hypoxia, although the effect of hypoxia was absent in high [K<sup>+</sup>]<sub>o</sub>. The lack of effect of traditional K<sup>+</sup> channel inhibitors, when applied singly or as a combination of glibenclamide plus BaCl<sub>2</sub>, may be explained by the hypothesis that more than one class of K<sup>+</sup> channel is involved, and so a combination of antagonists are required to block the hypoxia effect. Also, two-pore domain potassium (K2P) channels were not investigated in this study as there are no specific pharmacological inhibitors that can be used to dissect out the role of these channels. Moreover, the sensitivity of K<sup>+</sup> channels to inhibitors may be reduced during hypoxia (Venkatesh, Lamp & Weiss, 1991). If the pharmacological tools are rendered less effective during hypoxia, then knocking out/down individual K<sup>+</sup> channels using KO mice and siRNA might be useful to address some of these unanswered questions in future experiments.

Although cell culture is typically carried out at an atmospheric condition with an oxygen concentration of 20–21%, we are aware that cells *in vivo* reside under considerably lower oxygen tension, named physiologically relevant O<sub>2</sub> concentration, also known as physioxia (Carreau *et al.*, 2011). It has been reported that standard *in vitro* cultured cells experience a constant state of oxidative stress due to hyperoxic environment (Ferguson *et al.*, 2018). Therefore, most of the contemporary experiments using *in vitro* cell culture are conducted with oxidatively stressed cells. Culture in hyperoxic conditions has conferred a protective resilience to many kinds of oxidative stressors, which is thought to be an artefact arising from the current culture leading to poor translation of investigations from *in vitro* to *in vivo* (Ferguson *et al.*, 2018; Lin *et al.*, 2020). On the contrary, the extent of antioxidant enzyme activities and redox homeostasis are examined to be closer to *in vivo* conditions if the cells are pre-conditioned under physioxia (Pastore *et al.*, 2001; Shindo *et al.*, 1994). Evidence has also pointed to a distinct pattern of protein expression between the two, and microRNAs are strong regulators of the transcriptome under hypoxic and physioxic conditions (Carreau *et al.*, 2011). The different observations between hyperoxic and physioxic conditions should be well considered when looking at functional parameters in cells cultured *in vitro*, especially in experiments under various O<sub>2</sub> concentrations. Furthermore, a range of physiologically-relevant O<sub>2</sub> concentrations is suggested in order to reach a better understanding of the full scope of cellular responses as O<sub>2</sub> level is different within a tissue. Although we have tried a range of O<sub>2</sub> concentrations including 20%, 5% and 1% in this study, it is also crucial to conduct experiments in a variety of physiological O<sub>2</sub> concentrations to make further understanding of the underlying K<sub>ATP</sub> channel hypothesis. Moreover, suggestions have been made that chronic culture of mammalian cell culture

under physioxia before exposure to experimental stimuli should be incorporated into standard protocols, which have been proved to be relatively easy and inexpensive to achieve (Ferguson et al., 2018).

## CONCLUSIONS

Hypoxic vasodilation is crucial to the regulation of human coronary blood flow, but underlying cellular and molecular mechanisms are unknown. The current paper provides evidence that direct opening of K<sup>+</sup> channels of HCASMCs occurs in metabolic and hypoxic coronary vasodilation. Combined with the reports that altered Ca<sup>2+</sup> and K<sup>+</sup> channel function may be important in the coronary artery vascular changes seen in angina, diabetes and metabolic syndrome (Borbouse et al., 2010; Chutkow et al., 2002b; Dick & Tune, 2010; Miki et al., 2002; Miura et al., 2003), elucidation of detailed mechanisms underpinning hypoxic vasodilation will aid in development of therapeutic strategies to treat diseases associated with hypoxia/ischemia by inducing local vasodilation.

## ACKNOWLEDGEMENTS

We would like to thank Dr. Joanna Wardyn for her expert help with lentivirus and the Centre for Cellular Imaging, University of Liverpool, for confocal facilities.

## ADDITIONAL INFORMATION AND DECLARATIONS

### Funding

Current work was funded by the China Scholarship Council and University of Liverpool awarded to Mingming Yang and John Martin Quayle (201206090033), and the Fundamental Research Funds for the Central Universities awarded to Mingming Yang (2242019K40232). The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

### Grant Disclosures

The following grant information was disclosed by the authors:

China Scholarship Council and University of Liverpool awarded to Mingming Yang and John Martin Quayle: 201206090033.

The Fundamental Research Funds for the Central Universities awarded to Mingming Yang: 2242019K40232.

### Competing Interests

The authors declare there are no competing interests.

### Author Contributions

- Mingming Yang conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

- Caroline Dart and John M. Quayle conceived and designed the experiments, analyzed the data, authored or reviewed drafts of the paper, and approved the final draft.
- Tomoko Kamishima conceived and designed the experiments, performed the experiments, analyzed the data, prepared figures and/or tables, authored or reviewed drafts of the paper, and approved the final draft.

### Data Availability

The following information was supplied regarding data availability:

The raw measurements are available in the [Supplementary Files](#).

### Supplemental Information

Supplemental information for this article can be found online at <http://dx.doi.org/10.7717/peerj.10344#supplemental-information>.

## REFERENCES

- Alexander SPH, Mathie A, Peters JA. 2009.** Guide to receptors and channels (GRAC), 4th edition. *British Journal of Pharmacology* **158**:S1–S239  
[DOI 10.1111/j.1476-5381.2009.00499.x](https://doi.org/10.1111/j.1476-5381.2009.00499.x).
- Antcliff JF, Haider S, Proks P, Sansom MS, Ashcroft FM. 2005.** Functional analysis of a structural model of the ATP-binding site of the KATP channel Kir6.2 subunit. *Embo Journal* **24**:229–239 [DOI 10.1038/sj.emboj.7600487](https://doi.org/10.1038/sj.emboj.7600487).
- Apell HJ, Bersch B. 1987.** Oxonol VI as an optical indicator for membrane potentials in lipid vesicles. *Biochimica et Biophysica Acta/General Subjects* **903**:480–494  
[DOI 10.1016/0005-2736\(87\)90055-1](https://doi.org/10.1016/0005-2736(87)90055-1).
- Asano S, Bratz IN, Berwick ZC, Fancher IS, Tune JD, Dick GM. 2012.** Penitrem A as a tool for understanding the role of large conductance Ca(2+)/voltage-sensitive K(+) channels in vascular function. *Journal of Pharmacology and Experimental Therapeutics* **342**:453–460 [DOI 10.1124/jpet.111.191072](https://doi.org/10.1124/jpet.111.191072).
- Baczkó I, Giles WR, Light PE. 2004.** Pharmacological activation of plasma-membrane KATP channels reduces reoxygenation-induced Ca(2+) overload in cardiac myocytes via modulation of the diastolic membrane potential. *British Journal of Pharmacology* **141**:1059–1067 [DOI 10.1038/sj.bjp.0705702](https://doi.org/10.1038/sj.bjp.0705702).
- Berg J, Hung YP, Yellen G. 2009.** A genetically encoded fluorescent reporter of ATP:ADP ratio. *Nature Methods* **6**:161–166 [DOI 10.1038/nmeth.1288](https://doi.org/10.1038/nmeth.1288).
- Blatz AL, Magleby KL. 1987.** Calcium-activated potassium channels. *Trends in Neurosciences* **10**:463–467 [DOI 10.1016/0166-2236\(87\)90101-9](https://doi.org/10.1016/0166-2236(87)90101-9).
- Boedtker E, Praetorius J, Aalkjaer C. 2006.** NBCn1 (slc4a7) mediates the Na<sup>+</sup>-dependent bicarbonate transport important for regulation of intracellular pH in mouse vascular smooth muscle cells. *Circulation Research* **98**:515–523  
[DOI 10.1161/01.Res.0000204750.04971.76](https://doi.org/10.1161/01.Res.0000204750.04971.76).
- Borbouse L, Dick GM, Payne GA, Berwick ZC, Neeb ZP, Alloosh M, Bratz IN, Sturek M, Tune JD. 2010.** Metabolic syndrome reduces the contribution of K<sup>+</sup> channels to

- ischemic coronary vasodilation. *American Journal of Physiology-Heart and Circulatory Physiology* **298**:H1182–H1189 DOI [10.1152/ajpheart.00888.2009](https://doi.org/10.1152/ajpheart.00888.2009).
- Brown SES, Heming TA, Benedict CR, Bidani A. 1991.** ATP-sensitive Na<sup>+</sup>-H<sup>+</sup> antiport in type-II alveolar epithelial-cells. *American Journal of Physiology* **261**:C954–C963 DOI [10.1152/ajpcell.1991.261.6.C954](https://doi.org/10.1152/ajpcell.1991.261.6.C954).
- Carrasco AJ, Dzeja PP, Alekseev AE, Pucar D, Zingman LV, Abraham MR, Hodgson D, Bienengraeber M, Puceat M, Janssen E, Wieringa B, Terzic A. 2001.** Adenylate kinase phosphotransfer communicates cellular energetic signals to ATP-sensitive potassium channels. *Proceedings of the National Academy of Sciences of the United States of America* **98**:7623–7628 DOI [10.1073/pnas.121038198](https://doi.org/10.1073/pnas.121038198).
- Carreau A, El Hafny-Rahbi B, Matejuk A, Grillon C, Kieda C. 2011.** Why is the partial oxygen pressure of human tissues a crucial parameter? Small molecules and hypoxia. *Journal of Cellular and Molecular Medicine* **15**:1239–1253 DOI [10.1111/j.1582-4934.2011.01258.x](https://doi.org/10.1111/j.1582-4934.2011.01258.x).
- Chutkow WA, Pu J, Wheeler MT, Wada T, Makielski JC, Burant CF, McNally EM. 2002a.** Episodic coronary artery vasospasm and hypertension develop in the absence of Sur2 K(ATP) channels. *Journal of Clinical Investigation* **110**:203–208 DOI [10.1172/jci15672](https://doi.org/10.1172/jci15672).
- Chutkow WA, Pu J, Wheeler MT, Wada T, Makielski JC, Burant CF, McNally EM. 2002b.** Episodic coronary artery vasospasm and hypertension develop in the absence of Sur2 KATP channels. *Journal of Clinical Investigation* **110**:203–208 DOI [10.1172/JCI15672](https://doi.org/10.1172/JCI15672).
- Dart C, Standen NB. 1995.** Activation of ATP-dependent K<sup>+</sup> channels by hypoxia in smooth muscle cells isolated from the pig coronary artery. *Journal de Physiologie* **483**(Pt 1):29–39 DOI [10.1113/jphysiol.1995.sp020565](https://doi.org/10.1113/jphysiol.1995.sp020565).
- Dart C, Vaughan-Jones RD. 1992.** Na<sup>(+)</sup>-HCO<sub>3</sub><sup>-</sup> symport in the sheep cardiac Purkinje fibre. *Journal de Physiologie* **451**:365–385 DOI [10.1113/jphysiol.1992.sp019169](https://doi.org/10.1113/jphysiol.1992.sp019169).
- Daut J, Maier-Rudolph W, Von Beckerath N, Mehrke G, Gunther K, Goedel-Meinen L. 1990.** Hypoxic dilation of coronary arteries is mediated by ATP-sensitive potassium channels. *Science* **247**:1341–1344 DOI [10.1126/science.2107575](https://doi.org/10.1126/science.2107575).
- Davies LM, Purves GI, Barrett-Jolley R, Dart C. 2010.** Interaction with caveolin-1 modulates vascular ATP-sensitive potassium (KATP) channel activity. *Journal de Physiologie* **588**:3255–3266 DOI [10.1113/jphysiol.2010.194779](https://doi.org/10.1113/jphysiol.2010.194779).
- Dick GM, Tune JD. 2010.** Role of potassium channels in coronary vasodilation. *Experimental Biology and Medicine* **235**:10–22 DOI [10.1258/ebm.2009.009201](https://doi.org/10.1258/ebm.2009.009201).
- Duncker DJ, Bache RJ. 2008.** Regulation of coronary blood flow during exercise. *Physiological Reviews* **88**:1009–1086 DOI [10.1152/physrev.00045.2006](https://doi.org/10.1152/physrev.00045.2006).
- Dworetzky SI, Boissard CG, Lum-Ragan JT, McKay MC, Post-Munson DJ, Trojnecki JT, Chang CP, Gribkoff VK. 1996.** Phenotypic alteration of a human BK (hSlo) channel by hSlobeta subunit coexpression: changes in blocker sensitivity, activation/relaxation and inactivation kinetics, and protein kinase A modulation. *Journal of Neuroscience* **16**:4543–4550 DOI [10.1523/jneurosci.16-15-04543.1996](https://doi.org/10.1523/jneurosci.16-15-04543.1996).

- Epps DE, Wolfe ML, Groppi V. 1994.** Characterization of the steady-state and dynamic fluorescence properties of the potential-sensitive dye bis-(1, 3-dibutylbarbituric acid)trimethine oxonol (Dibac4(3)) in model systems and cells. *Chemistry and Physics of Lipids* **69**:137–150 DOI [10.1016/0009-3084\(94\)90035-3](https://doi.org/10.1016/0009-3084(94)90035-3).
- Ferguson DCJ, Smerdon GR, Harries LW, Dodd NJF, Murphy MP, Curnow A, Winyard PG. 2018.** Altered cellular redox homeostasis and redox responses under standard oxygen cell culture conditions versus physioxia. *Free Radical Biology and Medicine* **126**:322–333 DOI [10.1016/j.freeradbiomed.2018.08.025](https://doi.org/10.1016/j.freeradbiomed.2018.08.025).
- Flagg TP, Enkvetchakul D, Koster JC, Nichols CG. 2010.** Muscle KATP channels: recent insights to energy sensing and myoprotection. *Physiological Reviews* **90**:799–829 DOI [10.1152/physrev.00027.2009](https://doi.org/10.1152/physrev.00027.2009).
- Foster MN, Coetzee WA. 2016.** KATP channels in the cardiovascular system. *Physiological Reviews* **96**:177–252 DOI [10.1152/physrev.00003.2015](https://doi.org/10.1152/physrev.00003.2015).
- Francoobregon A, Urena J, Lopezbarneo J. 1995.** Oxygen-sensitive calcium channels in vascular smooth-muscle and their possible role in hypoxic arterial relaxation. *Proceedings of the National Academy of Sciences of the United States of America* **92**:4715–4719 DOI [10.1073/pnas.92.10.4715](https://doi.org/10.1073/pnas.92.10.4715).
- Gebremedhin D, Yamaura K, Harder DR. 2008.** Role of 20-HETE in the hypoxia-induced activation of Ca<sup>2+</sup>-activated K<sup>+</sup> channel currents in rat cerebral arterial muscle cells. *American Journal of Physiology-Heart and Circulatory Physiology* **294**:H107–H120 DOI [10.1152/ajpheart.01416.2006](https://doi.org/10.1152/ajpheart.01416.2006).
- Gluais P, Edwards G, Weston AH, Falck JR, Vanhoutte PM, Feletou M. 2005.** Role of SKCa and IKCa in endothelium-dependent hyperpolarizations of the guinea-pig isolated carotid artery. *British Journal of Pharmacology* **144**:477–485 DOI [10.1038/sj.bjp.0706003](https://doi.org/10.1038/sj.bjp.0706003).
- Hendgen-Cotta UB, Kelm M, Rassaf T. 2014.** Myoglobin's novel role in nitrite-induced hypoxic vasodilation. *Trends in Cardiovascular Medicine* **24**:69–74 DOI [10.1016/j.tcm.2013.06.006](https://doi.org/10.1016/j.tcm.2013.06.006).
- Imamura H, Nhat KP, Togawa H, Saito K, Iino R, Kato-Yamada Y, Nagai T, Noji H. 2009.** Visualization of ATP levels inside single living cells with fluorescence resonance energy transfer-based genetically encoded indicators. *Proceedings of the National Academy of Sciences of the United States of America* **106**:15651–15656 DOI [10.1073/pnas.0904764106](https://doi.org/10.1073/pnas.0904764106).
- Kiang JG, McKinney LC, Gallin EK. 1990.** Heat induces intracellular acidification in human A-431 cells - role of Na<sup>+</sup>-H<sup>+</sup> exchange and metabolism. *American Journal of Physiology* **259**:C727–C737 DOI [10.1152/ajpcell.1990.259.5.C727](https://doi.org/10.1152/ajpcell.1990.259.5.C727).
- Klapperstuck T, Glanz D, Klapperstuck M, Wohlrab J. 2009.** Methodological aspects of measuring absolute values of membrane potential in human cells by flow cytometry. *Cytometry Part A* **75A**:593–608 DOI [10.1002/cyto.a.20735](https://doi.org/10.1002/cyto.a.20735).
- Ko EA, Han J, Jung ID, Park WS. 2008.** Physiological roles of K<sup>+</sup> channels in vascular smooth muscle cells. *Journal of Smooth Muscle Research = Nihon Heikatsukin Gakkai kikanishi* **44**:65–81 DOI [10.1540/jsmr.44.65](https://doi.org/10.1540/jsmr.44.65).

- Kubo M, Quayle JM, Standen NB. 1997.** Angiotensin II inhibition of ATP-sensitive K<sup>+</sup> currents in rat arterial smooth muscle cells through protein kinase C. *Journal of Physiology-London* **503**:489–496 DOI [10.1111/j.1469-7793.1997.489bg.x](https://doi.org/10.1111/j.1469-7793.1997.489bg.x).
- Levick JR. 2010.** *An introduction to cardiovascular physiology*. London: Hodder Arnold.
- Li J, Shuai HY, Gylfe E, Tengholm A. 2013.** Oscillations of sub-membrane ATP in glucose-stimulated beta cells depend on negative feedback from Ca<sup>2+</sup>. *Diabetologia* **56**:1577–1586 DOI [10.1007/s00125-013-2894-0](https://doi.org/10.1007/s00125-013-2894-0).
- Lin Y, Li X, Fan C, Yang F, Hao D, Ge W, Gao Y. 2020.** Cardioprotective effects of rat adipose-derived stem cells differ under normoxic/physioxic conditions and are associated with paracrine factor secretion. *International Journal of Molecular Medicine* **45**:1591–1600 DOI [10.3892/ijmm.2020.4524](https://doi.org/10.3892/ijmm.2020.4524).
- Lynch FM, Austin C, Heagerty AM, Izzard AS. 2006.** Adenosine- and hypoxia-induced dilation of human coronary resistance arteries: evidence against the involvement of K(ATP) channels. *British Journal of Pharmacology* **147**:455–458 DOI [10.1038/sj.bjp.0706622](https://doi.org/10.1038/sj.bjp.0706622).
- Mackie AR, Byron KL. 2008.** Cardiovascular KCNQ (Kv7) potassium channels: physiological regulators and new targets for therapeutic intervention. *Molecular Pharmacology* **74**:1171–1179 DOI [10.1124/mol.108.049825](https://doi.org/10.1124/mol.108.049825).
- Madden JA, Keller PA, Kleinman JG. 2000.** Changes in smooth muscle cell pH during hypoxic pulmonary vasoconstriction: a possible role for ion transporters. *Physiological Research* **49**:561–566.
- Miki T, Suzuki M, Shibasaki T, Uemura H, Sato T, Yamaguchi K, Koseki H, Iwanaga T, Nakaya H, Seino S. 2002.** Mouse model of Prinzmetal angina by disruption of the inward rectifier Kir6.1. *Nature Medicine* **8**:466–472 DOI [10.1038/nm0502-466](https://doi.org/10.1038/nm0502-466).
- Misler S, Barnett DW, Gillis KD, Pressel DM. 1992.** Electrophysiology of stimulus-secretion coupling in human beta-cells. *Diabetes* **41**:1221–1228 DOI [10.2337/diabetes.41.10.1221](https://doi.org/10.2337/diabetes.41.10.1221).
- Miura H, Wachtel RE, Loberiza Jr FR, Saito T, Miura M, Nicolosi AC, Gutterman DD. 2003.** Diabetes mellitus impairs vasodilation to hypoxia in human coronary arterioles: reduced activity of ATP-sensitive potassium channels. *Circulation Research* **92**:151–158 DOI [10.1161/01.res.0000052671.53256.49](https://doi.org/10.1161/01.res.0000052671.53256.49).
- Mustafa AK, Sikka G, Gazi SK, Steppan J, Jung SM, Bhunia AK, Barodka VM, Gazi FK, Barrow RK, Wang R, Amzel LM, Berkowitz DE, Snyder SH. 2011.** Hydrogen sulfide as endothelium-derived hyperpolarizing factor sulphydrates potassium channels. *Circulation Research* **109**:1259–U1169 DOI [10.1161/circresaha.111.240242](https://doi.org/10.1161/circresaha.111.240242).
- Nelson MT, Quayle JM. 1995.** Physiological roles and properties of potassium channels in arterial smooth-muscle. *American Journal of Physiology-Cell Physiology* **268**:C799–C822 DOI [10.1152/ajpcell.1995.268.4.C799](https://doi.org/10.1152/ajpcell.1995.268.4.C799).
- Park WS, Han J, Kim N, Ko JH, Kim SJ, Earm YE. 2005.** Activation of inward rectifier K<sup>+</sup> channels by hypoxia in rabbit coronary arterial smooth muscle cells. *American Journal of Physiology-Heart and Circulatory Physiology* **289**:H2461–H2467 DOI [10.1152/ajpheart.00331.2005](https://doi.org/10.1152/ajpheart.00331.2005).

- Park WS, Son YK, Kim N, Ko JH, Kang SH, Warda M, Earm YE, Jung ID, Park YM, Han J. 2007. Acute hypoxia induces vasodilation and increases coronary blood flow by activating inward rectifier K<sup>+</sup> channels. *Pflugers Archiv-European Journal of Physiology* 454:1023–1030 DOI 10.1007/s00424-007-0269-4.
- Pastore A, Piemonte F, Locatelli M, Lo Russo A, Gaeta LM, Tozzi G, Federici G. 2001. Determination of blood total, reduced, and oxidized glutathione in pediatric subjects. *Clinical Chemistry* 47:1467–1469 DOI 10.1093/clinchem/47.8.1467.
- Quayle JM, Bonev AD, Brayden JE, Nelson MT. 1994. Calcitonin-gene-related peptide activated ATP-sensitive K<sup>+</sup> currents in rabbit arterial smooth-muscle via protein-kinase-A. *Journal of Physiology-London* 475:9–13 DOI 10.1113/jphysiol.1994.sp020045.
- Quayle JM, McCarron JG, Brayden JE, Nelson MT. 1993. Inward rectifier K<sup>+</sup> currents in smooth-muscle cells from rat resistance-sized cerebral-arteries. *American Journal of Physiology* 265:C1363–C1370 DOI 10.1152/ajpcell.1993.265.5.C1363.
- Quayle JM, Nelson MT, Standen NB. 1997. ATP-sensitive and inwardly rectifying potassium channels in smooth muscle. *Physiological Reviews* 77:1165–1232 DOI 10.1152/physrev.1997.77.4.1165.
- Quayle JM, Turner MR, Burrell HE, Kamishima T. 2006. Effects of hypoxia, anoxia, and metabolic inhibitors on KATP channels in rat femoral artery myocytes. *American Journal of Physiology-Heart and Circulatory Physiology* 291:H71–80 DOI 10.1152/ajpheart.01107.2005.
- Roy CS, Brown JG. 1880. The blood-pressure and its variations in the arterioles, capillaries and smaller veins. *Journal de Physiologie* 2:323–446.321 DOI 10.1113/jphysiol.1880.sp000068.
- Sampson LJ, Davies LM, Barrett-Jolley R, Standen NB, Dart C. 2007. Angiotensin II-activated protein kinase C targets caveolae to inhibit aortic ATP-sensitive potassium channels. *Cardiovascular Research* 76:61–70 DOI 10.1016/j.cardiores.2007.05.020.
- Sampson LJ, Hayabuchi Y, Standen NB, Dart C. 2004. Caveolae localize protein kinase A signaling to arterial ATP-sensitive potassium channels. *Circulation Research* 95:1012–1018 DOI 10.1161/01.RES.0000148634.47095.ab.
- Selivanov VA, Alekseev AE, Hodgson DM, Dzeja PP, Terzic A. 2004. Nucleotide-gated KATP channels integrated with creatine and adenylate kinases: amplification, tuning and sensing of energetic signals in the compartmentalized cellular environment. *Molecular and Cellular Biochemistry* 256-257:243–256 DOI 10.1023/b:mcbi.0000009872.35940.7d.
- Shimoda LA, Polak J. 2011. Hypoxia. 4. Hypoxia and ion channel function. *American Journal of Physiology. Cell Physiology* 300:C951–967 DOI 10.1152/ajpcell.00512.2010.
- Shindo Y, Witt E, Han D, Epstein W, Packer L. 1994. Enzymic and non-enzymic antioxidants in epidermis and dermis of human skin. *Journal of Investigative Dermatology* 102:122–124 DOI 10.1111/1523-1747.ep12371744.
- Standen NB, Quayle JM. 1998. K<sup>+</sup> channel modulation in arterial smooth muscle. *Acta Physiologica Scandinavica* 164:549–557 DOI 10.1046/j.1365-201X.1998.00433.x.
- Sturges NC, Ashford MLJ, Cook DL, Hales CN. 1985. The sulfonylurea receptor may be an ATP-sensitive potassium channel. *Lancet* 2:474–475.

- Tanaka Y, Meera P, Song M, Knaus HG, Toro L. 1997.** Molecular constituents of maxi KCa channels in human coronary smooth muscle: predominant  $\alpha$  + beta subunit complexes. *Journal de Physiologie* **502(Pt 3)**:545–557  
[DOI 10.1111/j.1469-7793.1997.545bj.x](https://doi.org/10.1111/j.1469-7793.1997.545bj.x).
- Tantama M, Hung YP, Yellen G. 2011.** Imaging intracellular pH in live cells with a genetically encoded red fluorescent protein sensor. *Journal of the American Chemical Society* **133**:10034–10037 [DOI 10.1021/ja202902d](https://doi.org/10.1021/ja202902d).
- Tantama M, Martínez-François JR, Mongeon R, Yellen G. 2013.** Imaging energy status in live cells with a fluorescent biosensor of the intracellular ATP-to-ADP ratio. *Nature Communications* **4**:2550 [DOI 10.1038/ncomms3550](https://doi.org/10.1038/ncomms3550).
- Tantama M, Yellen G. 2014.** Imaging changes in the cytosolic ATP-to-ADP ratio. *Methods in Enzymology* **547**:355–371 [DOI 10.1016/b978-0-12-801415-8.00017-5](https://doi.org/10.1016/b978-0-12-801415-8.00017-5).
- Totzeck M, Hendgen-Cotta UB, Luedike P, Berenbrink M, Klare JP, Steinhoff HJ, Semmler D, Shiva S, Williams D, Kipar A, Gladwin MT, Schrader J, Kelm M, Cossins AR, Rassaf T. 2012.** Nitrite regulates hypoxic vasodilation via myoglobin-dependent nitric oxide generation. *Circulation* **126**:325–334  
[DOI 10.1161/circulationaha.111.087155](https://doi.org/10.1161/circulationaha.111.087155).
- Traut TW. 1994.** Physiological concentrations of purines and pyrimidines. *Molecular and Cellular Biochemistry* **140**:1–22 [DOI 10.1007/bf00928361](https://doi.org/10.1007/bf00928361).
- Veech RL, Lawson JW, Cornell NW, Krebs HA. 1979.** Cytosolic phosphorylation potential. *Journal of Biological Chemistry* **254**:6538–6547.
- Venkatesh N, Lamp ST, Weiss JN. 1991.** Sulfonylureas, ATP-sensitive K<sup>+</sup> channels, and cellular K<sup>+</sup> loss during hypoxia, ischemia, and metabolic inhibition in mammalian ventricle. *Circulation Research* **69**:623–637 [DOI 10.1161/01.res.69.3.623](https://doi.org/10.1161/01.res.69.3.623).
- Von Beckerath N, Cyrus S, Dischner A, Daut J. 1991.** Hypoxic vasodilatation in isolated, perfused guinea-pig heart: an analysis of the underlying mechanisms. *Journal de Physiologie* **442**:297–319 [DOI 10.1113/jphysiol.1991.sp018794](https://doi.org/10.1113/jphysiol.1991.sp018794).
- Welsh DG, Jackson WF, Segal SS. 1998.** Oxygen induces electromechanical coupling in arteriolar smooth muscle cells: a role for L-type Ca<sup>2+</sup> channels. *American Journal of Physiology* **274**:H2018–2024 [DOI 10.1152/ajpheart.1998.274.6.H2018](https://doi.org/10.1152/ajpheart.1998.274.6.H2018).
- Wulff H, Castle NA. 2010.** Therapeutic potential of KCa3.1 blockers: recent advances and promising trends. *Expert Review of Clinical Pharmacology* **3**:385–396  
[DOI 10.1586/ecp.10.11](https://doi.org/10.1586/ecp.10.11).
- Yang M, Chadwick AE, Dart C, Kamishima T, Quayle JM. 2017.** Bioenergetic profile of human coronary artery smooth muscle cells and effect of metabolic intervention. *PLOS ONE* **12**:e0177951 [DOI 10.1371/journal.pone.0177951](https://doi.org/10.1371/journal.pone.0177951).
- Yoshida H, Feig JE, Morrissey A, Ghiu IA, Artman M, Coetzee WA. 2004.** K ATP channels of primary human coronary artery endothelial cells consist of a hetero-multimeric complex of Kir6.1, Kir6.2, and SUR2B subunits. *Journal of Molecular and Cellular Cardiology* **37**:857–869 [DOI 10.1016/j.yjmcc.2004.05.022](https://doi.org/10.1016/j.yjmcc.2004.05.022).