



Review Article

The role of Type III secretion system in the pathogenesis of *Pseudomonas aeruginosa* microbial keratitis

Justin J. Yang^{a,b}, Kai-Si Claire Tsuei^c, Elizabeth P. Shen^{c,d,e*}

^aDepartment of Chemistry, Trinity College of Arts and Sciences, Duke University, North Carolina, USA; ^bDepartment of Computer Science, Trinity College of Arts and Sciences, Duke University, North Carolina, USA; ^cDepartment of Ophthalmology, Taipei Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation, New Taipei, Taiwan; ^dCollege of Medicine, Tzu Chi University, Hualien, Taiwan; ^eDepartment of Ophthalmology, National Taiwan University Hospital and College of Medicine, National Taiwan University, Taipei, Taiwan

ABSTRACT

Pseudomonas aeruginosa is the most commonly isolated Gram-negative pathogen causing sight-threatening microbial keratitis (MK). Contact lens wear is the most significant risk factor associated with pseudomonal MK. Understanding the pathogenesis of MK due to *P. aeruginosa* and its interactions with contact lenses is crucial in preventing these often rapidly progressive and highly antibiotic-resistant infections. Bacterial virulence factor Type III secretion system (T3SS) has significant interplays between contact lens material, antibiotic sensitivity, disinfectant selectivity, and bacterial cell invasion. Depending on the T3SS exotoxins produced, *P. aeruginosa* strains are divided into cytotoxic or invasive strains. Cytotoxic strains are relatively resistant to commercial disinfectants, while invasive strains are more antibiotic resistant. Therefore, contact lens wearers are more predisposed to cytotoxic *P. aeruginosa* infections, and patients with trauma or previous surgery are more prone to infection by invasive strains. Previous studies with mutant *P. aeruginosa* strains unable to produce T3SS exotoxins were more susceptible to disinfectants and less able to adhere to soft contact lenses, indicating an essential role of T3SS in bacterial virulence. Invasion of *P. aeruginosa* intracellularly was found to be associated with control of scaffold protein IQ-domain GTPase-activating protein 1 (IQGAP1) and human corneal epithelial cell tight junctions. Knockdown of IQGAP1 strengthened tight junctions that prevented intracellular survival of invasive *P. aeruginosa* strains and enhanced corneal epithelial cell survival. These novel findings of the vital role of T3SS in the pathogenesis of pseudomonal MKs will provide new guidelines in both prevention and treatment of this common eye-blinding infection.

KEYWORDS: Contact lens, IQ-domain GTPase-activating protein 1, Microbial keratitis, *Pseudomonas aeruginosa*, Type III secretion system

Submission : 17-Feb-2021
Revision : 11-Mar-2021
Acceptance : 17-Mar-2021
Web Publication : 11-May-2021

INTRODUCTION

Microbial keratitis (MK) due to *Pseudomonas aeruginosa* causes severe ocular morbidity that may result in blindness if not treated promptly and appropriately [1]. *P. aeruginosa*, an opportunistic pathogen commonly found in our environment, is the most commonly isolated Gram-negative organism causing MK, particularly among contact lens wearers [2]. The incidence of this contact lens-related microbial keratitis (CLMK) is approximately 3.5–20.9 per 10,000 wearers, varying due to contact lens material and wearing schedules [3–5]. With young emmetropic individuals wearing cosmetic color-tinted contact lenses, the incidence of CLMK is unfortunately expected to increase [6,7].

Ongoing research to understand the complex interaction between contact lens material, the initiation mechanisms of

CLMK, and the bacterial–host immune response has continued to shed light into the perplex mechanisms causing MK. As our study and other previous studies have shown, contact lens material significantly affects the amount of bacterial adhesion [8–11]. Virulence factors of the bacterium may also influence its propensity to adhere [9,12–14]. Among the multitude of virulence factors, the Type III secretion system (T3SS), a contact-dependent protein secretion pathway secreting exotoxins, has been associated with significant host cell damage and thus clinical disease [15–17]. In this article, we review the present

*Address for correspondence:

Dr. Elizabeth P. Shen,
Department of Ophthalmology, Taipei Tzu Chi Hospital,
Buddhist Tzu Chi Medical Foundation, 289, Jianguo Road, Xindian
District, New Taipei, Taiwan.
E-mail: lizshen88@gmail.com

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Yang JJ, Tsuei KS, Shen EP. The role of Type III secretion system in the pathogenesis of *Pseudomonas aeruginosa* microbial keratitis. Tzu Chi Med J 2022; 34(1): 8–14.

Access this article online	
Quick Response Code: 	Website: www.tcmjmed.com
	DOI: 10.4103/tcmj.tcmj_47_21

understanding of the complex interactions between *P. aeruginosa* with contact lenses material in the initiation process of the pathogenesis of CLMK.

PATHOGENESIS OF *PSEUDOMONAS AERUGINOSA* MICROBIAL KERATITIS

Pseudomonal keratitis often presents as a rapidly progressive stromal ulceration with overlying epithelial defect. The stromal ulceration may lead to severe and often rapid stromal melting, resulting in corneal perforation and visual loss [18,19]. The development of CLMK commences with bacterial contamination and subsequent bacterial adhesion to contact lenses, rendering prolonged exposure of the cornea to microbial pathogens [Figure 1] [20-22]. Since *P. aeruginosa* is an opportunistic organism, the establishment of an infection requires a compromised state of the host. Trauma due to abrasion or hypoxic injury due to contact lens wear had been proposed to cause epithelial breakdown and thus allow ensuing pseudomonal invasion [12,23,24]. To increase the oxygen permeability to the cornea during contact lens wear, silicone hydrogel materials were developed and were found to have at least 6 times greater oxygen permeability than conventional hydrogel lenses [25]. Nevertheless, daily wear of silicone hydrogel contact lenses still had about 6 times greater risk of corneal infection than daily wear or daily disposable conventional soft contact lens [26]. Clearly, factors other than hypoxia must be involved. Therefore, besides the contact lens material itself, the interaction of bacterial virulence factors with host defenses or immunity had been previously studied.

PSEUDOMONAL TYPE III SECRETION SYSTEM

T3SS exoenzymes were found to be important virulence factors inducing *P. aeruginosa* keratitis [27-30]. The T3SS is a specialized protein export system that forms a needle-like complex between bacterial and host cells for the transport and secretion of four exotoxins, ExoS, ExoU, ExoT, and ExoY [Figure 2] [17]. The T3SS is encoded by 36 genes in five operons, which correspond to five functional components: the needle-like complex, the translocation apparatus, the regulatory proteins, the chaperones, and the effector proteins [17]. Structurally, the needle-like complex is a hollow filament about 60–120 nm long and 6–10 nm wide and serves as a tube through which secreted factors are transported [31]. The translocation apparatus is composed of proteins that form pores on the host's cell membrane. The needle-like complex connects with these pores, which then accept the effector proteins transported by the needle-like complex and deliver it across the host's cell membrane. A structural component of the needle complex, PscC protein is located on the outer membrane of *P. aeruginosa* and has been shown to be essential in effector transport [Figure 2] [17,32]. Mutations in the PscC protein result in loss of bacterial virulence and also T3SS exotoxin secretion [32-34]. The expression of the T3SS usually requires close cell contact *in vivo* or low calcium growth conditions *in vitro* [35-37]. Contamination of contact lenses with adhering bacteria thus provides a chance for relatively close contact with the corneal epithelium to induce the T3SS *in vivo*. After T3SS activation, the chaperone proteins which bind to effector proteins before secretion facilitate the delivery of effector proteins into the secretion system [38]. Chaperones remain in the bacterial cytoplasm after their effector protein





Figure 1: Proposed pathogenesis of pseudomonal keratitis. Complex interaction between bacterial virulence factor Type III secretion system, contact lens, and host immune response occurs during the initiation process of *Pseudomonas aeruginosa* keratitis. Understanding of these complex interactions may help to develop prevention strategies and target therapy



In contrast to ExoS, ExoU (74 kDa) is a phospholipase that causes host cell lysis within 1–2 h [42]. Analysis of its protein sequence reveals a patatin-like domain with phospholipase A₂ (PLA₂) activity between residues 107 and 357 [43]. PLA₂ enzymes hydrolyze the ester bond of phospholipids at the

ExoY is a 378-amino acid (42 kDa) adenylyl cyclase that also requires a host cell cofactor for activation, although the identity of this cofactor is currently unknown [51]. When ExoY was injected into mammalian cells, elevated intracellular cAMP concentrations and differential gene expression were

observed, which led to the disruption of the actin cytoskeleton, inhibition of host phagocytosis of bacteria, and increased endothelial permeability [17,51].

For reasons that are unclear, most strains do not carry all four genes encoding for the four effectors. In fact, while almost all strains of *P. aeruginosa* carry both *exoT* and *exoY* genes, most have only either *exoU* or *exoS*, not both [17]. As such, two genotypes of *P. aeruginosa* with different infection phenotypes can be distinguished by the secretion of either ExoU or ExoS. Strains that are ExoS positive and ExoU negative are known as invasive strains as they can invade into corneal epithelial cells [27,52-54]. On the other hand, strains that are ExoU positive and ExoS negative are known as cytotoxic strains because ExoU, a phospholipase, causes acute host cell lysis [53,55]. The invasive and cytotoxic phenotypes with their respective *exoS* and *exoU* genotypes were found to be mutually exclusive in nearly all strains [27].

PSEUDOMONAL TYPE III SECRETION SYSTEM AND CLINICAL MANIFESTATION OF MICROBIAL KERATITIS

From our investigation of invasive and cytotoxic strains isolated from MK cases over a 10-year period in Taiwan, the two phenotypes of *P. aeruginosa* showed significant differences both in clinical manifestation and in visual prognosis. Clinical *P. aeruginosa* isolates were mostly of cytotoxic strains among contact lens wearer, while invasive strains were significantly related to a history of trauma or previous surgery [9,56]. This was similarly noted among Australian isolates [57]. Cytotoxic strains were also more commonly seen in young females in contrast to invasive strains, which predominantly occurred in older males [56]. Initial presenting visual acuity for cytotoxic strains was also significantly better compared to MK due to invasive strains. This is mainly due to smaller infiltration size and less fulminant disease due to cytotoxic strains [56]. Invasive strains often had a greater infiltration size and depth accompanied by the presence of hypopyon therefore poorer final visual outcome and even higher treatment failure rate (Failure was defined as severe uncontrollable infection requiring therapeutic penetrating keratoplasty, evisceration, or enucleation) [56].

Antibiotic susceptibility varied according to the regions and between the two phenotypes. A greater proportion of cytotoxic strains from the USA, Australia, and India were more resistant to fluoroquinolones and aminoglycosides, although differences between phenotypes were not significant [58-60]. This is in contrast to isolates from Taiwan, which showed an increased antibiotic resistance to fluoroquinolones for invasive strains [56]. Cytotoxic strains from CLMK cases were all sensitive to commonly used aminoglycosides and fluoroquinolones [56]. These results suggest that analysis of *P. aeruginosa* phenotype may be helpful clinically in predicting visual prognosis and also suggesting clinical guidelines for the treatment of pseudomonal keratitis with some regional differences.

TYPE III SECRETION SYSTEM AND HOST INVASION

The effect of T3SS effectors on corneal epithelial cells differ

with their genotypes. Strains with the *exoS* genotype can invade and multiply within corneal epithelial cells both *in vitro* and *in vivo* [52,55]. It was found that functional T3SS was required for intracellular survival of *P. aeruginosa* within corneal epithelial cells [52]. Wild-type strains with functional T3SS can be seen replicating within plasma membrane blebs [61]. In contrast, the T3SS needle-complex *pscC* mutant could not form membrane blebs and thus had reduced intracellular survival [61]. Thus, these membrane blebs are essentially used as niches by *P. aeruginosa* for intracellular survival and replication [61]. Visualization of the membrane localization of intracellular *P. aeruginosa* can be seen with confocal microscopy using GFP-transformed strains [62]. Although preliminary work on how ExoS promotes intracellular survival had been done, the studies of ExoS on disruption of epithelial barrier function were mostly done on mucosal cells of the lung [63].

The effect of ExoS with a scaffold protein IQ-domain GTPase-activating protein 1 (IQGAP1) on the corneal epithelial barrier was recently published by ShenW *et al.* [64]. IQGAP1 co-localized with junctional proteins actin and zonular occludin 1 to induce changes in the corneal epithelial cell tight junction [64]. Knockdown of IQGAP1 enhanced tight junction formation with increased transepithelial resistance [64]. Since invasive strain *P. aeruginosa* invades human corneal epithelial cells by rapidly breaking down tight junctions, enhancement of tight junctions resulting from the knockdown of IQGAP1 can protect against *P. aeruginosa* invasion, leading to enhanced cell survival [64]. Viability of cells increased by 45% early in the infection [64]. After IQGAP1 knockdown, PAK was also less able to invade and survive within the corneal epithelial cells [64]. These novel findings suggested that T3SS enables invasiveness of *P. aeruginosa* through breakdown of tight junctions. Invasion of the bacterium into host cells protected the pathogen from host immune defenses and evaded antibiotics control. The poor clinical outcomes due to invasive pseudomonal strains may be partially due to this evasion tactic. Therefore, understanding the mechanism of pseudomonal invasion is essential in the development of new therapeutic targets, especially with rising antibiotic resistance.

TYPE III SECRETION SYSTEM AND CONTACT LENSES

The T3SS virulence factors not only influence clinical prognosis and bacterial antibiotic sensitivity, but it has also been shown to affect pseudomonal adherence to contact lenses. Previously, environmental and clinical isolates from lung, urinary tract, or burn wound infections were shown to be mostly invasive [65-67]. However, the higher prevalence or predisposition of finding cytotoxic genotype among contact lens wearers suggested that T3SS may be involved either directly or indirectly with adhesion to contact lens materials. By using low calcium-inducing conditions to compare the bacterial adhesions between wild-type and *pscC*-mutant strains, it was demonstrated that *pscC* mutants were unable to secrete T3SS exotoxins, irrespective of strain. These mutants showed significantly less adhesion to both conventional hydrogel and silicone hydrogel contact lenses compared to wild-type strains [9]. Compared to noninducing conditions, bacteria grown under T3SS-inducing

conditions also showed significantly greater bacterial adhesion to contact lenses, indicating that functional and active T3SS is essential for contact lens adhesion [9]. Although functional T3SS affects bacterial lens adhesion, genotype differences were not found. Therefore, other factors must be involved in the predisposition of cytotoxic genotype among CLMK.

Contact lens material had been shown to affect pseudomonal adherence. Regardless of strain, silicone hydrogels with plasma oxidation surface treatment tend to adhere greater bacteria [9]. Generally, galyfilcon silicone hydrogel lenses (lenses without surface treatment) had the least amount of bacterial adhesion compared to surface-treated silicone hydrogel and conventional hydrogels [9]. Thus, surface smoothness was proposed to affect bacterial adhesion. Cosmetic hydrogel lenses that easily dislodge color pigments were shown to adhere significantly more bacteria [68,69]. Scanning electron microscope of pigmented contact lenses showed a significant association of surface smoothness with pigment dislodgement [70]. Therefore, improper contact lens material or surface treatment may increase the propensity for bacterial adhesion onto lens surfaces.

TYPE III SECRETION SYSTEM AND DISINFECTANT SELECTIVITY

As previously mentioned, functional T3SS was required for bacterial adherence to contact lenses. However, there were no genotype differences in adhesion demonstrated by us and others [9,71]. Selectivity of cytotoxic strains by disinfectants or multipurpose solutions was thought to be a possible explanation. In 2001, Lakkis and Fleiszig found an increase in resistance to two contact lens chemical disinfectants [72]. This resistance was associated with acute cytotoxic activity to corneal epithelial cells [72]. Recently, we compared the sensitivity of the two genotypes of *P. aeruginosa* to four commercially available disinfectants (Renu Fresh, PureMoist, Replenish, and AoSept Plus). Cytotoxic strains were significantly more resistant to disinfectants, especially Renu Fresh, even at manufacturer's recommended disinfection time. The best disinfectant was AoSopt Plus, the hydrogen peroxide system, which was able to completely eradicate *P. aeruginosa* even at 25% of recommended disinfection time. Compared to standard wild-type strains, *pscC*-mutant strains were all susceptible to disinfectants, indicating that functional T3SS may contribute to bacterial disinfectant resistance. These cumulative findings suggest that the relative higher prevalence of cytotoxic strains with contact lens wearers may be due to the relative selectivity of marketed disinfectants. Careful selection of effective disinfectants should be advocated for proper lens hygiene.

CONCLUSION

P. aeruginosa is the most common pathogen causing vision-threatening corneal infections, especially in contact lens wearers. The rapid progressive course and poor clinical outcomes due to emerging antibiotic resistance stimulated extensive research in the role of bacterial virulence factor T3SS interactions with contact lenses. As extensive reviewed,

recognition of the importance of the various effects of T3SS exotoxins will better enable early prevention measures and novel treatment options to salvage the devastating sight-threatening complication due to *P. aeruginosa* infections.

Financial support and sponsorship

This article was supported by the funding from Taipei Tzu Chi Hospital, Buddhist Tzu Chi Medical Foundation (TCRD-TPE-110-20 and TCRD-TPE-110-47) and the Ministry of Science and Technology, Taiwan (MOST105-2314-B-002-167-MY3 and MOST106-2314-B-303-011MY3).

Conflicts of interest

There are no conflicts of interest.

REFERENCES

1. Stapleton F, Keay LJ, Sanfilippo PG, Katiyar S, Edwards KP, Naduvilath T. Relationship between climate, disease severity, and causative organism for contact lens-associated microbial keratitis in Australia. *Am J Ophthalmol* 2007;144:690-8.
2. Ormerod LD, Smith RE. Contact lens-associated microbial keratitis. *Arch Ophthalmol* 1986;104:79-83.
3. Cheng KH, Leung SL, Hoekman HW, Beekhuis WH, Mulder PG, Geerards AJ, et al. Incidence of contact-lens-associated microbial keratitis and its related morbidity. *Lancet* 1999;354:181-5.
4. Poggio EC, Glynn RJ, Schein OD, Seddon JM, Shannon MJ, Scardino VA, et al. The incidence of ulcerative keratitis among users of daily-wear and extended-wear soft contact lenses. *N Engl J Med* 1989;321:779-83.
5. Liesegang TJ. Contact lens-related microbial keratitis: Part I. *Epidemiol Cornea* 1997;16:125-31.
6. Steinemann TL, Fletcher M, Bonny AE, Harvey RA, Hamlin D, Zloty P, et al. Over-the-counter decorative contact lenses: Cosmetic or medical devices? A case series. *Eye Contact Lens* 2005;31:194-200.
7. Sauer A, Bourcier T; French Study Group for Contact Lenses Related Microbial Keratitis. Microbial keratitis as a foreseeable complication of cosmetic contact lenses: A prospective study. *Acta Ophthalmol* 2011;89:e439-42.
8. Henriques M, Sousa C, Lira M, Elisabete M, Oliveira R, Oliveira R, et al. Adhesion of *Pseudomonas aeruginosa* and *Staphylococcus epidermidis* to silicone-hydrogel contact lenses. *Optom Vis Sci* 2005;82:446-50.
9. Shen EP, Tsay RY, Chia JS, Wu S, Lee JW, Hu FR. The role of Type III secretion system and lens material on adhesion of *Pseudomonas aeruginosa* to contact lenses. *Invest Ophthalmol Vis Sci* 2012;53:6416-26.
10. Vijay AK, Zhu H, Ozkan J, Wu D, Masoudi S, Bandara R, et al. Bacterial adhesion to unworn and worn silicone hydrogel lenses. *Optom Vis Sci* 2012;89:1095-106.
11. Nilsson SE, Montan PG. The hospitalized cases of contact lens induced keratitis in Sweden and their relation to lens type and wear schedule: Results of a three-year retrospective study. *CLAO J* 1994;20:97-101.
12. Fleiszig SM, Efron N, Pier GB. Extended contact lens wear enhances *Pseudomonas aeruginosa* adherence to human corneal epithelium. *Invest Ophthalmol Vis Sci* 1992;33:2908-16.
13. Fletcher EL, Fleiszig SM, Brennan NA. Lipopolysaccharide in adherence of *Pseudomonas aeruginosa* to the cornea and contact lenses. *Invest Ophthalmol Vis Sci* 1993;34:1930-6.
14. Fletcher EL, Weissman BA, Efron N, Fleiszig SM, Curcio AJ, Brennan NA. The role of pili in the attachment of *Pseudomonas aeruginosa* to unworn hydrogel contact lenses. *Curr Eye Res* 1993;12:1067-71.
15. Engel J, Balachandran P. Role of *Pseudomonas aeruginosa* Type III effectors in disease. *Curr Opin Microbiol* 2009;12:61-6.

16. Yahr TL, Wolfgang MC. Transcriptional regulation of the *Pseudomonas aeruginosa* Type III secretion system. *Mol Microbiol* 2006;62:631-40.
17. Hauser AR. The Type III secretion system of *Pseudomonas aeruginosa*: Infection by injection. *Nat Rev Microbiol* 2009;7:654-65.
18. Cruz CS, Cohen EJ, Rapuano CJ, Laibson PR. Microbial keratitis resulting in loss of the eye. *Ophthalmic Surg Lasers* 1998;29:803-7.
19. Green M, Apel A, Stapleton F. Risk factors and causative organisms in microbial keratitis. *Cornea* 2008;27:22-7.
20. Cavanagh HD, Robertson DM, Petroll WM, Jester JV. Castroviejo lecture 2009: 40 years in search of the perfect contact lens. *Cornea* 2010;29:1075-85.
21. Fleiszig SM, Evans DJ. Pathogenesis of contact lens-associated microbial keratitis. *Optom Vis Sci* 2010;87:225-32.
22. Willcox MD. *Pseudomonas aeruginosa* infection and inflammation during contact lens wear: A review. *Optom Vis Sci* 2007;84:273-8.
23. Yanai R, Ko JA, Morishige N, Chikama T, Ichijima H, Nishida T. Disruption of zonula occludens-1 localization in the rabbit corneal epithelium by contact lens-induced hypoxia. *Invest Ophthalmol Vis Sci* 2009;50:4605-10.
24. Willcox MD, Holden BA. Contact lens related corneal infections. *Biosci Rep* 2001;21:445-61.
25. Holden BA. The Glenn A. Fry Award Lecture 1988: The ocular response to contact lens wear. *Optom Vis Sci* 1989;66:717-33.
26. Stapleton F, Keay L, Edwards K, Naduvilath T, Dart JK, Brian G, et al. The incidence of contact lens-related microbial keratitis in Australia. *Ophthalmology* 2008;115:1655-62.
27. Fleiszig SM, Wiener-Kronish JP, Miyazaki H, Vallas V, Mostov KE, Kanada D, et al. *Pseudomonas aeruginosa*-mediated cytotoxicity and invasion correlate with distinct genotypes at the loci encoding exoenzyme S. *Infect Immun* 1997;65:579-86.
28. Twining SS, Kirschner SE, Mahnke LA, Frank DW. Effect of *Pseudomonas aeruginosa* elastase, alkaline protease, and exotoxin A on corneal proteinases and proteins. *Invest Ophthalmol Vis Sci* 1993;34:2699-712.
29. O'Callaghan RJ, Engel LS, Hobden JA, Callegan MC, Green LC, Hill JM. *Pseudomonas keratitis*. The role of an uncharacterized exoprotein, protease IV, in corneal virulence. *Invest Ophthalmol Vis Sci* 1996;37:534-43.
30. Kessler E, Blumberg S. Specific inhibitors of *Pseudomonas aeruginosa* elastase as potential drugs for the treatment of *Pseudomonas keratitis*. *Antibiot Chemother* (1971) 1987;39:102-12.
31. Pastor A, Chabert J, Louwagie M, Garin J, Attree I. PscF is a major component of the *Pseudomonas aeruginosa* Type III secretion needle. *FEMS Microbiol Lett* 2005;253:95-101.
32. Lee VT, Smith RS, Tümmler B, Lory S. Activities of *Pseudomonas aeruginosa* effectors secreted by the Type III secretion system *in vitro* and during infection. *Infect Immun* 2005;73:1695-705.
33. Wolfgang MC, Lee VT, Gilmore ME, Lory S. Coordinate regulation of bacterial virulence genes by a novel adenylate cyclase-dependent signaling pathway. *Dev Cell* 2003;4:253-63.
34. McCaw ML, Lykken GL, Singh PK, Yahr TL. ExsD is a negative regulator of the *Pseudomonas aeruginosa* Type III secretion regulon. *Mol Microbiol* 2002;46:1123-33.
35. Yahr TL, Goranson J, Frank DW. Exoenzyme S of *Pseudomonas aeruginosa* is secreted by a Type III pathway. *Mol Microbiol* 1996;22:991-1003.
36. Frank DW. The exoenzyme S regulon of *Pseudomonas aeruginosa*. *Mol Microbiol* 1997;26:621-9.
37. Vallis AJ, Yahr TL, Barbieri JT, Frank DW. Regulation of ExoS production and secretion by *Pseudomonas aeruginosa* in response to tissue culture conditions. *Infect Immun* 1999;67:914-20.
38. Parsot C, Hamiaux C, Page AL. The various and varying roles of specific chaperones in Type III secretion systems. *Curr Opin Microbiol* 2003;6:7-14.
39. Zhang Y, Barbieri JT. A leucine-rich motif targets *Pseudomonas aeruginosa* ExoS within mammalian cells. *Infect Immun* 2005;73:7938-45.
40. Krall R, Zhang Y, Barbieri JT. Intracellular membrane localization of *Pseudomonas* ExoS and Yersinia YopE in mammalian cells. *J Biol Chem* 2004;279:2747-53.
41. Rolsma SL, Frank DW. *In vitro* assays to monitor the activity of *Pseudomonas aeruginosa* Type III secreted proteins. *Methods Mol Biol* 2014;1149:171-84.
42. Rabin SD, Hauser AR. Functional regions of the *Pseudomonas aeruginosa* cytotoxin ExoU. *Infect Immun* 2005;73:573-82.
43. Sato H, Frank DW, Hillard CJ, Feix JB, Pankhaniya RR, Moriyama K, et al. The mechanism of action of the *Pseudomonas aeruginosa*-encoded Type III cytotoxin, ExoU. *EMBO J* 2003;22:2959-69.
44. Phillips RM, Six DA, Dennis EA, Ghosh P. *In vivo* phospholipase activity of the *Pseudomonas aeruginosa* cytotoxin ExoU and protection of mammalian cells with phospholipase A2 inhibitors. *J Biol Chem* 2003;278:41326-32.
45. Rabin SD, Veesenmeyer JL, Biegging KT, Hauser AR. A C-terminal domain targets the *Pseudomonas aeruginosa* cytotoxin ExoU to the plasma membrane of host cells. *Infect Immun* 2006;74:2552-61.
46. Sato H, Feix JB, Frank DW. Identification of superoxide dismutase as a cofactor for the *Pseudomonas* Type III toxin, ExoU. *Biochemistry* 2006;45:10368-75.
47. Maresco AW, Baldwin MR, Barbieri JT. Ezrin/radixin/moesin proteins are high affinity targets for ADP-ribosylation by *Pseudomonas aeruginosa* ExoS. *J Biol Chem* 2004;279:38402-8.
48. Krall R, Schmidt G, Aktories K, Barbieri JT. *Pseudomonas aeruginosa* ExoT is a Rho GTPase-activating protein. *Infect Immun* 2000;68:6066-8.
49. Garrity-Ryan L, Kazmierczak B, Kowal R, Comolli J, Hauser A, Engel JN. The arginine finger domain of ExoT contributes to actin cytoskeleton disruption and inhibition of internalization of *Pseudomonas aeruginosa* by epithelial cells and macrophages. *Infect Immun* 2000;68:7100-13.
50. Sun J, Barbieri JT. *Pseudomonas aeruginosa* ExoT ADP-ribosylates CT10 regulator of kinase (Crk) proteins. *J Biol Chem* 2003;278:32794-800.
51. Yahr TL, Vallis AJ, Hancock MK, Barbieri JT, Frank DW. ExoY, an adenylate cyclase secreted by the *Pseudomonas aeruginosa* Type III system. *Proc Natl Acad Sci U S A* 1998;95:13899-904.
52. Fleiszig SM, Zaidi TS, Pier GB. *Pseudomonas aeruginosa* invasion of and multiplication within corneal epithelial cells *in vitro*. *Infect Immun* 1995;63:4072-7.
53. Fleiszig SM, Zaidi TS, Preston MJ, Grout M, Evans DJ, Pier GB. Relationship between cytotoxicity and corneal epithelial cell invasion by clinical isolates of *Pseudomonas aeruginosa*. *Infect Immun* 1996;64:2288-94.
54. Lee EJ, Cowell BA, Evans DJ, Fleiszig SM. Contribution of ExsA-regulated factors to corneal infection by cytotoxic and invasive *Pseudomonas aeruginosa* in a murine scarification model. *Invest Ophthalmol Vis Sci* 2003;44:3892-8.
55. Fleiszig SM, Zaidi TS, Fletcher EL, Preston MJ, Pier GB. *Pseudomonas aeruginosa* invades corneal epithelial cells during experimental infection. *Infect Immun* 1994;62:3485-93.
56. Shen EP, Hsieh YT, Chu HS, Chang SC, Hu FR. Correlation of *Pseudomonas aeruginosa* genotype with antibiotic susceptibility and clinical features of induced central keratitis. *Invest Ophthalmol Vis Sci* 2014;56:365-71.
57. Choy MH, Stapleton F, Willcox MD, Zhu H. Comparison of virulence factors in *Pseudomonas aeruginosa* strains isolated from contact lens- and non-contact lens-related keratitis. *J Med Microbiol* 2008;57:1539-46.
58. Khan M, Stapleton F, Summers S, Rice SA, Willcox MD. Antibiotic resistance characteristics of *Pseudomonas aeruginosa* isolated from keratitis in Australia and India. *Antibiotics (Basel)* 2020;9:600.
59. Borkar DS, Acharya NR, Leong C, Lalitha P, Srinivasan M,

- Oldenburg CE, et al. Cytotoxic clinical isolates of *Pseudomonas aeruginosa* identified during the Steroids for Corneal Ulcers Trial show elevated resistance to fluoroquinolones. *BMC Ophthalmol* 2014;14:54.
60. Zhu H, Conibear TC, Bandara R, Aliwarga Y, Stapleton F, Willcox MD. Type III secretion system-associated toxins, proteases, serotypes, and antibiotic resistance of *Pseudomonas aeruginosa* isolates associated with keratitis. *Curr Eye Res* 2006;31:297-306.
61. Angus AA, Lee AA, Augustin DK, Lee EJ, Evans DJ, Fleiszig SM. *Pseudomonas aeruginosa* induces membrane blebs in epithelial cells, which are utilized as a niche for intracellular replication and motility. *Infect Immun* 2008;76:1992-2001.
62. Tam C, LeDue J, Mun JJ, Herzmark P, Robey EA, Evans DJ, et al. 3D quantitative imaging of unprocessed live tissue reveals epithelial defense against bacterial adhesion and subsequent traversal requires MyD88. *PLoS One* 2011;6:e24008.
63. Soong G, Parker D, Magargee M, Prince AS. The Type III toxins of *Pseudomonas aeruginosa* disrupt epithelial barrier function. *J Bacteriol* 2008;190:2814-21.
64. Shen EP, Chen MR, Chen WL, Chu HS, Chen KL, Hu FR. Knockdown of IQGAP-1 enhances tight junctions and prevents *P. aeruginosa* invasion of human corneal epithelial cells. *Ocul Immunol Inflamm* 2020;28:876-83.
65. Feltman H, Schulert G, Khan S, Jain M, Peterson L, Hauser AR. Prevalence of Type III secretion genes in clinical and environmental isolates of *Pseudomonas aeruginosa*. *Microbiology (Reading)* 2001;147:2659-69.
66. Bradbury RS, Roddam LF, Merritt A, Reid DW, Champion AC. Virulence gene distribution in clinical, nosocomial and environmental isolates of *Pseudomonas aeruginosa*. *J Med Microbiol* 2010;59:881-90.
67. Lomholt JA, Poulsen K, Kilian M. Epidemic population structure of *Pseudomonas aeruginosa*: Evidence for a clone that is pathogenic to the eye and that has a distinct combination of virulence factors. *Infect Immun* 2001;69:6284-95.
68. Chan KY, Cho P, Boost M. Microbial adherence to cosmetic contact lenses. *Cont Lens Anterior Eye* 2014;37:267-72.
69. Shen EP, Chu HS, Hsieh YT, Chen WL, Chang SC, Hu FR. Analysis of *P. aeruginosa* disinfectant sensitivity and microbial adhesions to worn cosmetic contact lenses. *Cont Lens Anterior Eye* 2020;43:338-44.
70. Watanabe T, Uematsu M, Mohamed YH, Eguchi H, Imai S, Kitaoka T. Corneal erosion with pigments derived from a cosmetic contact lens: A case report. *Eye Contact Lens* 2018;44(Suppl 1):S322-5.
71. Choy MH, Stapleton F, Willcox MD, Zhu H. Comparison of virulence factors in *Pseudomonas aeruginosa* strains isolated from contact lens- and non-contact lens-related keratitis. *J Med Microbiol* 2008;57:1539-46.
72. Lakkis C, Fleiszig SM. Resistance of *Pseudomonas aeruginosa* isolates to hydrogel contact lens disinfection correlates with cytotoxic activity. *J Clin Microbiol* 2001;39:1477-86.