





Chemical Composition and Biological Effects of *Pistacia atlantica* Desf. Oleoresin Essential Oil

Azam Elyasi Ghahfarrokhi¹ , Mina Saeedi^{2,3}, Mahnaz Khanavi^{1,4*} , Somayeh Mojtavavi⁵, Farzad Kobarfard⁶, Mohammad Ali Faramarzi⁵

¹Department of Pharmacognosy, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran.

²Medicinal Plants Research Center, Faculty of Pharmacy, Tehran University of Medical Sciences, Tehran, Iran.

³Persian Medicine and Pharmacy Research Center, Tehran University of Medical Sciences, Tehran, Iran.

⁴Faculty of Land and Food Systems, University of British Columbia, Vancouver, BC, Canada.

⁵Department of Pharmaceutical Biotechnology, Faculty of Pharmacy and Biotechnology Research Center, Tehran University of Medical Sciences, Tehran, Iran.

⁶Department of Medicinal Chemistry, School of Pharmacy, Shahid Beheshti University of Medical Sciences, Tehran, Iran.

Abstract

Background and objectives: The oleoresin of *Pistacia atlantica* Desf. (known as “Baneh” in Iran) has been frequently used in traditional medicine for its medicinal properties. Herein, *P. atlantica* essential oil was investigated for its antimicrobial and α -glucosidase inhibitory activities since α -pinene which has been identified as the most abundant component in *Pistacia* genus oil, has demonstrated antimicrobial and anti- α -glucosidase properties. **Methods:** Fresh oleoresin was collected from Javanroud, Kermanshah, Iran and the essential oil was obtained by Clevenger-type apparatus. The chemical composition of essential oil was identified with GC/MS analysis and compared with those reported from various regions. The antimicrobial activity was evaluated against various strains (*Pseudomonas aeruginosa*, *Staphylococcus aureus*, *Escherichia coli*, *Candida albicans*, *Saccharomyces cerevisiae*, and *Lactobacilli* spp.) through MIC method. Also, its anti- α -glucosidase property and antioxidant activity by DPPH assay were investigated. **Results:** GC/MS analysis of the essential oil confirmed the presence of nineteen compounds and among them, α -pinene (64.8%) was identified as the major constituent. Also, β -pinene (5.7%) and *cis*-limonene oxide (4.5%) were relatively abundant. Our results revealed antimicrobial properties of the “Baneh” essential oil against various bacterial and fungal strains. Moreover, it demonstrated inhibitory activity toward α -glucosidase with IC₅₀ value of 41.5 ± 2.5 mg/mL compared with acarbose (IC₅₀= 0.5 ± 0.2 mg/mL). DPPH free-radical scavenging activity assay showed antioxidant activity with IC₅₀ value of 155.2 ± 1.4 mg/mL compared with quercetin (IC₅₀= 250.0 ± 0.0 μ g/mL). **Conclusion:** *Pistacia atlantica* oleoresin essential oil depicted satisfactory antimicrobial activity against both Gram-positive and Gram-negative strains. However, it demonstrated low antioxidant and α -glucosidase inhibitory effects.

Keywords: biological activity; chemical analysis; essential oil; oleoresin; *Pistacia atlantica*

Citation: Elyasi Ghahfarrokhi A, Saeedi M, Khanavi M, Mojtavavi S, Kobarfard F, Faramarzi MA. Chemical composition and biological effects of *Pistacia atlantica* Desf. oleoresin essential oil. Res J Pharmacogn. 2022; 9(3): 25–33.

*Corresponding author: khanavim@sina.tums.ac.ir

Introduction

The resiniferous wild pistachio tree belongs to *Pistacia* genus which comprises eleven deciduous species and is mainly distributed in the Mediterranean and Middle Eastern areas. *Pistacia atlantica* Desf., known as “Baneh” in Iran, has been traditionally used by indigenous people in the west of the Zagros Mountains (mountain range in western Iran) as a rich source of nutritional and medicinal constituents [1]. It has been widely utilized as an antiseptic, appetizer, phlegm purgative, astringent, laxative, and carminative drug [2]. However, biological properties of the plant have been considered in the recent studies, e. g. *P. atlantica* leaves extracts have depicted remarkable antidiabetic activity under in vitro and in vivo [3,4]. Repeated oral administration of the hexane extract of *P. atlantica* seeds has confirmed beneficial hypoglycemic effects on streptozotocine-induced diabetic mice after a two-week treatment [5].

Essential oils are secondary metabolites of the aromatic plants which have demonstrated antioxidant, antibacterial, antiviral, antifungal, and insecticidal activity [6,7]. It has been reported that *P. atlantica* oleoresin essential oil possesses various biological activities such as antibacterial, anti-inflammatory, antifungal, antioxidant and antidiabetic effects [2,8]. Oral administration of *P. atlantica* oleoresin essential oil has been reported to be safe up to higher doses (2000 mg/kg) [8]. Therefore, it is deemed to be safe to use in the food industry [9]. Various studies have also indicated the antibacterial activity of *P. atlantica* oleoresin essential oil against several foodborne pathogens, including *Staphylococcus aureus*, *Salmonella enteritidis*, *Bacillus cereus*, *Escherichia coli* and *Helicobacter pylori* [10,11].

The composition of essential oils is remarkably associated with geographical position, climate conditions, and the method of extraction [12,13]. Different studies revealed that *P. atlantica* oleoresin essential oil contained high amounts of monoterpenes such as α -pinene (25-78%) and β -pinene (3-16%). However, oxygenated monoterpenes and sesquiterpenes were found to be low [14]. In this study, the essential oil of freshly collected *P. atlantica* oleoresin was obtained and the chemical composition was determined with GC/MS analysis. Its antimicrobial activity was evaluated against different bacterial and fungal strains and

compared with those results reported previously for the same species from various regions. It should be noted that evaluation of the corresponding essential oil against *Lactobacilli* spp. is reported for the first time. Also, *P. atlantica* oleoresin essential oil was evaluated for its α -glucosidase inhibitory activity which was not considered for the same species from different regions.

Materials and Methods

Ethical considerations

The Ethics Committee of Tehran University of Medical Sciences approved this research by the code IR.TUMS.TIPS.REC.1397.111.

Chemicals

All chemicals including *p*-nitrophenyl glucopyranoside (*p*-NPG), 2,2-diphenyl-1-picrylhydrazyl radical (DPPH), quercetin, potassium dihydrogen phosphate, and dipotassium hydrogen phosphate as well as *Saccharomyces cerevisiae* α -glucosidase (EC 3.2.1.20, 20 U/mg) were purchased from Sigma-Aldrich.

Plant material

Pistacia atlantica oleoresin was harvested from Javanroud, Kermanshah Province, Iran (34°48'24"N 46°29'19"E) in July 2018. It was identified and deposited at the Herbarium of Faculty of Pharmacy, Tehran University of Medical Science, Tehran, Iran with the voucher specimen of PMP-890. The collected oleoresin was kept at 4 °C.

Bacterial strains

Antimicrobial activity of the essential oil was assessed against Gram-negative bacteria including *Escherichia coli* (ATCC 8739) and *Pseudomonas aeruginosa* (ATCC 9027), Gram-positive bacteria including *Staphylococcus aureus* (ATCC 6538), *Lactobacillus acidophilus* (ATCC 53671), *Lactobacillus casei* (ATCC 27139), *Lactobacillus reuteri* (ATCC 23272), *Lactobacillus rhamnosus* (ATCC 39595), and *Lactobacillus plantarum* (ATCC 14917) as well as yeasts including *Candida albicans* (ATCC 10231) and *Saccharomyces cerevisiae* (ATCC 1611). All microbial strains were provided by Iranian Research Organization for Science and Technology (IROST).

Preparation of the essential oil

The volatile fraction was obtained by hydrodistillation of (100 g) oleoresin for 3 h using Clevenger-type apparatus. The resulting essential oil was dried over anhydrous sodium sulfate and stored in a sealed glass vial at 4 °C.

GC/MS analysis

Chemical analysis of the sample was achieved on an Agilent 7890A GC instrument with a flame ionization detector (FID) accompanied with MS-5975C MSD (Agilent Technologies, USA). The silica capillary column was DB-5 Agilent (30 × 0.25 mm ID, film thickness: 0.25 µm; CA, USA) and pure helium (99.999%) was applied as the carrier gas at a flow rate of 1 mL/min. Also, injection volume was 1.0 µL with split ratio of 1:5. The initial oven temperature was held 5 min isothermal at 40 °C and increased to 250 °C at a rate of 50 °C/min for 25 min. Likewise, the MS system was set in electron ionization (EI) mode with a quadrupole detector at 70 eV ionization energy. The transfer line and ion source temperatures were set at 250 and 230 °C, respectively. To identify the chemical composition of essential oil, the Kovats retention indices of components were calculated using retention times of an *n*-alkane ladder (C₆-C₁₄) that was injected after essential oil and mass spectra were compared with those reported in the literature. Furthermore, the essential oil compositions were determined by comparing the fragmentation patterns of the peaks with libraries mass spectra (Wiley 7n and NIST05A). The fragmentation patterns of the mass spectra were also compared with those reported in the literature [15].

DPPH free-radical scavenging assay

The experiment was performed according to the literature with some modification [16]. Briefly, 1 mL of different concentrations of essential oil (3000, 1500, 750 µg/mL) was added to 2 mL solution of DPPH in methanol (1 mM). After incubation at 37 °C for 30 min, the absorbance was read at 517 nm. Quercetin was used at the concentrations of 250, 125, 63, and 32 µg/mL as the positive control. Inhibition percent which indicated the antioxidant activity was calculated by the following equation:

$$I\% = [(A_{\text{Blank}} - A_{\text{Sample}}) / A_{\text{Blank}}] \times 100$$

A_{blank} is the absorbance of the control reaction which contains all components without the tested sample and A_{sample} is the absorbance of the tested sample. The concentration that inhibited 50% of DPPH solution (IC₅₀) was calculated by plotting the inhibition percent against sample concentrations. The test was performed in triplicate and the IC₅₀ values were reported as means ± SD.

Evaluation of antimicrobial activity

To obtain the minimum inhibitory concentration (MIC), quantitative tests were performed employing microdilution broth method using 96 U-shaped well plates [17]. Different concentrations of the essential oil (1843, 921, 460, 230, 115, 58, 29, 14.5, 7.2 µg/mL) were prepared based on the literature to evaluate the activity against the bacteria including *Staphylococcus aureus* ATCC 6538, *Escherichia coli* ATCC 8739, *Pseudomonas aeruginosa* ATCC 9027, *Lactobacillus acidophilus* ATCC 53671, *Lactobacillus casei* ATCC 27139, *Lactobacillus reuteri* ATCC 23272, *Lactobacillus plantarum* ATCC 14917, and *Lactobacillus rhamnosus* ATCC 39595 as well as two species of fungi including *Saccharomyces cerevisiae* ATCC 1611 and *Candida albicans* ATCC 10231. Ciprofloxacin (8 µg/mL) and nystatin (30 µg/mL) were used as the standard antibacterial and antimycotic controls, respectively. For this purpose, the fungi and lactobacilli were cultured on Sabouraud dextrose and De Man, Rogosa and Sharpe (MRS) broth, respectively. Mueller-Hinton was also used for *S. aureus*, *E. coli*, and, *P. aeruginosa*. The essential oil samples at the aforementioned concentrations were added to 1×10⁶ CFU/mL of each microorganism. Positive control (cultured along with the microorganism without the essential oil) and negative control (cultured without the microorganism) were also examined. After 24 h of incubation at 37 °C for bacteria and 25 °C for fungal species, the microdilution plates were investigated for the absence or presence of visible growth in comparison to sample-free control well. Each experiment was repeated three times with similar results. The endpoint of MIC was the lowest concentration of the essential oil at which the test strain did not demonstrate visible growth.

α-Glucosidase inhibition assay

The assay was performed according to previously

reported methods with some modifications [18]. Briefly, a mixture of phosphate buffer saline (135 μ L, pH 6.8, 50 mM), solution of α -glucosidase (20 μ L), and the essential oil at different final concentrations in range of 1344-10.5 μ g/mL (20 μ L) was pre-incubated at 37 °C for 10 min in a 96-well plate. Then, *p*-NPG (25 μ L, 4 mM) was added to each well and the absorbance change was recorded at 400 nm within 20 min at 37 °C. The percent inhibition of all concentrations, control, and acarbose (positive control) was calculated using the following formula:

$$\% \text{ Inhibition} = [(\text{Abs control} - \text{Abs sample}) / \text{Abs control}] \times 100$$

IC₅₀ values were calculated from the nonlinear regression.

Results and Discussion

The *P. atlantica* oleoresin essential oil was obtained using a Clevenger-type apparatus by hydrodistillation of freshly collected oleoresin with 25% w/w yield. It was a viscous liquid in light-yellow colour, possessing a pungent odor and the density of 0.84 g/cm³. The composition of EO was determined using GC/MS analysis (Table 1). Nineteen compounds were identified, which accounted for 99.2% of the total composition. It should be noted that 75.1% of those compounds belonged to the monoterpenes and 24.1% were oxygenated monoterpenes. Among them, α -pinene (Table 1, entry 1) was the main constituent (64.8%). β -Pinene (5.7%), *cis*-limonene oxide (4.5%), *trans*-pinocarveol (3.3%), *trans*-verbenol (2.7%), α -campholenaldehyde (2.6%), and 8-hydroxy-*p*-cymene (2.3%) were also identified [19-34].

Comparing our results with those reported in the literature (Table 2) obtained from the same species in different parts of Iran, it was found that α -pinene is the most abundant compound; however, its quantity is depended on geographical region of the harvested plant. Accordingly, α -pinene was obtained in a wide range of 55.0-97.2 % [35-42]. The amount of α -pinene in *P. atlantica* oleoresin essential oil from different parts of Algeria was reported in the range of 25.4-79.8% [43,44]. It should be noted that chemical composition of the essential oil from the oleoresin grown in Iraq and Morocco demonstrated α -pinene of 57.1 and 42.9%, respectively [45,46].

Table 1. Chemical composition of *Pistacia atlantica* oleoresin essential oil, identified by GC/MS analysis

Entry	Compound	KI ^c	KI ^{ref}	Percentage
1	α -Pinene	973	965	64.8
2	Camphene	978	968	0.9
3	Sabinene	1002	991	0.9
4	β -Pinene	1020	999	5.7
5	<i>p</i> -Cymene	1030	1026	1.3
6	Limonene	1035	1031	1.5
7	Eucalyptol	1039	1039	0.6
8	2,3-Epoxy-pinane	1109	1103	1.0
9	α -Campholenaldehyde	1121	1122	2.6
10	<i>cis</i> -Limonene oxide	1137	1138	4.5
11	<i>trans</i> -Pinocarveol	1154	1149	3.3
12	<i>trans</i> -Verbenol	1158	1150	2.7
13	Pinocarvone	1174	1170	1.5
14	8-Hydroxy- <i>p</i> -cymene	1195	1194	2.3
15	Myrtenol	1207	1208	1.6
16	Verbenone	1221	1221	1.0
17	<i>trans</i> -Carveol	1227	1226	1.0
18	Carvone	1259	1257	1.0
19	Bornyl acetate	1293	1293	1.0
Monoterpenes				75.1
Oxygenated monoterpenes				24.1
Sesquiterpenes				-
Total identified				99.2

KI^c: Calculated Kovats Index, KI^{ref}: Reference Kovats Index

Table 2. Chemical constituents (%) of *Pistacia atlantica* oleoresin essential oil from different regions

Geographical region	*Iran	°Iran	Algeria	Iraq	Morocco
Components					
α -Pinene	64.8	55.0-97.2	25.4-79.8	57	42.9
β -Pinene	5.7	1.3-6.7	3-16.5	9.8	13.2
Limonene oxide	-	9	-	-	-
Myrcene	-	11.4	-	-	-
1,8-Cineol	0.6	0.6, 3.1	-	-	-
<i>trans</i> -Verbenol	2.7	2.7	-	3.8	-
<i>trans</i> -Pinocarveol	3.3	5.7	-	2.9	3.2
Carveol	-	2.2	-	-	-
Sabinene	1	1.2-4.5	-	-	-
<i>p</i> -Cymene	1.3	1.1, 1.2, 1.8	-	1.3	-
<i>p</i> -Cymenene	-	1.2, 1.4	-	-	-
Citral	-	5.7	-	-	-
Myrtenol	1.6	5.3	-	-	2.0
α -Terpinolene	-	1.3, 1.6, 1.8	-	-	-
Camphen	1	1.3, 1.4	-	1	-
Myrtenal	-	1.1, 1.4	-	1.7	-
Bornyl acetate	1	1.5	-	-	-
D-limonene	1.5	1.5, 10.1	5.1	-	2.0
Thujene	-	1.4	-	-	-
<i>E</i> -caryophyllene	-	2.6	-	-	-
Pinocarvone	1.5	-	5.5	-	-
Δ -2-Carene	-	1.1	-	-	-
<i>neo</i> -Isoverbenol	-	1.3, 1.4	-	-	-
<i>cis</i> -Limonene oxide	4.5	1.1-4.2	-	-	-
α -Phellandren	-	1.8	-	-	-
α -Campholenaldehyde	2.6	1.9	-	1.9	-

*Iran: The result of chemical analysis of *Pistacia atlantica* oleoresin essential oil collected from Javanroud; °Iran: The result of chemical analysis of *P. atlantica* oleoresin essential oil collected from other parts of Iran

The *P. atlantica* oleoresin essential oil showed in vitro antioxidant activity (>50%) at high dose of 2000 mg/mL with IC₅₀ value of 155.2±1.4 mg/mL compared with quercetin as the positive control (IC₅₀=250.0±0.0 µg/mL). This result was in accordance with identified compounds (Table 1), indicating a high level of monoterpenes.

The calculated minimum inhibitory concentration (MIC) values against examined strains were reported in Table 3. According to our results, the essential oil demonstrated antimicrobial activity against most microorganisms tested with MIC lower than 1 mg/mL. Among them, the essential oil showed the MIC value of 12.6 µg/mL against *L. acidophilus*. Also, it inhibited growth of *P. aeruginosa* and *C. albicans* with MIC value of 37.8 µg/mL. Antimicrobial activity of the essential oil against *S. aureus*, *E. coli*, *L. reuteri*, *L. plantarum*, and *L. rhamnosus* was found to be 115.2 µg/mL. However, it was not efficiently active toward *S. cerevisiae* and *L. casei* (MIC > 1 mg/mL).

Table 3. MIC values (µg/mL) of *Pistacia atlantica* oleoresin essential oil, ciprofloxacin, and nystatin against microbial strains

Bacteria/fungi	MIC (µg/mL)		
	EO*	Ciprofloxacin	Nystatin
<i>Staphylococcus aureus</i>	115.2	0.5	-
<i>Escherchia coli</i>	115.2	1.0	-
<i>Pseudomonae aeruginosa</i>	37.8	2.0	-
<i>Saccharomyces ceriviseae</i>	1034.5	-	30.0
<i>Candida albicans</i>	37.8	-	30.0
<i>Lactobacillus acidophilus</i>	12.6	1.0	-
<i>Lactobacillus casei</i>	1034.5	1.0	-
<i>Lactobacillus reuteri</i>	115.2	1.0	-
<i>Lactobacillus plantarum</i>	115.2	1.0	-
<i>Lactobacillus rhamnosus</i>	115.2	1.0	-

EO: essentialoil

Pistacia atlantica oleoresins grown in different geographical regions have demonstrated antibacterial and antifungal activity [10,47,48]. In this regard, high activity against Gram-positive bacteria has been significant. In our study, the more considerable activity of *P. atlantica* oleoresin essential oil was obtained against Gram-negative bacteria which are more resistant than Gram-positive type and can be considered for the respiratory infections. Also, good activity of the essential oil against *L. acidophilus* probably makes it applicable for oral health care. Antimicrobial activity of the essential oil seems to be associated with the presence of α -pinene. Based on the literature, this component has shown good activity against pathogenic yeasts or molds [49-52].

The antimicrobial activity of the essential oil can be explained by high hydrophobicity, which is directly correlated to log *P* (partitioning behavior of the lipophilic compounds in octanol/water) that targets the microbial cytoplasmic membrane [53]. Indeed, lipophilic compounds particularly those with log *P* higher than 3 possess high affinity for cell membranes, and their accumulation induces changes in membrane physico-chemical properties. In this respect, α -pinene, *p*-cymene, and β -pinene (monoterpenes) as the most abundant constituents of the EO, are lipophilic compounds and have log *P* values in the range of 3.1-4.1. Also, oxygenated monoterpenes which accounted for 24.1% of the essential oil could play a more or less important role in the antimicrobial activity. The presence of oxygenated terpenes in trace amount can also increase water solubility and the hydrogen bonding capacity, which are the main factors influencing MIC values of terpenoids against *P. aeruginosa* and *S. aureus* [54,55].

α -Glucosidase inhibitory activity of *P. atlantica* oleoresin essential oil as well as the major components (α -pinene and β -pinene), were evaluated and IC₅₀ values were obtained as 41.5±2.5, 0.2±3.2, and 0.49±1.4 mg/mL, respectively, compared with acarbose as the positive control (IC₅₀=0.50±0.2 mg/mL). The inhibitory activity may be associated with the presence of monoterpenes such as α -pinene and β -pinene which showed more potent activity than acarbose. However, the synergistic effect led to the reduction of activity.

In the case of inhibitory activity against α -glucosidase, which plays a key role in oligosaccharide hydrolysis, essential oils have been found as potent sources of natural inhibitors [56-59]. Some reports have also revealed that terpenes, as the main components of various plants, are capable of exerting blood glucose lowering effects and monoterpenes such as α -pinene, β -pinene, myrcene, 1,8-cineole, and sabinene have been reported as the potent inhibitors of α -amylase and α -glucosidase [60]. The essential oil of *Hertia cheirifolia* flowers indicated significant α -glucosidase inhibitory effect (IC₅₀=0.24±0.01, acarbose=0.28±0.01 mg/mL). Its composition was characterized by the predominance of α -pinene accounted for 70.4% of the oil [61]. However, in this study the presence of α -pinene as the major compound induced no satisfactory anti- α -glucosidase

activity. It seems that the inhibitory activity was affected by the whole components.

Conclusion

In this study, chemical composition and biological activities of the *P. atlantica* oleoresin essential oil which was collected from Javanroud, Kermanshah, Iran was investigated. According to the GC/MS analysis, α -pinene was detected as the most abundant component in the essential oil (64.8%). It showed broad-spectrum antibacterial activity against different gram-positive and gram-negative bacteria as well as two fungal strains, even against the multi-resistant bacteria, *P. aeruginosa*. It seems that this oil can be considered as a natural and potent antimicrobial agent. However, *P. atlantica* oleoresin essential oil depicted a weak α -glucosidase inhibitory activity in spite of the fact that its major component displayed potent activity.

Acknowledgments

This work was supported by grants from the Research Council of Tehran University of Medical Sciences with project No 97-03-33-40820.

Author contributions

Azam Elyasi performed the experiments and prepared the manuscript; Mina Saeedi participated in the characterization of components and preparation of the manuscript; Mahnaz Khanavi designed and supervised all project steps; Somayeh Mojtavavi contributed to the evaluation of biological activity; Farzad Kobarfard conducted GC/MS analysis; Mohammad Ali Faramarzi supervised biological assays.

Declaration of interest

The authors declare that there is no conflict of interest. The authors alone are responsible for the content of the paper.

References

- [1] Pourreza M, Shaw JD, Zangeneh H. Sustainability of wild pistachio (*Pistacia atlantica* Desf.) in Zagros forests. *For Ecol Manage*. 2008; 255(11): 3667–3671.
- [2] Ahmed ZB, Yousfi M, Viaene J, Dejaegher B, Demeyer K, Heyden YV. Four *Pistacia atlantica* subspecies (*atlantica*, *cabulica*, *kurdica* and *mutica*): a review of their botany,

ethnobotany, phytochemistry and pharmacology. *J Ethnopharmacol*. 2021; Article ID 113329.

- [3] Kasabri V, Afifi FU, Hamdan I. In vitro and in vivo acute antihyperglycemic effects of five selected indigenous plants from Jordan used in traditional medicine. *J Ethnopharmacol*. 2011; 133(2): 888–896.
- [4] Ahmed ZB, Yousfi M, Viaene J, Dejaegher B, Demeyer K, Mangelings D, Vander Heyden Y. Potentially antidiabetic and antihypertensive compounds identified from *Pistacia atlantica* leaf extracts by LC fingerprinting. *J Pharm Biomed Anal*. 2018; 5(149): 547–556.
- [5] Hashemnia M, Nikousefat Z, Yazdani-Rostam M. Antidiabetic effect of *Pistacia atlantica* and *Amygdalus scoparia* in streptozotocin-induced diabetic mice. *Comp Clin Pathol*. 2015; 24(6): 1301–1306.
- [6] Panikar S, Shoba G, Arun M, Sahayarayan JJ, Usha Raja Nanthini A, Chinnathambi A, Alharbi SA, Nasif O, Kim HJ. Essential oils as an effective alternative for the treatment of COVID-19: molecular interaction analysis of protease (Mpro) with pharmacokinetics and toxicological properties. *J Infect Public Health*. 2021; 14(5): 601–610.
- [7] Ayub M, Hanif M, Sarfraz R, Muhammad Shahid M. Biological activity of *Boswellia serrate* Roxb. oleo gum resin essential oil: effects of extraction by supercritical carbon dioxide and traditional methods. *Int J Food Prop*. 2018; 21(1): 808–820.
- [8] Memariani Z, Sharifzadeh M, Bozorgi M, Hajimahmoodi M, Farzaei MH, Gholami M, Siavoshi F, Saniee P. Protective effect of essential oil of *Pistacia atlantica* Desf. on peptic ulcer: role of α -pinene. *J Tradit Chin Med*. 2017; 37(1): 57–63.
- [9] Labdelli A, Zemour K, Simon V, Cerny M, Adda A, Merah O. *Pistacia atlantica* Desf., a source of healthy vegetable oil. *Appl Sci*. 2019; 9(12): 1–11.
- [10] Ghalem BR, Mohamed M. Essential oil from gum of *Pistacia atlantica* Desf: screening of antimicrobial activity. *Afr J Pharm Pharmacol*. 2009; 3(1): 13–15.
- [11] Benabderrahmane M, Benali M, Aouissat H, Jordán MJ. Antimicrobial activity of the essential oils of *Pistacia atlantica* Desf. from Algeria. 2009; 7(6): 304–308.
- [12] Biswas S, Koul M, Bhatnagar AK. Effect of

- salt, drought and metal stress on essential oil yield and quality in plants. *Nat Prod Commun.* 2011; 6(10): 1559–1564.
- [13] Locali-Pereira AR, Lopes NA, Menis-Henrique MEC, Janzantti NS, Nicoletti VR. Modulation of volatile release and antimicrobial properties of pink pepper essential oil by microencapsulation in single- and double-layer structured matrices. *Int J Food Microbiol.* 2020; Article ID 108890.
- [14] Asadollahzadeh H, Shamspur T. Chemical composition of the extracts of fruits of *Pistacia atlantica* Desf. from Kerman province in Iran. *J Essent Oil-Bear Plants.* 2013; 16(2): 243–246.
- [15] Adams RP. Identification of essential oil components by gas chromatography mass spectrometry. 4th ed. Carol Stream: Allured Publishing, 2007.
- [16] Kiashi F, Hadjiakhoondi A, Tofighi Z, Khanavi M, Ajani Y, Ahmadi Koulaei S, Yassa N. Compositions of essential oils and some biological properties of *Stachys laxa* Boiss. & Buhse and *S. byzantina* K. Koch. *Res J Pharmacogn.* 2021; 8(2): 5–15.
- [17] Balouiri M, Sadiki M, Ibnsouda SK. Methods for in vitro evaluating antimicrobial activity: a review. *J Pharm Analysis.* 2016; 6(2): 71–79
- [18] Saeedi M, Raeisi-Nafchi M, Sobhani S, Mirfazli SS, Zardkanlou M, Mojtavavi S, Faramarzi MA, Akbarzadeh T. Synthesis of 4-alkylaminoimidazo[1,2-a] pyridines linked to carbamate moiety as potent α -glucosidase inhibitors. *Mol Divers.* 2021; 25(4): 2399–2409.
- [19] Klesk K, Qian M. Preliminary aroma comparison of marion (*Rubus* spp. hyb) and evergreen (*R. laciniatus* L.) blackberries by dynamic headspace/OSME technique. *J Food Sci.* 2003; 68(2): 697–700.
- [20] Bylaite E, Meyer AS. Characterisation of volatile aroma compounds of orange juices by three dynamic and static headspace gas chromatography techniques. *Eur Food Res Technol.* 2006; 222(1): 176–184.
- [21] Miyazaki T, Plotto A, Goodner K, Gmitter FG. Distribution of aroma volatile compounds in tangerine hybrids and proposed inheritance. *J Sci Food Agric.* 2011; 91(3): 449–460.
- [22] Kim TH, Thuy NT, Shin JH, Baek HH, Lee HJ. Aroma-active compounds of miniature beefsteak plant (*Mosla dianthera* Maxim.). *J Agric Food Chem.* 2000; 48(7): 2877–2881.
- [23] Adams RP, González Elizondo MS, González Elizondo M, Slinkman E. DNA fingerprinting and terpenoid analysis of *Juniperus blancoi* var. *huehuentensis* (Cupressaceae), a new subalpine variety from Durango, Mexico. *Biochem Syst Ecol.* 2006; 34(3): 205–211.
- [24] Baccouri B, Ben Temime S, Campeol E, Cioni PL, Daoud D, Zarrouk M. Application of solid-phase microextraction to the analysis of volatile compounds in virgin olive oils from five new cultivars. *Food Chem.* 2007; 102(3): 850–856.
- [25] Loayza I, Abujder D, Aranda R, Jakupovic J, Collin G, Deslauriers H, Jean FI. Essential oils of *Baccharis salicifolia*, *B. latifolia* and *B. dracunculifolia*. *Phytochemistry.* 1995; 38(2): 381–389.
- [26] Senatore F, Napolitano F, Arnold NA, Bruno M, Herz W. Composition and antimicrobial activity of the essential oil of *Achillea falcata* L. (Asteraceae). *Flavour Fragr J.* 2005; 20(3): 291–294.
- [27] Choi HS. Character impact odorants of *Citrus Hallabong* [(*C. unshiu* Marcov \times *C. sinensis* Osbeck) \times *C. reticulata* Blanco] cold-pressed peel oil. *J Agric Food Chem.* 2003; 51(9): 2687–2692.
- [28] Courtois EA, Paine CE, Blandinieres PA, Stien D, Bessiere JM, Houel E, Baraloto C, Chave J. Diversity of the volatile organic compounds emitted by 55 species of tropical trees: a survey in French Guiana. *J Chem Ecol.* 2009; 35(11): 1349–1362.
- [29] Kowalski R, Wolski T. The chemical composition of essential oils of *Silphium perfoliatum* L. *Flavour Fragr J.* 2005; 20(3): 306–310.
- [30] Goodner KL, Mahattanatawee K, Plotto A, Sotomayor JA, Jordan MJ. Aromatic profiles of *Thymus hyemalis* and Spanish *T. vulgaris* essential oils by GC-MS/GC-O. *Ind Crops Prod.* 2006; 24(3): 264–268.
- [31] Hamm S, Bleton J, Connan J, Tchaplal A. A chemical investigation by headspace SPME and GC-MS of volatile and semi-volatile terpenes in various olibanum samples. *Phytochemistry.* 2005; 66(12): 1499–1514.
- [32] Mitiku SB, Sawamura M, Itoh T, Ukeda H. Volatile components of peel cold-pressed oils of two cultivars of sweet orange (*Citrus sinensis* (L.) Osbeck) from Ethiopia. *Flavour*

- Fragr J.* 2000; 15(4): 240–244.
- [33] Lawal OA, Oyedele AO. Chemical composition of the essential oils of *Cyperus rotundus* L. from South Africa. *Molecules.* 2009; 14(8): 2909–2917.
- [34] Mosayebi M, Amin G, Arzani H, Azarnivand H, Maleki M, Shafaghat A. Effect of habitat on essential oil of *Achillea filipendula* L. in Iran. *Asian J Plant Sci.* 2008; 7(8): 779–781.
- [35] Didehvar M, Ebadi M, Ayyari M. Qualitative and quantitative evaluation of *Pistacia atlantica* Desf. essential oil from thirteen natural habitats. *Iran J Horti Sci.* 2021; 52(2): 419–428.
- [36] Izadyar S, Babaei Y, Efhamisisi D. Effect of natural resin from wild pistachio trees on physical properties and durability of beech wood: alone and in combination with boric acid. *Drvna Ind.* 2020; 71(4): 379–388.
- [37] Peivastegan M, Rajabi M, Zaferani Arani H, Olya M, Atashi HA, Abolghasemi SH. Comparing the effects of oleoresin of *Pistacia atlantica* tree and diclofenac gel on the knee osteoarthritis improvement. *Shiraz E-Med J.* 2020; 21(10): 1–8.
- [38] Samedoroudi P, Mirzajani F, Aliahmadi A, Ghassempour A. Top-down thermal analysis versus bottom-up gas chromatography–mass spectrometry in an adulteration study of *Pistacia atlantica* Desf. oleoresin. *J Therm Anal Calorim.* 2016; 123: 2451–2457.
- [39] Najafi MBH, Farimani RH, Tavakoli J, Madayeni S. GC-MS analysis and antimicrobial activity of the essential oil of trunk exudates of *Pistacia atlantica* var. *mutica*. *Chem Nat Compd.* 2014; 50(2): 376–378.
- [40] Hosseini F, Adlgostar A, Sharifnia F. Antibacterial activity of *Pistacia atlantica* extracts on *Streptococcus mutans* biofilm. *Int Res J Biological Sci.* 2013; 2(2): 1–7.
- [41] Sharifi MS, Hazell SL. GC-MS analysis and antimicrobial activity of the essential oil of the trunk exudates from *Pistacia atlantica kurdica*. *J Pharm Sci Res.* 2011; 3(8): 1364–1367.
- [42] Delazar A, Reid RG, Sarker SD. GC-MS analysis of the essential oil from the oleoresin of *Pistacia atlantica* var. *mutica*. *Chem Nat Compd.* 2004; 40(1): 24–27.
- [43] Benhassaini H, Bendeddouche FZ, Mehdadi Z, Romane A. GC/MS analysis of the essential oil from the oleoresin of *Pistacia atlantica* Desf. subsp. *atlantica* from Algeria. *Nat Prod Commun.* 2008; 3(6): 929–932.
- [44] Benabderrahmane M, Aouissat M, Bueso MJ, Bouzidi A, Benali M. Chemical composition of essential oils from the oleoresin of *Pistacia atlantica* Desf. from Algeria. *J Biochem Int.* 2015; 2(4): 133–137.
- [45] Awla Shekhany H, Ahemad H. The study of chemical composition of gum in *Pistacia atlantica* in Erbil region. *Zanco J Pure Appl Sci.* 2018; 30(3): 26–32.
- [46] Barrero AF, Herrador MM, Arteaga JF, Akssira M, Mellouki F, Belgarrabe A, Blázquez MA. Chemical composition of the essential oils of *Pistacia atlantica* Desf. *J Essent Oil Res.* 2005; 17(1): 52–54.
- [47] Aksoy A, Duran N, Koksall F. In vitro and in vivo antimicrobial effects of mastic chewing gum against *Streptococcus mutans* and *mutans streptococci*. *Arch Oral Biol.* 2006; 51(6): 476–481.
- [48] Daifas DP, Smith JP, Blanchfield B, Sanders G, Austin JW, Koukoutisis J. Effects of mastic resin and its essential oil on the growth of proteolytic *Clostridium botulinum*. *Int J Food Microbiol.* 2004; 94(3): 313–322.
- [49] Leite AM, Lima EO, Souza EL, Melo Diniz MFF, Trajano VN, Medeiros IA. Inhibitory effect of beta-pinene, alpha-pinene and eugenol on the growth of potential infectious endocarditis causing Gram-positive bacteria. *Rev Bras Cienc Farm.* 2007; 43(1): 121–126.
- [50] Moreira ACP, Lima EO, Souza EL, Dingenen MAV, Trajano VN. Inhibitory effect of *Cinnamomum zeylanicum* Blume (Lauraceae) essential oil and β -pinene on the growth of dematiaceous moulds. *Braz J Microbiol.* 2007; 38(1): 33–38.
- [51] Lima IO, Oiviera RAG, Lima EO, Souza EL, Farias NP, Navarro DF. Inhibitory action of some phytochemicals on yeasts potentially causing of opportunistic infections. *Rev Bras Cienc Farm.* 2005; 41(2): 199–203.
- [52] Salehi B, Upadhyay S, Orhan IE, Jugran AK, Jayaweera SLD, Dias DA, Sharopov F, Taheri Y, Martins N, Baghalpour N, Cho WC, Sharifi-Rad J. Therapeutic potential of α - and β -pinene: a miracle gift of nature. *Biomolecules.* 2019; 9(11): 1–34.
- [53] Ben Arfa A, Combes S, Preziosi-Belloy L, Gontard N, Chalier P. Antimicrobial activity of carvacrol related to its chemical structure.

- Lett Appl Microbiol.* 2006; 43(2): 149–154.
- [54] Moumni S, Elaissi A, Trabelsi A, Merghni A, Chraief I, Jelassi B, Chemi R, Ferchichi S. Correlation between chemical composition and antibacterial activity of some Lamiaceae species essential oils from Tunisia. *BMC Complement Med Ther.* 2020; 20(1): 1–15.
- [55] Miladinović DL, Dimitrijević MV, Mihajilov-Krstev TM, Marković MS, Ćirić VM. The significance of minor components on the antibacterial activity of essential oil via chemometrics. *LWT.* 2021; Article ID 110305.
- [56] Wu T, Luo J, Xu B. In vitro antidiabetic effects of selected fruits and vegetables against glycosidase and aldose reductase. *Food Sci Nutr.* 2015; 3(6): 495–505.
- [57] Toyomizu M, Sugiyama S, Jin R, Nakatsu T. α -Glucosidase and aldose reductase inhibitors: constituents of cashew, *Anacardium occidentale*, nut shell liquids. *Phytother Res.* 1993; 7(3): 252–254.
- [58] Shihabudeen HMS, Priscilla DH, Thirumurugan K. Cinnamon extract inhibited the α -glucosidase activity and dampens postprandial glucose excursion in diabetic rats. *Nutr Metab.* 2011; 8(1): 1–11.
- [59] Jaradat NA, Al-Maharik N, Abdallah S, Shawahna R, Mosa A, Qtishat A. *Nepeta curviflora* essential oil: phytochemical composition, antioxidant, anti-proliferative and anti-migratory efficacy against cervical cancer cells, and α -glucosidase, α -amylase and porcine pancreatic lipase inhibitory activities. *Ind Crops Prod.* 2020; 158: 1–11.
- [60] Oboh G, Olasehinde TA, Ademosun AO. Inhibition of enzymes linked to type-2 diabetes and hypertension by essential oils from peels of orange and lemon. *Int J Food Prop.* 2017; 20(S1): 586–594.
- [61] Majouli K, Hlila MB, Hamdi A, Flamini G, Jannet HB, Kenani A. Antioxidant activity and α -glucosidase inhibition by essential oils from *Hertia cheirifolia* (L.). *Ind Crops Prod.* 2015; 82: 23–28.

Abbreviations

EO: essential oil; GC-MS: gas chromatography-mass spectrometry; MIC: minimum inhibitory concentration; DPPH: 2,2-diphenyl-1-picrylhydrazyl; IC₅₀: the half maximal inhibitory concentration; ATCC: American type culture collection; SD: standard deviation; *p*-NPG: *p*-nitrophenyl glucopyranoside