

## Explained: How to save our environment for our future generations through sustainable food systems?

**Parmeet Kaur**

Department of Dietetics

All India Institute of Medical Sciences, New Delhi, India

Email: [parjaskaur@yahoo.com](mailto:parjaskaur@yahoo.com)

**Date of Submission:** 13 Aug 2022 || **Date of Acceptance:** 3 Sep 2022

---

### Abstract

The food choices which we make in our daily lives are known to have a direct impact on our environment. Our planet earth has been getting warmer in recent years. Asia accounts for nearly half of global greenhouse gas emissions (GHGs). It is predicted that by 2050, a large number of regions in Asia will be exposed to deadly levels of GHGs. It is now time that we recognize the environmental impacts of the type and amount of food we eat considering the planet and our health. It is very important to address these issues because a growing body of scientific evidence suggests that our dietary habits are putting a lot of strain on the environment. Encountering the recent trends of overconsumption of discretionary foods (foods and beverages not required to provide nutrients the body needs) is key to aligning human and planetary health. Although the undesirable health effects of discretionary foods are an increased risk of obesity and chronic diseases, the environmental and broader sustainability impacts of these food products require more public awareness, especially since their dietary consumption has been increasing greatly in recent decades. Therefore, moving to more sustainable food choices, dietary patterns, and sustainable diets is required to strengthen global efforts to save our planet earth.

**Keywords:** sustainable food system, environment, discretionary foods

---

### How to Cite This Article:

Kaur Parmeet, Explained: How to save our environment for our future generations through sustainable food systems? *Annals of Ayurvedic Med.* 2022;11(3) 200-204, 10.5455/AAM.103794

---

### Introduction

A sustainable food system provides adequate and nutritious food without harmful environmental impacts. Earth's history is divided into a hierarchy of smaller portions of time, referred to as the geologic time scale. Anthropocene epoch is an unofficial unit of geologic time. It is referred to as the most recent period in Earth's history when human activity started having a notable influence on the planet's climate and ecosystems. It is still unclear whether the anthropocene started in the year 1800, with the industrial revolution when human activity created carbon and methane in the atmosphere. In 1945 radioactive particles were first released into the environment when humans tested and dropped the first atomic bombs.<sup>1</sup> Thereafter, the load of GHGs is continuously going on increasing due to untamed human

activities on the planet. In view of the COVID-19 pandemic, there was a brief stall in GHG in early 2020.<sup>2</sup> Intergovernmental panels on climate change (IPCC) has reported that the average surface temperature of the Earth will cross 1.5 degree Celsius in the next 20 years, and 2 degrees by the middle of the century. It is suggested that the countries in Asia and the Pacific should scale up and accelerate climate action by implementing justified strategies. Dietary choices are a key determinant of both health outcomes and environmental impacts.<sup>3</sup> Therefore, it is proposed to make the following dietary changes in our daily life to save our environment for our future generations:

### Follow traditional regional diets

The modern food systems in the last few decades have tremendously changed our dietary patterns. Trends of

global preference for highly processed foods, ready-to-eat foods and fast foods have further made a direct impact on the communities and the environment. The modern dietary trends negatively affect a trio of factors: health, agriculture, and the environment.

Nowadays, the link between dietary choices and carbon footprints is being recognized by more and more people. Dieticians can play a key role in the dissemination of information to the masses regarding attaining a sustainable food system, where food is purchased using techniques that protect the environment, public health, communities, natural ecosystems, and animals is needed. The distinct dietary patterns have markedly different environmental implications. The rice-based agricultural patterns are stated to have greater GHGEs than the wheat-based patterns, principally due to methane emissions from flooded rice crop production. Therefore, traditional regional diets prepared from locally available food ingredients should be encouraged to promote biodiversity.<sup>4</sup>

### **Safeguard natural ecosystem**

Climate change threatens soil health. The health of the soil is known to be connected with human and planetary health. Soil microbiome survives in soil ecosystems. These soil microbial communities are essential, for the cycling of carbon and other nutrients. Widespread use of inorganic fertilizers and pesticides affects soil ecology. Healthy soils favor soil carbon sequestration, and assist in removing carbon dioxide from the atmosphere and preserving it in a soil carbon pool through plants in form of soil organic carbon. However, the conversion of natural ecosystems to agriculture releases that carbon into the atmosphere. Additionally, reducing tillage, erosion control; organic amendments and cover crops can escalate soil organic carbon.<sup>5,6</sup>

### **Prefer plant based foods**

Food choices as a part of dietary patterns can make a significant impact on the environment. Scientific research has consistently shown that plant-based dietary patterns

are linked with lower environmental impact. Recent research of nine diets aligned with criteria for a healthful diet, specific to 140 countries found that shifts in diets to mostly plants based and low food chain animals such as forage fish, mollusks, and insects is advantageous. It was stated that the vegan diet had the greatest benefit. The next best eco-impact diets were a low food chain, two-thirds vegan, pescatarian, no dairy, no red meat, and vegetarian diet. Reducing the quantity of meat and one meatless day in a week was found to have the lowest advantage.<sup>7</sup>

The research findings further postulate that adopting any one of the common sustainable dietary patterns, ranging from vegan and vegetarian to traditional Mediterranean and New Nordic diets could result in reductions as high as 80% of GHGEs and land use, and 50% of water use. The findings from the Adventist health study and California state agricultural data showed that non-vegetarian diets used 2.9 times more water, 2.5 times more primary energy, 13 times more fertilizer, and 1.4 times more primary energy as compared to vegetarian dietary patterns. The consumption of beef has been documented as the major contributor. The EPIC-Oxford cohort study compared the GHGEs among meat eaters, fish eaters, vegetarians, and vegans in the United Kingdom. The values of carbon dioxide equivalent for vegetarians were 3.81, fish eaters, 3.91, and for low meat eaters were 4.67.<sup>8,9</sup>

According to the EAT-Lancet report, global consumption of fruits, vegetables, nuts, and legumes should be doubled and consumption of foods such as red meat and sugar should be reduced by more than 50%. A diet rich in plant-based foods with fewer animal source foods confers both health and environmental benefits.<sup>10</sup> However, adapting to dietary changes suggested by EAT-Lancet report would require higher income, nutritional assistance, and lower food prices for the diverse global population.<sup>11</sup>

Recently, methionine reduced diets have been associated to longevity in animal studies.<sup>12</sup> Animal sources of protein such as beef, lamb, pork, and eggs contain higher levels of methionine than plant-sourced proteins.<sup>13</sup> However, as

methionine is an essential amino acid for humans, the restriction could also increase the risk of deficiency.<sup>12</sup>

### **Integrate Crops and Livestock production**

Adopting integrated cropping and livestock production allows for better management of nutrient flows by recycling resources. In ecological livestock systems, manure is not a waste but serves as a valuable input that needs to be returned to soils and the crop waste serves as fodder.<sup>14, 15</sup>

### **Prefer Plant-Based Proteins**

Proteins vary broadly in terms of nutritional profile, digestibility and bioavailability, environmental implications, and consumer acceptance. Pulses are known to be among the most sustainable plant proteins. They are known to have the lowest carbon footprints, are drought tolerant, and enrich the soil through nitrogen-fixing bacteria which reduces the need for fertilizers. Pulses also promote food security in developing countries. It is estimated that the growing of pulses can help increase agricultural products which will be needed to feed the world by 2050. Animal foods require more energy to produce the same amount of protein than for high protein plant foods such as soy and pulses.<sup>16, 17</sup>

### **Trim down meat consumption**

Huge reductions are essential in meat-eating to avoid hazardous climate change. Countries, like the Netherlands, are advocating reducing red meat intake and increasing plant foods.<sup>18</sup> China, which dramatically increased meat consumption (its average intake of 13 kg per person per year in 1982 soared to 63 kg in 2016) along with affluence, has goals in its dietary guidelines to cut meat consumption in half.<sup>19</sup> British Dietetic Association's Environmentally Sustainable Diet (One Blue Dot) recommendations for the United Kingdom call for reducing red meat to less than 71 g per person per day, besides avoiding intake of processed meats and choosing high-protein plant foods, such as beans, lentils, soy, nuts, and seeds.<sup>20</sup>

### **Excuse yourself from Discretionary Foods**

Discretionary choices are foods or beverages high in saturated fat, added sugars, or salt. Discretionary food consumption is associated with an increased risk of obesity and chronic diseases. Consumption of these foods has increased steeply in high-income countries and the developing world. Eating healthful, nutritious foods, such as whole grains, legumes, vegetables, and fruits, is better for human health and the health of the planet.<sup>21</sup>

### **Monitor food related waste**

Food waste is not just a social and humanitarian concern it is also an environmental concern. If wasted food goes to the landfill and rots, it produces methane, a greenhouse gas that is even more potent than carbon dioxide. Today an estimated one-third of all the food produced in the world goes to waste. It could be enough calories to feed each undernourished person on the planet. Minimizing food packaging by skipping single-use, disposable straws, and cups and eating fast food less often is a sustainable strategy.<sup>22</sup>

### **Buy, grow and eat regional foods**

One of the most sustainable things people can do is to grow some food, by planting herbs, edible landscaping, shrubs, and trees in the home and the community. According to the World Watch Institute, food typically travels 1,500 to 2,500 miles from farm to plate. A study in Iowa tracked 2,211 miles for one carton of yogurt (milk, sugar, strawberries) to get to the processing plant. In a Swedish study, a typical Swedish breakfast of apple, bread, butter, cheese, coffee, cream, orange juice, and sugar traveled the circumference of the earth to get to the plate.<sup>23</sup> Eating locally produced foods not only create an economy for local producers but are also suitable for the environment.

### **Conclusion**

To conclude, it is time we recognize the environmental

impacts of the type and amount of food we eat considering the planet and our health. Environmentally sustainable diets should be based on the dietary recommendations of the country. Sustainable diet planning should focus on plant-based food sources of proteins such as beans, pulses/legumes, soya, nuts & seeds, and, whole grains. The consumption of red meat and processed meats should be minimized. Milk and its products need to be consumed in moderation. A variety of fish should be consumed and a policy on fish can be developed particularly for the coastal regions. Overconsumption of discretionary foods high in saturated fat added sugar and salt, and/or alcohol should be avoided to prevent not only obesity and chronic diseases but to consequently decrease carbon footprints. Avoiding consumption of air-freighted, pre-packaged, and prepared fruits and vegetables is another doable strategy. In place of drinking potable water from plastic bottles for hydration opt for tap water. Food waste is recyclable and hence must be used. Therefore, regulating our daily meals with sustainable food choices is a way forward to promote health and save our environment for our future generations.

## References

1. Perrin Selcer, "Anthropocene," Encyclopedia of the History of Science (March 2021), accessed 31 May 2022.
2. Klenert, D., Funke, F., Mattauch, L. *et al.* Five Lessons from COVID-19 for Advancing Climate Change Mitigation. *Environ Resource Econ* 76, 751–778 (2020).
3. IPCC, 2018: Global Warming of 1.5°C. An IPCC Special Report on the impacts of global warming of 1.5°C above pre-industrial levels and related global greenhouse gas emission pathways, in the context of strengthening the global response to the threat of climate change, sustainable development, and efforts to eradicate poverty [Masson-Delmotte, V., P. Zhai, H.-O. Pörtner, D. Roberts, J. Skea, P.R. Shukla, A. Pirani, W. Moufouma-Okia, C. Péan, R. Pidcock, S. Connors, J.B.R. Matthews, Y. Chen, X. Zhou, M.I. Gomis, E. Lonnoy, T. Maycock, M. Tignor, and T. Waterfield (eds.)].
4. M. H. Forouzanfar et al.; GBD 2013 Risk Factors Collaborators, Global, regional, and national comparative risk assessment of 79 behavioural, environmental and occupational, and metabolic risks or clusters of risks in 188 countries, 1990–2013: A systematic analysis for the global burden of disease study 2013. *Lancet* 386, 2287–2323 (2015).
5. International Food Information Council Foundation. 2019 Food & Health Survey.
6. <https://foodinsight.org/wp-content/uploads/2019/05/IFIC-Foundation-2019-Food-And-Health-Report-Final.pdf>.
7. Mejia NV, Reyes RP, Martinez Y, Cerritos R Implications Of The Western Diet For Agricultural Production, Health And Climate Change [Published Online December 20, 2018]. *Front Sustain Food Syst*. Doi: 10.3389/fsufs.2018.00088.
8. Joris P.C. Eekhout, Joris de Vente, Global impact of climate change on soil erosion and potential for adaptation through soil conservation, *Earth-Science Reviews*, Volume 226, 2022, 103921, ISSN 0012-8252, <https://doi.org/10.1016/j.earscirev.2022.103921>.
9. Kim BF, Santo RE Scatterday AP, Et al. Country– Specific Dietary Shifts To Mitigate Climate And Water Crises. *Glob Environ Change*. 2020;62:101926.
10. Saez - Almendros S, Obrador B, Bach-Faig A, Serra- Majem L. Environmental Footprints Of Mediterranean Versus Western Dietary Patients: Beyond The Health Benefits Of The Mediterranean Diet. *Environ Health*

2013;12:118.

11. Marlow HJ, Hayes WK, Soret S, Carter RL, Schwab ER, Sabate J. Diet And The Environment: Does What You Eat Matter? *Am J Clin Nutr.* 2009;89(5): 1699S-1703S.
12. Willett W, Rockström J, Loken B, Springmann M, Lang T, Vermeulen S, Garnett T, Tilman D, DeClerck F, Wood A, Jonell M. Food in the Anthropocene: the EAT–Lancet Commission on healthy diets from sustainable food systems. *The Lancet.* 2019 Feb 2;393(10170):447-92.
13. Sharma, Manika & Kishore, Avinash & Roy, Devesh & Joshi, Kuhu. (2020). A comparison of the Indian diet with the EAT-Lancet reference diet. *BMC public health.* 20. 812. 10.1186/s12889-020-08951-8.
14. Kitada, M., Ogura, Y., Monno, H., Xu, J., and Koya, D. 2021. Effect of methionine restriction on aging: Its relationship to oxidative stress. *Biomedicines*, 9: 130. <https://doi.org/10.3390/biomedicines9020130>. pmid:33572965. Google Scholar
15. Schmidt, J.A., Rinaldi, S., Scalbert, A., Ferrari, P., Achaintre, D., Gunter, M.J., et al. 2016. Plasma concentrations and intakes of amino acids in male meat-eaters, fish-eaters, vegetarians and vegans: a cross-sectional analysis in the EPIC-Oxford cohort. *Eur. J. Clin. Nutr.* 70: 306–312. <https://doi.org/10.1038/ejcn.2015.144>. pmid:26395436. Google Scholar
16. Food And Agriculture Organisation Of The United Nations; The Livestock's Long Shadow: Environmental Issues And Options. <https://www.fao.org/3/A0701E/a0701e.pdf>. Published 2006.
17. Pelletier N, Tyedmers P. Forecasting Potential Global Environmental Costs Of Livestock Production 2000-2050. *Proc Natl Acad Sci U S A.* 2010;107(43):18371-18374.
18. Sabate J, Soret S. Sustainability Of Plant-Based Diets: Back To The Future. *Am J Clin Nutr.* 2014;100 Suppl 1:476S-482S.
19. Oldways Whole Grains Council. Whole Grains: A Sustainable Food. [https://wholegrains-council.org/sites/default/files/atoms/files/WG\\_SustainableFood\\_infographic.pdf](https://wholegrains-council.org/sites/default/files/atoms/files/WG_SustainableFood_infographic.pdf).
20. Milman O, Leavenworth S. China's Plan To Cut Meat Consumption By 50% Cheered By Climate Campaigners. *The Guardian.* June 20, 2016. <https://www.theguardian.com/world/2016/jun/20/chinas-meat-consumption-climate-change>. Accessed October 14, 2020.
21. British Dietetic Association. One Blue Dot: Eating Patterns For Health And Environmental Sustainability: A Reference Guide For Dietitians. <https://www.bda.uk.com/uploads/assets/pdf>.
22. Hendrie, G., Baird, D., Ridoutt, B., Hadjikakou, M. and Noakes, M. (2016) 'Overconsumption of Energy and Excessive Discretionary Food Intake Inflates Dietary Greenhouse Gas Emissions in Australia', *Nutrients*, 8(11), pp. 690
23. Poore J, Nemecek T Reducing Food's Environmental impacts through producers and consumers. *Science.* 2018; 360(6392):987-992.

**Source of Support : Nil**  
**Conflict of Interest : None**