



RESEARCH ARTICLE

REVISED Perceived usefulness of receiving a potential smoking cessation intervention via mobile phones among smokers in Indonesia [version 2; peer review: 2 approved]

Mochammad Fadjar Wibowo ¹, Anil A. Kumar ², Surahyo Sumarsono ¹, Rashmi Rodrigues ²⁻⁴

¹Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

²St John's National Academy of Health Sciences, Bangalore, India

³Karolinska Institutet, Stockholm, Sweden

⁴Wellcome Trust/DBT India Alliance, New Delhi, India

v2 First published: 11 Jun 2019, 4:94
<https://doi.org/10.12688/wellcomeopenres.15135.1>
 Latest published: 19 Aug 2020, 4:94
<https://doi.org/10.12688/wellcomeopenres.15135.2>

Abstract

Background: The use of technology to support healthcare in Indonesia holds new promise in light of decreasing costs of owning mobile devices and ease of access to internet. However, it is necessary to assess end-user perceptions regarding mobile health interventions prior to its implementation. This would throw light on the acceptability of mobile phone communication in bringing about behavioral changes among the target Indonesian population. The aim of this study was to explore the perceived usefulness of receiving a potential smoking cessation intervention via mobile phones.

Methods: This is an exploratory cross-sectional study involving current and former adult tobacco smokers residing in Indonesia. Online advertisement and snowballing were used to recruit respondents. Data was collected using a web-based survey over a period of 4 weeks. Those willing to participate signed an online consent and were subsequently directed to the online questionnaire that obtained demographics, tobacco usage patterns, perceived usefulness of a mobile phone smoking cessation application and its design.

Results: A total of 161 people who smoked tobacco responded to the online survey. The mean age of the participants was 29.4. Of the 123 respondents, 102 were men. Prior experience with using a mobile phone for health communication (OR 3.6, $P=0.014$) and those willing to quit smoking (OR 5.1, $P=0.043$) were likely to perceive a mobile phone smoking cessation intervention as useful. A smartphone application was preferred over text messages, media messages or interactive voice response technology. Content consisting of motivational messages highlighting the methods and benefits of quitting smoking were requested.

Conclusion: People who smoke in Indonesia perceived receiving a

Open Peer Review

Reviewer Status

	Invited Reviewers	
	1	2
version 2		
(revision)	report	report
19 Aug 2020	↑	↑
version 1		
11 Jun 2019	report	report

1. **Mimi Nichter**, University of Arizona, Tucson, USA

2. **Judith McCool** , University of Auckland, Auckland, New Zealand

Any reports and responses or comments on the article can be found at the end of the article.

potential smoking cessation intervention via mobile phones as useful. A multi-component, personalized smartphone application was the desired intervention technique. Such an intervention developed and implemented within a public health program could help address the tobacco epidemic in Indonesia.

Keywords

mHealth, mobile phones, smoking cessation, Indonesia



This article is included in the [Wellcome Trust/DBT India Alliance](#) gateway.

Corresponding author: Mochammad Fadjar Wibowo (wibowo.fadjar@gmail.com)

Author roles: **Wibowo MF:** Conceptualization, Data Curation, Formal Analysis, Investigation, Methodology, Project Administration, Resources, Software, Writing – Original Draft Preparation; **Kumar AA:** Data Curation, Formal Analysis, Software, Validation, Visualization, Writing – Review & Editing; **Sumarsono S:** Conceptualization, Formal Analysis, Methodology, Project Administration, Resources, Supervision; **Rodrigues R:** Conceptualization, Formal Analysis, Methodology, Project Administration, Resources, Supervision, Validation, Writing – Review & Editing

Competing interests: No competing interests were disclosed.

Grant information: This study was supported by the Wellcome Trust through a Wellcome Trust/DBT India Alliance's Intermediate clinical and public health fellowship grant to RR [IA/CPHI/15/1/502042].

The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Copyright: © 2020 Wibowo MF *et al.* This is an open access article distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Wibowo MF, Kumar AA, Sumarsono S and Rodrigues R. **Perceived usefulness of receiving a potential smoking cessation intervention via mobile phones among smokers in Indonesia [version 2; peer review: 2 approved]** Wellcome Open Research 2020, 4:94 <https://doi.org/10.12688/wellcomeopenres.15135.2>

First published: 11 Jun 2019, 4:94 <https://doi.org/10.12688/wellcomeopenres.15135.1>

REVISED Amendments from Version 1

The version 2 includes major changes in the Introduction with Indonesian statistics such as current smartphone usage, cost of nicotine replacement therapies, cost of SMS/MMS, percentage of smokers who actually quit smoking after attempting to quit, and many other minor changes as suggested by peer reviewer reports.

Any further responses from the reviewers can be found at the end of the article

Introduction

With an estimated population of 266 million people in the year 2018, Indonesia is the fourth most populous country in the world¹ and the third highest number of tobacco smokers. Approximately 65% of adult males in Indonesia smoke tobacco². Of those who smoke tobacco, 61% start smoking before the age of 19 years³ with a 19% prevalence of smoking among Indonesian teenagers aged 13–15 years³.

The large revenue from cigarettes despite low taxes, the employment opportunities within the tobacco industry, the weak anti-tobacco legislations and public health campaigns against aggressive marketing, relatively low price and pervasive accessibility of tobacco products are considered to drive the tobacco epidemic in Indonesia^{4–6}. Indonesia is also the only country in the Asia Pacific that has not ratified the World Health Organizations (WHO) Framework Convention on Tobacco Control (FCTC) that addresses the demand and supply of tobacco and related products in order to promote health³.

This weak implementation of international regulations to minimize the marketing, distribution and sales of tobacco products along with prevailing economic, political and social factors have resulted in a high burden of tobacco-related morbidity and mortality in Indonesia⁴. As a consequence, >97 million non-smokers are exposed to second-hand tobacco smoke in the country⁷. In 2010, an estimated 12% of total deaths in Indonesia were the result of tobacco-related disease⁸. Greater than 3.5 million disability-adjusted life years (DALY) were also lost and an estimated 319 million to 1.2 billion USD was spent on healthcare for tobacco-related illnesses in Indonesia annually^{7,8}. Despite this large burden the support available to quit smoking, especially through the healthcare system, is limited⁸. This is evidenced by the fact that a third of the patients with TB attempting to quit relapsed into smoking six months after treatment⁹.

In Indonesia, nicotine replacement therapy such as nicotine patches, gums, and sprays are available without prescription in pharmacies but cost 7000–25000 IDR (0.0035–0.035 USD) per unit. Quitting with prescription medication such as varenicline, hypnotherapy or behavioral therapy costs 0.8–1 million IDR and are not very popular⁸. Further, most (71%) of those who want to quit smoking, attempt to quit without assistance while others may use traditional methods (herbal or medicinal plants), smokeless tobacco or counselling⁸. The costs of quitting

against the costs of cigarettes (12–24000 IDR/pack currently and 32–53000 IDR from 1st November 2019) probably influences both the quit strategy and the decision to continue smoking. The self-confidence of Indonesian physicians in providing smoking cessation counselling is reportedly low, with smoking cessation services offered only at a few healthcare facilities¹⁰ while no national toll-free quit-line is available¹¹.

Further, there are varied perceptions of the effects and complications of tobacco on health in Indonesia. For example; some patients with diabetes mellitus considered that they could smoke relatively lesser cigarettes (3/day) when compared with those who were healthy (12/day)¹² while others it did not know that smoking could complicate their illness¹³. Also, quitting smoking was considered an option only for those seriously ill which could be resumed on recovery¹². This prevailing scenario makes innovation and improvement in smoking cessation interventions in Indonesia a necessity.

The increasing use of mobile phones in resource poor settings and their adoption for healthcare delivery popularly known as mHealth¹⁴, provides an ideal opportunity to deliver smoking cessation interventions in any setting. mHealth supports a wide range of healthcare applications¹⁵ including clinical decision support and healthcare data collection¹⁶. Other mobile phone applications include behavior change interventions for medication adherence support and smoking cessation^{16,17}.

Using mobile phones in smoking cessation programs enables the personalization of smoking cessation support based on the quitter's background, time of the day or the location of the quitter. Short Messaging Service (SMS) Multimedia Messaging Service (MMS) (pictorial or messages or videos), live-voice calls and interactive voice response (IVR) technology that replace a human caller with a computer, provide motivation and counseling to those who want to quit smoking¹⁵. Such messages might use prompts (either text i.e.; SMS, picture i.e. MMS or voice i.e., calls and IVR) to encourage avoiding cigarettes, ashtrays, lighters, and environments where people usually smoke eg.; 'For the next 4 hours, stay away from cigs'. Additionally, messages also help identify the challenges to quitting and plans to overcome them. Prompts (either SMS or voice calls/IVR) to use telephone helplines and nicotine replacement therapy¹⁸, information regarding economic savings from quitting and nutrition are also useful.

In this regard, text messaging was effective for smoking cessation in New Zealand (personalized text messages to provide distraction, advise and support) and the United Kingdom (motivational messages and feedback focusing on their achievements), mobile phone applications, though not tested for efficacy in a randomized control trial (RCT), are known to reach smokers who are not seeking professional help^{18–23}.

A study from United Kingdom (UK) showed that motivational messages encouraged those wanting to quit smoking by focusing on their achievements. They also provided positive feedback, emphasized on the benefits of quitting, consequences of smoking and the process of quitting²⁴.

Personalized text messages were used to provide smoking cessation advice, support, and distraction from smoking in a study from New Zealand. Content covered symptoms expected on quitting, tips to avoid weight gain and improve nutrition, tips to cope with craving; advice to avoid smoking triggers; instructions on breathing exercises to perform instead of smoking and motivational support and distraction²².

Mobile phone penetration and mHealth development in Indonesia

The growth of mobile users in Indonesia is one of the fastest in Asia with a steady increase from 125 per 100 people in 2013 to 173 per 100 people in 2018²⁵. Given the improving internet accessibility and low cost of smartphones²⁶, with prices as low as 40 USD for a phone, smartphone penetration in Indonesia has reached 27% in 2018²⁷ and is predicted to reach 32% by 2022²⁸. The abundant use of mobile phones in Indonesia that parallels the tobacco epidemic in the country makes mobile phones deemed ideal for implementing smoking cessation interventions.

While mHealth is rapidly evolving in high income countries, scientific evaluation of mobile phone use for health care, particularly for smoking cessation interventions in Indonesia is still in inception^{29,30}. However, it is essential to first explore the acceptability and perceived usefulness of receiving a mobile based smoking cessation intervention among Indonesians who smoke prior to developing and testing such an intervention. We therefore, chose to determine the preferred mode of communication, potential content and communication characteristics of mobile phone-based smoking cessation interventions prior to developing such an intervention. To our knowledge this is the first study that has assessed the acceptability of mobile phone applications for smoking cessation in Indonesia.

Methods

This was an exploratory cross-sectional web-based survey conducted in Indonesia between March 23rd to April 21st 2015. As we did not have prior data on acceptability of mobile phone interventions in Indonesia, we did not estimate a sample size for the study.

For the survey, we developed a survey questionnaire and made it available via the internet for respondents to fill (Appendix A and B (Extended data³¹)). The questionnaire was 'face validated' for content and comprehension and was made available in the Indonesian language. The snowballing approach was used to distribute the questionnaire. For this, 25 potential respondents known to the first author were invited to participate in the study. On completing the survey questionnaire these respondents were requested to invite contacts who in turn were requested to invite their contacts and so forth. The respondents could access the questionnaire only on expressing consent to participate in the survey by clicking the AGREE button on the survey web page.

The survey was promoted via a [weblog](#) called [BerhentiMerokok.org](#) meaning "quit smoking". This website was

created to provide respondents information about the study, to enable respondents to refer the questionnaires to other potential participants and to communicate with the researcher. Another [website](#) and a Facebook page "Layanan Online Berhenti Merokok" ("Quit Smoking Online Information"), also promoted the survey (Appendix C (Extended data³¹)). The websites and Facebook page were further promoted via internet based smoking cessation campaigns and health promotion programs using websites, social media accounts and mobile applications using promotional banners (Appendix D (Extended data³¹)).

The questionnaire comprised four sections: (i) Introduction and informed consent (ii) demographic characteristics (iii) smoking status and smoking cessation aid seeking behavior (iv) mobile phone usage, perceived usefulness and preferences regarding mobile phone-based smoking cessation interventions. The survey included questions on the preferred mode of communication for the intervention delivery (SMS, voice calls, multimedia messaging, automated calls and smartphone applications), potential content and communication characteristics of a smoking cessation intervention delivered via mobile phones. The questionnaire was created using [Typeform](#) survey software and was made available online for data collection during the study period (Appendix E (Extended data³¹)).

During the four-weeks of data collection (March 23rd to April 21st 2015), 850 visitors had accessed the web-based survey. Of these, only 161 (19%) completed the survey and were included in the analysis. These respondents were current and former smokers, aged 18 years or older, residing in Indonesia for the past year. On average the participants' took 34:06 ($\pm 2:02$) minutes to complete the questionnaire.

Statistical analysis

Statistical analyses were performed using [SPSS](#) Version 22 for Windows. Complete case analysis was used to analyse the data. The variables were described using measures of central tendency and dispersion. Bivariate analysis (chi-square) was used to explore associations between perceived usefulness of receiving intervention and demographic variables, smoking status and mobile phone usage. Univariate logistic regression analyses were performed if the variable had more than three categories. Variables with p-values less than 0.25 were subsequently included in a multivariate regression model to identify the predictors of perceived usefulness of the intervention.

Ethics statement

Ethical clearance for the study (Ref: KE/FK/311/EC) was obtained from the Medical and Health Research Ethics Committee, Universitas Gadjah Mada, Yogyakarta, Indonesia, a state-owned university to which the researchers are affiliated (Appendix F (Extended data³¹)). Informed consent was obtained online prior to the survey by asking those willing to participate in the survey to click on an "AGREE" button online.

Results

Of the, 161 (19%) respondents who completed the survey, 47 (29%) used smartphones, 30 (19%) personal computers, and

24 (15%) used tablets. Respondents' locations represented 14 of the 34 provinces in Indonesia (see Underlying data³²).

Perceived usefulness of receiving a smoking cessation intervention via mobile phones

Overall, 116 (85%) of the respondents perceived that a potential smoking cessation intervention delivered via mobile phones was useful.

Socio-demographic characteristics. The socio-demography of the respondents is described in Table 1. The mean age of the respondents was 29.4 (± 7.11) years. Of the 123 respondents, 80 (65%) were aged < 30 years, 102 (83%) were men, 75 (61%) reported Indonesian as their primary language and 68/156 (44%) reported being literate in English. Most respondents were unmarried and had completed higher education. There were 96 (88%) respondents from urban areas. The respondents'

mean monthly expenditure was 4.7 million Indonesian Rupiah (IDR) (± 6.4 million) [USD 330 (± 450)]. There was no significant difference in the perceived usefulness of receiving smoking cessation intervention via mobile phones within different socio-demographic groups (Table 1 & Table 2).

Smoking status characteristics. Of the respondents, 111 (75%) were current smokers. Of these, 77 (52%) smoked daily. The mean age at which smoking was initiated was 16.55 (± 5.2) years. The mean duration of smoking was 8.5 (± 7) years while most were at a low or very low nicotine dependency as per the FTND. Participants reported smoking an average 9 (SD8) tobacco sticks/ day (range <1–32 tobacco sticks/ day) amounting to 3 pack years (range: 0.1–41 pack years).

Most current smokers (76, 68%) expressed their willingness to quit smoking and a majority (82, 74%) tried to quit in the

Table 1. Demographic profile of the participants.

Variables		Total	Female (n=22)	Male (n=102)	P value
Age (n=123)	Median (IQR)	27 (25-32) years	26.5 (23.75-30.25) years	27 (25-33.5) years	
	≥ 27 years	70 (57%)	11 (50%)	59 (58%)	
	<27 years	53 (43%)	11 (50%)	42 (41%)	0.470
Marital status (n=121)	Married	50 (41%)	6 (27%)	44 (43%)	
	Single	71 (59%)	16 (73%)	55 (54%)	0.139
Residence (n=123)	Rural	27 (22%)	4 (18%)	23 (23%)	
	Urban	96 (88%)	18 (82%)	78 (76%)	0.637
Education status (n=123)	High school and lower	17 (14%)	4 (18%)	13 (13%)	
	Undergraduate degree	79 (64%)	16 (73%)	63 (62%)	
	Postgraduate degree	27 (22%)	2 (9%)	25 (25%)	0.142
English Literacy (n=156)	No	88 (56%)	6 (27%)	48 (47%)	
	Yes	68 (44%)	16 (73%)	52 (51%)	0.076
Employment status (n=123)	Not gainfully employed	37 (30%)	6 (27%)	31 (30%)	
	Gainfully employed	86 (70%)	16 (73%)	70 (69%)	0.751
Income (in IDR) (n=120)	Median (IQR)	3000000 (2000000-5000000)	5000000 (2000000-7125000)	3000000 (1925000-5000000)	
	≥ 3000000 IDR	65 (54%)	15 (68%)	50 (49%)	
	<3000000 IDR	55 (46%)	7 (32%)	48 (47%)	0.144

IDR: Indonesian Rupiah. 1 USD = 14270 IDR as of August 5th, 2019.

Table 2. Demographic profile and its association with perceived usefulness of receiving smoking cessation intervention via mobile phone (N=122).

Variables		Total n (%)	Perceived as useful n (%)	P-value	Unadjusted OR (95% CI)
Age Median \pm SD (years)		27 \pm 7.11		0.887	0.995 (0.927-1.068)
Sex	Male	101 (82)	86 (85)		Referent
	Female	22 (18)	20 (91)	0.478	1.744 (0.369-8.247)
Marital Status	Married	49 (41)	41(84)		Referent
	Single	71 (59)	62 (87)	0.573	1.344 (0.479-3.768)
Primary Language	Indonesian	75 (61)	65 (87)		Referent
	Regional language	47 (39)	40 (85)	0.809	0.879 (0.310-2.495)
Education Status	High School and Lower	17 (14)	16 (94)		Referent
	Undergraduate Degree	79 (64)	65 (82)	0.279	0.313 (0.038-2.568)
	Postgraduate Degree	27 (22)	24 (89)	0.563	0.500 (0.048-5.242)
English Literacy	Illiterate	66 (50)	55 (85)		Referent
	Literate	67 (50)	58 (87)	0.602	1.289 (0.496-3.350)
Employment Status	Unemployed	36 (30)	33(92)		Referent
	Employed	86 (70)	72(84)	0.248	0.468 (0.126-1.738)
Monthly Expense Median \pm SD (in IDR)		3,000,000 \pm 6,358,018		0.253	1.000 (1.000-1.000)
Residence	Municipality	95 (78)	81 (85)		Referent
	Regency	27 (22)	24 (89)	0.631	1.383 (0.367-5.215)

IDR: Indonesian Rupiah. 1 USD = 14270 IDR as of August 5th, 2019.

past. Willingness to quit smoking was an important factor for perceived usefulness of an intervention. Details regarding smoking cessation methods used, the frequency of the health care provider enquiring about the smoking status, and the frequency of advice received to quit are described in [Table 3](#).

Of those who tried quitting smoking, 69 (91%) attempted quitting without assistance. Respondents who were willing to quit smoking were seven times more likely to perceive receiving a smoking cessation intervention via mobile phones as useful (OR=6.161, p-value=0.004) ([Table 3](#)).

Mobile phone usage patterns. Of the respondents, 154 (98%) used a smartphone and none of them shared their phones with others. Nearly all phone use was meant for personal reasons (153, 95%). Three-quarters (118, 77%) of the respondents reported being well acquainted with using mobile phones. Most respondents (140, 92%) had uninterrupted internet access via data services on their mobile phones.

Of the respondents, 95 (77%) used their mobile phone to communicate with others for health purposes. These respondents frequently communicated with physicians (33, 35%), health care

Table 3. Smoking status characteristics and its association with perceived usefulness of receiving smoking cessation intervention via mobile phone (N=136).

Characteristics		Frequency n (%)	Perceived usefulness n (%)	P- value	Unadjusted OR (95% CI)
Age at first smoke: Mean ± SD (years)		16.55 ± 5.2		0.989	1.001 (0.913-1.097)
Length of active smoking: Median ± SD (years)		7 ± 7.0		0.835	0.993 (0.932-1.059)
Smoking status	Daily smoker	77 (52)	61 (79)		Referent
	Occasional smoker	34 (23)	26 (76)	0.948	0.959 (0.271-3.395)
	Former smoker	38 (26)	29 (76)	0.243	0.535 (0.187-1.528)
Duration of abstinence: Median ± SD (years)		3 ± 6.9		0.605	0.978 (0.900-1.063)
Nicotine Dependence Level (Fagerstrom)	Very Low and Low	38 (59)	33 (89)		Referent
	Medium, High and Very high	27 (41)	23 (85)	0.632	0.697 (0.158-3.076)
Smoking cessation attempt within last 12 months	Yes	76 (76)	69 (91)		Referent
	No	24 (24)	18 (75)	0.045	0.304 (0.091-1.018)
Smoking cessation method used in the past 12 months	Quitting with assistance	7 (9)	5 (71)		Referent
	Quitting without assistance	69 (91)	58 (84)	0.569	1.933 (0.193-19.394)
Source of quit smoking advice	Motivation from others	35 (35)	30 (86)		Referent
	Self-motivation	65 (65)	57 (88)	0.779	1.118 (0.357-3.949)
Frequency being asked regarding smoking status by health provider	Every time	19 (17)	13 (68)		Referent
	Sometimes	45 (41)	38 (84)	0.397	2.019 (0.397-10.272)
	Never	45 (41)	35 (78)	0.625	1.462 (0.319-6.698)
Frequency being advised to quit smoking by health provider	Every time	14 (13)	9 (64)		Referent
	Sometimes	38 (35)	33 (87)	0.676	1.467 (0.243-8.854)
	Never	57 (52)	44 (77)	0.585	1.630 (0.282-9.412)
Willingness to Quit Smoking	No	13 (15)	8 (62)		Referent
	Yes	76 (85)	69 (91)	0.004	6.161 (1.579-24.033)

workers (13, 14%), family (60, 63%) and friends (56, 59%) for health. The content of these communications included request for advice regarding management of illness (51, 32%) and medication side effects (29, 18%), reporting symptoms (46, 29%), scheduling appointments (9, 6%), advising other regarding healthcare (27, 17%), and exchanging information regarding smoking cessation support (22, 14%).

Of the respondents, 85 (89%) who had used a mobile phone for health purposes perceived a potential smoking cessation intervention via mobile phones as useful (OR =3.598, p-value=0.014) (Table 4).

A multivariate logistic regression analysis of perceived usefulness of mobile phone smoking cessation interventions found only willingness to quit smoking as a predictor of perceived usefulness (Table 5).

Features of smoking cessation interventions via mobile phones preferred by respondents

Content and mode of communication. Of the respondents, 86 (62%) preferred a smartphone application as a potential smoking cessation intervention as opposed to 18% who were willing to have SMS or MMS for communication (Figure 1).

As for the content, motivational messages were the preferred content for mobile phone based smoking cessation interventions, followed by reasons for quitting and reminders about the the numbers of cigarettes that they could smoke each day as they approached their quit date (Figure 2).

Characteristics and features of communication. Two communication characteristics relevant for smoking cessation were explored, i.e., interactivity and personalization. Nearly half the respondents (65, 47%) preferred partially interactive communication, 41 (30%) preferred completely interactive communication and the rest (32, 23%) requested a non-interactive one-way communication. Personalization of content to their needs was a necessary feature for 126 (91%) respondents, while 84 (61%) requested interventions delivered at customized times.

Most respondents (55, 40%) wanted to receive smoking cessation communication on demand and throughout the day (37, 46%).

Potential features of the smartphone application for smoking cessation application requested are described in Figure 3. A calculator indicating the amount of money saved was the most popular followed by predicted lung performance and motivational messages.

Discussion

In Indonesia, an LMIC, the burden of tobacco smoking has risen from 59 million in 2000 to 70 million in 2010 and 73.6 million in 2015^{33,34} and continues to rise, increasing the financial burden due to tobacco related illnesses. The situation is compounded by the limited awareness regarding the hazards of smoking along with the minimalistic support available to quit. Currently, 10% of tobacco users quit tobacco annually³⁵. Further, literature from Indonesia shows that 66% of research


Table 4. Mobile phone ownership and usage patterns and its association with perceived usefulness of receiving a potential smoking cessation intervention via mobile phone (N=137).

Characteristics		Frequency n (%)	Perceived usefulness n (%)	P-value	Unadjusted OR (95% CI)
Mobile phone use proficiency*	Average	35 (23)	25 (71)		Referent
	Good	68 (44)	57 (84)	0.441	1.629 (0.471-5.629)
	Excellent	50 (33)	34 (68)	0.650	0.756 (0.226-2.531)
Monthly expense for mobile phone: Median ± SD (IDR)		150,000 ± 248,413		0.741	1.000 (1.000-1.000)
Main use of mobile internet	Browsing	62 (41)	46 (74)		Referent
	Text-messaging	49 (33)	35 (71)	0.459	0.685 (0.251-1.865)
	Accessing email and other applications	39 (24)	34 (87)	0.140	3.326 (0.675-16.392)
Alarm use	No	18 (12)	15 (83)		Referent
	Yes	132 (88)	101 (77)	0.283	0.337 (0.042-2.696)
Prior mobile phone use for health communication	No	29 (23)	21 (72)		Referent
	Yes	95 (77)	85 (89)	0.014	3.598 (1.240-10.441)

IDR: Indonesian Rupiah. 1 USD = 14270 IDR as of August 8th, 2019.

Table 5. Multivariate analysis of predictors of perceived usefulness (N =76).

Characteristics	Perceived usefulness (n)	Unadjusted OR 95% CI	Adjusted OR 95% CI
Employment status	No (33)		Referent
	Yes (72)	0.468 (0.126-1.738)	0.502 (0.080-3.173)
Monthly Expense Mean (std)		1.000 (1.000-1.000)	1.000 (1.000-1.000)
Smoking cessation attempt within last 12 months	Yes (69)		Referent
	No (18)	0.304 (0.091-1.018)	0.377 (0.087-1.641)
Willingness to quit smoking	No (8)		Referent
	Yes (69)	6.161 (1.579-24.033)	5.105 (1.051-24.808)
Prior mobile phone use for health-related communication	No (21)		Referent
	Yes (85)	3.598 (1.240-10.441)	1.799 (0.386-8.391)

**Figure 1. Potential content of the communication (n=140).**

participants in one study unsuccessfully attempted to quit smoking³⁶, while another study reported that 15% its participants had quit smoking³⁷. To address this issue innovative solutions that are acceptable to Indonesians wanting to quit is

essential. Therefore, given the current pervasiveness of mobile phone communication and its affordability we sought to explore the acceptability and design for a mobile phone smoking cessation intervention in Indonesia.




Figure 2. Preferred mode of communication for mobile phone based smoking cessation interventions (n=139). SMS: Short Message Service; MMS: Multimedia Messaging Service; IVR: Interactive Voice Response.




Figure 3. Preference of potential features of smartphone application for smoking cessation intervention (n=160).

Quitting with mobile phones

No universally effective intervention to address the tobacco epidemic exists. While willingness to quit smoking is a necessity, life-altering events also known as ‘teachable moments’ also lead to quitting^{38,39}. Behavior change interventions such as the counselling, self-help materials, physicians brief advice, telephone calls and pharmacotherapy are interventions commonly used in quitting⁴⁰. In addition, the rapid uptake of information technology (IT) has spurred innovative ways to support quitting via mobile phones.

Currently in Indonesia behavior change interventions are uncommon and when available are expensive. So also, is advice from physicians regarding quitting. Our study indicated that nearly half the participants did not receive any advice from their healthcare provider to quit, despite a reported desire to do so.

In this scenario, integrating mobile phones into the behavior learning theory (BLT)⁴¹ provides a theoretical model for mHealth interventions in smoking cessation. Based on BLT, quitting results from combined external antecedents or motivators (mHealth intervention) and internal antecedents (willingness to quit). Positive outcomes i.e., better health, money savings and better quality of life sustain quitting by reinforcing willingness and engagement with the intervention (Figure 4).

Perceived usefulness of a potential smoking cessation intervention via mobile phones

Studies globally have found smoking cessation intervention via mobile phones^{42–45} are feasible and acceptable to young people across different socio-economic groups. In our study, such interventions were more likely to be perceived as useful by respondents willing to quit smoking. An earlier study found smartphone applications were more frequently used by respondents who were willing to quit within 30 days¹⁹.

Though we did not find additional evidence, our study showed that respondents who had used a mobile phone for health-related communication perceived a smoking cessation intervention via mobile phones as useful. This was probably due to their experience and comfort with interventions delivered via mobile phones.

Features of smoking cessation interventions via mobile phones preferred by respondents

Mode of communication. Our study suggests that a smartphone application is the most preferred mode of communication for a potential smoking cessation intervention. A few respondents chose SMS, MMS, IVR or a combination of the three as the mode of communication. The larger percentage of respondents having access to the internet may explain this result. Given the improving internet accessibility and smartphone subscription in Indonesia²⁵, smartphone applications might be the most suitable mode of intervention for smoking cessation.

Further, literature showed that some of the widely used modes for delivery of health interventions via mobile phones were SMS and tele-counselling while MMS was not as widely used and tested as the other modes of communication¹⁶. SMS interventions are found effective in various behavioral change interventions such as diabetes self-management, weight loss management, physical activity, smoking cessation and medication adherence for antiretroviral therapy⁴⁶. SMS or text-message-based smoking cessation intervention is the only mobile phone-based intervention that was effective in randomized trials. Studies in the United Kingdom and New Zealand reported that text-message-based smoking cessation interventions are affordable, can be personalized, are age appropriate, and not location dependent^{18,22}. In Indonesia, no costs are incurred to receive a text message, while it costs IDR 360 IDR (3 cents) to send an individual message and when deployed in bulk, messaging would cost up to IDR 65,000 (USD 4.5) for




Figure 4. Behavior learning theory.

500 messages a month to be incurred by the intervention provider.

A study in New Zealand showed that an MMS-based smoking cessation intervention using video messages was effective. The results however were equivocal when a complex video messaging intervention was compared with simple general health videos that communicated general health messages. Video messaging was not considered economical in all socioeconomic groups, even in resource rich settings such as New Zealand⁴⁴. Another randomized trial that used multiple-component personalized counselling via telephone in high school students in the US showed an increase in abstinence rates⁴⁷. To deal with the problem of tobacco epidemics, many high-income countries have also established several tele-counselling interventions such as the “Quit Line” and incorporated it with the national health service. Such a quitline is not available in Indonesia, despite the large tobacco epidemic.

Smartphone applications are a promising medium to reach smokers across multiple nations. They have the potential to consolidate the advantages of smoking cessation interventions designed for use with or without the internet (i.e., computer based). Users can continue to access motivational features such as calculators for money saved per cigarette not smoked or information downloaded and saved within the applications from the internet. Mobile applications can be designed successfully to harness mobile phone features such as video, audio, interactive media and texting to promote engagement and constant motivation to quit smoking to the users¹⁹. Given the preference for smartphone applications for smoking cessation in our study, an application that uses pre-recorded audio and video based motivational messages could be useful. However, interventions designed should be contextual as the preference for smartphone applications over other forms mobile interventions, such as text messages, might vary globally⁴⁸.

Potential content. Our study found that motivational messages such as the benefits of quitting smoking and reminders about the users’ reason to quit smoking were preferred content for mobile phone-based smoking interventions in Indonesia. Motivation is the core of any smoking cessation intervention along with addressing barriers and benefits of quitting. The interventions also focus on providing cues to action and promoting self-efficacy and harness the theories of behavior change.

A study from the United Kingdom (UK) showed that motivational messages encouraged those wanting to quit smoking by focusing on their achievements^{18,24}. Personalized text messages were used to provide smoking cessation advice, support, and distraction from smoking in a study from New Zealand. Content had also covered symptoms expected on quitting, tips to avoid weight gain and improve nutrition, coping with craving; advice to avoid smoking triggers; instructions on breathing exercises to perform instead of smoking and motivational support and distraction²².

Communication characteristics. We explored two important characteristics namely, interactive communication and

personalized communication. Most respondents preferred to interact with a human facilitator and wanted personalized communications. They preferred receiving messages on demand or even throughout the day without a predetermined frequency. Though other studies have not discussed the timing of communication delivery, most interventions involved predetermined daily communication.

Although communication in the UK and New Zealand studies was an automated SMS, both interventions allowed participants to contact a quit line and speak to a counsellor at any time. Additionally, the intervention in New Zealand allowed the respondents to send free messages to friends and family in order to obtain support²². Earlier studies about the social network structure of large online communities for smoking cessation have shown a relationship between social network support for quitting and maintenance of abstinence⁴⁹. Higher levels of connectivity and positive social support are known associates of a greater quit rate and lower rate of relapse⁵⁰. Therefore, mobile-phone quit interventions should incorporate an interactive component to enable quitting in their design.

Both the UK and New Zealand interventions combined interactive and one-way communication. Although most text messages sent to the participants were push messages, the UK- study provided a “CRAVE” and “LAPSE” feature, where the participants could ask for additional messages¹⁸. Similarly, the New Zealand-based study provided a “txt crave” feature where participants could ask for additional messages during their moments of craving and the “txt quiz” feature where the participants could ask questions²².

Several studies have explored the effectiveness of personalised interventions for smoking cessation^{51–55}. Improved engagement and retention through mobile-based smoking cessation interventions in adolescents has been observed^{42,55,56}. Some studies used personalized messages^{18,42,44,54,57}. Participants sex, age, smoking history, goals, medical condition^{16,42,52} cultural and ethnic background^{53,58} are some factors used in personalising messages. The profound ethnic diversity of the Indonesian population, if considered, might increase the complexity of the intervention and costs for development⁵⁹.

Features currently available in mobile applications for smoking cessation

Smoking cessation applications are pervasive, some with exaggerated claims of effectiveness. Despite the large number of smartphone applications for smoking cessation^{20,60}, only a few are evidence-based²⁰ and are insufficient to stimulate self-motivation⁶⁰ to help quit smoking.

In 2012, an American-based survey analyzed 98 of the most popular smartphone applications for smoking cessation (available in English) downloaded via the iPhone and Android market. Popular applications had low levels of adherence to the U.S. Guidelines for Treating Tobacco Use and Dependence (GTTUD), with an average score of 12.9 of a possible 42 on the Adherence Index²⁰.

While the applications incorporated features such as instructiveness, user personalized advice to quit and assessment of current tobacco use, motivation through rewards, and quit plan assistance were missing. Additionally, advice for referral and follow-up were also missing⁶¹.

The list of potential smartphone features, in various combinations, for mobile phone interventions is exhaustive^{19,20,60}. One such feature is the interactive self-monitoring system that allows users to add their health data via questionnaires, texts, and audio or video recordings²⁰. These applications process, organize and graph this data to help users understand their progress. The data can help the users at every step in their quitting process, providing text information about quitting, showing the number of days users have been nicotine-free, providing logs to administer users' quit attempts and craving triggers along with sending them motivational messages and reminders¹⁹.

Some of the least explored features of mobile smoking cessation applications such as a calculator for predicting money saved from quitting and unsmoked cigarettes along with predicted lung function were features popular in our study. However, this may also be due to the structure of the questionnaire and the nuance of smartphone-based quitting applications in Indonesia.

A conceptual framework for designing mobile phone smoking cessation interventions

Based on the results we modified the conceptual framework for mHealth interventions by Rodrigues R (2014)⁶² to inform mHealth intervention design for smoking cessation (Figure 5). Such interventions should consider frequency, timing, personalization (tailoring) engagement and components (features, single or multiple) in their design. For example; an mHealth smoking cessation intervention could provide timed motivational

messages, distractions from craving, reinforcements such as graphic visualizations of money saved based on interactive data input from users. Further, the Cognitive-Affective Personality System (CAPS) model provides a possible mechanism to incorporate the intervention for behavior change^{63,64}. CAPS is a complex network of an individual's goals, beliefs, thoughts, feelings, self-regulatory standards, plans and competencies. An individual's thoughts and feelings are constantly changing. External stimuli through mobile phones (messages and prompts) can trigger these changes thereby influencing self-regulatory behavior.

Limitations

As this was a web-based study only those who were familiar with the internet were captured minimizing its generalizability to those familiar with information technology (IT). Nevertheless, as the proposed intervention is IT based, it captured the opinion of the beneficiaries that the intervention is likely to target. Also, as not all who accessed the questionnaire completed it, the numbers that were included in the analysis were low. However, despite the study's limited sample size and duration, information relevant to inform the design and piloting the mobile application was obtained. Item non-response and incomplete responses are known to affect the generalizability of the results of online surveys^{65,66}. As the reported levels of tobacco dependency were low, it is likely that the respondents were those who either had greater control over their smoking behavior or were more amenable to the idea of quitting. However, the level of nicotine dependence obtained may be questionable given the normalization of smoking in Indonesia^{5,67}. Further, social desirability bias also cannot be ruled out in the FTND as it is a self-report of dependence by the participant.

Given that quitting cold turkey is a popular method of smoking cessation we understand that our approach to smoking cessation reflects a dominance of interventionism. The proposed




Figure 5. Conceptual model for mHealth interventions in smoking cessation. This figure has been reproduced with permission from Rodrigues R (2014)⁶².

intervention should not be considered as a ‘one size fits all’ but rather one in a basket of solutions including behavioral therapy, pharmacotherapy or quitting cold turkey.

Conclusions

Our study showed that the Indonesian respondents to our survey perceived a potential smoking cessation intervention via mobile phones as useful. Perceived usefulness was associated with smokers’ willingness to quit smoking and their prior use of mobile phones for health-related communication. A multicomponent smartphone application personalized to time, frequency and content was desired. Such an application, if implemented, could be one in a basket of smoking cessation solutions offered within an organized program quit smoking programmes at schools, healthcare facilities and counseling centers could go a long way in addressing the tobacco epidemic in Indonesia.

Ethics and consent

Ethical approval was received from the Ethics Committee of Universitas Gadjah Mada, Indonesia (Ref: KE/FK/311/EC). Participants were fully informed of the study and consent was obtained prior to data collection.

Data availability

Underlying data

Harvard Dataverse: Perceived Usefulness of Receiving a Potential Smoking Cessation Intervention via Mobile Phones among Smokers in Indonesia. <https://doi.org/10.7910/DVN/N3QQE1>³²

This project contains the following underlying data:

- Main SPSS file.tab (SPSS file with underlying data)
- Raw data on MS Excel with codes and keys.xlsx (underlying data in Excel format)
- [Table 2 SPSS outputs.spv](#) (Data underlying Table 2)
- [Table 3 SPSS outputs.spv](#) (Data underlying Table 3)

- [Table 4 SPSS outputs.spv](#) (Data underlying Table 4)
- [Table 5 SPSS outputs.spv](#) (Data underlying Table 5)

Extended data

Harvard Dataverse: Perceived Usefulness of Receiving a Potential Smoking Cessation Intervention via Mobile Phones among Smokers in Indonesia. <https://doi.org/10.7910/DVN/EU6DZS>³¹

This project contains the following extended data:

- Appendix A - Questionnaire in English.pdf (Study questionnaire - English)
- Appendix B - Questionnaire in Indonesian Language - Kuisisioner Eksplorasi Penerimaan Penggunaan Ponsel untuk Dukungan Berhenti Merokok.docx (Study questionnaire - Indonesian)
- Appendix C - Promotion of survey on Facebook.png (Survey promotion via Facebook)
- Appendix D - Promotion of survey on Doctor Gratis mobile application, guetau.com and Twitter.png (Survey promotion via Doctor Gratis mobile application, guetau.com and Twitter)
- Appendix E - Survey on Typeform website.png (Image of Survey on Typeform)
- Appendix F - IEC and ethics clearance.pdf (Study consent form and approval document)

Data are available under the terms of the [Creative Commons Zero “No rights reserved” data waiver](#) (CC0 1.0 Public domain dedication).

Acknowledgements

The authors would like to thank all the participants in the study for their valuable time.

References

1. United Nations, Department of Economic and Social Affairs, Population Division: **World Population Prospects: The 2017 Revision, Volume II: Demographic Profiles**. ST/ESA/SER.A/400. 2017. [Reference Source](#)
2. World Health Organization: **Report on the global tobacco epidemic, 2017: monitoring tobacco use and prevention policies**. 2017. [Reference Source](#)
3. World Health Organization, Regional Office for South-East Asia: **Global Youth Tobacco Survey (GYTS): Indonesia Report, 2014**. New Delhi, India; 2015. [Reference Source](#)
4. Tandilittin H: **Civil Society and Tobacco Control in Indonesia: The Last Resort**. *Open Ethics J.* 2013; **7**: 11–18. [Publisher Full Text](#)
5. Nichter M, Padmawati S, Danardono M, et al.: **Reading culture from tobacco advertisements in Indonesia**. *Tob Control.* 2009; **18**(2): 98–107. [PubMed Abstract](#) | [Publisher Full Text](#)
6. Reynolds C: **Tobacco advertising in Indonesia: “the defining characteristics for success”**. *Tob Control.* 1999; **8**(1): 85–88. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
7. Barber S, Ahsan A, Adioetomo S, et al.: **Tobacco Economics in Indonesia**. 2008. [Reference Source](#)
8. Samlee Plianbanchang, Ministry of Health R of I, CDC Foundation, et al.: **Global Adult Tobacco Survey (GATS): Indonesia Report, 2011**. New Delhi: World Health Organization, Regional Office for South East Asia; 2012. [Publisher Full Text](#)
9. Ng N, Padmawati RS, Prabandari YS, et al.: **Smoking behavior among former tuberculosis patients in Indonesia: intervention is needed**. *Int J Tuberc Lung Dis.* 2008; **12**(5): 567–572. [PubMed Abstract](#)
10. Ng N, Prabandari YS, Padmawati RS, et al.: **Physician assessment of patient smoking in Indonesia: a public health priority**. *Tob Control.* 2007; **16**(3): 190–196. [PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
11. World Health Organization: **Report on the global tobacco epidemic, 2013**.

- Geneva, Switzerland.
[Reference Source](#)
12. Padmawati RS, Ng N, Prabandari YS, *et al.*: **Smoking among diabetes patients in Yogyakarta, Indonesia: cessation efforts are urgently needed.** *Trop Med Int Health.* 2009; **14**(4): 412–419.
[PubMed Abstract](#) | [Publisher Full Text](#)
 13. Ng N, Nichter M, Padmawati RS, *et al.*: **Bringing smoking cessation to diabetes clinics in Indonesia.** *Chronic Illn.* 2010; **6**(2): 125–135.
[PubMed Abstract](#) | [Publisher Full Text](#)
 14. WHO Global Observatory for eHealth, World Health Organization: **MHealth: new horizons for health through mobile technologies.** Geneva: World Health Organization; 2011; [cited 2019 Jan 4].
[Reference Source](#)
 15. Blaya JA, Fraser HS, Holt B: **E-health technologies show promise in developing countries.** *Health Aff (Millwood).* 2010; **29**(2): 244–251.
[PubMed Abstract](#) | [Publisher Full Text](#)
 16. Ghorai K, Akter S, Khatun F, *et al.*: **mHealth for Smoking Cessation Programs: A Systematic Review.** *J Pers Med.* 2014; **4**(3): 412–423.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 17. Cole-Lewis H, Kershaw T: **Text messaging as a tool for behavior change in disease prevention and management.** *Epidemiol Rev.* 2010; **32**(1): 56–69.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 18. Free C, Knight R, Robertson S, *et al.*: **Smoking cessation support delivered via mobile phone text messaging (txt2stop): a single-blind, randomised trial.** *Lancet.* 2011; **378**(9785): 49–55.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 19. BinDhim NF, McGeechan K, Trevena L: **Who Uses Smoking Cessation Apps? A Feasibility Study Across Three Countries via Smartphones.** *JMIR MHealth UHealth.* 2014; **2**(1): e4.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 20. Abrams LC, Lee Westmaas J, Bontemps-Jones J, *et al.*: **A content analysis of popular smartphone apps for smoking cessation.** *Am J Prev Med.* 2013; **45**(6): 732–736.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 21. Kong G, Ells DM, Camenga DR, *et al.*: **Text messaging-based smoking cessation intervention: a narrative review.** *Addict Behav.* 2014; **39**(5): 907–917.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 22. Rodgers A, Corbett T, Bramley D, *et al.*: **Do u smoke after txt? Results of a randomised trial of smoking cessation using mobile phone text messaging.** *Tob Control.* 2005; **14**(4): 255–261.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 23. Whittaker R, McRobbie H, Bullen C, *et al.*: **Mobile phone-based interventions for smoking cessation.** *Cochrane Database Syst Rev.* 2012; **11**: CD006611.
[PubMed Abstract](#) | [Publisher Full Text](#)
 24. Free C, Whittaker R, Knight R, *et al.*: **Txt2stop: a pilot randomised controlled trial of mobile phone-based smoking cessation support.** *Tob Control.* 2009; **18**(2): 88–91.
[PubMed Abstract](#) | [Publisher Full Text](#)
 25. International Telecommunication Union: **World Telecommunication/ICT Indicators database.** 2018.
[Reference Source](#)
 26. Global Business Guide: **An Overview of Indonesia's Telecommunication Sector.** Indonesia; 2012.
[Reference Source](#)
 27. Newzoo: **Global Mobile Market Report: Top 50 Countries/Markets by Smartphone Users and Penetration.** Amsterdam, The Netherlands; 2018.
[Reference Source](#)
 28. Statista: **The Statistics Portal. Smartphone penetration rate as share of the population in Indonesia from 2015 to 2022.** 2018.
[Reference Source](#)
 29. Chib A: **The Aceh Besar midwives with mobile phones project: Design and evaluation perspectives using the information and communication technologies for healthcare development model.** *J Comput Mediat Commun.* 2010; **15**(3): 500–525.
[Publisher Full Text](#)
 30. Aman YU, Lopulalan D, Budiman A, *et al.*: **2H2 Center: Praktik Cerdas dari Flores Timur.** Flores Timur: Indonesia; 2013.
 31. Wibowo MF, Kumar AA, Sumarsono S, *et al.*: **Perceived Usefulness of Receiving a Potential Smoking Cessation Intervention via Mobile Phones among Smokers in Indonesia.** Harvard Dataverse, V1. 2019.
<http://www.doi.org/10.7910/DV/N3Q6DZ5>
 32. Wibowo MF, Kumar AA, Sumarsono S, *et al.*: **Perceived Usefulness of Receiving a Potential Smoking Cessation Intervention via Mobile Phones among Smokers in Indonesia.** Harvard Dataverse. 2019.
<http://www.doi.org/10.7910/DV/N3Q6E1>
 33. Commar A, Prasad VK, Tursan d'Espaignet E, *et al.*: **WHO global report on trends in prevalence of tobacco smoking 2000–2025.** 2018.
[Reference Source](#)
 34. Zheng R, Marquez PV, Ahsan A, *et al.*: **Cigarette affordability in Indonesia: 2002–2017 (English).** WBG Global Tobacco Control Program Washington, D.C.: World Bank Group. 2018.
[Publisher Full Text](#)
 35. World Health Organization: **Regional Office for South-East Asia.** WNTD: Indonesia factsheet. World Health Organization. Regional Office for South-East Asia. 2018.
[Reference Source](#)
 36. Fawzani N, Triratnawati A: **Terapi Berhenti Merokok (Studi Kasus 3 Perokok Berat).** In *Jurnal Makara Kesehatan.* 2005; **9**(1): 15–22.
[Reference Source](#)
 37. Vebby Amellia Edwin and Besral: **Association of External Factors with Successful Smoking Cessation among Indonesian Adult** in The 2nd International Meeting of Public Health 2016 with theme “Public Health Perspective of Sustainable Development Goals: The Challenges and Opportunities in Asia-Pacific Region.” *KnE Life Sciences.* 2018; 27–36.
[Reference Source](#)
 38. Cohen DJ, Clark EC, Lawson PJ, *et al.*: **Identifying teachable moments for health behavior counseling in primary care.** *Patient Educ Couns.* 2011; **85**(2): e8–15.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 39. Black JH 3rd: **Evidence base and strategies for successful smoking cessation.** *J Vasc Surg.* 2010; **51**(6): 1529–1537.
[PubMed Abstract](#) | [Publisher Full Text](#)
 40. Samet JM, Yoon SY, World Health Organization, editors.: **Gender, women and the tobacco epidemic: Quitting smoking and beating nicotine addiction.** Geneva: World Health Organization; 2010.
[Reference Source](#)
 41. Munro S, Lewin S, Swart T, *et al.*: **A review of health behaviour theories: how useful are these for developing interventions to promote long-term medication adherence for TB and HIV/AIDS?** *BMC Public Health.* 2007; **7**: 104.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 42. Riley W, Obermayer J, Jean-Mary J: **Internet and mobile phone text messaging intervention for college smokers.** *J Am Coll Health.* 2008; **57**(2): 245–248.
[PubMed Abstract](#) | [Publisher Full Text](#)
 43. Whittaker R, Maddison R, McRobbie H, *et al.*: **A multimedia mobile phone-based youth smoking cessation intervention: findings from content development and piloting studies.** *J Med Internet Res.* 2008; **10**(5): e49.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 44. Whittaker R, Dorey E, Bramley D, *et al.*: **A theory-based video messaging mobile phone intervention for smoking cessation: randomized controlled trial.** *J Med Internet Res.* 2011; **13**(1): e10.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 45. Whittaker R, Merry S, Dorey E, *et al.*: **A development and evaluation process for mHealth interventions: examples from New Zealand.** *J Health Commun.* 2012; **17** Suppl 1: 11–21.
[PubMed Abstract](#) | [Publisher Full Text](#)
 46. Hall AK, Cole-Lewis H, Bernhardt JM: **Mobile text messaging for health: a systematic review of reviews.** *Annu Rev Public Health.* 2015; **36**: 393–415.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 47. Borland R, Balmford J, Benda P: **Population-level effects of automated smoking cessation help programs: a randomized controlled trial.** *Addiction.* 2013; **108**(3): 618–628.
[PubMed Abstract](#) | [Publisher Full Text](#)
 48. Ni Mhurchu PC, Te Morenga L, Tupai-Firestone R, *et al.*: **A co-designed mHealth programme to support healthy lifestyles in Māori and Pasifika peoples in New Zealand (OL@-OR@): a cluster-randomised controlled trial.** *Lancet Digit Health.* 2019; **1**(6): E298–E307.
[Publisher Full Text](#)
 49. Cobb NK, Graham AL, Abrams DB: **Social network structure of a large online community for smoking cessation.** *Am J Public Health.* 2010; **100**(7): 1282–1289.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 50. Muñoz RF, Barrera AZ, Delucchi K, *et al.*: **International Spanish/English Internet smoking cessation trial yields 20% abstinence rates at 1 year.** *Nicotine Tob Res.* 2009; **11**(9): 1025–1034.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 51. Sutton S, Smith S, Jamison J, *et al.*: **Study protocol for iQuit in Practice: a randomised controlled trial to assess the feasibility, acceptability and effectiveness of tailored web- and text-based facilitation of smoking cessation in primary care.** *BMC Public Health.* 2013; **13**: 324.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 52. Naughton F, Jamison J, Sutton S: **Attitudes towards SMS text message smoking cessation support: a qualitative study of pregnant smokers.** *Health Educ Res.* 2013; **28**(5): 911–922.
[PubMed Abstract](#) | [Publisher Full Text](#)
 53. Bramley D, Riddell T, Whittaker R, *et al.*: **Smoking cessation using mobile phone text messaging is as effective in Maori as non-Maori.** *N Z Med J.* 2005; **118**(1216): U1494.
[PubMed Abstract](#)
 54. Abrams LC, Ahuja M, Kodl Y, *et al.*: **Text2Quit: results from a pilot test of a personalized, interactive mobile health smoking cessation program.** *J Health Commun.* 2012; **17** Suppl 1: 44–53.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
 55. Peterson AV Jr, Kealey KA, Mann SL, *et al.*: **Group-randomized trial of a proactive, personalized telephone counseling intervention for adolescent smoking cessation.** *J Natl Cancer Inst.* 2009; **101**(20): 1378–1392.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

56. Obermayer JL, Riley WT, Asif O, *et al.*: **College smoking-cessation using cell phone text messaging.** *J Am Coll Health.* 2004; **53**(2): 71–78.
[PubMed Abstract](#) | [Publisher Full Text](#)
57. Brendryen H, Kraft P: **Happy ending: a randomized controlled trial of a digital multi-media smoking cessation intervention.** *Addiction.* 2008; **103**(3): 478–484; discussion 485–486.
[PubMed Abstract](#) | [Publisher Full Text](#)
58. Gould GS, McEwen A, Watters T, *et al.*: **Should anti-tobacco media messages be culturally targeted for Indigenous populations? A systematic review and narrative synthesis.** *Tob Control.* 2013; **22**(4): e7.
[PubMed Abstract](#) | [Publisher Full Text](#)
59. Arifin EN, Ananta A, Wilujeng Wahyu Utami DR, *et al.*: **Quantifying Indonesia's Ethnic Diversity: Statistics at National, Provincial, and District levels.** *Asian Popul Stud.* 2015; **11**(3): 233–256.
[Publisher Full Text](#)
60. Choi J, Noh GY, Park DJ: **Smoking cessation apps for smartphones: content analysis with the self-determination theory.** *J Med Internet Res.* 2014; **16**(2): e44.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
61. Abroms LC, Padmanabhan N, Thaweethai L, *et al.*: **iPhone apps for smoking cessation: a content analysis.** *Am J Prev Med.* 2011; **40**(3): 279–285.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)
62. Rodrigues RJ: **M-Health for antiretroviral treatment support: evidence from India.** [Stockholm]: Karolinska Institutet; 2014.
[Reference Source](#)
63. Mischel W, Ayduk O: **Self-Regulation in a Cognitive–Affective Personality System: Attentional Control in the Service of the Self.** *Self Identity.* 2002; **1**(2): 113–120.
[Publisher Full Text](#)
64. Kraft P, Drozd F, Olsen E: **Digital Therapy: Addressing Willpower as Part of the Cognitive-Affective Processing System in the Service of Habit Change.** In: Oinas-Kukkonen H, Hasle P, Harjumaa M, *et al.*, editors. *Persuas. Technol.* Berlin, Heidelberg: Springer Berlin Heidelberg; 2008; 177–188, [cited 2019 Apr 8].
[Publisher Full Text](#)
65. Denscombe M: **Item non-response rates: a comparison of online and paper questionnaires.** *Int J Soc Res Methodol.* 2009; **12**(4): 281–291.
[Publisher Full Text](#)
66. Bosniak M, Tuten TL: **Classifying Response Behaviors in Web-based Surveys.** *J Comput Mediat Commun.* 2001; **6**(3): JCMC636.
[Publisher Full Text](#)
67. Pérez-Ríos M, Santiago-Pérez MI, Alonso B, *et al.*: **Fagerstrom test for nicotine dependence vs heavy smoking index in a general population survey.** *BMC Public Health.* 2009; **9**: 493.
[PubMed Abstract](#) | [Publisher Full Text](#) | [Free Full Text](#)

Open Peer Review

Current Peer Review Status:  

Version 2

Reviewer Report 01 October 2020

<https://doi.org/10.21956/wellcomeopenres.16815.r40022>

© 2020 Nichter M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Mimi Nichter

School of Anthropology, University of Arizona, Tucson, AZ, USA

The authors have addressed the issues that were highlighted in the review.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: I am a medical anthropologist who has conducted extensive research on the development of tobacco cessation programs in Java, Indonesia.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Reviewer Report 21 September 2020

<https://doi.org/10.21956/wellcomeopenres.16815.r40023>

© 2020 McCool J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Judith McCool 

Department of Epidemiology and Biostatistics, School of Population Health, University of Auckland, Auckland, New Zealand

This revised version of the paper is considerably improved. All issues raised were addressed. This paper will add to the growing literature on mCessation; this one in respect to perceived usefulness of receiving text message support to quit smoking. In the interim since submission, there are likely to be several additional articles that would add value to this work, or to future interventions planned by the authors. Noting here that end-user perceptions of value is important but there is

an additional issue of sustainability of any mobile based or digital interventions, this is often outside the influence of the end-user and relies on the funder and implementers to commit to long-term delivery and evaluations.

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Global health, tobacco control, mass media communications, social media, behaviour change, Pacific region.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard.

Version 1

Reviewer Report 18 November 2019

<https://doi.org/10.21956/wellcomeopenres.16514.r36737>

© 2019 McCool J. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Judith McCool 

Department of Epidemiology and Biostatistics, School of Population Health, University of Auckland, Auckland, New Zealand

This paper adds practical, current information on what smokers in Indonesia want in terms of support to quit smoking. Tobacco use in Indonesia is a major global health challenge due to the scale of impact. It is the scale of the issue that lends itself to questioning whether mobile devices may boost the reach and accessibility for effective, evidence based smoking cessation support. In a context where most smokers quit without additional aids, it is important to question what mobile or other smoking cessation support is offered and what costs and benefits this adds over and above what is currently in place. This paper examines the perceived usefulness of mobile smoking cessation (mCessation) support for smokers. It is a credit to the research team that they have paused to ask smokers the question "how do you want your cessation support delivered" rather than going ahead with an costly mCessation pilot. Having said that, it is often stated, that "people do not know, what they don't know", in other words hypothetical interventions may not necessarily provide an accurate gauge of potential use or effectiveness of mobile interventions. However, this work also provides an insight into how mobile phones may be used to support smokers, in a low-middle income country. This information is likely to add value to other areas of work where mobile has been proven effective. It is also important to share information about the cultural variations in preferences for, and use of, mobile technologies. There is richer granularity in ITC use within countries than what is described in global mobile agency reports. Hence, this work is timely and will add value to others working in the field of mobile and digital health, as well as tobacco control.

General comments:

Introduction: Page 3. The statement regarding the high prevalence of tobacco use in Indonesia was attributed to the size of the population *and* the lack of public health campaigns addressing cessation. Might it also have something to do with tobacco industry interference, low taxation of tobacco and public awareness messages, smokefree environments, banning advertising and promotions? This statement gives the impression that the current rates are attributable to the size of the population and cessation information, exclusively.

Introduction Page 3. The WHO FCTC covers more than prevention of update of tobacco by adolescents and protection from second-hand cigarette smoke. It also covers cessation support and other measures that have evidence of impact. The statement on page X could be amended to reflect the broader intention of the WHO FCTC. The statement at the start of the following paragraph captures the challenges, but again does not make any reference to the weak regulation of the tobacco industry and marketing, distribution and sales of tobacco products. I note though that this paper is focused on the potential to increase cessation, but this is confounded by a supportive smoking environment.

Introduction, Page 3, paragraph 6. It was noted that the text messages were found to be effective (to support quit attempts) in New Zealand and the United Kingdom, smartphone android applications.... Is it the case that apps were trialled only on android phones? My interpretation is that they were similarly ineffective whether delivered on iPhone or android phones (Abroms *et al.*, 2013¹).

Methods. It is possible that having the survey promoted via a weblog that is for smoking cessation may have reduced the potential sample size given the study was about what people think they want for support for quitting, not about if they are trying to quit. This may, in addition to other factors, have reduced the final sample size.

Ethics statement, page 4. The ethics statement is included here and again at the end of the paper (on page 12), suggest presenting it once only.

Results: Smoking status characteristics; it was interesting to note that there were no differences in preference for receiving information via mobile device between socio-demographic groups. I also note that there was a low or very low level of dependency reported, despite the length of time that smoking was reported. Was this expected and is it explained in the discussion?

It was noted that most smokers in Indonesia quit without behavioural or NRT support. What proportion of those who attempt to quit, succeed. Is there any measure of these changes over time in terms of quitting? What were the methods that were used to support smokers to quit cold turkey? Is there any research on this that can shed light on the types of content that might be salient if delivered via mobile phone?

The finding that suggests a preference for smartphone apps over text message is unique and is inconsistent with other studies which have found little support for apps. I refer to a co-designed intervention using an app for behaviour change in New Zealand with indigenous and Pacific populations (Ni Mhurchu *et al.*, 2019²) among others.

Page 10. The statement that start "Only a few studies have explored the effectiveness of smartphone applications as a behaviour change intervention" is an important point but may need to be updated. There is an increasing number of publications coming through that question the value of apps in health and behaviour change.

Conclusion. Suggest revise the statement that a 'potential smoking cessation intervention...' was perceived as useful.

The final statement regarding the implementation of the app within an organised programme (via schools etc) was not a conclusion that came through strongly in the data. It might it be that any intervention, whether it is mobile device delivered or not, needs to complement a comprehensive tobacco control programme, backed by government support. Politics had not been mentioned in any detail until this point and seems out of place or underdeveloped at this stage.

Table 1. What does 'gainfully employed" mean - I am assuming this mean paid or salaried work?

Income level in IDR - could there be a proxy figure in US or Euro? It was noted earlier that the DALY for tobacco related illness was reported in USD.

Table 4. Column 1 - what does the *refer to on Mobile phone use proficiency*.

Figure 4. This figure does not substantively add to the paper. A brief description of the theory of behaviour change, and a reference would suffice.

References

1. Abroms LC, Lee Westmaas J, Bontemps-Jones J, Ramani R, et al.: A content analysis of popular smartphone apps for smoking cessation. *Am J Prev Med.* 2013; **45** (6): 732-6 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Ni Mhurchu C, Te Morenga L, Tupai-Firestone R, Grey J, et al.: A co-designed mHealth programme to support healthy lifestyles in Māori and Pasifika peoples in New Zealand (OL@-OR@): a cluster-randomised controlled trial. *The Lancet Digital Health.* 2019; **1** (6): e298-e307 [Publisher Full Text](#)

Is the work clearly and accurately presented and does it cite the current literature?

Yes

Is the study design appropriate and is the work technically sound?

Yes

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

Partly

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: Global health, tobacco control, mass media communications, social media, behaviour change, Pacific region.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 07 Aug 2020

Rashmi Rodrigues, St John's National Academy of Health Sciences, Bangalore, India

Dear Dr. McCool,

We very much appreciate and thank you for your valuable time towards reviewing our work. We have addressed your comments to the best of our ability and sincerely hope we have done justice to your suggestions on the report. The changes have been mentioned in a point-wise format as mentioned below, and a second version of the manuscript has been submitted to the journal.

We have added information on the sampling strategy, rephrased the table titles and removed associations for which adjusted odds ratios were unavailable among many other minor corrections, in line with the feedback received.

We hope the revised manuscript meets the standards to be considered for a formally approved status on Wellcome Open Research.

Thanking you again,

Sincerely,
Mochammad Fadjar Wibowo
Submitting Author
(on behalf of all the authors)

RESPONSE TO REVIEWERS:

EXTERNAL PEER-REVIEWER #1 (Judith McCool, BA, MPH, PhD)

Thank you once again for your comments. We have taken note of your detailed feedback on various sections in the manuscript and have addressed them to the best of our ability. Please find our point-counterpoint responses to your comments below:

- Introduction: Page 3. The statement regarding the high prevalence of tobacco use in Indonesia was attributed to the size of the population *and* the lack of public health campaigns addressing cessation. Might it also have something to do with tobacco industry interference, low taxation of tobacco and public awareness messages, smoke-free environments, banning advertising and promotions? This statement gives the impression that the current rates are attributable to the size of the population and cessation information, exclusively.

Response: We have now revised and reorganized the statement to read as follows (Introduction paragraph 2):

The large revenue from cigarettes despite low taxes, the employment opportunities within the tobacco industry, the weak anti-tobacco legislation and public health campaigns are considered to drive the tobacco epidemic in Indonesia.

- Introduction Page 3. The WHO FCTC covers more than prevention of update of tobacco by adolescents and protection from second-hand cigarette smoke. It also covers cessation support and other measures that have evidence of impact. The statement on page X could be amended to reflect the broader intention of the WHO FCTC.

Response: This statement has been amended as follows:

Indonesia is also the only country in the Asia Pacific that has not ratified the World Health Organizations (WHO) Framework Convention on Tobacco Control (FCTC) that addresses the demand and supply of tobacco and related products in order to promote health³.

- The statement at the start of the following paragraph captures the challenges, but again does not make any reference to the weak regulation of the tobacco industry and marketing, distribution and sales of tobacco products. I note though that this paper is focused on the potential to increase cessation, but this is confounded by a supportive smoking environment.

Response: Thank you very much for highlighting this. We have revised the sentence to read as follows:

This weak implementation of international regulations to minimize the marketing, distribution and sales of tobacco products along with prevailing economic, political and social factors have resulted in a high burden of tobacco-related morbidity and mortality in Indonesia ⁴

- Introduction, Page 3, paragraph 6. It was noted that the text messages were found to be effective (to support quit attempts) in New Zealand and the United Kingdom, smartphone android applications.... Is it the case that apps were trialled only on android phones? My interpretation is that they were similarly ineffective whether delivered on iPhone or android phones (Abroms *et al.*, 2013¹).

Response: Thank you very much for highlighting this. We have now revised our statement as follows:

In this regard, text messaging was effective for smoking cessation in New Zealand (personalized text messages to provide distraction, advise and support) and the United

Kingdom (motivational messages and feedback focusing on their achievements), **mobile phone applications**, though not tested for efficacy in a randomized control trial (RCT), are known to reach smokers who are not seeking professional help.

You will notice that we have changed the phrase **smartphone android applications to mobile phone applications**.

- Methods. It is possible that having the survey promoted via a weblog that is for smoking cessation may have reduced the potential sample size given the study was about what people think they want for support for quitting, not about if they are trying to quit. This may, in addition to other factors, have reduced the final sample size.

Response: Thank you for the suggestion. We did receive 800 odd responses, however as described in the results section. However, your comment has now been incorporated into the limitations of the study.

- Ethics statement, page 4. The ethics statement is included here and again at the end of the paper (on page 12), suggest presenting it once only.

Response: The duplicated ethical statement at the end of the article has been removed per your suggestion.

- Results: Smoking status characteristics; it was interesting to note that there were no differences in preference for receiving information via mobile device between socio-demographic groups. I also note that there was a low or very low level of dependency reported, despite the length of time that smoking was reported. Was this expected and is it explained in the discussion?

Response: Thank you once again. We do agree that there was no difference in the preference for receiving information between sociodemographic groups. However, as our study questionnaire did explore this further, we regret, that we are unable to comment on the reason for this at this stage.

We have now discussed the low level of dependency in the limitations of the study as follows:

As the reported levels of tobacco dependency were low, it is likely that the respondents were those who either had greater control over their smoking behavior or were more amenable to the idea of quitting. However, the level of nicotine dependence obtained may be questionable given the normalization of smoking in Indonesia.

- It was noted that most smokers in Indonesia quit without behavioural or NRT support. What proportion of those who attempt to quit, succeed. Is there any measure of these changes over time in terms of quitting? What were the methods that were used to support smokers to quit cold turkey? Is there any research on this that can shed light on the types of content that might be salient if delivered via mobile phone?

Response: Thanks for the suggestion. We have now incorporated the following in the 1st paragraph of our discussion:

Currently, 10% of tobacco users quit tobacco annually. [RR1] Further, literature from Indonesia shows that 66% of research participants in one study unsuccessfully attempted to quit smoking, [RR2] while another study reported that 15% its participants had quit smoking. [RR3]

Unfortunately, we did not explore the preference to quit smoking in our study as we primarily focused on quitting with mobile phones and providing support for the same over an extended period for those who would prefer this method. We do agree that this method is likely to work only if an individual is motivated enough to use it and that the choice of the method belongs to the individual. Given that quitting cold turkey is a popular method of smoking cessation we understand that there is a dominance of interventionism in our approach and have included this in our limitations as follows:

Given that quitting cold turkey is a popular method of smoking cessation we understand that our approach to smoking cessation reflects a dominance of interventionism. The proposed intervention should not be considered as a 'one size fits all' but rather one in a basket of interventions that includes interventions such as behavioral therapy, pharmacotherapy or quitting cold turkey.

- The finding that suggests a preference for smartphone apps over text message is unique and is inconsistent with other studies which have found little support for apps. I refer to a co-designed intervention using an app for behaviour change in New Zealand with indigenous and Pacific populations (Ni Mhurchu *et al.*, 2019²) among others.

Response: We agree that the preference to smart phone apps over text messages was found in this study. However, we believe that every population is unique, more so indigenous populations. Our attempt through this study was primarily to assess the acceptability of mobile phone interventions primarily for smoking cessation in Indonesia. The results therefore need to be viewed in this context and are therefore not expected to be generalizable to other populations unless they are similar.

We have incorporated your suggestion in the discussion section under "mode of communication" as follows:

However, interventions designed should be contextual as the preference for smartphone applications over other forms mobile interventions, such as text messages, might vary globally[RR4].

- Page 10. The statement that start "Only a few studies have explored the effectiveness of smartphone applications as a behaviour change intervention" is an important point but may need to be updated. There is an increasing number of publications coming through that question the value of apps in health and behaviour change.

Response: Thank you very much for highlighting this. We have rephrased this as follows, "Smartphone applications are a promising medium to reach smokers across multiple nations."

- Conclusion. Suggest revise the statement that a 'potential smoking cessation intervention...' was perceived as useful.

Response: Thank you for the suggestion. We have revised the statement as follows: Our study showed that the Indonesian respondents to our survey perceived a potential smoking cessation intervention via mobile phones as useful.

- The final statement regarding the implementation of the app within an organised programme (via schools etc) was not a conclusion that came through strongly in the data. It might be that any intervention, whether it is mobile device delivered or not, needs to complement a comprehensive tobacco control programme, backed by government support. Politics had not been mentioned in any detail until this point and seems out of place or underdeveloped at this stage.

Response: Thank you once again. We have revised our conclusion keeping in mind that we studied the acceptability of a mobile phone driven smoking cessation intervention as follows:

Our study showed that the Indonesian respondents to our survey perceived a potential smoking cessation intervention via mobile phones as useful. Perceived usefulness was associated with smokers' willingness to quit smoking and their prior use of mobile phones for health-related communication. A multicomponent smartphone application personalized to time, frequency and content was desired. Such an application, if implemented, could be one in a basket of smoking cessation solutions offered within an organized program quit smoking programmes at schools, healthcare facilities and counseling centers could go a long way in addressing the tobacco epidemic in Indonesia.

- Table 1. What does 'gainfully employed' mean - I am assuming this mean paid or salaried work?

Response: Yes, the term refers to an employee receiving steady work and payment from an employer.

- Income level in IDR - could there be a proxy figure in US or Euro? It was noted earlier that the DALY for tobacco related illness was reported in USD.

Response: A conversion for IDR and USD is now included in the table caption.

- Table 4. Column 1 - what does the *refer to on Mobile phone use proficiency*.

Response: This has been reworded to "Mobile phone usage skills".

- Figure 4. This figure does not substantively add to the paper. A brief description of the theory of behaviour change, and a reference would suffice.

Response: Thanks very much for the suggestion. However, we would like to keep the figure

as we feel that it pictorially depicts what we wish to express better.

Once again, we thank the reviewers immensely for their valuable comments. We hope we have addressed all the issues you have highlighted to your satisfaction.

Best wishes,
Authors

Competing Interests: No competing interests were disclosed.

Reviewer Report 23 July 2019

<https://doi.org/10.21956/wellcomeopenres.16514.r35764>

© 2019 Nichter M. This is an open access peer review report distributed under the terms of the [Creative Commons Attribution License](#), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.



Mimi Nichter

School of Anthropology, University of Arizona, Tucson, AZ, USA

This article provides information on the potential for smoking cessation interventions in Indonesia. This is an important topic as Indonesia has one of the highest prevalence of smoking among men in the world and is a country where phone use is extensive. Although this exploratory study is focused on the acceptability and perceived usefulness of a mobile phone smoking cessation app, it fails to consider the importance of incorporating the cultural context of smoking and perceived harm of tobacco use into the intervention. We need to consider the extent to which an intervention developed in New Zealand or the UK would have equal relevance to the Indonesian smoker. While this is beyond the focus of the current study, it should be discussed as critically important for the success of any mHealth intervention developed for LMIC.

1. Introduction: In relation to pharmaceuticals for nicotine addiction (patch, gum etc) - note that these medications, although available without prescription, are very expensive, and out of the realm of the possible for most Indonesians. (It would be interesting to know how much more they are than cigarettes, which are very inexpensive).
2. P. 4 Of the 850 visitors to the website, why did only 19% (n=161) complete the survey? This is a very low response rate. How long did the survey take to complete? Were respondents all from Java or from all over Indonesia? It is not clear from the discussion.
3. In one section, you mention 161 people as completing the survey, then in the socio-demographic data you note "Of the 123 respondents..." So how many people were there?
4. 83% of respondents were men; 17% were women—yet most studies to date show prevalence of women's smoking in Indonesia is about 2-3%, so please explain this

discrepancy.

5. In Table 1, provide IDR calculations to USD or to UK pounds so readers outside of Indonesia can interpret what the income figures mean.
6. Explain why only 75% of respondents were smokers. Why would the other 25% be interested in the intervention if they did not smoke? Were they ex-smokers who were concerned about relapse or people gathering information for family members?
7. Table 3: Do the authors have more detailed information on smoking status: daily smoker could be a very low level smoker (3 a day, or it could be 10 per day); occasional smoker could be defined differently by various people to mean once a week or twice a week or monthly, etc. The number of cigarettes smoked per day would be much more useful data, if available.
8. Consider eliminating the paragraph on what other uses the informants had for their phones (alarm function; texting, playing games...). It is not clear what relevance this has to the study at hand. Why is alarm use needed in Table 4? Why is the main use of mobile internet needed - it seems obvious that these are the features people want and use on a regular basis. If there is a reason for inclusion, please do make it clear.
9. Do people have to pay for text messages received in Indonesia or are they free, or does it depend on the plan? This might be important in people's decision to want text messages.
10. p. 7. Describe what is meant by "permitted number of cigarettes for a day". Who determines this? When is it determined? Is it in relation to number of cigarettes smoked per day prior to quitting?
11. Discussion, p. 7: the numbers cited in relation to DALYs are different than what is mentioned in introduction. Consider changing the first sentence to reflect the prevalence of smoking in Indonesia, and the lack of cessation services currently available.
12. With regards to knowledge about the harm of tobacco in Indonesia, the authors should review Padmawati, Ng *et al.* (2009)¹, in which it is noted that diabetes patients were believed to be able to smoke 3 cigarettes per day as relatively harmless while healthy people could smoke 12 cigarettes a day. See also Ng *et al.* (2010)². Also of use would be Nichter *et al.* (2009)³, for details about perceptions of smoking and quitting in Java. It would be useful for the reader to understand a bit more about Indonesia and people's attitudes toward quitting. A paragraph in the introduction would be sufficient and would highlight the reason for developing mobile phone apps for cessation and their potential in the country.
13. p. 7. Section on Quitting with mobile phones: Make it clear that at present behavior change interventions are not common in Indonesia; As your own data shows, few patients are getting a message to quit from their doctor.
14. Preferred mode of communication Figure 2: write out what SMS, MMS and IVR stand for.
15. Figure 2 & 3: Why does the n vary between figures?

16. Figure 3: Only a small percentage of your sample had smartphones, so what were they answering: what they would want in a smartphone if they had one?
17. Given that this study was done a few years ago, what is the prevalence of smartphone use now in Indonesia?
18. Explain early on what type of message would be delivered by SMS, MMS and IVR. How do they differ? Can you give examples from the intervention?
19. p. 10. It is unclear to the reader if any content for the proposed smoking cessation intervention has been developed or if the article is based solely on whether the audience would utilize an online intervention. Has thought been given or shared to the audience about the content of the intervention? Will it be based on an understanding of Indonesian smoker's behaviors and ideas about quitting or will the intervention be a translation of a program normed and developed in another country? If so, which country? Are there plans for the smoking intervention to be adapted for the Indonesian cultural context? Please note that even high level smokers in Indonesia do not think of themselves as "addicted,"—which clearly has implications for thinking about quitting.
20. p. 10. The authors state that few studies have explored use of smartphones for interventions. Hasn't the CDC in the US developed a large smoking cessation intervention? This is not a new phenomena. Further on p. 11, you review many of these interventions, so these sections need to be combined.
21. p. 10 The information about smartphone apps for a smoking cessation intervention should be moved into the introduction, as well as info on an MMS-based intervention. All of this information about development of apps in other countries should be noted in the beginning of the article so the reader understands a bit of the context.
22. p. 11 Move the information about the UK to the introduction. You should also note that messages to avoid environments where people smoke are extremely problematic in Indonesia where over 65% of men smoke and where few smoke free restrictions are in place.
23. p. 11 "the profound ethnic diversity of Indonesia...if considered..might increase the complexity and cost of the intervention"...Yes surely it would, but utilizing a 'one size fits all' approach does not seem like a recipe for success.
24. p. 12. Methodological Issues: Perhaps rename this as Limitations. The authors note the levels of tobacco dependency were low among respondents. It is possible that the reason for this is that the Fagerstrom as a measure of nicotine dependence in Indonesia is not very robust because smoking patterns are very different. This has been addressed in several publications of the Quit Tobacco International Project, conducted in Yogyakarta, the site of the present study.
25. p. 12 Conclusion: reword line one to indicate "Our study showed that among those smokers in Indonesia who responded to our survey..." rather than "our study showed that people in

Indonesia who smoke (which seems a bold claim for a small survey).

References

1. Padmawati RS, Ng N, Prabandari YS, Nichter M: Smoking among diabetes patients in Yogyakarta, Indonesia: cessation efforts are urgently needed. *Trop Med Int Health*. 2009; **14** (4): 412-9 [PubMed Abstract](#) | [Publisher Full Text](#)
2. Ng N, Nichter M, Padmawati RS, Prabandari YS, et al.: Bringing smoking cessation to diabetes clinics in Indonesia. *Chronic Illn*. 2010; **6** (2): 125-35 [PubMed Abstract](#) | [Publisher Full Text](#)
3. Nichter M, Nichter M, Padmawati S, Thresia CU, et al.: Anthropological contributions to the development of culturally appropriate tobacco cessation programs. *Anthropology and Public Health: Bridging Differences in Culture and Society*. 2009.

Is the work clearly and accurately presented and does it cite the current literature?

Partly

Is the study design appropriate and is the work technically sound?

Partly

Are sufficient details of methods and analysis provided to allow replication by others?

Partly

If applicable, is the statistical analysis and its interpretation appropriate?

Yes

Are all the source data underlying the results available to ensure full reproducibility?

No source data required

Are the conclusions drawn adequately supported by the results?

Partly

Competing Interests: No competing interests were disclosed.

Reviewer Expertise: I am a medical anthropologist who has conducted extensive research on the development of tobacco cessation programs in Java, Indonesia.

I confirm that I have read this submission and believe that I have an appropriate level of expertise to confirm that it is of an acceptable scientific standard, however I have significant reservations, as outlined above.

Author Response 07 Aug 2020

Rashmi Rodrigues, St John's National Academy of Health Sciences, Bangalore, India

Dear Dr. Nichter,

We very much appreciate and thank you for your valuable time on reviewing our work. We

have addressed your comments to the best of our ability and sincerely hope we have done justice to your suggestions on the report. The changes have been mentioned in a point-wise format as mentioned below, and a second version of the manuscript has been submitted to the journal.

We have added information on the sampling strategy, rephrased the table titles and removed associations for which adjusted odds ratios were unavailable among many other minor corrections, in line with the feedback received.

We hope the revised manuscript meets the standards to be considered for a formally approved status on Wellcome Open Research.

Thanking you again,

Sincerely,
Mochammad Fadjar Wibowo
Submitting Author
(on behalf of all the authors)

RESPONSE TO REVIEWERS:

EXTERNAL PEER-REVIEWER #1 (Mimi Nichter, PhD)

Thank you once again for your comments. We have taken note of your detailed feedback on various sections in the manuscript and have addressed them to the best of our ability. Please find our point-counterpoint responses to your comments below:

- Introduction: In relation to pharmaceuticals for nicotine addiction (patch, gum etc) - note that these medications, although available without prescription, are very expensive, and out of the realm of the possible for most Indonesians. (It would be interesting to know how much more they are than cigarettes, which are very inexpensive).

We have added this information under the 'Introduction' section paragraph 2 per your comment.

In Indonesia, nicotine replacement therapy such as nicotine patches, gums, and sprays are available without prescription in pharmacies but cost 7000-250,00 IDR (0.0035-0.035 USD) per unit. Quitting with prescription medication such as varenicline, hypnotherapy or behavioural therapy cost 0.8-1 million IDR and are not very popular ⁶. Further, most (71%) of those who want to quit smoking, attempt to quit without assistance while others may use traditional methods (herbal or medicinal plants), smokeless tobacco or counselling ⁶. The costs of quitting vs the costs of cigarettes (12-24000 IDR/ pack currently and 32-53000 IDR from 1st November 2019) probably influences both the quit strategy and the decision to continue smoking. Additionally, the self-confidence of Indonesian physicians in providing smoking cessation counselling is reportedly low, with smoking cessation services offered only at a few healthcare facilities ⁸. There is also no national toll-free quit-line ⁹. This prevailing scenario makes innovation and improvement in smoking cessation interventions in Indonesia a necessity.

- P. 4 Of the 850 visitors to the website, why did only 19% (n=161) complete the survey? This is a very low response rate. How long did the survey take to complete? Were respondents all from Java or from all over Indonesia? It is not clear from the discussion.

Answer:

Participants took 34:06 ($\pm 2:02$) minutes to complete the questionnaire. – this is now mentioned in the *last sentence of the 5th paragraph in the methods section* of the manuscript. The respondents' locations represented 14 of the 34 provinces in Indonesia- this was already mentioned in the *1st paragraph of the results section* of the manuscript.

- In one section, you mention 161 people as completing the survey, then in the socio-demographic data you note "Of the 123 respondents..." So how many people were there?

While 161 respondents completed the questionnaire, not all responded to every single question. Also, the questionnaire was designed such that respondents could skip questions based on their responses to previous questions, which changed the 'n' for various sections and questions of the questionnaire.

- 83% of respondents were men; 17% were women—yet most studies to date show prevalence of women's smoking in Indonesia is about 2-3%, so please explain this discrepancy.

While there were only 124 of the 161 respondents who provided complete demographic information, especially information on gender, we believe that the difference may be due to internet accessibility and receipt of information regarding the survey. Unlike the prevalence of smoking, the distribution of men and women accessing the internet and using social media (using which the questionnaire was made available) in Indonesia is approximately 50% each and the gender distribution in the study is a consequence of internet usage patterns.

- In Table 1, provide IDR calculations to USD or to UK pounds so readers outside of Indonesia can interpret what the income figures mean.

USD to IDR conversions have been added to Table 1, 2 and 4.

- Explain why only 75% of respondents were smokers. Why would the other 25% be interested in the intervention if they did not smoke? Were they ex-smokers who were concerned about relapse or people gathering information for family members?

Answer:

The questionnaire was administered to respondents to ever smoked. While 75% were current smokers, ex-smokers were required to respond to smokers 'hypothetically' in a scenario of the mobile phone intervention being made available when they were attempting to quit. We believe that they would be able to use their quitting experience to better advise us regarding an intervention that enabled quitting. Unfortunately, we did not obtain information on whether they were concerned about relapse.

We have mentioned this in the *5th paragraph* of the methods section of the manuscript as follows:

'These respondents were current and former smokers, aged 18 years or older, residing in Indonesia for the past year.'

- Table 3: Do the authors have more detailed information on smoking status: daily smoker could be a very low level smoker (3 a day, or it could be 10 per day); occasional smoker could be defined differently by various people to mean once a week or twice a week or monthly, etc. The number of cigarettes smoked per day would be much more useful data, if available.

We do agree with the reviewer that the number of cigarettes smoked would be more useful data.

The smoking status used in the analysis was obtained from the question i.e., the response to the question Do you currently smoke tobacco on a daily basis, less than daily, or not at all?

However, we also requested participants to report the number of cigarettes smoked/ day. Unfortunately, only 100 reported the number of cigarettes smoked/ day or week.

Participants reported smoking an average 9(SD8) cigarettes/ day, ranging from <1-32 cigarettes/ day. Hence, we used their self-report of smoking daily or occasionally or in the past as an indicator of their current smoking status. Most participants had smoked 3 pack years and ranged from 0.1-41 pack years. For the purpose of calculating pack years we assumed cigarette packs despite the different types of tobacco smoked.

We have now also reported the number of cigarettes smoked in a day in the results section under **Smoking status characteristics**.

- Consider eliminating the paragraph on what other uses the informants had for their phones (alarm function; texting, playing games...). It is not clear what relevance this has to the study at hand. Why is alarm use needed in Table 4? Why is the main use of mobile internet needed - it seems obvious that these are the features people want and use on a regular basis. If there is a reason for inclusion, please do make it clear.

Thank you for the suggestion - however, we would like to keep the paragraph as it suggests the other features that could be incorporated into the design of a mobile phone application for smoking cessation.

- Do people have to pay for text messages received in Indonesia or are they free, or does it depend on the plan? This might be important in people's decision to want text messages.

Receiving text messages in Indonesia is free irrespective of the plan. It however costs IDR 360 (3 cents) to send a message, while bulk messaging would cost upto IDR 65,000 (USD 4.5) for 500 messages a month. These costs would be incurred by the intervention provider.

- p. 7. Describe what is meant by "permitted number of cigarettes for a day". Who determines this? When is it determined? Is it in relation to number of cigarettes smoked per day prior to quitting?

Answer:

We have revised this sentence to read- *'reminders about the numbers of cigarettes that they could smoke each day as they approached their quit date'*

Studies have indicated that reduction in smoking is easier to achieve than complete cessation and once achieved could promote complete cessation. Reduction reverses neuroadaptation to tobacco and minimises the symptoms of withdrawal and the cravings experienced with sudden complete cessation, therefore a target is set each day in terms of minimising the cigarettes smoked until the quit date is reached.

West Rj et al Psychol Med. 1989 Nov; 19(4):981-5.

- Discussion, p. 7: the numbers cited in relation to DALYs are different than what is mentioned in introduction. Consider changing the first sentence to reflect the prevalence of smoking in Indonesia, and the lack of cessation services currently available.

Answer:

Thank you very much for highlighting this. The discussion however indicated global DALYs while the introduction indicated the DALYs lost in Indonesia due to smoking.

However, in the interest of minimizing confusion we have rewritten the first paragraph of the discussion to read as follows:

In Indonesia, an LMIC, the burden of tobacco smoking has risen from 59 million in 2000 to 70 million in 2010 and 73.6 million in 2015 and continues to rise, so does the amount spent on tobacco related illnesses. The situation is compounded by the limited awareness regarding the hazards of smoking along with the minimalistic support available to quit. Finding innovative solutions that are acceptable to Indonesians wanting to quit is therefore essential. Therefore, given the current pervasiveness of mobile phone communication and its minimal cost we sought to explore the acceptability and design for a mobile phone smoking cessation intervention in Indonesia.

- With regards to knowledge about the harm of tobacco in Indonesia, the authors should review Padmawati, Ng et al. (2009), in which it is noted that diabetes patients were believed to be able to smoke 3 cigarettes per day as relatively harmless while healthy people could smoke 12 cigarettes a day. See also Ng et al. (2010). Also, of use would be Nichter et al. (2009), for details about perceptions of smoking and quitting in Java. It would be useful for the reader to understand a bit more about Indonesia and people's attitudes toward quitting. A paragraph in the introduction would be sufficient and would highlight the reason for developing mobile phone apps for cessation and their potential in the country.

We agree with your thoughts on the baseline understanding of smoking and its effects among Indonesian subjects and have added a paragraph highlighting perceptions under the 'Introduction' section as follows:

Further, there are varied perceptions of the effects and complications of tobacco on health in Indonesia. For example; some patients with diabetes mellitus considered that they could smoke relatively lesser cigarettes/ day (3/day) when compared with those who were healthy (12/ day) [*Insert Ref D*], while others it did not know it could complicate their illness [*Insert Ref E*]. Also, reducing or quitting smoking was considered an option only for those seriously ill and could be resumed on recovery. [*Insert Ref D*]. This prevailing scenario makes innovation and improvement in smoking cessation interventions in Indonesia a necessity.

- p. 7. Section on Quitting with mobile phones: Make it clear that at present behavior change interventions are not common in Indonesia; As your own data shows, few patients are getting a message to quit from their doctor.

Answer:

Thank you once again for the suggestion. We have made the change and introduced the following paragraph under the section on 'Quitting with mobile phones'

Currently in Indonesia behaviour change interventions are uncommon and when available are expensive. So also, is advise from physicians regarding quitting. Our study indicated that nearly half the participants did not receive any advice from their healthcare provider to quit, despite a reported desire to do so.

- Preferred mode of communication Figure 2: write out what SMS, MMS and IVR stand

for.

Thank you for comment we have now expanded these abbreviations.

- Figure 2 & 3: Why does the n vary between figures?

We acknowledge that discrepancies in the numbers (n) between variables exist, these are due to missing data especially demographic, as respondents most probably chose not to respond to them. We have now discussed the discrepancy in the 'limitations section' of the manuscript.

- Figure 3: Only a small percentage of your sample had smartphones, so what were they answering: what they would want in a smartphone if they had one?

In the "Mobile phone usage patterns" section under "Results", we had stated that 154 (98%) of our respondents used a smartphone which reflects a majority. The minority of subjects who did not own a smartphone (7, 2%) answered the questions on desired features of an application had they owned a smartphone as you mentioned in your comment. We have thus made no changes here.

- Given that this study was done a few years ago, what is the prevalence of smartphone use now in Indonesia?

We have described the trend in mobile phone usage in Indonesia under the section '*Mobile phone penetration and mHealth development in Indonesia*' of the introduction as follows:

The growth of mobile users in Indonesia is one of the fastest in Asia with a steady increase from 125 per 100 people in 2013 to 173 per 100 people in 2018 ²⁰. Given the improving internet accessibility and low cost of smartphones ²¹, with prices as low as 40 USD for a phone, smartphone penetration in Indonesia has reached 27% in 2018 ²² and is predicted to reach 32% by 2022 ²³.

- Explain early on what type of message would be delivered by SMS, MMS and IVR. How do they differ? Can you give examples from the intervention?

Answer:

Thank you for the suggestion: we have now explained this in the introduction (7th paragraph) as follows:

Short Messaging Service (SMS) Multimedia Messaging Service (MMS) (pictorial messages), live-voice calls and interactive voice response (IVR) technology that replace a human caller with a computer, provide motivation and counseling to those who want to quit smoking ¹¹. Such messages use prompts (either text i.e.; SMS, picture i.e, MMS or voice i.e., calls and IVR) to encourage avoiding cigarettes, ashtrays, lighters, and environments where people usually smoke eg.; 'For the next 4 hours, stay away from cigs'. Additionally, messages also help identify the challenges to quitting and the plan to overcome them. Prompts to use telephone helplines and nicotine replacement therapy ¹⁷, economic savings from quitting and nutritional advice form the content of messages used to support quitting.

- p. 10. It is unclear to the reader if any content for the proposed smoking cessation intervention has been developed or if the article is based solely on whether the audience would utilize an online intervention. Has thought been given or shared to the audience about the content of the intervention? Will it be based on an understanding of Indonesian smoker's behaviors and ideas about quitting or will the intervention be a translation of a program normed and developed in another country? If so, which country? Are there plans for the smoking intervention to be adapted for the Indonesian cultural context? Please note that even high level smokers in Indonesia do not think of themselves as "addicted,"—which clearly has

implications for thinking about quitting.

We had not developed any content for the intervention when the study was initiated. Based on this study and available funding support further development of the intervention will happen for which intensive work into what content and its wording is acceptable to the Indonesian population will be identified and used. In this study we have aimed only to identify the kind of communication that is acceptable, its frequency and some basic content.

- p. 10. The authors state that few studies have explored use of smartphones for interventions. Hasn't the CDC in the US developed a large smoking cessation intervention? This is not a new phenomena. Further on p. 11, you review many of these interventions, so these sections need to be combined.

Thank you for highlighting this. Our aim was to indicate that such interventions have been used minimally in low-middle income contexts such as Indonesia. However, based on your suggestion we think it best to remove the paragraph from the manuscript. We however choose keep the sections as they are.

- p. 10 The information about smartphone apps for a smoking cessation intervention should be moved into the introduction, as well as info on an MMS-based intervention. All of this information about development of apps in other countries should be noted in the beginning of the article so the reader understands a bit of the context.

Answer: We have moved part of the content to the introduction as indicated in response 18

- p. 11 Move the information about the UK to the introduction. You should also note that messages to avoid environments where people smoke are extremely problematic in Indonesia where over 65% of men smoke and where few smoke free restrictions are in place.

We have moved the information on UK to the introduction as best as we can. We understand that messages to avoid environments where people smoke are extremely problematic in Indonesia where over 65% of men smoke. We will ensure that the intervention designed (based on available funding) will have an extensive situational analysis prior to design and deployment.

- p. 11 "the profound ethnic diversity of Indonesia...if considered..might increase the complexity and cost of the intervention"...Yes surely it would, but utilizing a 'one size fits all' approach does not seem like a recipe for success.

Thank you. We do not recommend a one size fits all approach. What we suggest is a menu of features within an application, identified based on potential user surveys. The beneficiaries/ users and then choose the features that best suit them. However, this will make the intervention complex and difficult to test scientifically.

- p. 12. Methodological Issues: Perhaps rename this as Limitations. The authors note the levels of tobacco dependency were low among respondents. It is possible that the reason for this is that the Fagerstrom as a measure of nicotine dependence in Indonesia is not very robust because smoking patterns are very different. This has been addressed in several publications of the Quit Tobacco International Project, conducted in Yogyakarta, the site of the present study.

Thank you!

We have renamed methodological issues as '*limitations*'

We have included your suggestion in the limitations as follows:

However, the level of nicotine dependence obtained may be questionable given the normalization of smoking in Indonesia. Further, social desirability bias also cannot be ruled out in the FTND as it is a self-report of dependence by the participant.

- p. 12 Conclusion: reword line one to indicate “Our study showed that among those smokers in Indonesia who responded to our survey...” rather than “our study showed that people in Indonesia who smoke (which seems a bold claim for a small survey).

This sentence has been reworded to reflect more clarity per your suggestion as follows: Our study showed that smokers in Indonesia who responded to our survey perceived receiving a potential smoking cessation intervention via mobile phones as useful.

Once again, we thank the reviewers immensely for their valuable comments. We hope we have addressed all the issues you have highlighted to your satisfaction.

Best wishes,
Authors

Competing Interests: No competing interests were disclosed.

