

RESEARCH

Open Access



Full-sized realistic 3D printed models of liver and tumour anatomy: a useful tool for the clinical medicine education of beginning trainees

Guoqiang Bao¹, Ping Yang¹, Jiangpu Yi², Shujia Peng¹, Jiahe Liang², Yajie Li³, Dian Guo³, Haoran Li¹, Kejun Ma³ and Zhenyu Yang^{1*}

Abstract

Background Simulation-based medical education (SBME) and three-dimensional printed (3DP) models are increasingly used in continuing medical education and clinical training. However, our understanding of their role and value in improving trainees' understanding of the anatomical and surgical procedures associated with liver surgery remains limited. Furthermore, gender bias is also a potential factor in the evaluation of medical education. Therefore, the aim of this study was to evaluate the educational benefits trainees receive from the use of novel 3DP liver models while considering trainees' experience and gender.

Methods Full-sized 3DP liver models were developed and printed using transparent material based on anonymous CT scans. We used printed 3D models and conventional 2D CT scans of the liver to investigate thirty trainees with various levels of experience and different genders in the context of both small group teaching and formative assessment. We adopted a mixed methods approach involving both questionnaires and focus groups to collect the views of different trainees and monitors to assess trainees' educational benefits and perceptions after progressing through different training programs. We used Objective Structured Clinical Examination (OSCE) and Likert scales to support thematic analysis of the responses to the questionnaires by trainees and monitors, respectively. Descriptive analyses were conducted using SPSS statistical software version 21.0.

Results Overall, a 3DP model of the liver is of great significance for improving trainees' understanding of surgical procedures and cooperation during operation. After viewing the personalized full-sized 3DP liver model, all trainees at the various levels exhibited significant improvements in their understanding of the key points of surgery ($p < 0.05$), especially regarding the planned surgical procedure and key details of the surgical procedures. More importantly, the trainees exhibited higher levels of satisfaction and self-confidence during the operation regardless of gender. However, with regard to gender, the results showed that the improvement of male trainees after training with the 3DP

*Correspondence:
Zhenyu Yang
yangzhenyu1023@fmmu.edu.cn

Full list of author information is available at the end of the article



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

liver model was more significant than that of female trainees in understanding and cooperation during the surgical procedure, while no such trend was found with regard to their understanding of the base knowledge.

Conclusion Trainees and monitors agreed that the use of 3DP liver models was acceptable. The improvement of the learning effect for practical skills and theoretical understanding after training with the 3DP liver models was significant. This study also indicated that training with personalized 3DP liver models can improve all trainees' presurgical understanding of liver tumours and surgery and males show more advantage in understanding and cooperation during the surgical procedure as compared to females. Full-sized realistic 3DP models of the liver are an effective auxiliary teaching tool for SBME teaching in Chinese continuing medical education.

Keywords 3D-printed models, Simulation-based medical education, Continuing medical education, Liver surgery

Introduction

In clinical medicine education, learning practical skills is an essential task during medical studies. The use of surgical simulation to reach specific target criteria significantly reduces operation times and improves the performance of surgeons [1]. Simulation-based medical education (SBME) is believed to be superior to the traditional style of medical education according to active and adult learning theories [2]. Furthermore, SBME has the potential to influence students' motivation and has already been implemented in many medical curricula [3]. In the daily practice of liver surgeons, a wide variety of theoretical, practical and psychosocial competencies must be mastered at a very high level, which can help a surgeon have the ability to perform difficult procedures as early as possible. Thus, the training of these competencies is a central component in the education of liver surgeons to impart what is required for their later practice [4].

With respect to SBME, virtual and augmented reality tools have the potential to offer a more comprehensive alternative to three-dimensional (3D) visualisation but lack the ability to provide tactile feedback [5, 6]. However, 3D printing has proven to be a revolutionary technology in medical education [7–9]. 3D printing, as an additive manufacturing technology based on rapid prototyping, facilitates the creation of patient-specific physical models with high precision [10]; thus, it has been deployed in various medical fields, such as medical education and preoperative planning [11–13]. Additionally, the growing need for tactile and haptic learning in medical education is leading to the increased use of 3D printing in SBME [14, 15]. 3DP models can have both realistic anatomy and textures derived from patients' individual CT and/or MRI images, thus facilitating the creation of patient-specific physical models with high precision; these models seem to be capable of satisfying the needs for tactile and spatial perception of human anatomical structures [16, 17]. The models can help trainees understand organ physiology, anatomy, tumour characteristics, and surgical procedures more accurately [18]. The application of 3DP models bridges the gap between two-dimensional (2D) imaging and realistic anatomy, as it accurately reproduces

anatomical structures and pathologies, thereby providing more tangible information than conventional imaging data [19]. 3DP models also appear to be a significantly more useful and cost-effective technique than traditional cadaveric models in medical education [20], such as those used in undergraduate dentistry training [4], urology residents [21], first-year medical students [22], craniofacial traumas [23], oral and cranio-maxillofacial surgery [24]. However, recent SBME changes regarding 3DP models have focused only on undergraduate medical education [25]. Continuing medical education after postgraduate education has been neglected and under evaluated.

As an intricately vascular architecture and stereoscopic space distribution, the liver is an intriguing yet complex multifunctional organ [26]. Despite technological improvements, hepatectomy remains a challenging operation associated with high complication and mortality, especially for young surgeons [27]. Couinaud's liver segment classification is currently used in liver surgery, but there are many anatomical variations and variables on tumour growth, prior operations, and regenerative growth [28]. Unlike conventional imaging data using 2D films, which can lead to confusion in the actual situation, 3DP liver models are capable of illustrating the relationship between the vascular and biliary channels [29]. 3D anatomical relationships of the vascular and biliary channels are crucial for performing accurate liver resections and predicting the actual size of hepatectomies [30, 31]. This approach may improve residents' and students' understanding of organ physiology, anatomy, tumour characteristics, and surgical procedures. Thus, it has the potential to gain increasing importance in the training of residents and students [32, 33], as well as in surgical decision-making and planning for complicated conditions and procedures [34].

According to some studies, gender bias poses a potential threat to the integrity of resident assessment in medical education [35, 36], and gender-related differences are more pronounced among medical students [37, 38], such as visual-spatial working memory [39], spatial learning ability in a virtual environment [40] and confidence [41]. To our knowledge, no study has investigated the

educational benefit of 3DP liver models with regard to trainees at different levels in their continuing medical education and trainees of different genders.

Study goals

Constructivism theory and quantitative post-positivist methods provided a solid theoretical foundation for simulation education in our study. Constructivism theory was proposed by Swiss psychologist Piaget and its core is student-centered, emphasizing students' active exploration, discovery and construction of the meaning of what they have learned [42, 43]. As the learning environment required by constructivism was strongly supported by the latest information technology achievements, the theory of constructivism has been increasingly combined with the medical education and clinical practice, thus becoming a guiding philosophy for deepening teaching reform in schools [44, 45]. Therefore, we hypothesized that 3DP liver models play a more important role in simulation-based continuing medical education after post-graduate education and may differ according to gender. Accordingly, the aim of the present study was to explore the depth and quality of the 3DP liver models used in the continuing medical education of these different trainees while considering the role of gender by applying SBME educational theory. Understanding the educational benefit of 3DP liver models in continuing medical education can generate evidence that can promote the reform of continuing medical education management programs. Specifically, the objectives of our research on the use of full-sized 3DP live models during trainee education training were as follows:

1) to identify and assess different levels of trainees' understanding of the relevant base knowledge between the different training models, which is a tool to help trainees better grasp abstract knowledge through spatial visualization.

2) to identify and assess different levels of trainees' understanding of the key points of surgery between the different training models.

3) to identify the helpfulness and satisfaction of various levels trainees, which were divided into different levels according to when they graduated, including interns (beginner), standardized training trainees

(not experienced) and professional training trainees (experienced).

4) to determine whether gender-related differences exist among trainees of continuing medical education, which refers to a specific form of continuing education that helps those in the medical field maintain competence and learn about new and developing areas of their field.

5) to evaluate monitors' level of satisfaction with the trainees' cooperation during the operation.

Materials and methods

Population and sampling

This was a cross-sectional, analytical and questionnaire-based study. As a preliminary exploratory study, we only surveyed all new trainees at a single centre in the current year, who were enrolled in continuing medical education after their undergraduate course at a large, academic, tertiary care hospital and a university-affiliated medical centre. After we estimated the eligible population in the 10 medical and surgical wards of three teaching tertiary referral hospitals located in the western regions of China, we assumed that 80% of the graduate students will participate and recruited in this study. Based on the predicted anticipated trainees, a minimum sample size of 30 was calculated using the online statistical program Openepi, taking into consideration a 90% confidence interval, 95% response distribution, 5% margin of error, and an estimated total population size of 70. All participants had completed their internal medicine clerkship rotation at the time of the survey. All participants were at various levels of their training, including interns (beginner), standardized training trainees (not experienced) and professional training trainees (experienced). In the first step, the trainees were divided into 3 groups according to their level of training and sorted by gender (Table 1). Second, all trainees were randomly assigned to one of two groups: one group that received 3D model training and one group that did not.

Educational context

Based on the syllabus, especially the relevant courses, the basics of liver imaging examinations as well as the physiology and anatomy of the healthy liver were reviewed jointly before continuing the hands-on training. All trainees were given multislice CT/MRI scans and received similar preoperative information on the patients' disease, tumour characteristics, planned surgery and related risk of complications. This information was delivered during a face-to-face consultation with the supervisors using CT/MRI scan images. Subsequently, the 3D printed models were presented randomly to the assigned trainers in each group. Finally, in this study, half of each group of trainers received a 3D model, while the other half

Table 1 Classification of the 30 trainees

| Group | Gender (Female/Male) | Level of Training | Qualification |
|----------|-------------------------|--|--------------------------------|
| A (n=8) | 2/6 | Beginner (interns) | No operation attended |
| B (n=10) | 4/6 | Not experienced (standardized training trainees) | No livers operated |
| C (n=12) | 2/10 | Experienced (professional training trainees) | Liver opera- tions assisted |

Table 2 Classification of Presentation Models

| Presentation Models | Description |
|---------------------|--|
| 2D | Conventional 2-D CT scans with or without MRI |
| 3D | Combination of the 3D printed liver models with conventional 2-D CT with or without MRI scan |

received only 2D images (Table 2). The training occurred 3 times a week, and each session lasted 40 min. After 4 weeks of continuous training, the trainees and the supervisors were required to perform the tasks listed in the questionnaires.

Two questionnaires were created to prospectively evaluate the trainees' level of preoperative knowledge and understanding. Their knowledge was first evaluated based on the previously delivered information and CT scan images, following which the 3D printed model was used to assess the improvement following the model presentation. Questionnaire #1 consisted of questions intended to evaluate two components of trainees' knowledge: (a) basic liver physiology and anatomy, (b) tumour characteristics and (c) questions involved in the planned surgical procedure ("Appendix 1, Table S1"). Questionnaire #2 investigated trainees' satisfaction using a 10-point Likert scale ("Appendix 2"). Questionnaire #3 was designed to enable supervisors to evaluate the performance of trainees ("Appendix 3"). The other questionnaires were used to evaluate the feedback provided by the trainees with regard to the training models using questions scored on a 10-point Likert scale ("Questionnaire 4, Appendix 4, Table S2").

3D printed model fabrication

Imaging data

Considering the cost of the printed model, the imaging data were collected from one patient who had been diagnosed with HCC as well as one healthy volunteer. The imaging data were scanned using a helical Philips iCT256 (Philips Electronics Co., Netherlands). The scanning parameters were 0.6-mm layer thickness, 0.6-mm layer spacing, 120 kV, 280 mAs, and 220 mm field of view (FOV). The contrast agent was iohexol injection 350 (Shanghai GE Pharmaceutical Co., LTD.) at a flow rate of 4.0 ml/s. The original image data were output in the standard digital imaging and communications in medicine (DICOM) format.

Mimics modeling

The DICOM data were analysed using Mimics to determine whether the scanning range was the anatomical site required by clinical practice. The segment parenchyma, including the artery, liver vein, portal vein, liver, gallbladder, and bile duct, were calculated and exported as STL files using Mimics. Then, procedures such as smoothing,

deleting and clipping the model, hollowing out with a certain wall thickness, and designing the connection structure were performed using Magics. A biomedical engineer working alongside surgeons completed model segmentation.

The liver digital model was printed using the PolyJet3D technology of the Stratasys J850 printer with VeroCyanV/VeroYellowV/VeroMagentaV/VeroPureWhite/VeroUltraClea as the model transparent material. According to the requirements, the printing temperature was adjusted between 72 and 76 °C, and the UV light wavelength was 365 nm. Photosensitive resin material with different colours was selected for printing different anatomical structures, i.e., the gallbladder (green), portal vein (violet), artery (red), and hepatic veins (blue), as shown in Figs. 1 and 2. Noncontact resin was used as the support material for jet moulding, and it was cleaned by high-speed water flushing. The minimum thickness of the printing layer can be up to 0.016 mm, and the printing accuracy can be up to 0.1 mm. It takes approximately 14 h to print a liver model by spraying multiple photosensitive resin materials of different colours onto the same plane, and the cost of printing materials is approximately \$260 USD, which can be reduced further as the number of prints increases.

Data analysis

The number of correct responses to the questions in Questions #1 and #4 was used as the endpoint. The selected number was used as the endpoint in Questionnaires #2 and #3. Number of correct response variables of their answers to specific questions and correlations between the different gender of each group are presented with absolute frequency and were compared with the Pearson's chi-square test or Fisher's exact test when appropriate. The median score of the evaluation was expressed as the median [Min-Max] and analyzed by Student's t test or Wilcoxon tests. Statistical significance was defined as a *P* value < 0.05 with a two-tailed test. Statistical analyses were performed using SPSS Statistics version 21.0 (IBM Corporation, Armonk, NY, USA).

Ethical statement

An informed consent form was provided to the trainees of each level and the supervisors involved in the survey, and the survey was conducted after ensuring that all members understood the importance of voluntary participation and acquiring written informed consent from each participant. In addition, this study was approved by the Ethics Committee of the Tangdu Hospital of Air Force Medical University (No. K202207-06) and conducted according to the standards stipulated in the Declaration of Helsinki. Moreover, in this study, the were collected from. Written consent also was obtained for




Fig. 1 The rigid liver 3D printing model of the healthy volunteer. The parenchyma was cast using photosensitive polymer in layers with ease of visualisation of the vascular structures: the venous structure (blue) and the artery tree (pink). **A:** Anterior view. **B:** Superior view. **C:** Digital 3D reconstruction of liver lobulation and internal duct structure. **D/E:** Digital 3D reconstruction shows liver segment features from the front. **F:** The rigid liver 3D printing model shows liver segment features with different colours




Fig. 2 Soft liver and tumour 3D printing model of one patient. The parenchyma was cast using photosensitive polymer in layers allowing an easy cut during preoperative planning. CT images of the cross-sectional venous phase showed the tumour (**A**) and based on CT images (**B** Relationship between hepatic vein and tumour, **C** Relationship between portal vein branch and tumour, **D** dorsal view, **E** Superior view); The rigid liver 3D printing model (**F**) and Intraoperative image of the patient (**G**: The resection line was marked with radiofrequency ablation needle, **H**: The right posterior branch of the portal vein was preserved, **I**: Photograph of resected specimen of tumour)

Table 3 Different levels of trainees' understanding assessment on the base knowledge between different training models

| Group ^a | Number of correct responses with different genders | | | | | | χ^2 | P |
|--------------------|--|--------|------|-------|--------|------|----------|-------|
| | 2D | | | 3D | | | | |
| | Total | Female | Male | Total | Female | male | | |
| A (n=4*5) | 14 | 3 | 11 | 16 | 4 | 12 | 0.533 | 0.465 |
| B (n=5*5) | 23 | 9 | 14 | 22 | 9 | 13 | 0.222 | 0.637 |
| C (n=6*5) | 27 | 4 | 23 | 28 | 5 | 23 | 0.218 | 0.640 |

a: According to Table 51, each trainee answers 5 questions about basic knowledge, so the total number of questions answered varies for each group: 20 for group A, 25 for group B and 30 for group C




Fig. 3 Number of correct responses on the baseline knowledge assessment across different groups with different genders

one patient and one healthy volunteer and their imaging data were anonymized.

Results

No improvement of trainees' understanding of the base knowledge

All the trainees had completed relatively standardized clinical theory courses and were given a systematic basic knowledge review together before the training. Regarding Questionnaire number 1, there was no significant difference in the number of correct responses between the different groups (Table 3). All the trainees showed good performance in answering the physiological and anatomical knowledge questions about the liver, and there was no significant difference between different training models and different genders (Fig. 3).

Significant improvement of trainees' understanding of the key points of surgery

The trainees were asked to identify the segment(s) in which the tumour was located. The combination of 3DP models with 2D CT scans led to an improvement in tumour allocation to the liver segment in all the groups ($p < 0.05$, respectively). The trainees with fundamental experience (Group B+C) showed better performance than the interns (Group A) the combination of 3DP model training ($p < 0.05$) (Table 4). In contrast, the trainees may have recognized a tumour situated within

segment VII but would fail to demonstrate exactly where the tumour is localized in a liver model.

The responses of trainers of different genders in each group were compared separately. We found that in group C, the 3DP model was more helpful in improving male trainees' understanding of the key points of surgical procedures ($p < 0.05$). However, this difference was not observed in the other two groups (Fig. 4).

The trainees were asked to draw their resection proposal line for each patient in the 3DP realistic liver model. During this test, the trainees were asked to give the minimal resection proposal, including the tumour, the safety margin, and the dependent liver tissue. Alternatively, the trainees could also proceed in a classic way by resecting the whole liver segment(s). We calculated the average percentage of the correct target area found and the number of answers that achieved more than 80% of the actual target area composed of the tumour, safety margin, and dependent liver tissue. The results are stratified according to the groups. A significant and measurable improvement could be demonstrated for the 3D presentations.

Evaluation of the field with help to the trainees

The major fields were analyzed in which 3DP models provided more help to the trainees. Although more trainees considered 3DP models helpful for their understanding of basic knowledge, including the liver itself and the tumour, no significant difference was found between 3DP models and 2D. In contrast, in the field associated with surgery and surgical details, 3DP models showed a significant advantage effect on the trainees, such as in planned surgical procedures ($p < 0.05$) and key details of the surgical procedures ($p < 0.05$) (Table 5; Fig. 5).

Evaluation of the monitors' satisfaction level with the trainees' cooperation in the operation

The control degree of basic knowledge and the assistance level during the operations were evaluated by the monitors. For the trainees given the hand-on 3DP model training, their supervisors gave them a better evaluation of their mastery degree of basic knowledge ($p < 0.05$). The supervisors also gave them better evaluation performance in assisted surgery ($p < 0.05$) (Table 6; Fig. 6).

Table 4 Different levels of trainees' understanding assessment of the key points of surgery between different training models

| Group ^a | Number of "No difficult responses" with different genders | | | | | | X ² | P |
|--------------------|---|--------|------|-------|--------|------|----------------|-------|
| | 2D | | 3D | | | | | |
| | Total | Female | Male | Total | Female | Male | | |
| A (n = 4*7) | 17 | 2 | 15 | 23 | 5 | 18 | 4.139 | 0.042 |
| B (n = 5*7) | 27 | 10 | 17 | 33 | 13 | 20 | 4.200 | 0.040 |
| C (n = 6*7) | 28 | 4 | 24 | 36 | 4 | 32 | 4.200 | 0.040 |

a: According to Table S1, each trainee answers 7 questions about the key points of surgery, so the total number of questions answered varies for each group: 28 for group A, 35 for group B and 42 for group C




Fig. 4 Number of "No difficult responses" on the key points of surgery of different groups with different genders

Table 5 Median score of the evaluation with help to the trainees

| Items | Median [Min-Max] score of trainee's responses | | t/Z | p |
|--|---|---------|-------|--------------------|
| | 2D | 3D | | |
| Basic liver physiology and anatomy | 5 [4–6] | 6 [5–7] | 3.333 | 0.068 |
| Disease and tumour characteristics | 6[4–7] | 6[5–7] | 1.154 | 0.283 ^a |
| planned surgical procedure | 5[4–7] | 6[6–8] | 3.968 | 0.046 |
| key details of the surgical procedures | 5[4–7] | 7[4–7] | 5.000 | 0.025 |

The score was given using a visual scale from 1 to 10 with 1 = "no help at all" and 10 = "of a great help" ("Appendix 2"). For each cognitive component, the score of the trainee's responses reflects the level of understanding

^a Wilcoxon tests

Evaluation of the trainees' satisfaction level on the training project

Finally, we analyzed the evaluation of the trainees on the training project. According to Questionnaire number 4, we found that the combination of 3DP model trainees had higher satisfaction with the teaching course in all groups ($p < 0.05$). For trainees with different experience levels, their teaching satisfaction was also significantly improved with the combination of 3DP liver models with conventional 2D scans (Table 7). Most trainees agreed with the usage of the 3DP model again and agreed with its usage as a training and testing tool. Furthermore, 73% (11/15) of trainees considered that they benefitted from the project technically. The satisfaction of trainers of different genders in each group was compared separately. The results showed that gender had no effect on satisfaction evaluation in the different training models across all trainers (Fig. 7).

Discussion

3D printing includes a wide range of technologies and applications in the fields of health care and medical education [46]. Successive improvements in 3D imaging and 3DP models have progressively led various surgical specialties to embrace these innovative technologies,




Fig. 5 Score of correct responses per trainee, conventional 2D CT scans and 3DP model presentation. Individual analysis of trainees' understanding improvement (number of correct responses) in four areas: A basic liver physiology and anatomy; B disease and tumour characteristics; C planned surgical procedure; D key details of the surgical procedures. A p value less than 0.05 was considered statistically significant

Table 6 The median score of the monitors' satisfaction level for the trainees

| Items | Median [Min-Max] score of monitors' satisfaction | | t/Z | p |
|---------------------------------|--|---------|-------|--------------------|
| | 2D | 3D | | |
| Master level of basic knowledge | 7[5–9] | 8[7–10] | 3.968 | 0.046 |
| Performance in assisted surgery | 4[2–7] | 6[3–9] | 5.000 | 0.025 ^a |

The score was given using a visual scale from 1 to 10 with 1 = "No satisfaction" and 10 = "Of great satisfaction" ("Appendix 3"). For each evaluation item, the score among the group of trainees reflects the monitors' satisfaction level

^a Wilcoxon tests

especially in the field of reconstructive surgery and SBME [47, 48]. SBME has recently received a great deal of attention due to the increasing use of 3D printing technology and 3DP models in continuing medical education and clinical training [49]. Understanding complex anatomical pathology and pre- and intraoperative sets represents a critical challenge for clinical medical education [50, 51]. 3DP models can provide precise and individualized 1:1 physical models that can help trainees easily understand actual liver tumour characteristics, including tumour size, depth, and location with respect to the arteriovenous and bile duct systems, thereby contributing to




Fig. 6 Score of correct responses per trainee, conventional 2D CT scans and 3DP model presentation. Individual analysis of monitors' satisfaction on two areas: **A** master's level of basic knowledge; **B** performance in assisted surgery; **A** p value less than 0.05 was considered statistically significant

preoperative surgical procedure simulation and intraoperative guidance [52].

First, in the present study, based on SBME theory, we successfully produced a full-sized physical 3DP liver model for continuing medical education evaluation using a solid transparent material, in which context the tumour and other features can be seen from the outside. Our subjective results indicated overall improvement and better performance in assisted surgery in terms of both test scores and questionnaires following the 3DP liver model training. The post-training test results of the 3D group were higher than those of the conventional 2D group. This finding is consistent with the results of previous studies [1, 49, 53]. Langridge et al. [54] used descriptive statistical methods to report the role of 3D printed models in surgical education. However, there was no difference in the mastery of basic knowledge between the two groups. Li et al. [55] did not find any significant difference ($p=0.0508$) in knowledge acquisition between 3DP models and manipulatable 3D imaging. In contrast, Lim et al. [56] found that 3DP models were superior to cadavers with regard to learning external cardiac anatomy. Further evidence is thus required to determine which aspects of the 3DP models facilitate learning and whether these models could provide a superior alternative to cadavers in traditional anatomical teaching [54]. A unique feature of this study is that both self-assessment by questionnaire and rating by monitors were used to evaluate the effect of training.

Second, in addition to facilitating knowledge acquisition with regard to anatomy, 3DP liver models can also be used for surgical procedures, as they convey the

anatomical relationships between organs and the surrounding tissues [57]. Accordingly, our results demonstrate that the use of 3DP liver models can significantly improve all trainees' understanding of the key points of surgery, especially for experienced male trainees. This finding further reflects the value of using 3DP liver models in continuing medical education and clinical training [58]. The evaluation of the questionnaires also showed that the participants were very satisfied overall with training on the 3DP models. Participants of all levels of experience agreed that they would benefit from such training opportunities in the future, thereby confirming the acceptance of this type of training. It was also found that providing trainees with full-sized realistic 3DP models of the liver is superior to 2D anatomy teaching methods, which is consistent with the findings of previous studies [32, 33]. The use of 3D printing technology to develop surgical simulation models and training devices has proven to offer valuable resources to clinical trainees in surgical training exercises, contribute to understanding surgical procedures, and improve surgical performance in terms of enhancing training quality and reducing learning curves [34, 59].

Finally, we investigated the gender of the trainees as an influencing factor of the training effect, and the results showed that after training with the 3DP liver models, male trainees performed better in terms of their understanding of base knowledge than female trainees, especially in the group of interns; however, there were no significant differences in the group of trainees in standardized training and the group of trainees in professional training. Vaccarezza M [60] reported that both

Table 7 Evaluation of the trainees' satisfaction level with the training project

| Group ^a | Number of correct responses with different genders | | | | | | χ ² | p |
|--------------------|--|--------|------|-------|--------|------|----------------|-------|
| | 2D | | | 3D | | | | |
| | Total | Female | Male | Total | Female | Male | | |
| A (n = 4*4) | 9 | 2 | 7 | 14 | 3 | 11 | 3.865 | 0.049 |
| B (n = 5*4) | 12 | 5 | 7 | 18 | 7 | 11 | 4.800 | 0.028 |
| C (n = 6*44) | 18 | 3 | 15 | 22 | 4 | 18 | 4.547 | 0.033 |

a: According to Table S2, each trainee answers 4 questions about the training project, so the total number of questions answered varies for each group: 16 for group A, 20 for group B and 24 for group C

male and female students spent less time answering questions regarding base knowledge on the spine models in the 3D group than did a conventional group. The different results found in the above research may be due to the variations in trainees experience and organs. The 3DP model was more helpful in improving male trainees' understanding of the key points of surgical procedures and performed better in terms of cooperation with the surgical procedure. Although the number of participants included in this study was small, the results are consistent with those reported by previous studies [41, 61–62]. These findings may be due to the fact that female trainees exhibited less procedural experience and lower confidence with regard to performing procedural skills than their male counterparts despite their equal or superior performance [62–64]. In a study of fourth-year medical students, females reported overall lower levels of experience and confidence performing technical skills than males [41]. Moreover, some study also indicated that gender was an independent factor affecting spatial learning ability, a male advantage in visual-spatial working memory [39] and show faster spatial learning in a virtual environment as compared to females [40].

Compared with the recently developed multiuser virtual reality application [65], 3DP liver models offer different advantages with regard to improving resection planning for vascular reconstruction localization of resection plains, thus reducing the risks for devascularization and complications and facilitating intraoperative detection of small and deeply located tumours [66]. The study reflected the potential benefits that 3DP liver models offer in clinical medicine education compared to traditional educational tools. The results proved that 3DP models are highly suitable for training purposes in this field [67–69]. Skill-specific simulation training and other interventions may improve skill development in medical students in light of the obstacles they face when developing proficiency in the clinical setting [70].

Limitations

However, both this study and the 3D printed training model have certain limitations. First, with regard to the design and statistical analysis of the study, the nonrandomized controlled trial and the self-reported nature of the data are the two biggest limitations. Meanwhile, this study only included thirty trainees, a sufficiently robust sample size has not been achieved, the number of participants in each group was relatively small, and the levels of anatomical education exhibited by the participants in the studies were not assessed prior to using the models, which may have introduced bias; Thus, a large-scale trial is necessary in the future. The results and findings of the study presented here should be viewed as preliminary data to support a larger and more comprehensive




Fig. 7 Number of correct responses of the trainees' satisfaction level on the training project of different groups with different genders

study featuring high-level medical training. This research could be enhanced by the inclusion of more longitudinal follow-up of participants who are exposed to 3DP models to assess their long-term skill acquisition.

Second, existing simulator 3DP models have several limitations. The most common is the material properties of the liver analogue are not physically accurate, and 3DP models have been unable to fully simulate the differences in deformation elasticity and biomechanical characteristics of human tissues. Therefore, the loss of these characteristics may limit the model's ability to reflect both the mechanical characteristics and surface smoothness of normal tissue and may significantly affect the perceptions of and feedback provided by trainees. Furthermore, although these models replicate the procedure well, there are still few liver disease models; for example, the background of liver cirrhosis could not be effectively illustrated by the current printed models. 3D printing technology can only capture major vasculature structures [71]. Small vessels and perforators cannot be illustrated with current technology. In addition, most importantly, only a few typical liver models were printed in this study, and more complex models were not presented because of their high costs [72, 73]; since such models are an emerging technology, this situation is also a barrier to widespread clinical adoption.

Conclusion

We have successfully created a liver 3DP model to help improve the learning effect for standardized residency training trainees and junior residents in terms of practical skills and theoretical understanding. This study supports the claim that training with 3DP models could facilitate all trainees' presurgical understanding of liver tumours and surgery. Meanwhile, 3DP liver models of

real production can serve as an effective auxiliary teaching tool for SBME teaching of Chinese standardized residency training and continuing medical education.

Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12909-023-04535-3>.

Supplementary Material 1

Acknowledgements

The authors would like to thank the students involved in the study.

Author contributions

Contributions: (I) Conception and design: Zhenyu Yang; (II) Administrative support: Zhenyu Yang and Kejun Ma; (III) Provision of study materials: Jiangpu Yi, Guoqiang Bao and Haoran Li; (IV) Collection and assembly of data: Dian Guo, Jiahe Liang and Yajie Li; (V) Data analysis and interpretation: Shujia Peng and Ping Yang; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Funding

This study was funded by the innovation programs of the Second Affiliated Hospital of Air Force Medical University (No. 2020XKPT010 and No. 2020XKPT013).

Data availability (ADM)

All data generated or analyzed during this study are included in this published article and its supplementary information files.

Declarations

Ethics approval and consent to participate

The authors are accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved. All trainees received oral information and written informed consent about the objectives and methodology of the study. All CT images were anonymized before use for 3D modeling. This study was approved by the Ethics Committee of the Tangdu Hospital of Air Force Medical University (No. K202207-06) and conducted according to the standards of the Declaration of Helsinki.

Consent for publication

Not applicable.

Competing interests

All authors declare that they have no competing interests.

Author details

¹Department of General Surgery, The Second Affiliated Hospital of Air Force Medical University, Xi'an, Shaanxi 710038, China

²3D Printing Research Center of Tangdu Hospital, Air Force Medical University, Xi'an, Shaanxi, China

³Xi'an Ma Ke Medical Technology Ltd, Room 21516, Block C, Chaoyang International Plaza, Xi'an, Shaanxi, China

Received: 17 November 2022 / Accepted: 24 July 2023

Published online: 15 August 2023

References

1. Seymour NE, Gallagher AG, Roman SA et al. Virtual reality training improves operating room performance: results of a randomized, double-blinded study. *Ann Surg*. 2002; 236(4): 458 – 63; discussion 463–4.
2. Akaiki M, Fukutomi M, Nagamune M, et al. Simulation-based medical education in clinical skills laboratory. *J Med Invest*. 2012;59(1–2):28–35.
3. Moll-Khosrawi P, Zöllner C, Cronje JS, Schulte-Uentrop L. The effects of simulation-based education on medical students' motivation. *Int J Med Educ*. 2021;12:130–5.
4. Richter M, Peter T, Rüttermann S, Sader R, Seifert LB. 3D printed versus commercial models in undergraduate conservative dentistry training. *Eur J Dent Educ*. 2022;26(3):643–51.
5. Rochlen LR, Putnam E, Levine R, Tait AR. Mixed reality simulation for peripheral intravenous catheter placement training. *BMC Med Educ*. 2022;22(1):876.
6. Goharnejad S, Goharnejad S, Hajesmaeel-Gohari S, Bahaadinbeigy K. The usefulness of virtual, augmented, and mixed reality technologies in the diagnosis and treatment of attention deficit hyperactivity disorder in children: an overview of relevant studies. *BMC Psychiatry*. 2022;22(1):4.
7. Lawson S, Reid J, Morrow M, Gardiner K. Simulation-based education and human factors training in Postgraduate Medical Education: A Northern Ireland Perspective. *Ulster Med J*. 2018;87(3):163–7.
8. Tenegra JC, Hoffman MR, Mueller Luckey GS, DiLalla LF, Ledford C. Simulation-Based Medical Education in Family Medicine Residencies: a CERA study. *Fam Med*. 2022;54(4):264–9.
9. Liu C, Liu W, Jiao M, et al. A combined behavioural economics- and simulation-based medical education to promote effectiveness among medical residents in coping with workplace violence in Northern China: a quasi-experimental study. *BMC Public Health*. 2022;22(1):1090.
10. Michiels C, Jambon E, Sarrazin J, et al. [Comprehensive review of 3D printing use in medicine: comparison with practical applications in urology]. *Prog Urol*. 2021;31(12):762–71.
11. Brunner BS, Thierij A, Jakob A, et al. 3D-printed heart models for hands-on training in pediatric cardiology - the future of modern learning and teaching. *GMS J Med Educ*. 2022;39(2):Doc23.
12. Bangeas P, Tsioukas V, Papadopoulos VN, Tsoulfas G. Role of innovative 3D printing models in the management of hepatobiliary malignancies. *World J Hepatol*. 2019;11(7):574–85.
13. Su W, Xiao Y, He S, Huang P, Deng X. Three-dimensional printing models in congenital heart disease education for medical students: a controlled comparative study. *BMC Med Educ*. 2018;18(1):178.
14. Comeau M, Goudie C, Murphy D, Fowler E, Dubrowski A. The Use of Silicone Vaginal Repair Models as an Adjunct to Mannequins for Simulation Training in sexual assault clinical learning for obstetrics and Gynecology Medical residents. *Cureus*. 2020;12(3):e7410.
15. Joo MW, Lee YS, Chung YG, Lee HK. Sarcomas in Teachers using Three-Dimensional Printers: a report of three patients and literature review. *Clin Orthop Surg*. 2022;14(2):310–7.
16. Schmauss D, Haerberle S, Hagl C, Sodan R. Three-dimensional printing in cardiac surgery and interventional cardiology: a single-centre experience. *Eur J Cardiothorac Surg*. 2015;47(6):1044–52.
17. Xie Y, Wu G, Liang Y, Fan G. Three-Dimensional Physical Model in Urologic Cancer. *Front Surg*. 2022;9:757337.
18. Loke YH, Harahsheh AS, Krieger A, Olivieri LJ. Usage of 3D models of tetralogy of Fallot for medical education: impact on learning congenital heart disease. *BMC Med Educ*. 2017;17(1):54.
19. Sutherland J, Belec J, Sheikh A, et al. Applying modern virtual and augmented reality Technologies to Medical images and models. *J Digit Imaging*. 2019;32(1):38–53.
20. Nestic D, Schaefer BM, Sun Y, Saulacic N, Sailer I. 3D Printing Approach in Dentistry: the future for personalized oral soft tissue regeneration. *J Clin Med*. 2020. 9(7).
21. Yamazaki M, Takayama T, Fujita A, et al. 3D printed kidney model could be an important educational tool for residents. *Asian J Endosc Surg*. 2023;16(2):197–202.
22. Hammerton C, Yip S, Manobharath N, Myers G, Sturrock A. Are 3D printed models acceptable in assessment. *Clin Teach*. 2022;19(3):221–8.
23. Nicot R, Druelle C, Schlund M, et al. Use of 3D printed models in student education of craniofacial traumas. *Dent Traumatol*. 2019;35(4–5):296–9.
24. Meglioli M, Naveau A, Macaluso GM, Catros S. 3D printed bone models in oral and cranio-maxillofacial surgery: a systematic review. *3D Print Med*. 2020;6(1):30.
25. Maggio LA, Tannery NH, Chen HC, ten Cate O, O'Brien B. Evidence-based medicine training in undergraduate medical education: a review and critique of the literature published 2006–2011. *Acad Med*. 2013;88(7):1022–8.
26. Török G, Erdei Z, Lilienberg J, Apáti Á, Homolya L. The importance of transporters and cell polarization for the evaluation of human stem cell-derived hepatic cells. *PLoS ONE*. 2020;15(1):e0227751.
27. Cai J, Jiang G, Liang Y, Xie Y, Zheng J, Liang X. Safety and effectiveness evaluation of a two-handed technique combining harmonic scalpel and laparoscopic Peng's multifunction operative dissector in laparoscopic hemihepatectomy. *World J Surg Oncol*. 2021;19(1):198.
28. van Leeuwen MS, Fernandez MA, van Es HW, Stokking R, Dillon EH, Feldberg MA. Variations in venous and segmental anatomy of the liver: two- and three-dimensional MR imaging in healthy volunteers. *AJR Am J Roentgenol*. 1994;162(6):1337–45.
29. Frericks BB, Caldarone FC, Nashan B, et al. 3D CT modeling of hepatic vessel architecture and volume calculation in living donated liver transplantation. *Eur Radiol*. 2004;14(2):326–33.
30. Lamadé W, Glombitza G, Fischer L, et al. The impact of 3-dimensional reconstructions on operation planning in liver surgery. *Arch Surg*. 2000;135(11):1256–61.
31. Raichurkar KK, Lochan R, Jacob M, Asthana S. The Use of a 3D Printing Model in planning a Donor Hepatectomy for living Donor Liver transplantation: first in India. *J Clin Exp Hepatol*. 2021;11(4):515–7.
32. Smith B, Dasgupta P. 3D printing technology and its role in urological training. *World J Urol*. 2020;38(10):2385–91.
33. Knoedler M, Feibus AH, Lange A, et al. Individualized physical 3-dimensional kidney tumor models constructed from 3-dimensional printers result in Improved Trainee Anatomic understanding. *Urology*. 2015;85(6):1257–61.
34. Fan G, Li J, Li M, et al. Three-dimensional physical model-assisted planning and Navigation for Laparoscopic partial nephrectomy in patients with endophytic renal tumors. *Sci Rep*. 2018;8(1):582.
35. Klein R, Julian KA, Snyder ED, et al. Gender Bias in Resident Assessment in Graduate Medical Education: review of the literature. *J Gen Intern Med*. 2019;34(5):712–9.
36. Klein R, Koch J, Snyder ED, et al. Association of gender and Race/Ethnicity with Internal Medicine In-Training examination performance in Graduate Medical Education. *J Gen Intern Med*. 2022;37(9):2194–9.
37. Ali A, Subhi Y, Ringsted C, Konge L. Gender differences in the acquisition of surgical skills: a systematic review. *Surg Endosc*. 2015;29(11):3065–73.
38. Sanghvi R. Gender perspectives in medical education. *Indian J Med Ethics*. 2019;4(2):148–53.
39. Voyer D, Voyer SD, Saint-Aubin J. Sex differences in visual-spatial working memory: a meta-analysis. *Psychon Bull Rev*. 2017;24(2):307–34.
40. Piber D, Nowacki J, Mueller SC, Wingenfeld K, Otte C. Sex effects on spatial learning but not on spatial memory retrieval in healthy young adults. *Behav Brain Res*. 2018;336:44–50.
41. Barr J, Graffeo CS. Procedural experience and confidence among Graduating Medical Students. *J Surg Educ*. 2016;73(3):466–73.
42. Brown DK, Fosnight S, Whitford M, et al. Interprofessional education model for geriatric falls risk assessment and prevention. *BMJ Open Qual*. 2018;7(4):e000417.

43. Hu X, Liu L, Xu Z, et al. Creation and application of war trauma treatment simulation software for first aid on the battlefield based on undeformed high-resolution sectional anatomical image (chinese visible human dataset). *BMC Med Educ*. 2022;22(1):498.
44. Ghadiri ZND, Fardanesh H. Investigation of the effect of computer simulation-based learning based on constructivism in removing physics misconception. *Interdiscip J Virtual Learn Med Sci*. 2015;6(4):e11997.
45. Li JLY, Bao H. An empirical study of the virtual Simulation System Teaching Method in NC Machining. *Int J Technol Human Interact*. 2020;16(3):109–23.
46. Anagnostopoulos S, Gallos P, Zoulias E, Fotos N, Mantas J. 3D Digital Printing in Healthcare: Technologies, Applications and Health Issues. *Stud Health Technol Inform*. 2022;295:394–7.
47. Cohen A, Laviv A, Berman P, Nashef R, Abu-Tair J. Mandibular reconstruction using stereolithographic 3-dimensional printing modeling technology. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod*. 2009;108(5):661–6.
48. Duncan JM, Nahas S, Akhtar K, Daurka J. The Use of a 3D printer in Pre-operative Planning for a patient requiring Acetabular reconstructive surgery. *J Orthop Case Rep*. 2015;5(1):23–5.
49. Ye Z, Dun A, Jiang H, et al. The role of 3D printed models in the teaching of human anatomy: a systematic review and meta-analysis. *BMC Med Educ*. 2020;20(1):335.
50. Patera E, Rust PA. Creation of 3D anatomical models illustrating an intact and centrally torn triangular fibrocartilage complex for patient education prior treatment. *Ann Anat*. 2022;240:151854.
51. Holdsworth C, Skinner EH, Delany CM. Using simulation pedagogy to teach clinical education skills: a randomized trial. *Physiother Theory Pract*. 2016;32(4):284–95.
52. Javan R, Herrin D, Tangestanipoor A. Understanding spatially Complex Segmental and Branch anatomy using 3D Printing: liver, lung, prostate, coronary arteries, and Circle of Willis. *Acad Radiol*. 2016;23(9):1183–9.
53. Jones DB, Sung R, Weinberg C, Korelitz T, Andrews R. Three-Dimensional modeling May improve Surgical Education and Clinical Practice. *Surg Innov*. 2016;23(2):189–95.
54. Langridge B, Momin S, Coumbe B, Woin E, Griffin M, Butler P. Systematic review of the Use of 3-Dimensional Printing in Surgical Teaching and Assessment. *J Surg Educ*. 2018;75(1):209–21.
55. Li Z, Li Z, Xu R, et al. Three-dimensional printing models improve understanding of spinal fracture—A randomized controlled study in China. *Sci Rep*. 2015;5:11570.
56. Lim KH, Loo ZY, Goldie SJ, Adams JW, McMenamin PG. Use of 3D printed models in medical education: a randomized control trial comparing 3D prints versus cadaveric materials for learning external cardiac anatomy. *Anat Sci Educ*. 2016;9(3):213–21.
57. Facco G, Massetti D, Coppa V, et al. The use of 3D printed models for the pre-operative planning of surgical correction of pediatric hip deformities: a case series and concise review of the literature. *Acta Biomed*. 2022;92(6):e2021221.
58. Chandran J, Balakrishnan N, Sreenivasagan S. Awareness on three-dimensional printing of orthodontic appliances among dental students. *J Adv Pharm Technol Res*. 2022;13(Suppl 2):S63–S567.
59. Lemarteleur V, Peycelon M, Sablayrolles JL, Plaisance P, El-Ghoneimi A, Ceccaldi PF. Realization of Open Software Chain for 3D modeling and Printing of Organs in Simulation Centers: Example of Renal Pelvis Reconstruction. *J Surg Educ*. 2021;78(1):232–44.
60. Vaccarezza M, Papa V. 3D printing: a valuable resource in human anatomy education. *Anat Sci Int*. 2015;90(1):64–5.
61. Ryan JF, Istl AC, Luhoway JA, et al. Gender disparities in Medical Student Surgical Skills Education. *J Surg Educ*. 2021;78(3):850–7.
62. Gude T, Finset A, Anvik T, et al. Do medical students and young physicians assess reliably their self-efficacy regarding communication skills? A prospective study from end of medical school until end of internship. *BMC Med Educ*. 2017;17(1):107.
63. Madrazo L, Lee CB, McConnell M, Khamisa K. Self-assessment differences between genders in a low-stakes objective structured clinical examination (OSCE). *BMC Res Notes*. 2018;11(1):393.
64. Lind DS, Rekkas S, Bui V, Lam T, Beierle E, Copeland EM 3. Competency-based student self-assessment on a surgery rotation. *J Surg Res*. 2002;105(1):31–4.
65. Boedecker C, Huettl F, Saalfeld P, et al. Using virtual 3D-models in surgical planning: workflow of an immersive virtual reality application in liver surgery. *Langenbecks Arch Surg*. 2021;406(3):911–5.
66. Huber T, Huettl F, Tripke V, Baumgart J, Lang H. Experiences with three-dimensional Printing in Complex Liver surgery. *Ann Surg*. 2021;273(1):e26–7.
67. Boshra M, Godbout J, Perry JJ, Pan A. 3D printing in critical care: a narrative review. *3D print Med*. 2020. 6(1): 28.
68. He J, Wang J, Pang Y, et al. Bioprinting of a hepatic tissue model using Human-Induced Pluripotent Stem Cell-derived hepatocytes for Drug-Induced Hepatotoxicity evaluation. *Int J Bioprint*. 2022;8(3):581.
69. Li L, Zhang K, Wang R, et al. A study of three-dimensional reconstruction and printing models in two cases of soft tissue sarcoma of the thigh. *Int J Comput Assist Radiol Surg*. 2021;16(9):1627–36.
70. Luhoway JA, Ryan JF, Istl AC, et al. Perceived Barriers to the development of Technical Skill Proficiency in Surgical Clerkship. *J Surg Educ*. 2019;76(5):1267–77.
71. Ryan JR, Almefty KK, Nakaji P, Frakes DH. Cerebral aneurysm clipping surgery Simulation using patient-specific 3D Printing and Silicone Casting. *World Neurosurg*. 2016;88:175–81.
72. Perica ER, Sun Z. A systematic review of Three-Dimensional Printing in Liver Disease. *J Digit Imaging*. 2018;31(5):692–701.
73. Yang T, Lin S, Xie Q, et al. Impact of 3D printing technology on the comprehension of surgical liver anatomy. *Surg Endosc*. 2019;33(2):411–7.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.