

POSTER PRESENTATION

Open Access

Frovatriptan vs almotriptan for treatment of menstrual migraine: a double-blind, randomized, cross-over, multicenter Italian study

M Bartolini^{1*}, MA Giamberardino², C Lisotto³, P Martelletti⁴, D Moscato⁴, B Panascia⁵, L Savi⁶, LA Pini⁷, G Sances⁸, P Santoro⁹, G Zanchin¹⁰, S Omboni¹¹, MD Ferrari¹², B Fierro¹³, F Brighina¹³

From The European Headache and Migraine Trust International Congress
London, UK. 20-23 September 2012

Objective

To compare the efficacy and safety of frovatriptan and almotriptan in women with menstrually related migraine (IHS Classification of Headache disorders) enrolled in a multicenter, randomized, double blind, cross-over study.

Methods

Patients received frovatriptan 2.5 mg or almotriptan 12.5 mg in a randomized sequence: after treating 3 episodes of migraine in no more than 3 months with the first treatment, the patient switched to the other treatment.

Results

67 of the 96 female patients of the intention-to-treat population of the main study had regular menstrual cycles and were thus included in this subgroup analysis. 77 migraine attacks classified as related to menses were treated with frovatriptan and 78 with almotriptan. Rate of pain relief at 2- and 4-hrs was 36% and 53% for frovatriptan and 41% and 50% for almotriptan ($p=NS$ between treatments). Rate of pain free at 2- and 4-hrs was 19% and 47% with frovatriptan and 29% and 54% for almotriptan ($p=NS$). At 24-hrs, 62% of frovatriptan- and 67% of almotriptan-treated patients had pain relief, while 60% vs. 67% were pain free ($p=NS$). Recurrence at 24-hrs was significantly ($p<0.05$) lower with frovatriptan (8% vs. 21% almotriptan). This was the case also at 48-hrs (9% vs. 24%, $p<0.05$).

Conclusions

Frovatriptan was as effective as almotriptan in the immediate treatment of menstrually related migraine

attacks. However, it showed a more favorable sustained effect, as shown by a lower rate of migraine recurrence.

Author details

¹Clinica Neurologica, Ospedali Riuniti, Università Politecnica delle Marche, Ancona, Italy. ²Dipartimento di Medicina e Scienze dell'Invecchiamento, Università "G. D'Annunzio", Chieti, Italy. ³Ospedale Civile San Vito al Tagliamento, Pordenone, Italy. ⁴UOS Cefalea, Ospedale S. Andrea, Università La Sapienza, Roma, Italy. ⁵Centro Cefalee, A.O. Universitaria Vittorio Emanuele, Catania, Italy. ⁶Department of Neurology, University of Torino, Italy. ⁷Centro Cefalee, Università degli Studi di Modena e Reggio Emilia, Reggio Emilia, Italy. ⁸Centro Cefalee, IRCCS Neurologico C. Mondino, Pavia, Italy. ⁹Clinica Neurologica, Ospedale S. Gerardo, Monza, Italy. ¹⁰Department of Neurology, University of Padova, Italy. ¹¹Italian Institute of Telemedicine, Varese, Italy. ¹²Leiden Centre for Translational Neuroscience, Department of Neurology, Leiden University Medical Centre, Netherlands. ¹³Department of Neurology and Psychiatry, University of Palermo, Palermo, Italy.

Published: 21 February 2013

Reference

1. Bartolini M, Giamberardino MA, Lisotto C, Martelletti P, Moscato D, Panascia B, Savi L, Pini LA, Sances G, Santoro P, Zanchin G, Omboni S, Ferrari MD, Brighina F, Fierro B: **Frovatriptan versus almotriptan for acute treatment of menstrual migraine: analysis of a double-blind, randomized, cross-over, multicenter, Italian, comparative study.** *J Headache Pain* 2012.

doi:10.1186/1129-2377-14-S1-P192

Cite this article as: Bartolini et al.: Frovatriptan vs almotriptan for treatment of menstrual migraine: a double-blind, randomized, cross-over, multicenter Italian study. *The Journal of Headache and Pain* 2013 **14**(Suppl 1):P192.

¹Clinica Neurologica, Ospedali Riuniti, Università Politecnica delle Marche, Ancona, Italy

Full list of author information is available at the end of the article