

RESEARCH NOTE

Open Access



# Difference in risk of preterm and small-for-gestational-age birth depending on maternal occupations in Japan

Tasuku Okui<sup>1\*</sup>

## Abstract

**Objectives** In this study, an association between the mother's occupations with preterm and small-for-gestational-age (SGA) births was investigated using national data in Japan, and individual-level birth data from the Report of Vital Statistics: Occupational and Industrial Aspects in the 2015 fiscal year were used. Preterm and SGA birth rates were calculated for each of infant characteristics, and relative risk of each type of maternal occupations (categorized into 12 types) for the outcomes was estimated using a log binomial regression model.

**Results** Data of 997,600 singleton births were analyzed. Among maternal occupations, preterm birth rate was highest among carrying, cleaning, packaging, and related workers (5.65%) and lowest among security workers (4.24%). SGA birth rate was highest among manufacturing process workers (5.91%) and lowest among security workers (4.00%). We found significantly elevated risks for preterm birth among manufacturing process workers compared with unemployed mothers, and significantly elevated risks for SGA birth compared with unemployed mothers were observed among sales workers, service workers, and manufacturing process workers. In contrast, security workers had a significantly decreased risk for SGA birth compared with unemployed mothers.

**Keywords** Small-for-gestational-age, Preterm birth, Japan, Maternal occupations

## Introduction

Preterm birth and small-for-gestational-age (SGA) in infants is a major adverse birth outcome. SGA rate differs largely depending on regions and tends to be high in developing countries [1, 2]. Although preterm birth rate globally stands at approximately 10% [2], the rate in Japan is much lower [3]. However, preterm birth is a risk factor for perinatal and infant mortality [4, 5], and adversely effects neurodevelopment or the incidence of

cardiovascular diseases [6]. Infants with SGA, which is the other adverse birth outcome and a focus of epidemiological studies [7, 8], have an elevated risk of delayed development or childhood mortality [9–11]. Therefore, prevention of these adverse birth outcomes is a basic medical and societal need.

It is known that some characteristics of mothers, such as maternal age, smoking habits, and maternal socioeconomic status, are risk factors of adverse birth outcomes [12–14]. In Japan, epidemiological studies have shown that the mother's pre-pregnancy body mass index (BMI) and the maternal level of education were risk factors for preterm and SGA births [15–18]. In addition, occupation is another major socioeconomic characteristic of parents. The association between maternal occupation

\*Correspondence:

Tasuku Okui  
task10300@gmail.com

<sup>1</sup>Medical Information Center, Kyushu University Hospital, Fukuoka city, Japan



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

and outcomes, such as infant mortality or low birth weight, has been investigated in Japan [19, 20]. However, there has been no systematic study for the relationship between the mother's occupations and preterm and SGA births, using nationwide government statistics data in Japan. One study on maternal occupations in Sweden showed that "mechanics and iron and metalware workers" had a higher risk of giving birth to infants with SGA [21], and difference was also investigated and observed also in Korea and the United States [22, 23]. By identifying occupations that confer high risk for these adverse birth outcomes to parents also in Japan, we might be able to provide targeted support to people with those occupations at their workplace or in their communities.

To address this goal, we used the Vital Statistics data in Japan to investigate the association between maternal occupations and preterm and SGA births.

## Main text

### Methods

#### *Data source and data preprocessing*

We obtained nationwide individual-level data of the Report of Vital Statistics: Occupational and Industrial Aspects in the 2015 fiscal year from the Japanese Ministry of Health, Labor, and Wealth. Use of the data for research was permitted by the Ministry based on Article 33 of the Statistics Act. The survey for the report is conducted every 5 years, and the data include infants which were born between April 1, 2015 and March 31, 2016. The Report corresponds to the Vital Statistics with occupational and industry information. Vital Statistics is a survey by the Japanese government, and information from the birth certificates of all over the country are gathered by the Ministry of Health, Labor, and Wealth. It is mandatory for a parent to submit birth certificates of their children to a municipality within 14 days after childbirth. The birth certificate is a self-administered questionnaire, while health-related information, such as birthweight and gestational age, are written by physicians or midwives at the time of childbirth. The same data have been analyzed in previous studies in Japan [19, 24], and the maternal occupations at the time of birth were surveyed. We used the birth data, which included maternal age, maternal nationality, wedlock status, sex, birth weight, number of fetuses, gestational age, parity, household occupation, and mother's occupation for each infant. Categories for maternal occupation in the data were 12 types, which was based on the Standard Occupational Classification for Japan. Household occupation means type of main occupation for each household, which is classified as farmer, self-employed worker, full-time worker 1, full-time worker 2, others, and unemployed. These six types of household occupations are the

choices listed in the birth certificate for decades, and the parents must choose one from them.

Preterm birth was defined as gestational age below 37 weeks. SGA is often defined as a birth weight below the 10th percentile of the sex- and gestational age-specific standard reference value [8, 25]. In Japan, previous studies of SGA births generally used a sex- parity- and gestational age-specific standard reference for birthweight based on a Japanese neonatal anthropometric chart [7, 15, 26, 27]. Following this precedent, we defined SGA infants as those with birth weight below the 10th percentile of the same standard reference. For this study, we excluded multiples, and restricted the analysis to singleton infants. In addition, infants who were born between 22 and 41 weeks were used for the analysis of SGA births because the standard reference for birthweight in Japan covers only gestational ages of those periods [26].


#### Statistical analysis

We tallied the number of births, preterm births, and SGA births and preterm and SGA birth rates for each infant characteristic. We defined six disjoint groups for maternal age ( $\leq 19$  years, 20–24 years, 25–29 years, 30–34 years, 35–39 years, and  $\geq 40$  years), three groups for gestational age ( $< 32$  weeks, 32–36 weeks, and  $\geq 37$  weeks), and three groups for birth weight ( $< 1500$  g, 1500–2499 g,  $\geq 2500$  g). Parity for infants was categorized into primiparous and multiparous. Maternal nationality was classified into Japanese and non-Japanese. Chi-square test was conducted between each of the birth characteristics and each of the outcomes. In addition, the relative risk of preterm and SGA birth for each type of occupation was calculated using a log binomial regression model with other characteristics as covariates. The log binomial regression model is a method used for calculating the relative risks for binary data [28, 29]. Unemployed persons were used as a reference in the regression analysis for maternal occupations because the number of births was the largest. We reported the relative risk, its 95% confidence interval, and the p-value.

Complete-case analysis was conducted in the chi-square test and the regression analysis for dealing with missing data. In addition, multiple imputation was used as a sensitivity analysis for dealing with missing data in the regression analysis [30]. All statistical analyses were performed by using the statistical software R (version 4.6.1) [31]. In addition, statistics shown in this study were analyzed by the author using the data provided from the Ministry of Health, Labor and Wealth, and those are different from statistics published by the Ministry.

## Results

Figure 1 shows the flowchart of selecting birth data for analysis.



**Fig. 1** Flowchart of selecting birth data

Table 1 shows number of live births, preterm births, and SGA births and rates of preterm and SGA births for each of the characteristics. The greatest number of births was registered for unemployed persons among maternal occupations. The preterm and SGA rates were particularly high for mothers of age 40 and older. Among maternal occupations, preterm birth rate was highest among carrying, cleaning, packaging, and related workers (5.65%) and lowest among security workers (4.24%).

SGA birth rate was highest among manufacturing process workers (5.91%) and lowest among security workers (4.00%). Most of the characteristics such as maternal occupations were statistically significantly associated with the birth outcomes.

Table 2 shows analysis results of regression analysis showing relative risk of maternal occupations for preterm and SGA birth. We found significantly elevated risks for preterm birth among manufacturing process workers

**Table 1** Number of live births, preterm births, and SGA births and rates of preterm and SGA births for each of the characteristics

|  | Preterm birth         |        |          |                      | SGA birth                                  |        |          |                      |
|--|-----------------------|--------|----------|----------------------|--|--------|----------|----------------------|
|  | Number of live births | Number | Rate (%) | p-value <sup>a</sup> | Number of infants born between 22–41 weeks | Number | Rate (%) | p-value <sup>a</sup> |
| Total  | 997,600               | 46,685 | 4.68     |                      | 995,473                                    | 51,603 | 5.18     |                      |
| Maternal age group                                 |                       |        |          | < 0.001              |  |        |          | < 0.001              |
| Under 20 years                                     | 11,907                | 659    | 5.53     |                      | 11,882                                     | 624    | 5.25     |                      |
| 20–24 years  | 84,431                | 3,581  | 4.24     |                      | 84,208                                     | 4,422  | 5.25     |                      |
| 25–29 years  | 259,444               | 10,230 | 3.94     |                      | 258,890                                    | 13,185 | 5.09     |                      |
| 30–34 years  | 362,287               | 16,076 | 4.44     |                      | 361,535                                    | 18,223 | 5.04     |                      |
| 35–39 years  | 225,745               | 12,401 | 5.49     |                      | 225,282                                    | 11,939 | 5.30     |                      |
| 40 years or more                                   | 53,786                | 3,738  | 6.95     |                      | 53,676                                     | 3,210  | 5.98     |                      |
| Maternal nationality                               |                       |        |          | < 0.001              |  |        |          | < 0.001              |
| Japanese   | 974,010               | 45,429 | 4.66     |                      | 971,954                                    | 50,932 | 5.24     |                      |
| Non-Japanese                                       | 23,590                | 1,256  | 5.32     |                      | 23,519                                     | 671    | 2.85     |                      |
| Sex  |                       |        |          | < 0.001              |  |        |          | 0.765                |
| Female   | 486,483               | 19,836 | 4.08     |                      | 485,398                                    | 25,123 | 5.18     |                      |
| Male   | 511,117               | 26,849 | 5.25     |                      | 510,075                                    | 26,480 | 5.19     |                      |
| Parity   |                       |        |          | < 0.001              |  |        |          | < 0.001              |
| Primiparous  | 475,620               | 21,609 | 4.54     |                      | 473,912                                    | 25,378 | 5.36     |                      |
| Multiparous  | 521,980               | 25,076 | 4.80     |                      | 521,561                                    | 26,225 | 5.03     |                      |
| Wedlock status                                     |                       |        |          | < 0.001              |  |        |          | < 0.001              |
| In wedlock   | 972,910               | 44,949 | 4.62     |                      | 970,864                                    | 49,944 | 5.14     |                      |
| Out of wedlock                                     | 24,690                | 1,736  | 7.03     |                      | 24,609                                     | 1,659  | 6.74     |                      |
| Gestational age                                    |                       |        |          | < 0.001              |  |        |          | < 0.001              |
| < 32 weeks   | 5,672                 | 5,672  | 100.00   |                      | 5,666                                      | 1,173  | 20.70    |                      |
| 32–36 weeks  | 41,013                | 41,013 | 100.00   |                      | 41,013                                     | 3,636  | 8.87     |                      |
| ≥ 37 weeks   | 950,915               | 0      | 0.00     |                      | 948,794                                    | 46,794 | 4.93     |                      |
| Birth weight                                       |                       |        |          | < 0.001              |  |        |          | < 0.001              |
| 1499 g or less                                     | 5,841                 | 5,786  | 99.06    |                      | 5,835                                      | 2,337  | 40.05    |                      |
| 1500–2499 g  | 75,561                | 23,763 | 31.45    |                      | 75,546                                     | 29,988 | 39.70    |                      |
| ≥ 2500 g   | 916,198               | 17,136 | 1.87     |                      | 914,092                                    | 19,278 | 2.11     |                      |
| Household occupation                               |                       |        |          | < 0.001              |  |        |          | < 0.001              |
| Farmer   | 12,789                | 608    | 4.75     |                      | 12,755                                     | 642    | 5.03     |                      |
| Self-employed worker                               | 71,471                | 3,414  | 4.78     |                      | 71,301                                     | 3,662  | 5.14     |                      |
| Full-time worker 1 <sup>b</sup>                    | 332,563               | 15,703 | 4.72     |                      | 331,852                                    | 17,859 | 5.38     |                      |
| Full-time worker 2 <sup>c</sup>                    | 454,301               | 20,672 | 4.55     |                      | 453,433                                    | 22,660 | 5.00     |                      |
| Others   | 84,339                | 3,867  | 4.59     |                      | 84,131                                     | 4,301  | 5.11     |                      |
| Unemployed   | 20,160                | 1,277  | 6.33     |                      | 20,106                                     | 1,309  | 6.51     |                      |
| Missing  | 21,977                | 1,144  | 5.21     |                      | 21,895                                     | 1,170  | 5.34     |                      |
| Maternal occupation                                |                       |        |          | 0.001                |  |        |          | < 0.001              |
| Unemployed persons                                 | 535,812               | 25,135 | 4.69     |                      | 534,739                                    | 27,325 | 5.11     |                      |
| Administrative and managerial workers              | 5,140                 | 254    | 4.94     |                      | 5,129                                      | 245    | 4.78     |                      |
| Professional and engineering workers               | 149,002               | 6,800  | 4.56     |                      | 148,662                                    | 7,613  | 5.12     |                      |
| Clerical workers                                   | 127,405               | 5,915  | 4.64     |                      | 127,135                                    | 6,727  | 5.29     |                      |
| Sales workers                                      | 32,548                | 1,495  | 4.59     |                      | 32,467                                     | 1,776  | 5.47     |                      |
| Service workers                                    | 64,555                | 3,092  | 4.79     |                      | 64,385                                     | 3,475  | 5.40     |                      |
| Security workers                                   | 2,904                 | 123    | 4.24     |                      | 2,901                                      | 116    | 4.00     |                      |
| Agriculture, forestry, and fishery workers         | 4,132                 | 183    | 4.43     |                      | 4,124                                      | 197    | 4.78     |                      |
| Manufacturing process workers                      | 16,569                | 883    | 5.33     |                      | 16,543                                     | 977    | 5.91     |                      |
| Transport and machine operating workers            | 1,549                 | 74     | 4.78     |                      | 1,547                                      | 67     | 4.33     |                      |
| Construction and mining workers                    | 1,893                 | 85     | 4.49     |                      | 1,888                                      | 91     | 4.82     |                      |
| Carrying, cleaning, packaging, and related workers | 1,718                 | 97     | 5.65     |                      | 1,712                                      | 78     | 4.56     |                      |

**Table 1** (continued)

|   | Number of live births | Preterm birth |          |                      | Number of infants born between 22–41 weeks | SGA birth |          |                      |
|---|-----------------------|---------------|----------|----------------------|--|-----------|----------|----------------------|
|   |                       | Number        | Rate (%) | p-value <sup>a</sup> |  | Number    | Rate (%) | p-value <sup>a</sup> |
| Workers engaged in an unclassified occupation | 12,833                | 555           | 4.32     |                      | 12,800                                     | 656       | 5.13     |                      |
| Missing                                       | 41,540                | 1,994         | 4.80     |                      | 41,441                                     | 2,260     | 5.45     |                      |

SGA, small-for-gestational-age. a. p-value indicates the result of chi-square test for an association between an outcome and each variable. b. Full-time worker 1 means household of a full-time worker of a company or private shop (except for public offices) who has 1–99 employees. c. Full-time worker 2 means household of a board member or of a full-time worker who does not correspond with the full-time worker 1.

compared with unemployed mothers, and significantly elevated risks for SGA birth compared with unemployed mothers were observed among sales workers, service workers, and manufacturing process workers. In contrast, security workers had a significantly decreased risk for SGA birth compared with unemployed mothers. In addition, the relative risk was the highest level among manufacturing process workers both for preterm and SGA births.

Supplementary Table 1 shows analysis results of regression analysis showing relative risk of maternal occupations for preterm and SGA birth using multiple imputation. Similar results were obtained as the main analysis, while a significant risk ratio for SGA birth was not observed for sales and service workers.

## Discussion

An association between the adverse birth outcomes and maternal occupations was revealed from an analysis of nationwide Vital Statistics data in Japan. Specifically, we have found that the risk of preterm and SGA births varied depending on maternal occupations. We now discuss the possible reasons for those associations.

Regarding the mother's occupation, it was suggested that the risk for manufacturing process workers was the highest. The elevated risk for adverse birth outcomes among manufacturing process workers might indirectly reflect the prevalence of lower socioeconomic status among Japanese female manufacturing process workers. It is known that, in Japan, the proportion of university graduates is relatively low and the proportion of junior high school or elementary school graduates is relatively high for manufacturing process workers in the occupational classifications among women aged 20–49 years old [32]. The association of maternal education level or socioeconomic status with preterm birth and SGA is known [33, 34]. Specifically, lower socioeconomic status of the mother is positively associated with higher rate of preterm and SGA births in Japan [16, 18]. Furthermore, lower socioeconomic status is associated with the presence of medical risk factors, such as high smoking rate and low utilization of prenatal care, for adverse birth outcomes in Japan [35, 36]. However, the risk was

not elevated among other types of workers whose educational level were relatively low, such as transport and machine operating workers, indicating that education level and socioeconomic status are not sufficient to fully explain the observations. Instead, other factors, such as employment status or occupation-specific factors could, in part, explain the elevated risk for preterm and SGA births among manufacturing process workers or other workers (service workers and sales workers). For example, long standing time at work or frequent night shifts are also known to be associated with the adverse birth outcomes [37–39], and it might have affected the results. In addition, security workers had the lowest risk for SGA births. As one possibility, physical characteristics of those workers might be related to the results. It is known that maternal height or BMI affects SGA births, and lower height or BMI is associated with higher risk of SGA birth [40–42]. Generally, adequate body weight and height are required for becoming security workers, such as police officers and self-defense officers, and it might have favorable effects on birth outcomes. An association of maternal occupations with socioeconomic and physical characteristics is needed to be surveyed in order to identify the reasons for the difference in risks.

One strength of this study is that we used a large set of nationwide birth data for the analysis, and the study results represent nationwide trends in Japan. In particular, the association between maternal occupations and SGA and preterm birth is demonstrated for the first time using nationwide Vital Statistics. We specifically highlighted a higher risk of preterm birth and SGA for mothers employed as manufacturing process workers. This finding has important implications. First, future studies are needed to investigate the causes for the observed association. Second, there is a need for implementing effective prevention measures to lower the risk of adverse birth outcomes. Adjusting the work load or improving working conditions might be needed to lower those risks for pregnant manufacturing process workers. A call for attention in working places or prenatal care for pregnant women with those occupations might help. Finally, providing health guidance targeted to husbands of pregnant

**Table 2** Analysis results of regression analysis showing relative risk of maternal occupations for preterm and SGA birth

|  | Preterm birth        |         | SGA birth            |         |
|--|----------------------|---------|----------------------|---------|
|  | RR (95% CI)          | p-value | RR (95% CI)          | p-value |
| Maternal age group                                 |                      |         |                      |         |
| Under 20 years                                     | 1.084 (0.999, 1.177) | 0.053   | 0.932 (0.858, 1.013) | 0.098   |
| 20–24 years  | 0.911 (0.877, 0.945) | < 0.001 | 0.991 (0.958, 1.026) | 0.620   |
| 25–29 years  | 0.880 (0.858, 0.903) | < 0.001 | 0.992 (0.970, 1.015) | 0.491   |
| 30–34 years  | Reference            |         | Reference            |         |
| 35–39 years  | 1.232 (1.204, 1.262) | < 0.001 | 1.052 (1.028, 1.077) | < 0.001 |
| 40 years or more                                   | 1.558 (1.504, 1.614) | < 0.001 | 1.171 (1.128, 1.216) | < 0.001 |
| Maternal nationality                               |                      |         |                      |         |
| Japanese   | Reference            |         | Reference            |         |
| Non-Japanese                                       | 1.096 (1.034, 1.161) | 0.002   | 0.532 (0.492, 0.575) | < 0.001 |
| Sex  |                      |         |                      |         |
| Female   | Reference            |         | Reference            |         |
| Male   | 1.288 (1.264, 1.312) | < 0.001 | 1.003 (0.986, 1.020) | 0.739   |
| Parity   |                      |         |                      |         |
| Primiparous  | Reference            |         | Reference            |         |
| Multiparous  | 1.004 (0.986, 1.024) | 0.645   | 0.930 (0.914, 0.947) | < 0.001 |
| Wedlock status                                     |                      |         |                      |         |
| In wedlock   | Reference            |         | Reference            |         |
| Out of wedlock                                     | 1.398 (1.318, 1.482) | < 0.001 | 1.240 (1.169, 1.315) | < 0.001 |
| Household occupation                               |                      |         |                      |         |
| Farmer   | 1.062 (0.974, 1.158) | 0.173   | 1.017 (0.935, 1.106) | 0.694   |
| Self-employed worker                               | 1.013 (0.977, 1.050) | 0.496   | 1.029 (0.994, 1.065) | 0.110   |
| Full-time worker 1 <sup>a</sup>                    | 1.046 (1.024, 1.067) | < 0.001 | 1.083 (1.062, 1.105) | < 0.001 |
| Full-time worker 2 <sup>b</sup>                    | Reference            |         | Reference            |         |
| Others   | 1.024 (0.989, 1.059) | 0.183   | 1.032 (0.999, 1.066) | 0.058   |
| Unemployed   | 1.182 (1.107, 1.262) | < 0.001 | 1.246 (1.169, 1.329) | < 0.001 |
| Maternal occupation                                |                      |         |                      |         |
| Unemployed persons                                 | Reference            |         | Reference            |         |
| Administrative and managerial workers              | 0.975 (0.863, 1.100) | 0.678   | 0.909 (0.802, 1.030) | 0.135   |
| Professional and engineering workers               | 0.979 (0.953, 1.005) | 0.112   | 1.003 (0.978, 1.029) | 0.806   |
| Clerical workers                                   | 0.975 (0.948, 1.003) | 0.076   | 1.026 (0.999, 1.053) | 0.060   |
| Sales workers                                      | 0.985 (0.935, 1.037) | 0.564   | 1.062 (1.013, 1.113) | 0.013   |
| Service workers                                    | 1.031 (0.993, 1.070) | 0.106   | 1.043 (1.007, 1.080) | 0.017   |
| Security workers                                   | 0.951 (0.798, 1.134) | 0.578   | 0.817 (0.684, 0.977) | 0.027   |
| Agriculture, forestry, and fishery workers         | 0.913 (0.782, 1.066) | 0.249   | 0.956 (0.824, 1.110) | 0.557   |
| Manufacturing process workers                      | 1.156 (1.082, 1.234) | < 0.001 | 1.178 (1.107, 1.254) | < 0.001 |
| Transport and machine operating workers            | 1.029 (0.823, 1.288) | 0.801   | 0.847 (0.669, 1.072) | 0.167   |
| Construction and mining workers                    | 0.941 (0.763, 1.160) | 0.570   | 0.952 (0.779, 1.164) | 0.631   |
| Carrying, cleaning, packaging, and related workers | 1.173 (0.965, 1.426) | 0.108   | 0.878 (0.705, 1.094) | 0.247   |
| Workers engaged in an unclassified occupation      | 0.901 (0.828, 0.981) | 0.017   | 0.987 (0.912, 1.067) | 0.736   |

SGA, small-for-gestational-age; RR, relative risk; CI, confidence interval. a. Full-time worker 1 means household of a full-time worker of a company or private shop (except for public offices) who has 1–99 employees. b. Full-time worker 2 means household of a board member or of a full-time worker who does not correspond with the full-time worker 1.

women whose occupations have higher risk of adverse birth outcomes might be also effective.

### Limitations

One limitation of this study is the lack of data on women's behaviors or physical characteristics, such as smoking status, BMI, utilization of prenatal care, chronic diseases, and working hours, that are known risk factors for adverse birth outcomes. Another limitation is the

indirect inference of maternal educational level or physical characteristics because of the absence of such data in the dataset we analyzed. Socioeconomic status indicators of a mother, such as educational level or household income were not available in the data, and we could not fully differentiate the effect of maternal occupations from that of maternal socioeconomic status. Thus, to properly account for those factors that the survey does not collect,

an epidemiological study designed to target those factors needs to be conducted in the future.

#### List of abbreviations

|     |                           |
|-----|---------------------------|
| SGA | small-for-gestational-age |
| RR  | relative risk             |
| CI  | confidence interval       |
| BMI | body mass index           |

#### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13104-023-06539-0>.

Supplementary Material 1

#### Acknowledgements

None.

#### Author contributions

Conceptualization: TO. Data curation: TO. Formal analysis: TO. Methodology: TO. Funding acquisition: TO. Writing- original draft: TO. Writing - review & editing: TO.

#### Funding

This study was supported by JSPS KAKENHI Grant Number JP22K17372.

#### Data Availability

The data that support the findings of this study are available from the Ministry of Health, Labour, and Welfare in Japan. However, restrictions apply to the availability of these data, which were provided under license for the current study.

#### Declarations

##### Ethics approval and consent to participate

This study was conducted on the basis of the Statistics Act with the permission of the Ministry of Health, Labour, and Welfare in Japan, and it was approved by the ethical committee of Kyushu University (No. 22221-00). Informed consent was not required because we used the official statistics data that were provided from the government in accordance with the law. All the works were conducted in compliance with relevant laws and guidelines.

##### Consent for publication

Not applicable.

##### Competing interests

None.

Received: 12 March 2023 / Accepted: 26 September 2023

Published online: 05 October 2023

#### References

- Lee AC, Katz J, Blencowe H, Cousens S, Kozuki N, Vogel JP, Adair L, Baqui AH, Bhutta ZA, Caulfield LE, Christian P, Clarke SE, Ezzati M, Fawzi W, Gonzalez R, Huybregts L, Kariuki S, Kolsteren P, Lusingu J, Marchant T, Meriadi M, Mongkolkeha A, Mullany LC, Ndirangu J, Newell ML, Nien JK, Osrin D, Roberfroid D, Rosen HE, Sania A, Silveira MF, Tielsch J, Vaidya A, Willey BA, Lawn JE, Black RE. ChERG SGA-Preterm Birth Working Group. National and regional estimates of term and preterm babies born small for gestational age in 138 low-income and middle-income countries in 2010. *Lancet Glob Health*. 2013;1:e26–36. [https://doi.org/10.1016/S2214-109X\(13\)70006-8](https://doi.org/10.1016/S2214-109X(13)70006-8).
- Chawanpaiboon S, Vogel JP, Moller AB, Lumbiganon P, Petzold M, Hogan D, Landoulsi S, Jampathong N, Kongwattanakul K, Laopaiboon M, Lewis C, Ratanakanokchai S, Teng DN, Thinkhamrop J, Watananirun K, Zhang J, Zhou W, Gülmezoglu AM. Global, regional, and national estimates of levels of preterm birth in 2014: a systematic review and modelling analysis. *Lancet Glob Health*. 2019;7:e37–e46. [https://doi.org/10.1016/S2214-109X\(18\)30451-0](https://doi.org/10.1016/S2214-109X(18)30451-0).
- Sakata S, Konishi S, Ng CFS, Watanabe C. Preterm birth rates in Japan from 1979 to 2014: analysis of national vital statistics. *J Obstet Gynaecol Res*. 2018;44:390–6. <https://doi.org/10.1111/jog.13460>.
- Yamaoka Y, Morisaki N, Noguchi H, Takahashi H, Tamiya N. Comprehensive assessment of risk factors of cause-specific infant deaths in Japan. *J Epidemiol*. 2018;28:307–14.
- Sugai MK, Gilmour S, Ota E, Shibuya K. Trends in perinatal mortality and its risk factors in Japan: analysis of vital registration data, 1979–2010. *Sci Rep*. 2017;7:46681. <https://doi.org/10.1038/srep46681>.
- Luu TM, Rehman Mian MO, Nuyt AM. Long-term impact of preterm birth: neurodevelopmental and physical health outcomes. *Clin Perinatol*. 2017;44:305–14. <https://doi.org/10.1016/j.clp.2017.01.003>.
- Shibata M, Ogawa K, Kanazawa S, Kawasaki M, Morisaki N, Mito A, Sago H, Horikawa R, Arata N. Association of maternal birth weight with the risk of low birth weight and small-for-gestational-age in offspring: a prospective single-center cohort study. *PLoS ONE*. 2021;16:e0251734. <https://doi.org/10.1371/journal.pone.0251734>.
- He H, Miao H, Liang Z, Zhang Y, Jiang W, Deng Z, Tang J, Liu G, Luo X. Prevalence of small for gestational age infants in 21 cities in China, 2014–2019. *Sci Rep*. 2021;11:7500. <https://doi.org/10.1038/s41598-021-87127-9>.
- Takeuchi A, Yorifuji T, Takahashi K, Nakamura M, Kageyama M, Kubo T, Ogino T, Doi H. Neurodevelopment in full-term small for gestational age infants: a nationwide Japanese population-based study. *Brain Dev*. 2016;38:529–37. <https://doi.org/10.1016/j.braindev.2015.12.013>.
- Ludvigsson JF, Lu D, Hammarström L, Cnattingius S, Fang F. Small for gestational age and risk of childhood mortality: a Swedish population study. *PLoS Med*. 2018;15:e1002717. <https://doi.org/10.1371/journal.pmed.1002717>.
- Sania A, Smith ER, Manji K, Duggan C, Masanja H, Kisenge R, Msamanga G, Urassa W, Fawzi W. Neonatal and infant mortality risk associated with preterm and small for gestational age births in Tanzania: individual level pooled analysis using the Intergrowth Standard. *J Pediatr*. 2018;192:66–72. <https://doi.org/10.1016/j.jpeds.2017.09.007>.
- Londero AP, Rossetti E, Pittini C, Cagnacci A, Driul L. Maternal age and the risk of adverse pregnancy outcomes: a retrospective cohort study. *BMC Pregnancy Childbirth*. 2019;19:261. <https://doi.org/10.1186/s12884-019-2400-x>.
- Bushnik T, Yang S, Kaufman JS, Kramer MS, Wilkins R. Socioeconomic disparities in small-for-gestational-age birth and preterm birth. *Health Rep*. 2017;28:3–10.
- Shah NR, Bracken MB. A systematic review and meta-analysis of prospective studies on the association between maternal cigarette smoking and preterm delivery. *Am J Obstet Gynecol*. 2000;182:465–72. [https://doi.org/10.1016/S0002-9378\(00\)70240-7](https://doi.org/10.1016/S0002-9378(00)70240-7).
- Iwama N, Obara T, Ishikuro M, Murakami K, Ueno F, Noda A, Onuma T, Matsuzaki F, Hoshiai T, Saito M, Metoki H, Sugawara Y, Jaegashi N, Kuriyama S. Risk scores for predicting small for gestational age infants in Japan: the TMM birth cohort study. *Sci Rep*. 2022;12:8921. <https://doi.org/10.1038/s41598-022-12892-0>.
- Fujiwara T, Ito J, Kawachi I. Income inequality, parental socioeconomic status, and birth outcomes in Japan. *Am J Epidemiol*. 2013;177:1042–52. <https://doi.org/10.1093/aje/kws355>.
- Tamura N, Hanaoka T, Ito K, Araki A, Miyashita C, Ito S, Minakami H, Cho K, Endo T, Sengoku K, Ogasawara K, Kishi R. Different risk factors for very low birth weight, term-small-for-gestational-age, or preterm birth in Japan. *Int J Environ Res Public Health*. 2018;15:369. <https://doi.org/10.3390/ijerph15020369>.
- Okui T. Analysis of an association between preterm birth and parental educational level in Japan using national data. *Child (Basel)*. 2023;10:342. <https://doi.org/10.3390/children10020342>.
- Suzuki Y, Senda Y, Honjo K. Maternal occupation and infant mortality in Japan: insights from the vital statistics (Occupational and Industrial Aspects). *Nihon Koshu Eisei Zasshi*. 2021;68(10):669–76. <https://doi.org/10.11236/jph.20-151>.
- Okui T, Ochiai M, Nakashima N. An association between maternal occupations and low birth weight infants in Japan from 1995 to 2015. *Int J Environ Res Public Health*. 2021;18:8040. <https://doi.org/10.3390/ijerph18158040>.
- Li X, Sundquist J, Sundquist K. Parental occupation and risk of small-for-gestational-age births: a nationwide epidemiological study in Sweden. *Hum Reprod*. 2010;25:1044–50. <https://doi.org/10.1093/humrep/deq004>.



22. Park MJ, Son M, Kim YJ, Paek D. Social inequality in birth outcomes in Korea, 1995–2008. *J Korean Med Sci*. 2013;28(1):25–35. <https://doi.org/10.3346/jkms.2013.28.1.25>. Epub 2013 Jan 8. PMID: 23341708; PMCID: PMC3546100.
23. Savitz DA, Olshan AF, Gallagher K. Maternal occupation and pregnancy outcome. *Epidemiology*. 1996;7:269–74. <https://doi.org/10.1097/00001648-199605000-00009>.
24. Wada K, Gilmour S. Inequality in mortality by occupation related to economic crisis from 1980 to 2010 among working-age Japanese males. *Sci Rep*. 2016;6:22255. <https://doi.org/10.1038/srep22255>.
25. Shittu AAT, Kumar BP, Okafor U, Berkelhamer SK, Goniewicz ML, Wen X. Changes in e-cigarette and cigarette use during pregnancy and their association with small-for-gestational-age birth. *Am J Obstet Gynecol*. 2022;226:730.e1–730.e10.
26. Itabashi K, Miura F, Uehara R, Nakamura Y. New Japanese neonatal anthropometric charts for gestational age at birth. *Pediatr Int*. 2014;56:702–8. <https://doi.org/10.1111/ped.12331>.
27. Hayashi I, Takakura K, Yamaguchi K, Sumitomo M, Suzuki M, Sumitomo A, Minato S, Nose Y, Nagai N, Sakane N. Association between socioeconomic status and small-for-gestational-age in Japan: a single center retrospective cohort study. *J Obstet Gynaecol Res*. 2020;46:110–8. <https://doi.org/10.1111/jog.14069>.
28. Chen W, Qian L, Shi J, Franklin M. Comparing performance between log-binomial and robust Poisson regression models for estimating risk ratios under model misspecification. *BMC Med Res Methodol*. 2018;18:63. <https://doi.org/10.1186/s12874-018-0519-5>.
29. Urquia ML, Qiao Y, Ray JG, Liu C, Hjert A. Birth outcomes of foreign-born, native-born, and mixed couples in Sweden. *Paediatr Perinat Epidemiol*. 2015;29:123–30. <https://doi.org/10.1111/ppe.12179>.
30. van Buuren S, Groothuis-Oudshoorn K. Mice: multivariate imputation by chained equations in R. *J Stat Softw*. 2013;45:1–67.
31. R Core Team. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. [cited 2023 March 13]. <https://www.R-project.org/>.
32. Ministry of Internal Affairs and Communications. The Census. [cited 2023 March 13]. <https://www.e-stat.go.jp/stat-search/files?page=1&itoukei=00200521>.
33. Mortensen LH, Lauridsen JT, Diderichsen F, Kaplan GA, Gissler M, Andersen AM. Income-related and educational inequality in small-for-gestational age and preterm birth in Denmark and Finland 1987–2003. *Scand J Public Health*. 2010;38:40–5. <https://doi.org/10.1177/1403494809353820>.
34. Ruiz M, Goldblatt P, Morrison J, Kukla L, Švancara J, Riitta-Järvelin M, Taanila A, Saurel-Cubizolles MJ, Lioret S, Bakoula C, Veltsista A, Porta D, Forastiere F, van Eijdsen M, Vrijkotte TG, Eggesbø M, White RA, Barros H, Correia S, Vrijheid M, Torrent M, Rebagliato M, Larrañaga I, Ludvigsson J, Olsen Faresjö Å, Hryhorczuk D, Antipkin Y, Marmot M, Pikhart H. Mother's education and the risk of preterm and small for gestational age birth: a DRIVERS meta-analysis of 12 European cohorts. *J Epidemiol Community Health*. 2015;69:826–33. <https://doi.org/10.1136/jech-2014-205387>.
35. Tomioka K, Kurumatani N, Saeki K. The Association between education and smoking prevalence, independent of occupation: a nationally representative survey in Japan. *J Epidemiol*. 2020;30:136–42. <https://doi.org/10.2188/jea.JE20180195>.
36. Osawa E, Kodama T. Regional socio-environmental characteristics associated with inadequate prenatal care during pregnancy: an ecological study of 47 prefectures in Japan. *BMC Pregnancy Childbirth*. 2021;21:619. <https://doi.org/10.1186/s12884-021-04100-0>.
37. Mozurkewich EL, Luke B, Avni M, Wolf FM. Working conditions and adverse pregnancy outcome: a meta-analysis. *Obstet Gynecol*. 2000;95:623–35. [https://doi.org/10.1016/s0029-7844\(99\)00598-0](https://doi.org/10.1016/s0029-7844(99)00598-0).
38. Saurel-Cubizolles MJ, Zeitlin J, Lelong N, Papiernik E, Di Renzo GC, Bréart G, Europop Group. Employment, working conditions, and preterm birth: results from the Europop case-control survey. *J Epidemiol Community Health*. 2004;58:395–401. <https://doi.org/10.1136/jech.2003.008029>.
39. Croteau A, Marcoux S, Brisson C. Work activity in pregnancy, preventive measures, and the risk of delivering a small-for-gestational-age infant. *Am J Public Health*. 2006;96:846–55. <https://doi.org/10.2105/AJPH.2004.058552>.
40. Goto E. Dose-response associations of maternal height and weight with small for gestational age: a meta-analysis. *Eur J Clin Nutr*. 2020;74:106–11. <https://doi.org/10.1038/s41430-019-0432-y>.
41. Goetzinger KR, Cahill AG, Macones GA, Odibo AO. The relationship between maternal body mass index and tobacco use on small-for-gestational-age infants. *Am J Perinatol*. 2012;29:153–8. <https://doi.org/10.1055/s-0031-1284224>.
42. Kozuki N, Katz J, Lee AC, Vogel JP, Silveira MF, Sania A, Stevens GA, Cousens S, Caulfield LE, Christian P, Huybregts L, Roberfroid D, Schmiedelow C, Adair LS, Barros FC, Cowan M, Fawzi W, Kolsteren P, Merialdi M, Mongkolkeha A, Saville N, Victora CG, Bhutta ZA, Blencowe H, Ezzati M, Lawn JE, Black RE. Short maternal stature increases risk of small-for-gestational-age and preterm births in low- and middle-income countries: individual participant data meta-analysis and population attributable fraction. *J Nutr*. 2015;145:2542–50. <https://doi.org/10.3945/jn.115.216374>. Child Health Epidemiology Reference Group Small-for-Gestational-Age/Preterm Birth Working Group.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.