

POSTER PRESENTATIONS

Open Access

Strain specific protective immunity induced by pre-erythrocytic infection of toque monkeys with *Plasmodium cynomolgi*

Wathsala Wijayalath¹, Sandie Cheesman³, Jagath Rajakaruna¹, Sudath Weerasinghe¹, Gamini Kapilananda¹, Kamal Gamage², Kamal Perera², Shiroma Handunnetti⁴, Richard Carter³, Sisira Pathirana^{1*}

From Parasite to Prevention: Advances in the understanding of malaria
Edinburgh, UK. 20-22 October 2010

Strain specific protective immunity (SSPI) was investigated using two strains of *Plasmodium cynomolgi*, Pc746 and PcCeylon, in toque monkey. Two groups of monkeys were immunized against either Pc746 (n=5) or PcCeylon (n=4), by giving bites with 2-4 sporozoite-infected *Anopheles tessellates* mosquitoes per monkey. Primary blood infection was prevented by simultaneous chloroquine cover and secondary, hypnozoite-induced, blood infection was prevented by treating the monkeys one month later with primaquine. The two immunized groups and a group of unimmunized monkeys (n=4) were given a mixed-strain sporozoite challenge infection, 140 and 100 days respectively after immunization. Parasite DNA was collected for 5-8 consecutive days after parasitaemia reached 0.05% or above. The proportions of the two parasite strains in these samples were quantified using a PyrosequencingTM assay based on SNPs in MSP1 and CSP genes. In subsequent blood infections in immunized monkeys the earliest recorded proportion of parasites of an immunizing strain was significantly lower than its proportion in monkeys immunized against a heterologous strain (P=0.014 and 0.027 for the two SNPs) and the proportions of an immunizing strain tended to decline during the period of sample collection. These results show that a parasite strain specific protective immunity to *P. cynomolgi* was induced following a sporozoite induced pre-erythrocytic infection. This immunity may have been directed against the liver stages or against the blood stage parasites, or against both

Acknowledgements

Financial support by NSF, Sri Lanka (research grant -RG/2005/HS/14 and scholarship - NSF/SCH/2004/08) and NRC, Sri Lanka 05-66 is acknowledged. SC and RC were supported by a Wellcome Trust Programme Grant.

Author details

¹Malaria Research Unit, Dept of Parasitology, Faculty of Medicine, University of Colombo, Colombo, Sri Lanka. ²Animal House, Faculty of Medicine, University of Colombo, Colombo, Sri Lanka. ³School of Biological Sciences, Institute of Immunology & Infection Research, University of Edinburgh, UK. ⁴Institute of Biochemistry, Molecular Biology and Biotechnology, University of Colombo, Colombo, Sri Lanka.

Published: 20 October 2010

doi:10.1186/1475-2875-9-S2-P58

Cite this article as: Wijayalath et al.: Strain specific protective immunity induced by pre-erythrocytic infection of toque monkeys with *Plasmodium cynomolgi*. *Malaria Journal* 2010 **9**(Suppl 2):P58.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit



¹Malaria Research Unit, Dept of Parasitology, Faculty of Medicine, University of Colombo, Colombo, Sri Lanka
Full list of author information is available at the end of the article