

RESEARCH

Open Access



# Mycothioliol maintains the homeostasis and signalling of nitric oxide in *Streptomyces coelicolor* A3(2) M145

Tomoki Yoshizumi<sup>1</sup>, Yukiko Shibui<sup>1</sup>, Minoru Kogo<sup>1</sup>, Sota Honma<sup>1</sup>, Shinsaku Ito<sup>1</sup>, Shunsuke Yajima<sup>1</sup> and Yasuyuki Sasaki<sup>1\*</sup>

## Abstract

**Background** Previous studies have revealed a nitric oxide (NO) metabolic cycle in which NO, nitrate (NO<sub>3</sub><sup>-</sup>), and nitrite (NO<sub>2</sub><sup>-</sup>) circulate. The NO produced in this cycle serves as a signalling molecule that regulates actinorhodin (ACT) production via the DevS/DevR NO-dependent two-component system (TCS) in *Streptomyces coelicolor* A3(2) M145. However, the mechanisms involved in the regulation of NO signalling in *S. coelicolor* have not yet been elucidated. Mycothiol (MSH), a thiol molecule produced by *Actinomyces*, is involved in the defence mechanisms against oxidative stress. Therefore, this study focused on the correlation between intracellular NO and MSH levels.

**Results** To investigate the interaction of MSH with endogenously produced NO, we generated an *S. coelicolor* A3(2) strain deficient in MSH biosynthesis. This mutant strain exhibited a decrease in low-molecular-weight S-nitrosothiols and intracellular NO levels during culture compared to those of the wild-type strain. Moreover, the mutant strain exhibited reduced activity of the DevS/DevR TCS, a regulator of NO homeostasis and ACT production, from the early stage of culture, along with a decrease in ACT production compared to those of the wild-type strain.

**Conclusions** This study suggests that MSH maintains intracellular NO homeostasis by forming S-nitrosomycothiol, which induces NO signalling. Finally, we propose a metabolic model in which MSH from endogenously produced NO facilitates the maintenance of both NO homeostasis and signalling in *S. coelicolor* A3(2) M145.

**Keywords** Nitric oxide, Actinobacteria, Mycothiol, Intracellular homeostasis, Signal transduction, Secondary metabolism

## Background

Nitric oxide (NO) is a highly reactive and diffusible molecule that can easily permeate the cell membrane [1]. Since Palmer first reported its activity in 1987, NO has been recognised as a signalling molecule that regulates various physiological functions in most organisms [2].

NO mediates cell-to-cell communication and regulates several metabolic pathways by interacting with transition metals (e.g., Fe and Cu [3]), thiol groups [4], free-radical oxygen [5], and unsaturated fatty acids [6]. However, increased levels of NO exert cytotoxic effects owing to its high reactivity. NO reacts with O<sub>2</sub><sup>-</sup> to form peroxynitrite (ONOO<sup>-</sup>), which leads to intracellular dysregulation due to the oxidation of a wide range of cellular molecules such as proteins, lipids, and nucleic acids [7]. Thus, cells must strictly regulate the internal NO concentration to ensure their viability.

\*Correspondence:

Yasuyuki Sasaki

y1sasaki@nodai.ac.jp

<sup>1</sup> Department of Bioscience, Faculty of Life Sciences, Tokyo University of Agriculture, Sakuragaoka, Tokyo, Setagaya-Ku 156-8502, Japan



© The Author(s) 2023. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

Several studies have reported mechanisms involved in the regulation of intracellular NO levels. NO production by enzymes such as nitric oxide synthase and nitrite reductase, which reduce nitrite to NO, depends on the transient expression of genes [8, 9]. To protect cells from NO toxicity, flavohaemoglobin (Fhb) and nitric oxide reductase eliminate cellular NO by converting it into nitrate and nitrous oxide, respectively [10, 11]. Additionally, low-molecular-weight (LMW) thiol compounds such as glutathione (GSH) and thionein modulate the NO concentration by generating S-nitrosothiol [12, 13]. These small molecules can therefore protect cells against nitrosylative stress and regulate the S-nitrosylation of proteins and their corresponding functions.

*Streptomyces* species are gram-positive soil-dwelling filamentous bacteria that exhibit a complex life cycle similar to that of fungi. These bacteria produce many commercially available secondary metabolites, including anticancer agents [14], agrochemicals [15], and antibiotics [16]. Recently, we reported an NO metabolic cycle in which NO, nitrate, and nitrite are circulated by Fhb and nitrate reductase (Nar) in *Streptomyces coelicolor* A3(2) M145; the NO produced in this cycle serves as a signaling molecule to regulate morphological differentiation and secondary metabolism in bacteria [17]. In addition, the production of NO and actinorhodin (ACT; a blue-pigmented antibiotic) were demonstrated to be regulated by the NO-dependent two-component regulatory system (TCS) DevS/DevR in *Streptomyces coelicolor* A3(2) M145 [18]. The sensor protein DevS binds NO and phosphorylates DevR, a cognate response regulator. Phosphorylated DevR then activates the expression of its own genes (*devR*, *narG2*, and *actII-ORF4*). In addition, DevS senses NO elevation during cell growth and inactivates its autophosphorylation activity, thereby inhibiting the NO synthesis pathway. Thus, DevS/R functions as an autoregulatory system to maintain NO homeostasis in *S. coelicolor* A3(2) M145 [18].

Mycothiols (MSH) is an *Actinomycetes*-specific LMW thiol compound consisting of an acetylated cysteine

linked to glucosamine and inositol [19]. Additionally, MSH acts as an antioxidant buffer by forming mycothione, which is reduced by disulphide reductase [20]. Similar to GSH, MSH reacts with NO to form S-nitroso-mycothiol (MSNO) and mitigates reactive nitrogen species-induced damage [21]. Four enzymes [MshA (SCO4204), MshB (SCO5126), MshC (SCO1663), and MshD (SCO4151)] are involved in the MSH biosynthesis pathway in *S. coelicolor* A3(2) M145 [22]. Although the correlation between LMW thiol compounds and oxidative stress has been extensively studied, the relationship between MSH and intracellular NO signaling in *S. coelicolor* A3(2) M145 remains unclear. In the present study, we investigated the relationship between MSH and endogenously produced NO in *S. coelicolor* A3(2) M145 cells.

## Results


### Effect of MSH deficiency on cell growth

Previous studies have reported that MSH eliminates reactive oxygen and nitrogen species generated in cells [20, 21]. However, the relationship between MSH expression and endogenous NO production remains unclear. A *Streptomyces coelicolor* A3(2) M145 strain lacking MSH synthesis-related genes (*mshA*, *mshC*, and *mshD*) cannot synthesize MSH [22]. Thus, we deleted *mshC* by homologous recombination using the plasmid pGM160 (Fig. 1a) to obtain a model strain incapable of MSH biosynthesis.

To evaluate the effect of the loss of MSH biosynthesis ability in the cells, we comparatively investigated the colony areas and wet weights of the wild-type (WT) M145 strain and MSH-deficient  $\Delta mshC$  strain cultured on yeast extract-malt extract-glutamine (YEME-gln) agar medium using ImageJ. Only relatively small differences in colony area and wet weight were found between strains after 72 and 168 h of incubation, respectively (Fig. 1b, c). Furthermore, only slight differences in colony area and wet weight were also found between the *nar* deletion strain ( $\Delta nar$  strain), with reduced NO production ability, and the *nar* and *mshC*

(See figure on next page.)

**Fig. 1** Effect of mycothiol (MSH) deficiency on colony growth. **a** Construction of the *mshC* disruptant strain. The plasmid pGM160, including the 2,000 nucleotides upstream and downstream of *mshC* flanking an *aac(3)IV* gene, was transformed into *Streptomyces coelicolor* A3(2) M145. The double-crossover event is shown schematically. **b** The colony area of strains M145 (blue bars),  $\Delta mshC$  (red bars), and  $\Delta mshC::mshC$  (yellow bars) cultured for the indicated durations. Data are presented as mean and standard deviation ( $n=9$ ). \*\*\* $p < 0.001$ ; NS, not significant (Student's t-test). **c** The wet weight of M145 (blue bars),  $\Delta mshC$  (red bars), and  $\Delta mshC::mshC$  (yellow bars) strains cultured for the indicated durations. Data are presented as mean and standard deviation values from three independent experiments. \*\*\* $p < 0.001$ ; \* $p < 0.05$ . NS, not significant (Student's t-test). **d** The colony area of  $\Delta nar$  (blue bars) and  $\Delta nar/mshC$  (red bars) strains cultured for the indicated durations. Data are presented as mean and standard deviation ( $n=9$ ) from three independent experiments. **e** The wet weight of  $\Delta nar$  (blue bars) and  $\Delta nar/mshC$  (red bars) strains cultured for the indicated durations. Data are presented as the mean and standard deviation values from three independent experiments. \*\*\* $p < 0.001$ ; \*\* $p < 0.01$ ; NS, not significant (Student's t-test)



**Fig. 1** (See legend on previous page.)

double-deletion strains ( $\Delta nar/mshC$  strain) (Fig. 1d, e). These results suggested that MSH has a relatively small effect on the growth of *S. coelicolor* A3(2) M145, regardless of the presence or absence of endogenously produced NO.

**Deficiency of MSH synthesis decreases the intracellular NO level in *S. coelicolor* A3(2)M145**

Because *hmpA* (encoding Fhb) is regulated by the NO-dependent transcription factor NsrR, the gene expression level is reflective of the intracellular NO concentration


[23]. To verify the effect of MSH synthesis deficiency on endogenous NO levels, we examined the expression levels of *hmpA* in the M145 and  $\Delta mshC$  strains. The transcription levels of *hmpA* increased in a time-dependent manner in both strains but were maintained at lower levels in the  $\Delta mshC$  strain compared to those in the M145 strain (Fig. 2a).

DevS/R has been reported to regulate NO synthesis in a dose-dependent manner to maintain NO homeostasis [18]. Therefore, we further examined NO homeostasis in the  $\Delta mshC$  strain by evaluating the expression of the DevS/R TCS regulon *devR*. The transcription level of *devR* in the  $\Delta mshC$  strain was decreased compared

to that in the M145 strain during culture (Fig. 2b). These results suggest that the lack of MSH suppressed NO production in the  $\Delta mshC$  strain during the early stages of culture. This was supported by two exogenous NO donors inducing the expression of *hmpA* in the  $\Delta mshC$  strain (Fig. 2c).

#### MSH deletion decreases LMW S-nitrosothiol levels

Since previous studies demonstrated that S-nitrosothiols are formed by thiol groups that bind to NO in the presence of electron acceptors in cells [24, 25], we verified LMW S-nitrosothiol levels in the M145,  $\Delta mshC$ , and  $\Delta nar$  strains in a time-dependent manner. The




**Fig. 2** Mycothiol (MSH) gene deletion decreases intracellular nitric oxide (NO) levels. **a** Reverse-transcription quantitative polymerase chain reaction (RT-qPCR) analysis of the relative expression levels of *hmpA* in the total RNA extracted from strains M145 (blue bars),  $\Delta mshC$  (red bars), and  $\Delta mshC::mshC$  (yellow bars) cultured for the indicated durations. Data are presented as the mean and standard deviation values from six independent experiments. \*\* $p < 0.01$ ; \* $p < 0.05$ ; NS, not significant (Student's *t*-test). **b** Relative mRNA expression levels of *devR* in the total RNA extracted from strains M145 (blue bars),  $\Delta mshC$  (red bars), and  $\Delta mshC::mshC$  (yellow bars) cultured for the indicated durations. Data are presented as the mean and standard deviation values from three independent experiments. \*\* $p < 0.01$ , \* $p < 0.05$  (Student's *t*-test). **c** Relative mRNA expression levels of *hmpA* in the total RNA extracted from M145 and  $\Delta mshC$  strains. After the strains were cultured for 72 h, they were treated with 1 ml water (control; blue bars), 100  $\mu\text{M NO}_2^-$  for 1 h (red bars), or 1  $\mu\text{M NOC5}$  solution (yellow bars); incubated for 30 min; and subjected to RT-qPCR analysis. Data are presented as mean and standard deviation values from three independent experiments. \* $p < 0.05$ ; NS, not significant (Dunnett's test)

LMW S-nitrosothiol levels increased from 96 to 144 h but decreased after 144 h of culture in the M145 strain (Fig. 3). This is consistent with the results of a previous study demonstrating time-dependent changes in NO production in vivo [18]. Intracellular NO levels in *S. coelicolor* A3(2) M145 increased during the middle phase of culture and decreased during the late phase. In contrast, the LMW S-nitrosothiol levels in the  $\Delta mshC$  and  $\Delta nar$  strains were lower than those in the M145 strain between 96 and 144 h (Fig. 3). These results indicated that the amount of LMW S-nitrosothiols is


positively correlated with intracellular NO levels in *S. coelicolor* A3(2) M145.

**MSH is needed for optimal ACT synthesis**

A previous study demonstrated that the DevS/R TCS modulated ACT synthesis by regulating *actII-ORF4* expression levels [18]. Since we found that the lack of MSH affected cellular NO levels, ACT production and *actII-ORF4* expression levels were examined in the M145 and  $\Delta mshC$  strains. Minimal ACT production (Fig. 4a) and downregulated *actII-ORF4* expression (Fig. 4b) were



**Fig. 3** Defect of mycothiol (MSH) synthesis decreases low-molecular-weight (LMW) S-nitrosothiol (SNO) levels. M145 (blue bars),  $\Delta mshC$  (red bars), and  $\Delta nar$  (yellow bars) strains were cultured for the indicated durations. After deproteinisation, SNO was converted to  $NO_2^-$  using  $Hg^{2+}$  as a catalyst. SNO levels were measured using the Saville–Griess assay. Concentrations below the detection limit (500 nM) were recorded as the detection limit concentration. Data are presented as the mean and standard deviation values from three or four independent experiments. \*\*\* $p < 0.001$ ; \* $p < 0.05$ ; NS, not significant (Tukey–Kramer test)



**Fig. 4** Mycothiol (MSH) is needed for optimal actinorhodin (ACT) synthesis. **a** Spores were inoculated onto yeast extract-malt extract-glutamine agar medium with a toothpick and cultured for 96 h at 30 °C. **b** M145 (blue bar),  $\Delta mshC$  (red bar), and  $\Delta mshC::mshC$  (yellow bar) strains were cultured under the same conditions as those in (a). Total RNA was extracted from each strain to measure the relative expression level of *actII-ORF4* using reverse transcription-quantitative polymerase chain reaction. Data are presented as mean and standard deviation values from three independent experiments. \*\*\* $p < 0.001$  (Student’s *t*-test)

found in the  $\Delta mshC$  strain inoculated on YEME-gln agar and cultured for 96 h. These results suggest that the lack of MSH negatively affects NO signalling and ACT production in *S. coelicolor* A3(2) M145 cells.

Thus, MSH is essential for maintaining intracellular NO homeostasis and for mediating NO signalling in *S. coelicolor* A3(2) M145.

## Discussion

As NO is highly reactive, its cellular concentration must be strictly regulated to exert its physiological functions. The results of the present study indicate that MSH-mediated intracellular NO homeostasis facilitates optimal NO signalling in bacteria. The GSH synthesis pathway has not been detected in *Actinomycetes*; however, members of this bacterial genus produce MSH as their major redox buffer and coenzyme [26–28]. MSH-deficient mutants of *Mycobacterium smegmatis* exhibit increased sensitivity to exogenous NO [29], reactive oxygen species [30], alkali agents, and antibiotics [31]. *Mycobacterium tuberculosis* relies on MSH-dependent systems to infect macrophages [32]. *Corynebacterium glutamicum* metabolises aromatic compounds that require MSH as a coenzyme [33]. Thus, *Actinomycetes* use MSH to adapt to their environments. In addition, an MSH-defective strain of *S. coelicolor* A3(2) was reported to exhibit high sensitivity to hydrogen peroxide [34].

While being statistically significant, the differences in cell growth between the M145 and  $\Delta mshC$  strains remained relatively small during culture. However, the lack of MSH due to *mshC* deletion disrupted intracellular NO homeostasis. These results suggest that MSH acts not only as a defence molecule against reactive oxygen species but also as a mediator involved in maintaining intracellular NO homeostasis.

Although previous studies have revealed that bacteria endogenously produce NO and possess regulatory mechanisms for NO-mediated metabolism, these are transient adaptive responses to environmental changes [35–38]. *Streptomyces coelicolor* A3(2) M145 scavenges endogenously produced NO via MSH, in addition to Fhb-mediated NO metabolism. NO signalling can start gradually as intracellular NO levels overcome the metabolisable levels of both mechanisms. Therefore, *S. coelicolor* A3(2) M145 possesses an advanced mechanism for regulating intracellular NO homeostasis.

We further found that the ACT production level in the  $\Delta mshC$  strain was lower than that in the WT strain. The DevS/R activity in the  $\Delta mshC$  strain was also lower than that in the WT strain. ACT production is regulated by NO signalling via the DevS/R TCS [18]. Therefore, the dysregulation of ACT production in the  $\Delta mshC$  strain

can be explained by the upregulation of intracellular NO-induced DevS/R repression in the absence of MSH.

Thiols bind directly to NO under aerobic conditions [39] or in the presence of electron acceptors such as NAD<sup>+</sup>, metals, and metalloproteins in vivo [24, 25]. GSH forms S-nitrosoglutathione (GSNO) with the intracellular NO produced by nitric oxide synthase to maintain the life cycle of mammalian cells [40]. GSNO is a stable NO-carrying molecule in both plant and animal cells that regulates protein activity by transferring an NO group to cysteines [41–43]. LMW S-nitrosothiols function as donors of NO and nitrosonium ions and are involved in cellular signalling [44, 45]. S-Nitrosothiols are required for cell survival. In particular, GSNO interacts with specific thiol groups of proteins to produce high-molecular-weight S-nitrosothiols in a process called S-transnitrosation [46] and regulates the secondary metabolite content through protein S-nitrosylation [47].

LMW S-nitrosothiols were detected in the WT M145 strain. In contrast, LMW S-nitrosothiols were almost undetectable in the  $\Delta mshC$  strain. This suggests that MSH interacts with endogenously produced NO and regulates the intracellular NO concentration through MSNO formation; however, we did not identify MSNO in the present study. In the future, it will be important to elucidate the mechanisms by which LMW S-nitrosothiols, including MSH, affect NO metabolism in actinomycetes.

The LMW S-nitrosothiol levels were not significantly different between the M145 and  $\Delta mshC$  strains at 192 h, suggesting that the amount of other LMW thiols increased instead of MSH and that these thiols were involved in maintaining intracellular NO levels in the late stages of culture. This is consistent with the results of a previous study, which demonstrated that the amounts of other thiol compounds increased in an MSH-deficient strain of *S. coelicolor* A3(2) [34]. However, the levels of LMW S-nitrosothiols in the *mshC* strain were lower than those in the M145 strain at both 96 and 144 h. MSH deficiency inactivated the DevS/R TCS from the early stages, indicating that MSH is the main LMW thiol for maintaining NO homeostasis in the early stages of culture, which is necessary for signalling secondary metabolism in *S. coelicolor* A3(2) M145. *Actinomycetes* synthesise ergothioneine, an LMW thiol produced by Actinobacteria, in addition to MSH [48]. Thus, to understand the homeostatic regulation of NO in *Actinomycetes*, the roles of other thiols in NO homeostasis must be considered.

This study suggests that intracellular NO homeostasis, which must be strictly maintained for optimal secondary metabolism, depends on MSH, a novel component that regulates NO signalling. MSH deficiency decreased ACT production. Elucidation of the regulatory mechanism of secondary metabolism will contribute to the efficient

improvement of fermentation products. Additionally, *M. tuberculosis* has a similar form of NO metabolism [49, 50]. Therefore, identifying the role of MSH in maintaining intracellular NO homeostasis will enable the development of clinical therapeutic strategies against *M. tuberculosis* infections. Future studies should focus on the regulation of NO homeostasis through various mechanisms in several organisms, which can provide novel perspectives on NO biology and improvement of human health.

## Conclusions

This study suggests that MSH regulates intracellular NO homeostasis by promoting MSNO formation. MSH-mediated maintenance of NO homeostasis, which is dependent on intracellular NO levels, is important for ACT production in *S. coelicolor* A3(2) M145 under NO-producing conditions (Fig. 5).

## Methods


### Bacterial strains, plasmids, and culture conditions

The *S. coelicolor* A3(2) M145 parental strain and all mutants used in this study are listed in Table 1. To induce

sporulation, the strains were cultured on mannitol soya flour agar (2% mannitol, 2% soya flour, and 2% agar) at 30 °C. Additionally, the strains were incubated for each experiment in YEME-gln agar medium (0.3% yeast extract, 0.3% malt extract, 0.5% bacto-peptone, 1% glucose, 50 mM L-glutamine, and 2% agar, pH 7.2) at 30 °C. As NO easily diffuses and affects strains grown in the same incubator, the  $\Delta nar$  and  $\Delta nar/mshC$  strains were incubated independently. Detailed culture conditions are described in the figure legends. *Escherichia coli* DH5 $\alpha$  and *E. coli* HST04 *dam*<sup>-</sup>/*dcm*<sup>-</sup> (TaKaRa Bio) strains were used as hosts for cloning and cultured in Luria–Bertani broth (1% tryptone, 0.5% yeast extract, and 0.5% NaCl) at 37 °C.

### Construction of the $\Delta mshC$ and complementation strains

Two fragments corresponding to the upstream and downstream regions of *mshC* (*sco1663*) and a fragment corresponding to the apramycin-resistance cassette [51] flanked by flippase recognition target sites were polymerase chain reaction (PCR)-amplified with PrimeSTAR GXL DNA polymerase (TaKaRa Bio) using specific primer sets (Table 2), along with the introduction of



**Fig. 5** Proposed model of intracellular nitric oxide (NO) homeostasis in *Streptomyces coelicolor* A3(2) M145. Flavohaemoglobin (Fhb) and nitrate reductase (Nar) mediate the circulation of NO, NO<sub>3</sub><sup>-</sup>, and NO<sub>2</sub><sup>-</sup>. DevS/R is activated or inactivated depending on intracellular NO levels. NO directly regulates actinorhodin synthesis through the DevS/R two-component regulatory system, which modulates *actII-ORF4* expression. DevR also regulates *nar2* expression and NO production. Intracellular NO homeostasis is maintained through Fhb metabolism and its interaction with mycothiol (MSH) (S-nitrosylation) to form S-nitrosomycothiol (MSNO)

**Table 1** Strains used in this study

<i>Streptomyces coelicolor</i> A3(2) M145 strain	Genotype and/or characteristics	Reference or source
M145 (wild type)	Lacks native plasmids, SCP1 and SCP2	[17]
$\Delta nar$	M145 SCO6535::hyg/SCO0216::scar/SCO4947::scar ( <i>narGG2G3</i> removed)	[17]
$\Delta mshC$	M145 SCO1663::aac(3)IV ( <i>mshC</i> removed)	This study
$\Delta mshC::mshC$	M145 SCO1663::aac(3)IV/pku460-SCO1163 (complementation of <i>mshC</i> )	This study
$\Delta nar/mshC$	M145 SCO6535::hyg/SCO0216::scar/SCO4947::scar/SCO1663::aac(3)IV ( <i>narGG2G3</i> and <i>mshC</i> removed)	This study

**Table 2** Primers used in this study

Primer name	Sequence (5' → 3')	Purpose
hrdB F	GCATGCTCTTCTGGACCTCAT	qPCR
hrdB R	TGGAGAACTTGTAGCCCTTGGTGTA	qPCR
hmpA F	AACAGTCCGTTCCCGTGGT	qPCR
hmpA R	CGAAGAGCTTGCGGTAGAACAG	qPCR
actII-ORF4 F	AAAGGAATATCGCGCACCTGGAAG	qPCR
actII-ORF4 R	GTTCCGGAATCATCGGCCCTATTC	qPCR
devR F	ACGACGAACCGGACATCACC	qPCR
devR R	CGTCGTCGAAGGAGGTCAGC	qPCR
mshA F	GTGAGCCAGTACGTCAGCAG	qPCR
mshA R	AGCTCCACGATGTAGACGTTC	qPCR
mshB F	TCGCGAAGGTCTACTGGAAC	qPCR
mshB R	CCTTCTCGAACGGCAGTCC	qPCR
mshC F	CGTACGACGCCACCCACAT	qPCR
mshC R	TAGTGGACCTGCCGCTTGGT	qPCR
mshD F	TTGACCGGCTGGACCTG	qPCR
mshD R	TGAGCGACCCCTGTTCGG	qPCR
ΔmshC upstream F	AAGCTTTAAACAGGCGTACAGACAGAAGAAAGA	Disruption
ΔmshC upstream R	GAATTCACCCCGATCGACAAGTACCG	Disruption
ΔmshC downstream F	GAATTCGCCGAACCTGGGGAACGAG	Disruption
ΔmshC downstream R	AAGCTTCATGTCATGAGCCTAACCCGGA	Disruption
FRT drug resi F	CTCGAGAAGCTTATTCCGGGGATCCGTCGACC	Disruption
FRT drug resi R	CTCGAGAAGCTTGTAGGCTGGAGCTGCTTC	Disruption
comp mshC F	CTCGAGGAATCCCGCTGAAGGACGGCGCA	Complementation
comp mshC R	CTCGAGAAGCTTTTACAGGCCACGCCCAGC	Complementation

qPCR quantitative polymerase chain reaction

*EcoRI* and *HindIII* restriction sites. The two fragments were ligated into the *SmaI*-digested pUC19 plasmid using Ligation High Ver. 2 (TOYOBO). The two products were transformed into *E. coli* DH5α competent cells (TaKaRa Bio). The plasmids were purified from *E. coli* cells, digested with *EcoRI* and *HindIII*, and the two fragments were ligated into the *EcoRI*-digested pKU450 plasmid using Ligation High Ver. 2. The resulting plasmids were transformed into *E. coli* DH5α competent cells. The plasmids were then purified. After digestion of the apramycin-resistance cassettes and plasmids with *HindIII*, the two fragments were ligated using Ligation High Ver. 2. The resulting products were transformed into *E. coli* DH5α competent cells. The cosmids were purified, digested with *EcoRI*, and ligated with *EcoRI*-digested pGM160 using Ligation High Ver. 2. The resulting construct was introduced into *E. coli* HST04 *dam*<sup>-</sup>/*dcm*<sup>-</sup> competent cells (TaKaRa Bio) to obtain non-methylated cosmids. Purified cosmids were introduced into the *S. coelicolor* A3(2) M145 parental strain or each mutant. The drug-resistant recombinant strains were screened. Genomic DNA was checked for DNA recombination by PCR with the appropriate primer sets (Table 2).

The pKU460 plasmid was used for genetic complementation of the knockout mutants. The complementary cassette was obtained by PCR using the complementary primer set (Table 2) and ligated to the *SmaI*-digested pUC19 plasmid using Ligation High Ver. 2. The ligation product was introduced into *E. coli* DH5α competent cells, and the cosmid pUC19-*mshC* was purified and ligated with pKU460. The ligation product pKU460-*mshC* was introduced into *E. coli* HST04 *dam*<sup>-</sup>/*dcm*<sup>-</sup> competent cells to obtain a non-methylated cosmid. The obtained cosmid was then introduced into the Δ*mshC* strain. The complementary mutant was verified by PCR using the appropriate primer sets (Table 2) and a complementation study.

#### Measurement of colony areas

Spores were incubated on the YEME-gln agar medium at intervals of 1.8 cm (total of nine spots, 3×3 squares) using a sterilised toothpick. After incubating at 30 °C, a line of colonies (three colonies) including the central colony were photographed under a microscope (OLYMPUS) at a 7× multiplier, and then the nine colony areas for

three plates were measured using ImageJ (Version 1.53t, 24 August 2022 upgrade).

#### Measurement of cell wet weight

Spores were incubated on YEME-gln agar medium at intervals of 1.8 cm (total of nine spots, 3×3 squares) using a sterilised toothpick. After incubating at 30 °C, all colonies were collected in a microtube while carefully removing the agar medium with a toothpick and then the cell wet weight was measured using a precision balance.

#### S-nitrosothiol quantification

Spores were incubated on YEME-gln agar medium at intervals of 1.8 cm (total of nine spots, 3×3 squares) using a sterilised toothpick. After incubating at 30 °C, all colonies were collected in a 10 ml conical tube while carefully removing the agar medium with a toothpick.

S-nitrosothiol was measured using the Griess assay [52]. Cells were lysed in extraction buffer (0.1 M Tris-HCl, pH 7.2) on ice using sonication. The lysate was centrifuged at 20,000×g and 4 °C for 15 min. Protein concentrations were measured using Bradford Protein assay kit (BIO-RAD) [53]. To precipitate proteins, the cell-free extract was incubated with acetone at -20 °C for 15 min. The samples were then centrifuged at 20,000×g and 4 °C for 10 min. Acetone was removed using an evaporator. The sample was incubated with an equivalent volume of buffer A (0.5 M HCl, 1% sulphanilamide) or buffer B (0.5 M HCl, 1% sulphanilamide, 0.2% HgCl<sub>2</sub>) for 15 min at 20–25 °C, followed by incubation with an equal volume of Griess reagent [0.5 M HCl, 0.02% *N*-(1-naphthyl)-ethylenediamine dihydrochloride] for 15 min. S-nitrosothiol was quantified by measuring the absorbance of the reaction mixture at 540 nm. A standard curve was constructed using different concentrations (0, 1, 5, 10, 15, and 20 μM) of GSNO in 50 μM NaNO<sub>2</sub>. Concentrations below the detection limit (500 nM) [54].

#### RNA extraction and reverse transcription-quantitative PCR (RT-qPCR)

Spores were inoculated on YEME-gln agar medium at intervals of 1.8 cm (total of nine spots, 3×3 squares) using a sterilised toothpick. After incubating at 30 °C, all colonies were collected in a microtube while carefully removing the agar medium with a toothpick. Total RNA was purified from cells using the ReliaPrep RNA Cell Miniprep System (Promega). The isolated RNA was reverse-transcribed into complementary DNA using ReverTra Ace RT-qPCR Master Mix with gDNA Remover (TOYOBO), which was used as a template for qPCR analysis. The primers used for RT-qPCR analysis are listed in Table 2. qPCR was performed using

the Thermal Cycler Dice Real-Time System (TaKaRa Bio). The PCR mixture (20 μL) contained 1 μg cDNA, 10 pmol of an appropriate primer set (Table 2), and THUNDERBIRD SYBR qPCR Mix (TOYOBO) or KOD SYBR qPCR Mix (TOYOBO). The expression levels of the target genes were normalised to those of *hrdB*, a housekeeping gene.

#### ACT production and RNA extraction

The strains were inoculated in a triangular shape in the YEME-GLN medium. The ACT-derived blue pigment was visualised using a camera (Canon). Total RNA was extracted from the cells using the ReliaPrep RNA Cell Miniprep System (Promega) by the same technique as described above.

#### Statistical analysis and reproducibility

All bars in figures represent the mean values obtained from at least three independent biological experiments and error bars represent the standard deviation. The detailed sample number (*n*) is provided in the legend of each figure. Differences in means between strains or conditions were analysed using Student's *t*-test, Dunnett's test, or the Tukey-Kramer test, depending on the comparison items. Differences were considered statistically significant at *p* < 0.05.

#### Abbreviations

ACT	Actinorhodin
Fhb	Flavo-haemoglobin
GSH	Glutathione
GSNO	S-nitrosoglutathione
LMW	Low molecular weight
MSH	Mycotoxin
MSNO	S-nitrosomycotoxin
Nar	Nitrate reductase
NO	Nitric oxide
PCR	Polymerase chain reaction
RT-qPCR	Reverse transcription-quantitative polymerase chain reaction
TCS	Two-component system
WT	Wild type
YEME-gln	Yeast extract-malt extract-glutamine

#### Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s12866-023-03036-z>.

Additional file 1.

#### Acknowledgements

Not applicable.

#### Authors' contributions

T.Y., Y.S., M.K., S.H., S.I., S.Y., and Y.S. designed the study. T.Y., M.K. and Y.S. carried out experiments and analyzed data. T.Y. prepared the manuscript. All authors approved final draft.

### Funding

This work was supported in part by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science (C; Grant No. 20K05811) and a research project from the Tokyo University of Agriculture.

### Availability of data and materials

The dataset supporting the conclusions of this study is included in the article and an additional file (Additional file 1). All data will be made available from the corresponding author upon request.

### Declarations

#### Ethics approval and consent to participate

Not applicable.

#### Consent for publication

Not applicable.

#### Competing interests

The authors declare no competing interests.

Received: 8 December 2022 Accepted: 1 October 2023

Published online: 05 October 2023

### References

1. Feelisch M, Stamler J. *Methods in nitric oxide research*. Wiley-Blackwell; 1996.
2. Palmer RM, Ferrige AG, Moncada S. Nitric oxide release accounts for the biological activity of endothelium-derived relaxing factor. *Nature*. 1987;327:524–6. <https://doi.org/10.1038/327524a0>.
3. Ford PC, Lorkovic IM. Mechanistic aspects of the reactions of nitric oxide with transition-metal complexes. *Chem Rev*. 2002;102:993–1018. <https://doi.org/10.1021/cr0000271>.
4. Gaston B. Nitric oxide and thiol groups. *Biochim Biophys Acta*. 1999;1411:323–33. [https://doi.org/10.1016/s0005-2728\(99\)00023-7](https://doi.org/10.1016/s0005-2728(99)00023-7).
5. Rubbo H, Darley-Usmar V, Freeman BA. Nitric oxide regulation of tissue free radical injury. *Chem Res Toxicol*. 1996;9:809–20. <https://doi.org/10.1021/tx960037q>.
6. Hogg N, Kalyanaram B. Nitric oxide and lipid peroxidation. *Biochim Biophys Acta*. 1999;1411:378–84. [https://doi.org/10.1016/s0005-2728\(99\)00027-4](https://doi.org/10.1016/s0005-2728(99)00027-4).
7. Pacher P, Beckman JS, Liaudet L. Nitric oxide and peroxynitrite in health and disease. *Physiol Rev*. 2007;87:315–424. <https://doi.org/10.1152/physrev.00029.2006>.
8. Aktan F. iNOS-mediated nitric oxide production and its regulation. *Life Sci*. 2004;75:639–53. <https://doi.org/10.1016/j.lfs.2003.10.042>.
9. Reutov VP, Sorokina EG. NO-synthesis and nitrite-reductase components of nitric oxide cycle. *Biochemistry (Mosc)*. 1998;63:874–84.
10. Gardner PR, Gardner AM, Martin LA, Salzman AL. Nitric oxide dioxygenase: an enzymic function for flavohemoglobin. *Proc Natl Acad Sci U S A*. 1998;95:10378–83. <https://doi.org/10.1073/pnas.95.18.10378>.
11. Zumft WG. Cell biology and molecular basis of denitrification. *Microbiol Mol Biol Rev*. 1997;61:533–616. <https://doi.org/10.1128/mmb.61.4.533-616.1997>.
12. Smith BC, Marletta MA. Mechanisms of S-nitrosothiol formation and selectivity in nitric oxide signaling. *Curr Opin Chem Biol*. 2012;16:498–506. <https://doi.org/10.1016/j.cbpa.2012.10.016>.
13. Zhou S, Narukami T, Masuo S, Shimizu M, Fujita T, Doi Y, et al. NO-inducible nitrosothionein mediates NO removal in tandem with thioredoxin. *Nat Chem Biol*. 2013;9:657–63. <https://doi.org/10.1038/nchembio.1316>.
14. Noomnual S, Thasana N, Sungkeeree P, Mongkolsuk S, Loprasert S. Streptanoate, a new anticancer butanoate from *Streptomyces* sp. DC3. *J Antibiot (Tokyo)*. 2016;69:124–7. <https://doi.org/10.1038/ja.2015.95>.
15. Ishiyama T, Hara I, Matsuoka M, Sato K, Shimada S, Izawa R, et al. Studies on the preventive effect of kasugamycin on rice blast. *J Antibiot (Tokyo)*. 1965;18:115–9.
16. Watve MG, Tickoo R, Jog MM, Bhole BD. How many antibiotics are produced by the genus *Streptomyces*? *Arch Microbiol*. 2001;176:386–90. <https://doi.org/10.1007/s002030100345>.
17. Sasaki Y, Oguchi H, Kobayashi T, Kusama S, Sugiura R, Moriya K, et al. Nitrogen oxide cycle regulates nitric oxide levels and bacterial cell signaling. *Sci Rep*. 2016;6:22038. <https://doi.org/10.1038/srep22038>.
18. Honma S, Ito S, Yajima S, Sasaki Y. Nitric oxide signaling for actinorhodin production in *Streptomyces coelicolor* A3(2) via the DevS/R Two-Component System. *Appl Environ Microbiol*. 2021;87:e0048021. <https://doi.org/10.1128/AEM.00480-21>.
19. Newton GL, Buchmeier N, Fahey RC. Biosynthesis and functions of mycothiol, the unique protective thiol of Actinobacteria. *Microbiol Mol Biol Rev*. 2008;72:471–94. <https://doi.org/10.1128/MMBR.00008-08>.
20. Ung KS, Av-Gay Y. Mycothiol-dependent mycobacterial response to oxidative stress. *FEBS Lett*. 2006;580:2712–6. <https://doi.org/10.1016/j.febslet.2006.04.026>.
21. Imber M, Pietrzyk-Brzezinska AJ, Antelmann H. Redox regulation by reversible protein S-thiolation in Gram-positive bacteria. *Redox Biol*. 2019;20:130–45. <https://doi.org/10.1016/j.redox.2018.08.017>.
22. Park JH, Cha CJ, Roe JH. Identification of genes for mycothiol biosynthesis in *Streptomyces coelicolor* A3(2). *J Microbiol*. 2006;44:121–5.
23. Tucker NP, Le Brun NE, Dixon R, Hutchings MI. There's NO stopping NsrR, a global regulator of the bacterial NO stress response. *Trends Microbiol*. 2010;18:149–56. <https://doi.org/10.1016/j.tim.2009.12.009>.
24. Gow AJ, Buerk DG, Ischiropoulos H. A novel reaction mechanism for the formation of S-nitrosothiol in vivo. *J Biol Chem*. 1997;272:2841–5. <https://doi.org/10.1074/jbc.272.5.2841>.
25. Bosworth CA, Toledo JC, Zmijewski JW, Li Q, Lancaster JR. Dinitrosyliron complexes and the mechanism(s) of cellular protein nitrosothiol formation from nitric oxide. *Proc Natl Acad Sci U S A*. 2009;106:4671–6. <https://doi.org/10.1073/pnas.0710416106>.
26. Fahey RC, Brown WC, Adams WB, Worsham MB. Occurrence of glutathione in bacteria. *J Bacteriol*. 1978;133:1126–9. <https://doi.org/10.1128/jb.133.3.1126-1129.1978>.
27. Newton GL, Fahey RC, Cohen G, Aharonowitz Y. Low-molecular-weight thiols in streptomycetes and their potential role as antioxidants. *J Bacteriol*. 1993;175:2734–42. <https://doi.org/10.1128/jb.175.9.2734-2742.1993>.
28. Jothivasan VK, Hamilton CJ. Mycothiol: synthesis, biosynthesis and biological functions of the major low molecular weight thiol in actinomycetes. *Nat Prod Rep*. 2008;25:1091–117. <https://doi.org/10.1039/b616489g>.
29. Miller CC, Rawat M, Johnson T, Av-Gay Y. Innate protection of *Mycobacterium smegmatis* against the antimicrobial activity of nitric oxide is provided by mycothiol. *Antimicrob Agents Chemother*. 2007;51:3364–6. <https://doi.org/10.1128/AAC.00347-07>.
30. Newton GL, Unson MD, Anderberg SJ, Aguilera JA, Oh NN, delCardayre SB, et al. Characterization of *Mycobacterium smegmatis* mutants defective in 1-d-myo-inosityl-2-amino-2-deoxy-alpha-d-glucopyranoside and mycothiol biosynthesis. *Biochem Biophys Res Commun*. 1999;255:239–44. <https://doi.org/10.1006/bbrc.1999.0156>.
31. Rawat M, Newton GL, Ko M, Martinez GJ, Fahey RC, Av-Gay Y. Mycothiol-deficient *Mycobacterium smegmatis* mutants are hypersensitive to alkylating agents, free radicals, and antibiotics. *Antimicrob Agents Chemother*. 2002;46:3348–55. <https://doi.org/10.1128/AAC.46.11.3348-3355.2002>.
32. Sassetti CM, Rubin EJ. Genetic requirements for mycobacterial survival during infection. *Proc Natl Acad Sci U S A*. 2003;100:12989–94. <https://doi.org/10.1073/pnas.2134250100>.
33. Feng J, Che Y, Milse J, Yin YJ, Liu L, Rückert C, et al. The gene ncgl2918 encodes a novel maleylpyruvate isomerase that needs mycothiol as cofactor and links mycothiol biosynthesis and gentisate assimilation in *Corynebacterium glutamicum*. *J Biol Chem*. 2006;281:10778–85. <https://doi.org/10.1074/jbc.M513192200>.
34. Nakajima S, Satoh Y, Yanashima K, Matsui T, Dairi T. Ergothioneine protects *Streptomyces coelicolor* A3(2) from oxidative stresses. *J Biosci Bioeng*. 2015;120:294–8. <https://doi.org/10.1016/j.jbiosc.2015.01.013>.
35. Corker H, Poole RK. Nitric oxide formation by *Escherichia coli*. Dependence on nitrite reductase, the NO-sensing regulator Fnr, and flavohemoglobin Hmp. *J Biol Chem*. 2003;278:31584–92. <https://doi.org/10.1074/jbc.M303282200>.

36. Renner ED, Becker GE. Production of nitric oxide and nitrous oxide during denitrification by *Corynebacterium nephridii*. *J Bacteriol.* 1970;101:821–6. <https://doi.org/10.1128/jb.101.3.821-826.1970>.
37. Silva WM, Carvalho RD, Soares SC, Bastos IF, Folador EL, Souza GH, et al. Label-free proteomic analysis to confirm the predicted proteome of *Corynebacterium pseudotuberculosis* under nitrosative stress mediated by nitric oxide. *BMC Genomics.* 2014;15:1065. <https://doi.org/10.1186/1471-2164-15-1065>.
38. Bodenmiller DM, Spiro S. The *yjeB* (*nsrR*) gene of *Escherichia coli* encodes a nitric oxide-sensitive transcriptional regulator. *J Bacteriol.* 2006;188:874–81. <https://doi.org/10.1128/JB.188.3.874-881.2006>.
39. Kolesnik B, Palten K, Schrammel A, Stessel H, Schmidt K, Mayer B, et al. Efficient nitrosation of glutathione by nitric oxide. *Free Radic Biol Med.* 2013;63:51–64. <https://doi.org/10.1016/j.freeradbiomed.2013.04.034>.
40. Ferrini ME, Simons BJ, Bassett DJ, Bradley MO, Roberts K, Jaffar Z. S-nitrosoglutathione reductase inhibition regulates allergen-induced lung inflammation and airway hyperreactivity. *PLoS ONE.* 2013;8: e70351. <https://doi.org/10.1371/journal.pone.0070351>.
41. Corpas FJ, Alché JD, Barroso JB. Current overview of S-nitrosoglutathione (GSNO) in higher plants. *Front Plant Sci.* 2013;4:126. <https://doi.org/10.3389/fpls.2013.00126>.
42. Broniowska KA, Diers AR, Hogg N. S-nitrosoglutathione. *Biochim Biophys Acta.* 1830;2013:3173–81. <https://doi.org/10.1016/j.bbagen.2013.02.004>.
43. Zaffagnini M, Fermani S, Marchand CH, Costa A, Sparla F, Rouhier N, et al. Redox homeostasis in photosynthetic organisms: novel and established thiol-based molecular mechanisms. *Antioxid Redox Signal.* 2019;31:155–210. <https://doi.org/10.1089/ars.2018.7617>.
44. Arnelle DR, Stamler JS. NO<sup>+</sup>, NO, and NO<sup>-</sup> donation by S-nitrosothiols: implications for regulation of physiological functions by S-nitrosylation and acceleration of disulfide formation. *Arch Biochem Biophys.* 1995;318:279–85. <https://doi.org/10.1006/abbi.1995.1231>.
45. Lipton AJ, Johnson MA, Macdonald T, Lieberman MW, Gozal D, Gaston B. S-nitrosothiols signal the ventilatory response to hypoxia. *Nature.* 2001;413:171–4. <https://doi.org/10.1038/35093117>.
46. Hogg N. The biochemistry and physiology of S-nitrosothiols. *Annu Rev Pharmacol Toxicol.* 2002;42:585–600. <https://doi.org/10.1146/annurev.pharmtox.42.092501.104328>.
47. Liu R, Zhu T, Chen X, Wang Z, Yang Z, Ren A, et al. GSNOR regulates ganoderic acid content in *Ganoderma lucidum* under heat stress through S-nitrosylation of catalase. *Commun Biol.* 2022;5:32. <https://doi.org/10.1038/s42003-021-02988-0>.
48. Fahey RC. Glutathione analogs in prokaryotes. *Biochim Biophys Acta.* 2013;1830:3182–98. <https://doi.org/10.1016/j.bbagen.2012.10.006>.
49. Cunningham-Bussel A, Zhang T, Nathan CF. Nitrite produced by *Mycobacterium tuberculosis* in human macrophages in physiologic oxygen impacts bacterial ATP consumption and gene expression. *Proc Natl Acad Sci U S A.* 2013;110:E4256–65. <https://doi.org/10.1073/pnas.1316894110>.
50. Cunningham-Bussel A, Bange FC, Nathan CF. Nitrite impacts the survival of *Mycobacterium tuberculosis* in response to isoniazid and hydrogen peroxide. *Microbiol Open.* 2013;2:901–11. <https://doi.org/10.1002/mbo3.126>.
51. Gust B, Challis GL, Fowler K, Kieser T, Chater KF. PCR-targeted *Streptomyces* gene replacement identifies a protein domain needed for biosynthesis of the sesquiterpene soil odor geosmin. *Proc Natl Acad Sci U S A.* 2003;100(4):1541–6. <https://doi.org/10.1073/pnas.0337542100>. Epub 2003 Jan 31.
52. Thalineau E, Truong HN, Berger A, Fournier C, Boscardi A, Wendehenne D, et al. Cross-regulation between N metabolism and nitric oxide (NO) signaling during plant immunity. *Front Plant Sci.* 2016;7:472. <https://doi.org/10.3389/fpls.2016.00472>.
53. Bradford MM. A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Anal Biochem.* 1976;7(72):248–54. <https://doi.org/10.1006/abio.1976.9999>.
54. Gow A, Doctor A, Mannick J, Gaston B. S-Nitrosothiol measurements in biological systems. *J Chromatogr B Anal Technol Biomed Life Sci.* 2007;851(1–2):140–51. <https://doi.org/10.1016/j.jchromb.2007.01.052>. Epub 2007 Feb 25.

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more [biomedcentral.com/submissions](https://biomedcentral.com/submissions)

