

One-pot synthesis of 2-oxa-7-azaspiro[4.4]nonane-8,9-diones using Mn(III)-based oxidation of 4-acylpyrrolidine-2,3-diones [☆]



Thanh-Truc Huynh ^a, Van-Ha Nguyen ^b, Hiroshi Nishino ^{c,*}

^a Department of Chemistry, Graduate School of Science and Technology, Kumamoto University, Kurokami, Kumamoto 860-8555, Japan

^b Department of Chemistry, Dalat University, 1 Phu Dong Thien Vuong St., Dalat, Viet Nam

^c Department of Chemistry, Faculty of Science, Kumamoto University, Kurokami 2-39-1, Chûou-Ku, Kumamoto 860-8555, Japan

ARTICLE INFO

Article history:

Received 7 June 2017

Revised 27 July 2017

Accepted 1 August 2017

Available online 3 August 2017

Keywords:

Oxaazaspiro[4.4]nonanedione

Oxidation

1,1-Diarylethenes

4-Acylpyrrolidine-2,3-diones

Manganese(III) acetate

Cycloaddition

ABSTRACT

2-Oxa-7-azaspiro[4.4]nonane-8,9-diones were newly synthesized in good yields by the Mn(III)-based reaction of a mixture of 1,1-diarylethenes and 4-acylpyrrolidine-2,3-diones. Under the stated reaction conditions, the pyrrolidinedione ring remained intact and became one of the two rings of the 2-oxa-7-azaspiro[4.4]nonanedione scaffold. The procedure was simple and the product was easily separated. The structure determination and the mechanism for the formation of the 2-oxa-7-azaspiro[4.4]nonanediones were also discussed.

© 2017 Elsevier Ltd. All rights reserved.

Introduction

The Mn(III)-based oxidation is a powerful tool in organic synthesis, and new applications and protocols have been continuously reported.^{1,2} Tetrahydrofuran derivatives are found in many natural compounds, some of which derived from tetrose, pentose, hexose, and glycoside, have a significant biological importance.³ Pyrrolidinediones have displayed an interesting biological activity and been used as an inhibitor of aldose reductase⁴ and endothelin receptor antagonists.⁵ Pyrrolidinediones⁶ were also used as a versatile reagent for the preparation of β -lactams.⁷ We previously reported a unique synthesis of spirodi- γ -lactones,^{8a} spirodioxanes,^{8b} trioxaspiro compounds,^{8c} dioxatricyclic^{8d} and oxa-aza-tricyclic compounds,^{8e} spirofurans,^{8f} and aza-spiro compounds^{8g} using manganese(III) acetate dihydrate, Mn(OAc)₃·2H₂O. In the course of our study, we found a simple and straightforward route based on the Mn(III) oxidation for the synthesis of new pyrrolidine-2,3-diones **I** with an ethenyl group substituted at the α -position of the carbonyl group (Scheme 1).^{9,10} At that time, we anticipated producing a spiro compound (path a), but the deprotonation was fast under the stated

conditions (path b). Based on these results, we postulated if the presence of a keto-carbonyl group at the C-4 position of the pyrrolidine-2,3-dione instead of an ester could allow the cyclization to produce a spiro bicyclic compound such as a furan connected through the α -carbon of the pyrrolidinediones. We then attempted to verify this idea using the Mn(III)-based oxidation of 1,1-diarylethenes with 4-acylpyrrolidine-2,3-diones as the starting material. As a result, the cyclization proceeded at the carbonyl oxygen and new 1-exomethylene-2-oxa-7-azaspiro[4.4]nonane-8,9-dione derivatives were produced in good yields as expected.


Results and discussion

The pyrrolidine-2,3-diones **2a–e**^{9,11} were prepared by the condensation of 2,4-dioxoalkanoates¹² with *N*-benzyl or *N*-methyl-methanimines.¹³ The pyrrolidinediones were purified by silica gel column chromatography, then recrystallization. With the starting material of the pyrrolidinediones **2a–e** in hand, we commenced the Mn(III)-based oxidation in the presence of 1,1-disubstituted alkenes. 1,1-Bis(4-methylphenyl)ethene (**1a**) and 1-benzyl-4-isobutyrylpyrrolidine-2,3-dione (**2a**) were first selected and the reaction using Mn(OAc)₃·2H₂O was carried out in glacial acetic acid at 70 °C. Since the oxidant was consumed in 16 min, the reaction was quenched and the mixture was worked up. Gratifyingly, the desired 2-oxa-7-azaspiro compound **3aa** was obtained in 38% isolated yield (Scheme 2 and Table 1, Entry 1).


[☆] Dedicated to Dr. Kazu Kurosawa, Professor Emeritus of Kumamoto University, on his 80th birthday.

* Corresponding author.

E-mail address: nishino@sci.kumamoto-u.ac.jp (H. Nishino).



Scheme 1. Mn(III)-based reaction of alkene with 4-hydroxy-5-oxo-2,5-dihydropyrrole-3-carboxylate.



Scheme 2. Mn(III)-based reaction of alkenes **1a–e** with 4-acylpyrrolidine-2,3-diones **2a–e**.

When a similar reaction was performed at reflux temperature, the reaction was finished in 2 min and the yield of **3aa** was significantly improved (Entry 2). We then optimized the reaction and the maximum yield of **3aa** was 90% (Entry 4). The structure of **3aa** was determined by IR, ^1H NMR, ^{13}C NMR, and ^{13}C DEPT spectra, a 2D NMR study and elemental analysis. The presence of the dimethylmethylene group and C-5 spiro carbon was confirmed by the NMR spectrum, and the HMBC experiment was also in good agreement with the structure of 7-benzyl-3,3-bis(4-methylphenyl)-1-(propan-2-ylidene)-2-oxa-7-azaspiro[4.4]nonane-8,9-dione (Fig. 1).¹⁴

Having succeeded in the synthesis of the anticipated spiro compound **3aa**, we turned our attention to a similar reaction using the other 1,1-diarylethenes **1b–e**. The reaction was conducted under similar conditions and the corresponding 2-oxa-7-azaspiro[4.4]nonane-8,9-diones **3ba–ea** were obtained in good yields (Table 1, Entries 5–8). The use of 4-propionylpyrrolidine-2,3-dione **2b** instead of **2a** also gave the 2-oxa-7-azaspiro compound **3ab** in 81% yield (Entry 9). Although the reaction of 5-phenyl-4-propionyl- **2c** and the 4-butyryl-5-phenyl-pyrrolidine-2,3-diones **2d**, both bearing a phenyl group at the C-5 position of the pyrrolidine-




Fig. 1. Important ^1H and ^{13}C chemical shifts (left) and HMBC study of **3aa** (right).





Fig. 2. Important ^1H chemical shifts, NOE (left) and HMBC study (right) of **3ad** (upper) and **3ae** (lower).

Table 1
Mn(III)-based reaction of 1,1-diarylethenes **1a–e** with pyrrolidine-2,3-diones **2a–e**.^a

Entry	Ethene/ 1	Pyrrolidinedione/ 2	1:2 :Mn(OAc) ₃ ^b	Temp/°C	Time/min	3 /Yield/% ^c
1	1a : Ar = 4-Me-C ₆ H ₄	2a : R ¹ = R ² = Me, R ³ = H, R ⁴ = Bn	1:1.5:3	70	16	3aa (38)
2	1a	2a	1:1.5:3	reflux	2	3aa (67)
3	1a	2a	1:2:3	reflux	3	3aa (87)
4	1a	2a	1:3:5	reflux	3	3aa (90)
5	1b : Ar = Ph	2a	1:3:5	reflux	3	3ba (87)
6	1 : Ar = 4-F-C ₆ H ₄	2a	1:3:5	reflux	3	3ca (74)
7	1 : Ar = 4-Cl-C ₆ H ₄	2a	1:2:3	reflux	3	3da (68)
8	1e : Ar = 4-MeO-C ₆ H ₄	2a	1:2:3	reflux	3	3ea (60)
9	1a : Ar = 4-Me-C ₆ H ₄	2b : R ¹ = Me, R ² = R ³ = H, R ⁴ = Bn	1:2:3	reflux	3	3ab (81)
10	1a	2c : R ¹ = R ⁴ = Me, R ² = H, R ³ = Ph	1:3:5	reflux	3	3ac (83)
11	1a	2d : R ¹ = Et, R ² = H, R ³ = Ph, R ⁴ = Me	1:3:5	reflux	3	3ad (80)
12	1a	2e : R ¹ = R ² = R ⁴ = Me, R ³ = Ph	1:3:5	reflux	3	3ae (60) ^d

^a The reaction of ethene **1** (1 mmol) was carried out in acetic acid (15 mL).

^b Molar ratio.

^c Isolated yield based on the ethene **1**.

^d 1-Hydroxy-1-isopropyl-3,3-bis(4-methylphenyl)-7-methyl-6-phenyl-2-oxa-7-azaspiro[4.4]nonane-8,9-dione (**4**) was also isolated in 22% yield.

dione, led to a similar result (Entries 10 and 11), the reaction using 4-isobutyryl-1-methyl-5-phenylpyrrolidine-2,3-dione (**2e**) resulted in the decreased yield of the desired product (Entry 12). However, after a thorough chromatographic separation, 1-hydroxy-1-isopropyl-3,3-bis(4-methylphenyl)-7-methyl-6-phenyl-2-oxa-7-azaspiro[4.4]nonane-8,9-dione (**4**) was also isolated in 22% yield probably due to the addition of water during the reaction (vide infra).¹⁵ The ¹H NMR spectrum of the 6-phenyl-2-oxa-7-azaspiro[4.4]nonane-8,9-diones **3ac**, **3ad**, and **3ae** deserves comments.^{16–18} The *ortho*-protons of the phenyl group appeared around δ 6.3 as a broad singlet because of the rotational barrier of the phenyl group by the C-1 exomethylene group (Fig. 2). Simultaneously, the H-4 proton (ca. δ 3.4) was deshielded by the ring current effect of one of the C-3 aryl groups. When the NMR spectrum of **3ae** was taken in CDCl₃ at 50 °C, the broad peak of the phenyl protons became sharp (See Supplementary data). In addition, the H-10 sp² proton (δ 3.37) of **3ac** and **3ad**, and the C-11 methyl group (δ 0.67) of **3ae** were shielded by the anisotropic effect of the C-9 carbonyl group. In the case of the by-product **4**, the anisotropic effect of the C-9 carbonyl group was extremely strong toward one of the methyl groups of the isopropyl group, showing δ 0.00 (3H, d, J = 6.7 Hz) (Fig. 3).¹⁵ Fortunately, we got a single crystal of **3ae** from chloroform and finally established the structure by an X-ray crystallographic measurement (Fig. 4 and supplementary data).¹⁹

It was reported that the Mn(III)-enolate complex formation is the rate-determining step in the Mn(III)-based reaction of the α -alkyl-substituted 1,3-dicarbonyl compounds with alkenes.^{1c,20} In this case, a similar enolization of **2** with Mn(OAc)₃ would occur during the first stage, producing complex **A** (Scheme 3). Complex **A** is electron deficient, thus an electron-rich alkene **2** should be easily oxidized to give radical **B**, which would be further oxidized to produce the carbocation **C**. The cation **C** would spontaneously cyclize with the carbonyl oxygen and undergo subsequent β -proton elimination that produces the 2-oxa-7-azaspiro compounds **3**. The formation of the stable tertiary carbocation **C** is crucial for the next *O*-cyclization to produce **3**. In fact, the reaction using styrene and terminal alkenes such as 1-hexene was complicated. When 4-isobutyryl-1-methyl-5-phenylpyrrolidine-2,3-dione (**2e**) was subjected to the oxidation, the desired 1-exomethylene-2-oxa-7-azaspiro compound **3ae** was mainly produced along with the hydroxy-2-oxa-7-azaspiro compound **4** formed by nucleophilic addition of water to the intermediate cation **D** due to relief of the steric hindrance.

In conclusion, our initial forecast was proved to be correct by the fact that the 2-oxa-7-azaspiro[4.4]nonane-8,9-dione derivatives **3** containing tetrahydrofuran and 2,3-pyrrolidinedione rings could be successfully synthesized in good yields by the Mn(III) oxidation of a mixture of 1,1-diarylethenes **1** and 4-acylpyrrolidine-2,3-diones **2**. The reaction was straightforward, the reaction time was significantly short, and the procedure was simple to





Fig. 4. Crystal structure of **3ae**.



Scheme 3. Plausible mechanism for the formation of **3** and **4**.


obtain the desired product **3**. In addition, the structures of the products **3** and **4** were well characterized by spectroscopic methods including the X-ray single crystal analysis of **3ae**, and the mechanism for the formation of the products **3** and **4** was logically interpreted by the Mn(III)-based oxidation chemistry.^{1,2,20,21} Further optimization of the reactions listed in Table 1, application of the reaction using the pyrrolidinediones **2** bearing other substituents, and bioassay of the pyrrolidinediones **2** and the products **3** for antibacterial, antiviral, bactericidal, insecticidal, herbicidal activities are currently underway.

Acknowledgments

This research was supported by a Grant-in-Aid for Scientific Research (C), No. 25410049, from the Japan Society for the Promotion of Science – Japan. We also acknowledge Nissan Chemical Industries, Ltd., and the Astellas Foundation for Research on Metabolic Disorders – Japan for their financial support.

A. Supplementary data

Supplementary data (experimental detail, spectroscopic data of the products **3ba**, **3ca**, **3da**, **3ea**, **3ab**, and the copies of ¹H NMR, ¹³C NMR, DEPT, COSY, NOESY, HMQC, and HMBC spectra for all



the compounds **3** and **4**, and X-ray brief report of **3ae**.) associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.tetlet.2017.08.002>.

References

- (a) de Klein WJ. In: Mijis WJ, de Jonge CRH, eds. *Organic Syntheses by Oxidation with Metal Compounds*. New York: Plenum Press; 1986:261–314;
(b) . Reviews for the Mn(III)-based oxidation: Iqbal J, Bhatia B, Nayyar NK. *Chem Rev*. 1994;94:519–564;
(c) Snider BB. *Chem Rev*. 1996;96:339–363;
(d) Melikyan G. *Org React*. 1997;49:627–675.
- (a) Recent reviews for the Mn(III)-based oxidation: Demir AS, Emrullahoglu M. *Current. Org Synth*. 2007;4:321–351;
(b) Pan X-Q, Zou J-P, Zhang W. *Mol. Divers.*. 2009;13:421–438;
(c) Burton JW. Manganese(III) Acetate, CAN, and Fe(III) Salts in Oxidative Radical Chemistry. In: Chatgililoglu C, Studer A, eds. *Encyclopedia of Radicals in Chemistry, Biology and Materials*. New York: Wiley; 2012:901–941;
(d) Jahn U. *Top Curr Chem*. 2012;320:121–190;
(e) Mondal M, Bora U. *RSC Adv*. 2013;3:18716–18754;
(f) Wang G-W, Li F-BJ. *Nanosci Nanotech.*. 2007;7:1162–1175;
(g) Nishino H. Manganese(III)-Based Peroxidation of Alkenes to Heterocycles. In: Eguchi S, ed. *Topics in Heterocyclic Chemistry, Bioactive Heterocycles I*. Berlin: Springer; 2006:39–76;
(h) Terent'ev AO, Borisov DA, Vil' VA, Dembitsky VM. *Beilstein J Org Chem*. 2014;10:34–114.
- Eicher T, Hauptmann S. *The Chemistry of Heterocycles: Structure, Reactions, Syntheses, and Applications*. 2nd ed. Wiley; 2004.
- (a) Balendiran GK, Martin H-J, El-Hawari Y, Maser E. *Chem Biol Int*. 2009;178:134–137;
(b) Srivastava SK, Ramana KV, US Pat Appl Publ, 2009, US 20090270490 A1 20091029.;
(c) Verma M, Martin H-J, Haq W, O'Connor TR, Maser E, Balendiran GK. *Eur. J. Pharm.*. 2008;584:213–221(d) Goering BK, Ph.D. Dissertation, Cornell University, 1995.
- Haslam E. *Shikimic Acid Metabolism and Metabolites*. New York: John Wiley & Sons; 1993.
- (a) Vaughan WR, Peters LR. *J Org Chem*. 1953;18:382–392;
(b) Wasserman HH, Ivest JL. *J Org Chem*. 1985;50:3573–3580.
- (a) Buchanan JG, Sable HZ. In: Thyagarajan BS, ed. *Selective Organic Transformations*, Vol. 2. New York: Wiley-Interscience; 1972:1–95;
(b) Nishikawa T, Kajii S, Isobe M. *Chem Lett*. 2004;33:440–441;
(c) Nubbemeyer U. *Top Curr Chem*. 2001;216:125–196;
(d) Ley SV, Cox LR, Meek G. *Chem Rev*. 1996;96:423–442.
- (a) Ito N, Nishino H, Kurosawa K. *Bull Chem Soc Jpn*. 1983;56:3527–3528;
(b) Qian C-Y, Yamada T, Nishino H, Kurosawa K. *Bull Chem Soc Jpn*. 1992;65:1371–1378;
(c) Yamada T, Iwahara Y, Nishino H, Kurosawa K. *J Chem Soc, Perkin Trans 1*. 1993;609–616;
(d) Nishino H, Hashimoto H, Korp JD, Kurosawa K. *Bull Chem Soc Jpn*. 1995;68:1999–2009;
(e) Nishino H, Ishida K, Hashimoto H, Kurosawa K. *Synthesis*. 1996;888–896;
(f) Fujino R, Nishino H. *Synthesis*. 2005;731–740;
(g) Tsubusaki T, Nishino H. *Tetrahedron*. 2009;65:9448–9459.
- (a) Nguyen V-H, Nishino H, Kurosawa K. *Tetrahedron Lett*. 1997;38:1773–1776;
(b) Nguyen V-H, Nishino H, Kurosawa K. *Heterocycles*. 1998;48:465–480.
- Chowdhury FA, Nishino H, Kurosawa K. *Heterocycles*. 1999;51:575–591.
- Emerson DW, Titus RL, Jones MD. *J Heterocycl Chem*. 1998;35:611–617.
- (a) DeNinno MP, Andrews M, Bell AS, et al. *Bioorg Med Chem Lett*. 2009;19:2537–2541;
(b) Galenko AV, Khlebnikov AF, Novikov MS, Avdontceva MS. *Tetrahedron*. 2015;71:1940–1951.
- (a) Sugimoto H, Nakamura S, Ohwada T. *Adv Synth Catal*. 2007;349:669–679;
(b) Campbell KN, Sommers AH, Campbell BK. *J Am Chem Soc*. 1944;66:82.
- The structure of **3aa** was determined by spectroscopic methods, a 2D NMR study and elemental analysis. 7-Benzyl-3,3-bis(4-methylphenyl)-1-(propan-2-ylidene)-2-oxa-7-azaspiro[4.4]nonane-8,9-dione (**3aa**): yellow needles (from chloroform/hexane); mp 182.0–182.5 °C; IR (CHCl₃) 1762.8 (—CO—), 1714.6 (—CON—); ¹H NMR (500 MHz, CDCl₃) δ 7.34–7.30 (3H, m, arom H), 7.22–7.20 (2H, m, arom H), 7.13–7.11 (4H, m, arom H), 7.07–7.03 (4H, m, arom H), 4.56 (1H, d, J = 14.4 Hz, H_a-CH), 4.52 (1H, d, J = 14.4 Hz, HC-H_b), 3.04 (1H, d, J = 11.8 Hz, H_a-6), 2.84 (1H, d, J = 11.8 Hz, H_b-6), 2.83 (1H, d, J = 12.4 Hz, H_a-4), 2.80 (1H, d, J = 12.4 Hz, H_b-4), 2.31 (3H, s, Me), 2.26 (3H, s, Me), 1.80 (3H, s, Me-12), 1.15 (3H, s, Me-11); ¹³C NMR (125 MHz, CDCl₃) δ 199.5 (C-9), 158.4 (C-8), 149.0 (C-1), 140.6, 140.0, 137.3, 137.0, 134.1 (arom C), 129.1 (2C), 128.84 (2C), 128.82 (2C), 128.5 (2C), 128.3, 125.6 (2C), 125.4 (2C) (arom CH), 102.6 (C-10), 88.1 (C-3), 53.6 (C-6), 53.2 (C-4), 51.7 (C-5), 48.3 (CH₂), 20.93 (Me), 20.90 (Me), 18.9 (Me-12), 17.5 (Me-11); FAB HRMS (acetone/NBA): calcd for C₃₁H₃₂NO₃ 466.2382 (M+H); found 466.2365. Anal Calcd for C₃₁H₃₁NO₃·1/2H₂O: C, 78.45; H, 6.58; N, 2.95. Found: C, 78.57; H, 6.73; N, 2.92.
- The structure of **4** was determined by spectroscopic methods, an HMQC study and elemental analysis (Fig. 3). 1-Hydroxy-1-isopropyl-3,3-bis(4-methylphenyl)-7-methyl-6-phenyl-2-oxa-7-azaspiro[4.4]nonane-8,9-dione (**4**): colorless cubes (from chloroform); mp 218.5–219.5 °C; IR (KBr) 3196 (OH), 1689.5 (C=O, N=C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.44 (2H, d, J = 8.1 Hz, arom H), 7.40 (2H, d, J = 8.2 Hz, arom H), 7.31–7.26 (3H, m, arom H), 7.14 (2H, J = 8.2 Hz, arom H), 7.12 (2H, m, arom H), 7.07 (2H, d, J = 8.1 Hz, arom H), 5.45 (1H, s, OH; exchanged by D₂O), 4.26 (1H, s, H-6), 3.55 (1H, d, J = 13.4 Hz, HC-H_a-4), 3.32 (1H, d, J = 13.4 Hz, HC-H_b-4), 2.53 (3H, s, Me-7), 2.40 (1H, sep, J = 6.7 Hz, H-10), 2.29 (3H, s, Me), 2.26 (3H, s, Me), 0.71 (3H, d, J = 6.7 Hz, Me-12), 0.00 (3H, d, J = 6.7 Hz, Me-11); ¹³C NMR (125 MHz, CDCl₃) δ 212.9 (C-9), 170.4 (C-8), 143.2, 143.0, 137.0, 136.7, 134.7 (arom C), 129.2 (2C), 128.9 (3C), 128.8 (2C), 128.7 (2C), 125.6 (2C), 125.0 (2C) (arom CH), 106.4 (C-1), 88.9 (C-3), 67.8 (C-6), 66.1 (C-5), 47.5 (C-4), 39.7 (C-10), 28.7 (Me-7), 21.0 (Me), 20.9 (Me), 20.1 (Me-12), 17.2 (Me-11); FAB HRMS (acetone/NBA): calcd for C₃₁H₃₄NO₄ 484.2488 (M+H); found 484.2501. Anal Calcd for C₃₁H₃₃NO₄·1/4H₂O: C, 76.28; H, 6.92; N, 2.87. Found: C, 76.55; H, 7.03; N, 2.84.
- The structure of **3ac** was determined by spectroscopic methods, a 2D NMR study and elemental analysis. 1-Ethylidene-3,3-bis(4-methylphenyl)-7-methyl-6-phenyl-2-oxa-7-azaspiro[4.4]nonane-8,9-dione (**3ac**): colorless needles (from chloroform/hexane); mp 182–183 °C; IR (CHCl₃) ν 1767 (C=O), 1715 (N=C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.41 (2H, d, J = 8.2 Hz, ArH), 7.27 (2H, d, J = 8.2 Hz, ArH), 7.21 (2H, d, J = 8.0 Hz, ArH), 7.18 (3H, t, J = 7.4 Hz, ArH), 7.06 (2H, d, J = 8.0 Hz, ArH), 6.32 (2H, br, s, ArH), 4.40 (1H, s, H-6), 3.37 (1H, q, J = 6.9 Hz, H-10), 3.32 (1H, d, J = 12.7 Hz, H_a-4), 2.92 (1H, d, J = 12.7 Hz, H_b-4), 2.90 (3H, s, N-Me), 2.42 (3H, s, Me), 2.27 (3H, s, Me), 1.44 (3H, t, J = 6.9 Hz, Me); ¹³C NMR (125 MHz, CDCl₃) δ 198.5 (C-9), 160.0 (C-8), 150.0 (C-1), 142.2, 140.8, 137.6, 137.1, 135.0 (arom C), 129.3 (2C), 128.9 (2C), 128.6, 128.3 (2C), 126.0 (2C), 125.2 (4C) (arom CH), 100.1 (C-10), 87.9 (C-3), 70.1 (C-6), 60.9 (C-5), 53.0 (C-4), 30.9 (N-Me), 21.1, 20.9 (Me), 10.7 (Me); FAB HRMS (acetone/NBA): calcd for C₃₀H₃₀NO₃ 452.2226 (M+H); found 452.2225. Anal Calcd for C₃₀H₂₉NO₃·4/5H₂O: C, 77.33; H, 6.62; N, 3.01. Found: C, 77.08; H, 6.37; N, 2.98.
- The structure of **3ad** was determined by spectroscopic methods, a 2D NMR study and elemental analysis. 3,3-Bis(4-methylphenyl)-7-methyl-6-phenyl-1-propylidene-2-oxa-7-azaspiro[4.4]nonane-8,9-dione (**3ad**): colorless needles (from chloroform/hexane); mp 188–189 °C; IR (CHCl₃) ν 1767 (C=O), 1717 (N=C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.42 (2H, d, J = 8.2 Hz, ArH), 7.27 (2H, d, J = 8.2 Hz, ArH), 7.21 (2H, d, J = 8.3 Hz, ArH), 7.17 (3H, t, J = 7.5 Hz, ArH), 7.05 (2H, d, J = 8.3 Hz, ArH), 6.31 (2H, br, s, ArH), 4.42 (1H, s, H-6), 3.37 (1H, dd, J = 6.7, 6.3 Hz, H-10), 3.35 (1H, d, J = 12.8 Hz, H_a-4), 2.90 (1H, d, J = 12.8 Hz, H_b-4), 2.90 (3H, s, N-Me), 2.43 (3H, s, Me), 2.26 (3H, s, Me), 2.07 (1H, m, H-CH), 1.86 (1H, m, HC-H), 0.55 (3H, t, J = 7.5 Hz, Me); ¹³C NMR (125 MHz, CDCl₃) δ 198.5 (C-9), 160.0 (C-8), 149.1 (C-1), 142.2, 140.8, 137.5, 137.0, 134.9 (arom C), 129.3 (2C), 128.9 (2C), 128.5, 128.3 (2C), 126.0 (2C), 125.1 (4C) (arom CH), 107.5 (C-10), 87.9 (C-3), 70.1 (C-6), 60.1 (C-5), 52.8 (C-4), 30.8 (N-Me), 21.1, 20.8 (Me), 18.7 (CH₂), 13.7 (Me); FAB HRMS (acetone/NBA): calcd for C₃₁H₃₂NO₃ 466.2382 (M+H); found 466.2377.
- The structure of **3ae** was determined by spectroscopic methods, a 2D NMR study and elemental analysis. 3,3-Bis(4-methylphenyl)-7-methyl-6-phenyl-1-(propan-2-ylidene)-2-oxa-7-azaspiro[4.4]nonane-8,9-dione (**3ae**): colorless microcrystals (from chloroform/hexane); mp 174–175 °C; IR (CHCl₃) ν 1765 (C=O), 1717 (N=C=O); ¹H NMR (500 MHz, CDCl₃) δ 7.53 (2H, d, J = 8.3 Hz, ArH), 7.27 (2H, d, J = 8.5 Hz, ArH), 7.21 (2H, m, ArH), 7.19 (2H, d, J = 8.3 Hz, ArH), 7.15 (1H, t, J = 8.03 Hz, ArH), 7.05 (2H, d, J = 8.3 Hz, ArH), 6.50 (2H, br, s, ArH), 4.55 (1H, s, H-6), 3.42 (1H, d, J = 12.8 Hz, H_a-4), 3.00 (3H, s, N-Me), 2.89 (1H, d, J = 12.8 Hz, H_b-4), 2.39 (3H, s, Me), 2.26 (3H, s, Me), 1.58 (3H, s, Me-12), 0.67 (3H, s, Me-11); ¹³C NMR (125 MHz, CDCl₃) δ 197.4 (C-9), 160.0 (C-8), 143.9 (C-1), 143.7, 141.2, 137.6, 136.8, 135.0 (arom C), 129.3 (2C), 128.9 (4C), 128.3, 127.8 (2C), 126.1 (2C), 125.0 (2C) (arom CH), 107.0 (C-10), 86.8 (C-3), 69.5 (C-6), 60.2 (C-5), 54.7 (C-4), 31.2 (N-Me), 21.0, 20.9 (Me), 20.5 (Me-11), 17.5 (Me-12); Anal Calcd for C₃₁H₃₁NO₃: C, 79.97; H, 6.71; N, 3.01. Found: C, 79.73; H, 6.78; N, 3.07.
- X-ray crystal data of **3ae** (Fig. 4): Empirical Formula C₃₁H₃₁NO₃; Formula Weight 465.59; Crystal Color, Habit colorless, block; Crystal Dimensions 0.370 × 0.327 × 0.187 mm; Crystal System triclinic; Lattice Type Primitive; Lattice Parameters a = 9.6559(4) Å, b = 9.9138(5) Å, c = 14.9614(8) Å, α = 75.345(2)°, β = 69.807(2)°, γ = 78.722(1)°, V = 1291.2(1) Å³; Space Group P-1 (#2); Z value 2; D_{calc} 1.197 g/cm³; F₀₀₀ 496.00; μ(MoKα) 0.763 cm⁻¹; R₁ (I > 2.0σ(I)) 0.0688; R (All reflections) 0.1444; wR₂ (All reflections) 0.2614; Goodness of Fit Indicator 1.131.
- Snider BB. *Tetrahedron*. 2009;65:10738–10744.
- Cossy J, Bouzide A, Leblanc C. *J Org Chem*. 2000;65:7258–7265.