

Harnack inequalities for functional SDEs with multiplicative noise and applications[☆]

Feng-Yu Wang^{a,b,*}, Chenggui Yuan^b

^a School of Mathematical Sciences, Beijing Normal University, Beijing 100875, China

^b Department of Mathematics, Swansea University, Singleton Park, SA2 8PP, UK

Received 28 December 2010; received in revised form 15 June 2011; accepted 8 July 2011

Available online 23 July 2011

Abstract

By constructing a new coupling, the log-Harnack inequality is established for the functional solution of a delay stochastic differential equation with multiplicative noise. As applications, the strong Feller property and heat kernel estimates w.r.t. quasi-invariant probability measures are derived for the associated transition semigroup of the solution. The dimension-free Harnack inequality in the sense of Wang (1997) [14] is also investigated.

© 2011 Elsevier B.V. All rights reserved.

MSC: 60H10; 47G20

Keywords: Harnack inequality; Functional solution; Delay SDE; Strong Feller property; Heat kernel

1. Introduction

The dimension-free Harnack inequality introduced in [14] has become a useful tool in the study of diffusion semigroups, in particular, for the uniform integrability, contractivity properties, and estimates on heat kernels, see e.g. [2,1,4,9–11,15,16,19,20] and the references within. Recently, by using coupling arguments, the dimension-free Harnack inequality has been established in [18] for stochastic differential equations (SDEs) with multiplicative noise, and

[☆] Supported in part by WIMCS, SRFDP and the Fundamental Research Funds for the Central Universities.

* Corresponding author at: School of Mathematical Sciences, Beijing Normal University, Beijing 100875, China.

E-mail addresses: wangfy@bnu.edu.cn, F.-Y.Wang@swansea.ac.uk (F.-Y. Wang), C.Yuan@swansea.ac.uk (C. Yuan).

in [7] for stochastic differential delay equations (SDDEs) with additive noise. The aim of this paper is to extend these existed results to the functional solution of SDDEs with multiplicative noise. Due to the double difficulty caused by delay and non-constant diffusion coefficient, both couplings constructed in [18,7] are no longer valid. Under a reasonable assumption (see (A) below), we will construct a successful coupling which leads to an explicit log-Harnack inequality of the functional solution (see Theorem 1.1 below). This weaker version of Harnack inequality was introduced in [12,17] for elliptic diffusion processes, and it is powerful enough to imply some regularity properties of the semigroup such as the strong Feller property and heat kernel estimates w.r.t. quasi-invariant probability measures (see Corollary 1.2 below). The dimension-free Harnack inequality in the sense of [14] is also derived (see Theorem 4.1 below).

Let $r_0 > 0$ be fixed, and let $\mathcal{C} = C([-r_0, 0]; \mathbb{R}^d)$ be equipped with the uniform norm $\|\cdot\|_\infty$. Let $\mathcal{B}_b(\mathcal{C})$ be the set of all bounded measurable functions on \mathcal{C} . Let $B(t)$ be a d -dimensional Brownian motion on a complete filtered probability space $(\Omega, \{\mathcal{F}_t\}_{t \geq 0}, \mathbb{P})$, and let

$$\begin{aligned}\sigma &: [0, \infty) \times \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}^d \otimes \mathbb{R}^d, \\ Z &: [0, \infty) \times \mathbb{R}^d \times \Omega \rightarrow \mathbb{R}^d, \\ b &: [0, \infty) \times \mathcal{C} \times \Omega \rightarrow \mathbb{R}^d\end{aligned}$$

be progressively measurable, locally bounded in the first variable and continuous w.r.t. the second variable. Consider the following delay SDE on \mathbb{R}^d :

$$dX(t) = \{Z(t, X(t)) + b(t, X_t)\} dt + \sigma(t, X(t)) dB(t), \quad X_0 \in \mathcal{C}, \quad (1.1)$$

where for each $t \geq 0$, $X_t \in \mathcal{C}$ is fixed as $X_t(u) = X(t+u)$, $u \in [-r_0, 0]$. Let $\|\cdot\|$ and $\|\cdot\|_{HS}$ be the operator norm and the Hilbert–Schmidt norm for $d \times d$ -matrices respectively.

To ensure the existence, uniqueness, non-explosion, and further regular properties of the solution, we make use of the following assumption:

(A) σ is invertible, $\sup_{\Omega} \{|\sigma(t, 0)| + |Z(t, 0)| + |b(t, 0)|\}$ is locally bounded in t , and for any $T > 0$ there exist constants $K_1, K_2 \geq 0, K_3 > 0$ and $K_4 \in \mathbb{R}$ such that

$$\begin{aligned}(A1) \quad & |\sigma(t, \eta(0))^{-1} \{b(t, \xi) - b(t, \eta)\}| \leq K_1 \|\xi - \eta\|_\infty, \quad t \in [0, T], \xi, \eta \in \mathcal{C}; \\ (A2) \quad & |(\sigma(t, x) - \sigma(t, y))| \leq K_2 (1 \wedge |x - y|), \quad t \in [0, T], x, y \in \mathbb{R}^d; \\ (A3) \quad & |\sigma(t, x)^{-1}| \leq K_3, \quad t \geq 0, x \in \mathbb{R}^d; \\ (A4) \quad & \|\sigma(t, x) - \sigma(t, y)\|_{HS}^2 + 2\langle x - y, Z(t, x) - Z(t, y) \rangle \leq K_4 |x - y|^2, \quad t \in [0, T], x, y \in \mathbb{R}^d\end{aligned}$$

hold almost surely.

Obviously, (A) is satisfied if uniformly in $t \geq 0$ and $\omega \in \Omega$: σ is bounded and Lipschitz continuous on \mathbb{R}^d with bounded σ^{-1} , b is Lipschitz continuous on \mathcal{C} and ∇Z is bounded above on \mathbb{R}^d .

We remark that in [7] σ is assumed to be the unit matrix and (A4) holds for non-positive K_4 , so that (A) holds for $K_2 = 0, K_3 = 1$ and K_1 being the Lipschitz constant of b . Moreover, it is easy to see that (A) is satisfied provided σ is uniformly invertible, and σ, b, Z are Lipschitz continuous w.r.t. the second variable uniformly in the first and third variables. Therefore, our framework is much more general.

Moreover, if (A) holds then for any \mathcal{F}_0 -measurable X_0 , the Eq. (1.1) has a unique strong solution and the solution is non-explosive, see [13, Theorem 2.3]. We note that although this result was proved for the time-homogenous case, its proof works also for the non-homogenous case if the conditions therein hold uniformly w.r.t. $t \in [0, T]$ for any $T > 0$.

To indicate the dependence of the solution on the initial data, for any $\xi \in \mathcal{C}$ we shall use $X^\xi(t)$ and X_t^ξ respectively to denote the solution and the functional solution to the equation with $X_0 = \xi$. We shall investigate the Harnack inequality and applications for the family of Markov operators $(P_t)_{t \geq 0}$ on $\mathcal{B}_b(\mathcal{C})$ given by

$$P_t f(\xi) = \mathbb{E} f(X_t^\xi), \quad t \geq 0, f \in \mathcal{B}_b(\mathcal{C}).$$

We note that due to the delay, the solution $X(t)$ is not Markovian. But when Z, b and σ are deterministic, the functional solution X_t is a strong Markov process.

Theorem 1.1. Assume (A) for $T > r_0$. Then the log-Harnack inequality

$$P_T \log f(\eta) \leq \log P_T f(\xi) + H_T(\xi, \eta), \quad \xi, \eta \in \mathcal{C}$$

holds for $f \in \mathcal{B}_b(\mathcal{C})$ with $f \geq 1$ and

$$H_T(\xi, \eta) := \inf_{s \in (0, T-r_0]} \left\{ \frac{2K_3^2 K_4 |\xi(0) - \eta(0)|^2}{1 - e^{-K_4 s}} + 3K_1^2 \left\{ \frac{r_0}{2} + 4s \left(1 + K_2^2 K_3^2 \right) \right\} \right. \\ \left. \times e^{3K_2^2 (K_1^2 s + 4)s} \|\xi - \eta\|_\infty^2 \right\}.$$

Consequently, for any $T > r_0$, P_T is strong Feller, i.e. $P_T \mathcal{B}_b(\mathcal{C}) \subset C_b(\mathcal{C})$, the set of bounded continuous functions on \mathcal{C} .

It is easy to see that the log-Harnack inequality only holds for $T > r_0$. Indeed, if the inequality holds for some $T \in (0, r_0]$ then by taking $f(\xi) = (1 + |\xi(-r_0)| \wedge n)^n$ and letting $n \rightarrow \infty$, the inequality implies that

$$\log(1 + |\eta(T - r_0)|) \leq \log(1 + |\xi(T - r_0)|)$$

holds for all $\xi, \eta \in \mathcal{C}$, which is however impossible.

Next, we present some consequences of the above log-Harnack inequality for heat kernels of P_T w.r.t. a quasi-invariant probability measure μ .

Definition 1.1. Let (E, \mathcal{F}) be a measurable space with $\mathcal{B}_b(E)$ the set of all bounded measurable functions, let μ be a probability measure on E , and let P be a bounded linear operator on $\mathcal{B}_b(E)$.

- (i) μ is called quasi-invariant of P , if μP is absolutely continuous w.r.t. μ , where $(\mu P)(A) := \mu(P1_A)$, $A \in \mathcal{F}$. If $\mu P = \mu$ then μ is called an invariant probability measure of P .
- (ii) A measurable function p on E^2 is called the kernel of P w.r.t. μ , if

$$Pf = \int_E p(\cdot, y) f(y) \mu(dy), \quad f \in \mathcal{B}_b(E).$$

Corollary 1.2. Assume (A). Let $t > r_0$ and μ be a quasi-invariant probability measure of P_t . Then:

- (1) P_t has a kernel p_t w.r.t. μ .
- (2) The kernel p_t satisfies the entropy inequality

$$\int_{\mathcal{C}} p_t(\xi, \cdot) \log \frac{p_t(\xi, \cdot)}{p_t(\eta, \cdot)} d\mu \leq H_t(\xi, \eta), \quad \xi, \eta \in \mathcal{C},$$

where we set $r \log \frac{r}{s} = 0$ if $r = 0$ and $r \log \frac{r}{s} = \infty$ if $r > 0$ and $s = 0$.

(3) The kernel p_t satisfies

$$\int_{\mathcal{C}} p_t(\xi, \cdot) p_t(\eta, \cdot) d\mu \geq \exp[-H_t(\xi, \eta)], \quad \xi, \eta \in \mathcal{C}.$$

(4) P_t has at most one invariant probability measure, and if it has, the kernel of P_t w.r.t. the invariant probability measure is strictly positive.

Note that if P_t is symmetric w.r.t. μ , then $\int_{\mathcal{C}} p_t(\xi, \cdot) p_t(\eta, \cdot) d\mu = p_{2t}(\xi, \eta)$ so that (3) provides a Gaussian type lower bound for the heat kernel. Moreover, if μ is an invariant probability measure of P_t , then (2) gives an entropy-cost inequality as in [12, Corollary 1.2(3)]. More precisely, letting P_t^* be the adjoint operator of P_t in $L^2(\mu)$, for any $f \geq 0$ with $\mu(f) = 1$, one has

$$\mu((P_t^* f) \log P_t^* f) \leq \inf_{\pi \in \mathcal{C}(\mu, f\mu)} \int_{\mathcal{C} \times \mathcal{C}} H_t(\xi, \eta) \pi(d\xi, d\eta), \quad t > 0,$$

where $\mathcal{C}(\mu, f\mu)$ is the set of all couplings for μ and $f\mu$. The right hand side of the above inequality is called the transportation-cost between μ and $f\mu$ with cost function H_t . Finally, we note that the uniqueness of the invariant probability measure has been investigated in [8] for SDDEs in terms of the asymptotic coupling property.

To conclude this section, let us present an existence result of the quasi-invariant measure (see [6] for existence of the invariant probability measure).

Proposition 1.3. Assume (A) and let σ and Z be deterministic and time-independent. If $x \mapsto -\|\sigma(x)\|_{HS}^2 - \langle x, Z(x) \rangle$ is a compact function, i.e. $\{x \in \mathbb{R}^d : -\|\sigma(x)\|_{HS}^2 - \langle x, Z(x) \rangle \leq r\}$ is a compact set for any constant $r > 0$, then $\{P_t\}_{t \geq 0}$ has a quasi-invariant probability measure, i.e. the measure is quasi-invariant for all $P_t, t \geq 0$.

The Proof of Theorem 1.1 is presented in Section 2 while those of Corollary 1.2 and Proposition 1.3 are addressed in Section 3. Finally, in Section 4 we investigate the dimension-free Harnack inequality in the sense of [14].

2. Proof of Theorem 1.1

According to [18, Proposition 2.3], the claimed log-Harnack inequality implies the strong Feller property of P_T , see also Proposition 3.1(1) below. So, we only have to prove the desired log-Harnack inequality. To make the proof easy to follow, let us first explain the main idea of the argument.

Let $T > r_0$ and $t_0 \in (0, T - r_0]$ be fixed. Let $X(s)$ solve (1.1) with $X_0 = \xi$. For $\gamma \in C^1([0, t_0])$ such that $\gamma(r) > 0$ for $r \in [0, t_0]$ and $\gamma(t_0) = 0$, let $Y(t)$ solve the equation

$$\begin{aligned} dY(t) = & \left\{ Z(t, Y(t)) + b(t, X_t) + \frac{1_{\{t < t_0\}}}{\gamma(t)} \sigma(t, Y(t)) \sigma(t, X(t))^{-1} (X(t) - Y(t)) \right\} dt \\ & + \sigma(t, Y(t)) dB(t), \quad Y_0 = \eta. \end{aligned} \quad (2.1)$$

The key point of our coupling is that $X(t)$ and $Y(t)$ will move together from time t_0 on, so that $X_T = Y_T$. To this end, we add the drift term

$$\frac{1_{\{t < t_0\}}}{\gamma(t)} \sigma(t, Y(t)) \sigma(t, X(t))^{-1} (X(t) - Y(t)) dt$$

to force $Y(t)$ to meet $X(t)$ at time t_0 . In order to dominate the non-trivial martingale part of $X(t) - Y(t)$, the force has to be infinitely strong near by t_0 , for this we need $\gamma(t_0) = 0$. More precisely, as in [18] we shall take

$$\gamma(t) = \frac{2-\theta}{K_4} \left(1 - e^{(t-t_0)K_4}\right), \quad t \in [0, t_0]$$

for a parameter $\theta \in (0, 2)$. In this case, we have

$$2 + \gamma'(t) - K_4\gamma(t) = \theta, \quad t \in [0, t_0]. \quad (2.2)$$

Moreover, to ensure these two process moving together after the coupling time (i.e. the first meeting time), they should solve the same equation from that time on. This is the reason why we have to follow the line of [7] to take the delay term in (2.1) by using X_t rather than Y_t . In this case both $X(t)$ and $Y(t)$ solve the equation

$$\begin{aligned} d\Xi(t) = & \left\{ Z(t, \Xi(t)) + b(t, X_t) + \frac{1_{\{t < t_0\}}}{\gamma(t)} \sigma(t, \Xi(t)) \sigma(t, X(t))^{-1} (X(t) - \Xi(t)) \right\} dt \\ & + \sigma(t, \Xi(t)) dB(t), \end{aligned}$$

so that by the uniqueness of the solution which is easy to verify from (A), they move together from the coupling time on. Since the additional drift is singular at time t_0 , it is only clear that $Y(t)$ is well solved before time t_0 . To solve $Y(t)$ for all $t \in [0, T]$, we need to reformulate the equation by using a new Brownian motion determined by the Girsanov transform induced by the coupling.

Let

$$\phi_t = \sigma(t, Y(t))^{-1} \{b(t, Y_t) - b(t, X_t)\} - \frac{1_{\{t < t_0\}}}{\gamma(t)} \sigma(t, X(t))^{-1} (X(t) - Y(t)), \quad t \geq 0.$$

From (A) it is easy to see that

$$R_t := \exp \left[\int_0^t \langle \phi_s, dB(s) \rangle - \frac{1}{2} \int_0^t |\phi_s|^2 ds \right]$$

is a martingale for $t \in [0, t_0]$. We shall further prove that

(i) $\{R_t\}_{t \geq 0}$ is a well-defined martingale.

Whence (i) is confirmed, by the Girsanov theorem, under probability $d\mathbb{Q}_T := \mathbb{R}_T d\mathbb{P}$ the process

$$\tilde{B}_t := B_t - \int_0^t \langle \phi_s, ds \rangle, \quad t \in [0, T]$$

is a d -dimensional Brownian motion and (2.1) reduces to

$$dY(t) = \{Z(t, Y(t)) + b(t, Y_t)\} dt + \sigma(t, Y(t)) d\tilde{B}(t), \quad t \in [0, T], Y_0 = \eta. \quad (2.3)$$

Therefore, (2.1) has a unique solution $\{Y(t)\}_{t \in [0, T]}$ under the probability \mathbb{Q}_T , and

$$P_T f(\eta) = \mathbb{E}_{\mathbb{Q}_T} f(Y_T) = \mathbb{E}[R_T f(Y_T)]. \quad (2.4)$$

Next, we shall prove that

(ii) The coupling time $\tau := \inf\{t \geq 0 : X(t) = Y(t)\} \leq t_0$, \mathbb{Q}_T -a.s..

From (1.1) and (2.1) we see that for $t \geq \tau$, the two processes $X(t)$ and $Y(t)$ solve the same equation, because the additional drift term disappears as soon as $X(t) = Y(t)$. By the uniqueness

of the solution to (1.1) we have $X(t) = Y(t)$ for $t \geq \tau$. Combining this with (ii) and noting that $t_0 \leq T - r_0$, we conclude that $X_T = Y_T$, \mathbb{Q}_T -a.s.. So, by the Young inequality and (2.4), we arrive at (see [3])

$$P_T \log f(\eta) = \mathbb{E}[R_T \log f(Y_T)] = \mathbb{E}[R_T \log f(X_T)] \leq \log P_T f(\xi) + \mathbb{E}R_T \log R_T.$$

Therefore, to complete the proof it remains to show that

$$(iii) \mathbb{E}[R_T \log R_T] \leq \frac{2K_3^2 K_4 |\xi(0) - \eta(0)|^2}{1 - e^{-K_4 t_0}} + 3K_1^2 \left\{ \frac{r_0}{2} + t_0(1 + K_2^2 K_3^2) \right\} e^{3K_2^2(K_1^2 t_0 + 4)t_0} \|\xi - \eta\|_\infty^2.$$

In the remainder of the section, we will prove the above claimed (i)–(iii) respectively.

2.1. Proofs of (i)

The key result of this subsection is the following.

Proposition 2.1. Assume (A). Then for any $t \in [0, t_0]$,

$$\mathbb{E}[R_t \log R_t] \leq \frac{2K_3^2 K_4 |\xi(0) - \eta(0)|^2}{\theta(2 - \theta)(1 - e^{-K_4 t_0})} + \frac{12t K_1^2(1 + K_2^2 K_3^2) e^{3K_2^2(K_1^2 t + 4)t}}{\theta^2} \|\xi - \eta\|_\infty^2.$$

Proof. Let $t \in (0, t_0)$ be fixed. Then $\{\tilde{B}(s)\}_{s \leq t}$ is a d -dimensional Brownian motion under the probability $d\mathbb{Q}_t := R_t d\mathbb{P}$, so that (A1) and (A3) imply

$$\begin{aligned} \mathbb{E}[R_t \log R_t] &= \mathbb{E}_{\mathbb{Q}_t} \log R_t = \mathbb{E}_{\mathbb{Q}_t} \left\{ \int_0^t \langle \phi_s, d\tilde{B}(s) \rangle + \frac{1}{2} \int_0^t |\phi_s|^2 ds \right\} \\ &= \frac{1}{2} \int_0^t \mathbb{E}_{\mathbb{Q}_t} |\phi_s|^2 ds \\ &\leq K_1^2 \int_0^t \mathbb{E}_{\mathbb{Q}_t} \|Y_s - X_s\|_\infty^2 ds + K_3^2 \int_0^t \frac{1}{\gamma(s)^2} \mathbb{E}_{\mathbb{Q}_t} |X(s) - Y(s)|^2 ds \\ &=: I_1 + I_2. \end{aligned} \quad (2.5)$$

To estimate I_1 and I_2 , let us reformulate Eq. (1.1) using the new Brownian motion $\tilde{B}(s)$:

$$dX(s) = \{Z(s, X(s)) + b(s, X_s) + \sigma(s, X(s))\phi_s\} ds + \sigma(s, X(s))d\tilde{B}(s), \quad s \leq t.$$

Since

$$\begin{aligned} &\sigma(s, X(s))\phi_s + b(s, X_s) - b(s, Y_s) \\ &= \{\sigma(s, X(s)) - \sigma(s, Y(s))\} \sigma(s, Y(s))^{-1} (b(s, Y_s) - b(s, X_s)) - \frac{X(s) - Y(s)}{\gamma(s)}, \end{aligned}$$

the equation reduces to

$$\begin{aligned} dX(s) &= \left\{ \{\sigma(s, X(s)) - \sigma(s, Y(s))\} \sigma(s, Y(s))^{-1} (b(s, Y_s) - b(s, X_s)) \right. \\ &\quad \left. + Z(s, X(s)) + b(s, Y_s) - \frac{X(s) - Y(s)}{\gamma(s)} \right\} ds + \sigma(s, X(s))d\tilde{B}(s), \\ &\quad s \leq t. \end{aligned} \quad (2.6)$$

Combining this with (2.3) and using the Itô formula, we obtain from (A1), (A2) and (A4) that

$$\begin{aligned} d|X(s) - Y(s)|^2 &\leq 2 \left\langle X(s) - Y(s), (\sigma(s, X(s)) - \sigma(s, Y(s))) d\tilde{B}(s) \right\rangle \\ &\quad + \left\{ 2K_1 K_2 \|X_s - Y_s\|_\infty |X(s) - Y(s)| + \left(K_4 - \frac{2}{\gamma(s)} \right) |X(s) - Y(s)|^2 \right\} ds, \\ s &\leq t. \end{aligned} \quad (2.7)$$

Since it is easy to see that

$$K_4 \leq \frac{2}{\gamma(0)} \leq \frac{2}{\gamma(s)},$$

it follows that

$$\begin{aligned} d|X(s) - Y(s)| &\leq \left\langle \frac{X(s) - Y(s)}{|X(s) - Y(s)|}, (\sigma(s, X(s)) - \sigma(s, Y(s))) d\tilde{B}(s) \right\rangle \\ &\quad + K_1 K_2 \|X_s - Y_s\|_\infty ds, \quad s \leq t. \end{aligned} \quad (2.8)$$

Let

$$M(s) = \int_0^s \left\langle \frac{X(r) - Y(r)}{|X(r) - Y(r)|}, (\sigma(r, X(r)) - \sigma(r, Y(r))) d\tilde{B}(r) \right\rangle, \quad s \leq t,$$

which is a martingale under \mathbb{Q}_t . By (A2) and the Doob inequality we have

$$\mathbb{E}_{\mathbb{Q}_t} \sup_{r \in [0, s]} M(r)^2 \leq 4K_2^2 \int_0^s \mathbb{E}_{\mathbb{Q}_t} \|X_r - Y_r\|_\infty^2 dr, \quad s \leq t.$$

Combining this with (2.8) we obtain

$$\mathbb{E}_{\mathbb{Q}_t} \|X_s - Y_s\|_\infty^2 \leq 3\|\xi - \eta\|_\infty^2 + 3K_2^2(K_1^2 s + 4) \int_0^s \mathbb{E}_{\mathbb{Q}_t} \|X_r - Y_r\|_\infty^2 dr, \quad s \leq t.$$

By the Gronwall lemma, this implies that

$$\mathbb{E}_{\mathbb{Q}_t} \|X_s - Y_s\|_\infty^2 \leq 3\|\xi - \eta\|_\infty^2 e^{3K_2^2(K_1^2 s + 4)s}, \quad s \leq t. \quad (2.9)$$

On the other hand, let

$$d\tilde{M}(s) = \frac{2}{\gamma(s)} \left\langle X(s) - Y(s), (\sigma(s, X(s)) - \sigma(s, Y(s))) d\tilde{B}(s) \right\rangle, \quad s \leq t,$$

which is a martingale under \mathbb{Q}_t . It follows from (2.2) and (2.7) that

$$\begin{aligned} d \frac{|X(s) - Y(s)|^2}{\gamma(s)} - d\tilde{M}(s) &\leq \left\{ \frac{2K_1 K_2}{\gamma(s)} \|X_s - Y_s\|_\infty |X(s) - Y(s)| + \frac{K_4 \gamma(s) - 2 - \gamma'(s)}{\gamma(s)^2} |X(s) - Y(s)|^2 \right\} ds \\ &\leq \left\{ \frac{2K_1 K_2}{\gamma(s)} \|X_s - Y_s\|_\infty |X(s) - Y(s)| - \frac{\theta |X(s) - Y(s)|^2}{\gamma(s)^2} \right\} ds, \quad s \leq t. \end{aligned} \quad (2.10)$$

Combining this with (2.9), we obtain

$$\begin{aligned} h(t) &:= \int_0^t \frac{\mathbb{E}_{\mathbb{Q}_t} |X(s) - Y(s)|^2}{\gamma(s)^2} ds \\ &\leq \frac{|\xi(0) - \eta(0)|^2}{\theta \gamma(0)} + \frac{2K_1 K_2}{\theta} h(t)^{1/2} \left(\int_0^t \mathbb{E}_{\mathbb{Q}_t} \|X_s - Y_s\|_\infty^2 ds \right)^{1/2} \\ &\leq \frac{|\xi(0) - \eta(0)|^2}{\theta \gamma(0)} + \frac{h(t)}{2} + \frac{6K_1^2 K_2^2 t}{\theta^2} e^{3K_2^2 (K_1^2 t + 4)t} \|\xi - \eta\|_\infty^2. \end{aligned}$$

Therefore,

$$\int_0^t \frac{\mathbb{E}_{\mathbb{Q}_t} |X(s) - Y(s)|^2}{\gamma(s)^2} ds \leq \frac{2|\xi(0) - \eta(0)|^2}{\theta \gamma(0)} + \frac{12K_1^2 K_2^2 t e^{3K_2^2 (K_1^2 t + 4)t}}{\theta^2} \|\xi - \eta\|_\infty^2. \quad (2.11)$$

Substituting this and (2.9) into (2.5), we complete the proof. \square

Proof of (i). According to Proposition 2.1, $\{R_t\}_{t \in [0, t_0]}$ is a uniformly integrable continuous martingale. So, by the martingale convergence theorem,

$$R_{t_0} = \lim_{t \uparrow t_0} R_t \quad (2.12)$$

exists and $\{R_t\}_{t \in [0, t_0]}$ is again a uniformly integrable martingale. In particular, $\tilde{B}(t)$ is a d -dimensional Brownian motion under $d\mathbb{Q}_{t_0} := R_{t_0} d\mathbb{P}$ such that $Y(t)$ can be solved from (2.3) for $t \in [0, t_0]$. To solve the equation for $t > t_0$, let us use the filtered probability space $(\Omega, \{\mathcal{F}_t\}_{t \geq t_0}, \mathbb{Q}_{t_0})$. Since $\{B(t) - B(t_0)\}_{t \geq t_0}$ is independent of R_{t_0} , it is easy to see that $\{B(t)\}_{t \geq t_0}$ is a d -dimensional Brownian motion on this probability space. Moreover, due to (2.9), X_{t_0} and Y_{t_0} are \mathcal{F}_{t_0} -measurable random variables on \mathcal{C} with

$$\mathbb{E}_{\mathbb{Q}_{t_0}} \|X_{t_0} - Y_{t_0}\|_\infty^2 \leq 3\|\xi - \eta\|_\infty^2 e^{3K_2^2 (K_1^2 t_0 + 4)t_0}. \quad (2.13)$$

Therefore, by (A2) and (A4), starting from Y_{t_0} at time t_0 the Eq. (2.1) has a unique solution $\{Y(t)\}_{t \geq t_0}$, and by the Itô formula and (A4),

$$\begin{aligned} d|X(t) - Y(t)|^2 &\leq 2\langle X(t) - Y(t), (\sigma(t, X(t)) - \sigma(t, Y(t))) dB(t) \rangle \\ &\quad + K_4 |X(t) - Y(t)|^2 dt, \quad t \geq t_0. \end{aligned}$$

Combining this with (A2) and noting that $\{B(t)\}_{t \geq t_0}$ is a \mathbb{Q}_{t_0} -Brownian motion, we obtain from the Burkholder–Davies–Gundy inequality that

$$\begin{aligned} &\mathbb{E}_{\mathbb{Q}_{t_0}} \left(\sup_{s \in [t_0, t]} |X(s) - Y(s)|^2 \middle| \mathcal{F}_{t_0} \right) \\ &\leq |X(t_0) - Y(t_0)|^2 + 2K_2 \left(\int_{t_0}^t \mathbb{E}_{\mathbb{Q}_{t_0}} \left(|X(s) - Y(s)|^2 \middle| \mathcal{F}_{t_0} \right) ds \right)^{1/2} \\ &\quad + K_4 \int_{t_0}^t \mathbb{E}_{\mathbb{Q}_{t_0}} \left(|X(s) - Y(s)|^2 \middle| \mathcal{F}_{t_0} \right) ds \\ &\leq \|X_{t_0} - Y_{t_0}\|_\infty^2 + 1 + K \int_{t_0}^t \mathbb{E}_{\mathbb{Q}_{t_0}} \left(\sup_{r \in [t_0, s]} |X(r) - Y(r)|^2 \middle| \mathcal{F}_{t_0} \right) ds \end{aligned}$$

holds for some constant $K > 0$ and all $t \geq t_0$. Combining this with (2.13) we arrive at

$$\mathbb{E}_{\mathbb{Q}_{t_0}} \sup_{s \in [t_0, t]} \|X_s - Y_s\|_\infty^2 \leq \left(\mathbb{E}_{\mathbb{Q}_{t_0}} \|X_{t_0} - Y_{t_0}\|_\infty^2 + 1 \right) e^{K(t-t_0)} < \infty, \quad t \geq t_0.$$

Since

$$|\phi_t| = \left| \sigma(t, Y(t))^{-1} (b(t, Y_t) - b(t, X_t)) \right| \leq K_1 \|X_t - Y_t\|_\infty, \quad t \geq t_0,$$

this implies that

$$\frac{R_t}{R_{t_0}} = \exp \left[\int_{T-r}^t \langle \phi_s, dB(s) \rangle - \frac{1}{2} \int_{t_0}^t |\phi_s|^2 ds \right], \quad t \geq t_0$$

is a \mathbb{Q}_{t_0} -martingale, and thus, for $t > s \geq t_0$ and $A \in \mathcal{F}_s$,

$$\mathbb{E}(R_t 1_A) = \mathbb{E}_{\mathbb{Q}_{t_0}} \left\{ 1_A \frac{R_t}{R_{t_0}} \right\} = \mathbb{E}_{\mathbb{Q}_{t_0}} \left\{ 1_A \frac{R_s}{R_{t_0}} \right\} = \mathbb{E}(R_s 1_A).$$

This means that $\{R_t\}_{t \geq t_0}$ is a \mathbb{P} -martingale, and thus, $\{R_t\}_{t \geq 0}$ is a well-defined \mathbb{P} -martingale as claimed since $\{R_t\}_{t \in [0, t_0]}$ is already a martingale. \square

2.2. Proof of (ii)

Since $\{R_t\}_{t \in [0, T]}$ is a martingale, for any $t \in [0, t_0)$ the inequality (2.11) holds for \mathbb{Q}_T in place of \mathbb{Q}_t . Therefore,

$$\mathbb{E}_{\mathbb{Q}_T} \int_0^{t_0} \frac{|X(t) - Y(t)|^2}{\gamma(t)^2} dt < \infty. \quad (2.14)$$

This implies that $\tau \leq t_0$, \mathbb{Q}_T -a.s.. Indeed, since $t \mapsto X(t)$ and $t \mapsto Y(t)$ are continuous \mathbb{Q}_T -a.s., there exists $\Omega_0 \subset \Omega$ with $\mathbb{Q}_T(\Omega_0) = 1$ such that for any $\omega \in \Omega_0$, $X(t)(\omega)$ and $Y(t)(\omega)$ are continuous in t . If $\omega \in \Omega_0$ such that $\tau(\omega) > t_0$, then

$$\inf_{t \in [0, t_0]} |X(t) - Y(t)|(\omega) > 0,$$

so that

$$\int_0^{t_0} \frac{|X(t) - Y(t)|^2(\omega)}{\gamma(t)^2} dt \geq \inf_{t \in [0, t_0]} |X(t) - Y(t)|(\omega) \int_0^{t_0} \frac{dt}{\gamma(t)^2} = \infty.$$

This means that

$$\mathbb{Q}_T(\tau > t_0) \leq \mathbb{Q}_T \left(\int_0^{t_0} \frac{|X(t) - Y(t)|^2}{\gamma(t)^2} dt = \infty \right)$$

which equals to zero according to (2.14).

2.3. Proof of (iii)

Since $\{R_t\}_{t \geq 0}$ is a martingale, by the Girsanov theorem $\{\tilde{B}(t)\}_{t \in [0, T]}$ is Brownian motion under \mathbb{Q}_T . Then

$$\begin{aligned}
\mathbb{E}[R_T \log R_T] &= \frac{1}{2} \mathbb{E}_{\mathbb{Q}_T} \int_0^T |\phi_t|^2 dt \\
&= \frac{1}{2} \mathbb{E}_{\mathbb{Q}_{t_0}} \int_0^{t_0} |\phi_t|^2 dt + \frac{1}{2} \mathbb{E}_{\mathbb{Q}_T} \int_{t_0}^T |\phi_t|^2 dt \\
&= \mathbb{E}[R_{t_0} \log R_{t_0}] + \frac{1}{2} \mathbb{E}_{\mathbb{Q}_T} \int_{t_0}^T |\phi_t|^2 dt.
\end{aligned} \tag{2.15}$$

Since $\tau \leq t_0$ and $X(t) = Y(t)$ for $t \geq \tau$, we have $X(t) = Y(t)$ for $t \geq t_0$. So, it follows from (A1) that

$$\int_{t_0}^T |\phi_t|^2 dt \leq K_1^2 \int_{t_0}^T \|X_t - Y_t\|_\infty^2 dt \leq K_1^2 r_0 \|Y_{t_0} - X_{t_0}\|_\infty^2.$$

Combining this with (2.9), which also holds for $t = s = t_0$ by (2.12) and the Fatou lemma, we arrive at

$$\frac{1}{2} \mathbb{E}_{\mathbb{Q}_T} \int_{t_0}^T |\phi_t|^2 dt \leq \frac{3K_1^2 r_0}{2} e^{3K_2^2(K_1^2 t_0 + 4)t_0} \|\xi - \eta\|_\infty^2.$$

Substituting this into (2.15) and noting that (2.12) and Proposition 2.1 with $\theta = 1$ imply

$$\mathbb{E}[R_{t_0} \log R_{t_0}] \leq \frac{2K_3^2 K_4 |\xi(0) - \eta(0)|^2}{1 - e^{-K_4 t_0}} + 12t_0 K_1^2 (1 + K_2^2 K_3^2) e^{3K_2^2(K_1^2 t_0 + 4)t_0} \|\xi - \eta\|_\infty^2,$$

we prove (iii).

3. Proofs of Corollary 1.2 and Proposition 1.3

Proof of Corollary 1.2. According to Theorem 1.1, we have

$$e^{P_T f}(\xi) \leq \{P_T e^f(\eta)\} e^{\Psi_T(\|\xi - \eta\|_\infty)}, \quad \xi, \eta \in \mathcal{C}, T > r_0, f \in \mathcal{B}_b(\mathcal{C}), f \geq 0$$

holds for some continuous function Ψ_T with $\lim_{r \rightarrow 0} \Psi_T(r) = 0$. So, the desired assertions follow immediately from the following more general result for $\Phi(r) = e^r$. \square

Proposition 3.1. Let (E, \mathcal{F}) be the Borel measurable space of a topology space E , P a Markov operator on $\mathcal{B}_b(E)$, and μ a quasi-invariant probability measure of P . Let $\Phi \in C^1([0, \infty))$ be an increasing function with $\Phi'(1) > 0$ and $\Phi(\infty) := \lim_{r \rightarrow \infty} \Phi(r) = \infty$, such that

$$\Phi(Pf(x)) \leq \{P\Phi(f)(y)\} e^{\Psi(x,y)}, \quad x, y \in E, f \in \mathcal{B}_b(E), f \geq 0 \tag{3.1}$$

holds for some measurable non-negative function Ψ on E^2 .

- (1) If $\lim_{y \rightarrow x} \{\Psi(x, y) + \Psi(y, x)\} = 0$ holds for all $x \in E$, then P is strong Feller.
- (2) P has a kernel p w.r.t. μ , so that any invariant probability measure of P is absolutely continuous w.r.t. μ .
- (3) P has at most one invariant probability measure and if it has, the kernel of P w.r.t. the invariant probability measure is strictly positive.
- (4) The kernel p of P w.r.t. μ satisfies

$$\int_E p(x, \cdot) \Phi^{-1} \left(\frac{p(x, \cdot)}{p(y, \cdot)} \right) d\mu \leq \Phi^{-1}(e^{\Psi(x,y)}), \quad x, y \in E,$$

where $\Phi^{-1}(\infty) := \infty$ by convention.

(5) If $r\Phi^{-1}(r)$ is convex for $r \geq 0$, then the kernel p of P w.r.t. μ satisfies

$$\int_E p(x, \cdot) p(y, \cdot) d\mu \geq e^{-\Psi(x, y)}, \quad x, y \in E.$$

Proof. (1) Let $f \in \mathcal{B}_b(E)$ be positive. Applying (3.1) to $1 + \varepsilon f$ in place of f for $\varepsilon > 0$, we have

$$\Phi(1 + \varepsilon Pf(x)) \leq \{P\Phi(1 + \varepsilon f)(y)\}e^{\Psi(x, y)}, \quad x, y \in E, \varepsilon > 0.$$

By the Taylor expansion this implies

$$\Phi(1) + \varepsilon \Phi'(1)Pf(x) + o(\varepsilon) \leq \{\Phi(1) + \varepsilon \Phi'(1)Pf(y) + o(\varepsilon)\}e^{\Psi(x, y)} \quad (3.2)$$

for small $\varepsilon > 0$. Letting $y \rightarrow x$ we obtain

$$\varepsilon Pf(x) \leq \varepsilon \liminf_{y \rightarrow x} Pf(y) + o(\varepsilon).$$

Thus, $Pf(x) \leq \lim_{y \rightarrow x} Pf(y)$ holds for all $x \in E$. On the other hand, letting $x \rightarrow y$ in (3.2) gives $Pf(y) \geq \limsup_{x \rightarrow y} Pf(x)$ for any $y \in E$. Therefore, Pf is continuous.

(2) To prove the existence of kernel, it suffices to prove that for any $A \in \mathcal{F}$ with $\mu(A) = 0$ we have $P1_A \equiv 0$. Applying (3.1) to $f = 1 + n1_A$, we obtain

$$\Phi(1 + nP1_A(x)) \int_E e^{-\Psi(x, y)} \mu(dy) \leq \int_E \Phi(1 + n1_A(y))(\mu P)(dy), \quad n \geq 1. \quad (3.3)$$

Since $\mu(A) = 0$ and μ is quasi-invariant for P , we have $1_A = 0$, μP -a.s.. So, it follows from (3.3) that

$$\Phi(1 + nP1_A(x)) \leq \frac{\Phi(1)}{\int_E e^{-\Psi(x, y)} \mu(dy)} < \infty, \quad x \in E, n \geq 1.$$

Since $\Phi(1 + n) \rightarrow \infty$ as $n \rightarrow \infty$, this implies that $P1_A(x) = 0$ for all $x \in E$.

Now, for any invariant probability measure μ_0 of P , if $\mu(A) = 0$ then $P1_A \equiv 0$ implies that $\mu_0(A) = \mu_0(P1_A) = 0$. Therefore, μ_0 is absolutely continuous w.r.t. μ .

(3) We only need to prove that the kernel of P w.r.t. an invariant probability measure μ_0 is strictly positive, which implies the uniqueness of the invariant measure according to e.g. [5, Proposition 4.1.1 and Theorem 4.1.2]. To this end, it suffices to show that for any $x \in E$ and $A \in \mathcal{F}$, $P1_A(x) = 0$ implies that $\mu_0(A) = 0$. Since $P1_A(x) = 0$, applying (3.1) to $f = 1 + nP1_A$ we obtain

$$\Phi(1 + nP1_A(y)) \leq \{P\Phi(1 + n1_A)(x)\}e^{\Psi(y, x)} = \Phi(1)e^{\Psi(y, x)}, \quad y \in E, n \geq 1.$$

Letting $n \rightarrow \infty$ we conclude that $P1_A \equiv 0$ and hence, $\mu_0(A) = \mu_0(P1_A) = 0$.

(4) Applying (3.1) to

$$f = n \wedge \Phi^{-1}\left(\frac{p(x, \cdot)}{p(y, \cdot)}\right)$$

and letting $n \rightarrow \infty$, we obtain the desired inequality.

(5) Let $r\Phi^{-1}(r)$ be convex for $r \geq 0$. By the Jensen inequality we have

$$\int_E p(x, \cdot) \Phi^{-1}(p(x, \cdot)) d\mu \geq \Phi^{-1}(1).$$

So, applying (3.1) to

$$f = n \wedge \Phi^{-1}(p(x, \cdot))$$

and letting $n \rightarrow \infty$, we obtain

$$\int_E p(x, \cdot) p(y, \cdot) d\mu \geq e^{-\Psi(x,y)} \Phi \left(\int_E p(x, \cdot) \Phi^{-1}(p(x, \cdot)) d\mu \right) \geq e^{-\Psi(x,y)}. \quad \square$$

Proof of Proposition 1.3. Consider the Itô SDE without delay:

$$d\tilde{X}(t) = Z(\tilde{X}(t))dt + \sigma(\tilde{X}(t))dB(t). \quad (3.4)$$

The equation has a unique strong solution which is a strong Markov process. Since $-\|\sigma(x)\|_{HS}^2 - \langle x, Z(x) \rangle$ is a compact function, it is standard that the process has an (indeed unique, due to the ellipticity) invariant probability measure μ_0 , so that with initial distribution μ_0 the process is stationary. Let μ be the distribution of the \mathcal{C} -valued random variable

$$\{\tilde{X}(r_0 + u)\}_{u \in [-r_0, 0]},$$

where $\tilde{X}(0)$ has distribution μ_0 . Then the \mathcal{C} -valued Markov process $\{X_t\}_{t \geq 0}$ with

$$X_t(u) := X(t+u) := \tilde{X}(t+r_0+u), \quad u \in [-r_0, 0]$$

has an invariant probability measure μ . Let $\bar{B}(t) = B(t+r_0) - B(r_0)$, we have

$$dX(t) = Z(X(t))dt + \sigma(X(t))d\bar{B}(t), \quad t \geq 0. \quad (3.5)$$

As before, let $X^\xi(t)$ be the solution of this equation with $X_0 = \xi$. To formulate $P_t f(\xi)$ using X_t^ξ , we take

$$\tilde{B}^\xi(t) = \bar{B}(t) + \int_0^t \sigma(X^\xi(s))^{-1} b(s, X_s^\xi) ds.$$

Then (3.5) implies that

$$dX^\xi(t) = \{Z(X^\xi(t)) + b(t, X_t^\xi)\}dt + \sigma(X^\xi(t))d\tilde{B}^\xi(t), \quad t \geq 0.$$

By (A) it is easy to see that

$$R_t^\xi := \exp \left[\int_0^t \langle b(s, X_s^\xi), \sigma(X^\xi(s)) d\bar{B}(s) \rangle - \frac{1}{2} \int_0^t |\sigma(X^\xi(s))^{-1} b(s, X_s^\xi)|^2 ds \right]$$

is a martingale, and by the Girsanov theorem for any $T > 0$, $\{\tilde{B}^\xi(t)\}_{t \in [0, T]}$ is a Brownian motion under probability $d\mathbb{Q}_T^\xi := R_T^\xi d\mathbb{P}$. Therefore,

$$P_T f(\xi) = \mathbb{E}[R_T^\xi f(X_T^\xi)], \quad T \geq 0, f \in \mathcal{B}_b(\mathcal{C}). \quad (3.6)$$

Since μ is an invariant probability measure of X_t , for any μ -null set A we have

$$\int_{\Omega \times \mathcal{C}} 1_A(X_T^\xi(\omega)) (\mathbb{P} \times \mu)(d\omega, d\xi) = \mu(A) = 0.$$

Combining this with (3.6) we obtain

$$(\mu P_T)(A) = \int_{\mathcal{C}} P_T 1_A d\mu = \int_{\Omega \times \mathcal{C}} \left\{ 1_A(X_T^\xi(\omega)) R_T^\xi(\omega) \right\} (\mathbb{P} \times \mu)(d\omega, d\xi) = 0.$$

Therefore, μ is a quasi-invariant probability measure of P_T . \square

4. The Harnack inequality

In this section we aim to establish the Harnack inequality with a power $p > 1$ in the sense of [14]:

$$P_T f(\eta) \leq \{P_T f^p(\xi)\}^{1/p} \exp[\Phi_p(T, \xi, \eta)], \quad f \geq 0, T > r_0, \xi, \eta \in \mathcal{C} \quad (4.1)$$

for some positive function Φ_p on $(r_0, \infty) \times \mathcal{C}^2$. As shown in [18] for the case without delay, we will have to assume that $p > (1 + K_2 K_3)^2$. In this case, letting

$$\lambda_p = \frac{1}{2(\sqrt{p} - 1)^2},$$

the set

$$\Theta_p := \left\{ \varepsilon \in (0, 1) : \frac{(1 - \varepsilon)^4}{2(1 + \varepsilon)^3 K_2^2 K_3^2} \geq \lambda_p \right\}$$

is non-empty. Let

$$W_\varepsilon(\lambda) = \max \left\{ \frac{8(1 + \varepsilon)r_0 K_1^3 K_2 \lambda \{4(1 + \varepsilon)r_0 K_1 K_2 \lambda + \varepsilon\}}{\varepsilon^2}, \frac{2(1 + \varepsilon)^2 \lambda}{\varepsilon^2}, \right. \\ \left. \frac{(1 + \varepsilon)^3 K_1^2 K_2^2 K_3^2 \lambda}{8\varepsilon^2(1 - \varepsilon)^3} \right\},$$

and

$$s_\varepsilon(\lambda) = \frac{\sqrt{K_1^2 + 2W_\varepsilon(\lambda)} - K_1}{4W_\varepsilon(\lambda)K_2}, \quad \varepsilon \in (0, 1), \lambda > 0.$$

The following result provides an explicit Harnack inequality. Since to derive this inequality we have to use the Hölder inequality for several times which leads to the loss of sharpness, when $\sigma = I$ our inequality does not cover the one derived in [7].

Theorem 4.1. Assume (A) for $T > r_0$. For any $p > (1 + K_2 K_3)^2$, the Harnack inequality (4.1) holds for

$$\Phi_p(T, \xi, \eta) := \frac{\sqrt{p} - 1}{\sqrt{p}} \inf_{\varepsilon \in \Theta_p} \inf_{s \in (0, s_\varepsilon(\lambda_p) \wedge (T - r_0)]} \left\{ \frac{\varepsilon}{2(1 + \varepsilon)} + \frac{16K_2^2 s^2 W_\varepsilon(\lambda_p)}{1 - 4K_1 K_2 s} \right. \\ \left. + \frac{\lambda_p(1 + \varepsilon)^2 K_3^2 K_4 |\xi(0) - \eta(0)|^2}{2\varepsilon(1 - \varepsilon)^2(1 + 2\varepsilon)(1 - e^{-K_4 s})} + (K_1^2 r_0 \lambda_p + 2s W_\varepsilon(\lambda_p)) \|\xi - \eta\|_\infty^2 \right\}.$$

Consequently, there exists a decreasing function $C : ((1 + K_2 K_3)^2, \infty) \rightarrow (0, \infty)$ such that (4.1) holds for

$$\Phi_p(T, \xi, \eta) = C(p) \left\{ 1 + \frac{|\xi(0) - \eta(0)|^2}{T - r_0} + \|\xi - \eta\|_\infty^2 \right\}.$$

Proof. (a) We first observe that the second assertion is a consequence of the first. Indeed, for any $q > (1 + K_2 K_3)^2$, we take $(\varepsilon, s) = (\varepsilon_q, s_q(T))$ for a fixed $\varepsilon_q \in \Theta_q$ and $s_q(T) :=$

$s_{\varepsilon_q}(\lambda_q) \wedge (T - r_0)$. By the definition of Φ_q , there exists two positive constants $c_1(q)$ and $c_2(q)$ such that

$$\begin{aligned}\Phi_q(T, \xi, \eta) &\leq c_1(q) \left(1 + \|\xi - \eta\|_\infty^2 + \frac{|\xi(0) - \eta(0)|^2}{c_2(q) \wedge (T - r_0)} \right) \\ &\leq \frac{c_1(q)(1 + c_2(q))}{c_2(q)} \left(1 + \|\xi - \eta\|_\infty^2 + \frac{|\xi(0) - \eta(0)|^2}{T - r_0} \right), \\ &\quad T > r_0, \xi, \eta \in \mathcal{C}.\end{aligned}$$

So, for any $p > (1 + K_2 K_3)^2$ and any $q \in ((1 + K_2 K_3)^2, p]$, by the first assertion and using the Jensen inequality, we obtain

$$\begin{aligned}P_T f(\eta) &\leq (P_T f^q)^{1/q}(\xi) \exp \left[\frac{c_1(q)(1 + c_2(q))}{c_2(q)} \left(1 + \|\xi - \eta\|_\infty^2 + \frac{|\xi(0) - \eta(0)|^2}{T - r_0} \right) \right] \\ &\leq (P_T f^p)^{1/p}(\xi) \exp \left[\frac{c_1(q)(1 + c_2(q))}{c_2(q)} \left(1 + \|\xi - \eta\|_\infty^2 + \frac{|\xi(0) - \eta(0)|^2}{T - r_0} \right) \right].\end{aligned}$$

Therefore, the second assertion holds for

$$C(p) = \inf_{q \in ((1 + K_2 K_3)^2, p]} \frac{c_1(q)(1 + c_2(q))}{c_2(q)}$$

which is decreasing in p .

(b) To prove the first assertion, let us fix $\varepsilon \in \Theta_p$ and $t_0 \in (0, s_\varepsilon(\lambda_p) \wedge (T - r_0)]$. We shall make use of the coupling constructed in Section 2 for $\theta = 2(1 - \varepsilon)$. Since $t_0 \leq T - r_0$ and $X(t) = Y(t)$ for $t \geq t_0$, we have $X_T = Y_T$ and

$$P_T f(\eta) = \mathbb{E}[R_T f(Y_T)] = \mathbb{E}[R_T f(X_T)] \leq (P_T f^p(\xi))^{1/p} (\mathbb{E} R_T^{p/(p-1)})^{(p-1)/p}. \quad (4.2)$$

By the definition of R_T and \mathbb{Q}_T , we have

$$\begin{aligned}\mathbb{E} R_T^{p/(p-1)} &= \mathbb{E}_{\mathbb{Q}_T} R_T^{1/(p-1)} = \mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{1}{p-1} \int_0^T \langle \phi_t, d\tilde{B}(t) \rangle + \frac{1}{2(p-1)} \int_0^T |\phi_t|^2 dt \right] \\ &= \mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{1}{p-1} \int_0^T \langle \phi_t, d\tilde{B}(t) \rangle - \frac{\sqrt{p}+1}{2(p-1)^2} \int_0^T |\phi_t|^2 dt \right. \\ &\quad \left. + \frac{p+\sqrt{p}}{2(p-1)^2} \int_0^T |\phi_t|^2 dt \right] \\ &\leq \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{\sqrt{p}+1}{p-1} \int_0^T \langle \phi_t, d\tilde{B}(t) \rangle - \frac{(\sqrt{p}+1)^2}{2(p-1)^2} \int_0^T |\phi_t|^2 dt \right] \right)^{1/(1+\sqrt{p})} \\ &\quad \times \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{(\sqrt{p}+1)(p+\sqrt{p})}{2(p-1)^2 \sqrt{p}} \int_0^T |\phi_t|^2 dt \right] \right)^{\sqrt{p}/(\sqrt{p}+1)} \\ &= \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_p \int_0^T |\phi_t|^2 dt \right] \right)^{\sqrt{p}/(\sqrt{p}+1)}.\end{aligned}$$

Combining this with (4.2), we obtain

$$P_T f(\eta) \leq (P_T f^p)^{1/p}(\xi) \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_p \int_0^T |\phi_t|^2 dt \right] \right)^{(\sqrt{p}-1)/\sqrt{p}}.$$

Therefore, to prove the first assertion, it suffices to show that

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_p \int_0^T |\phi_t|^2 dt \right] &\leq \exp \left[\frac{\varepsilon}{2(1+\varepsilon)} + \frac{\lambda_p(1+\varepsilon)^2 K_3^2 K_4 |\xi(0) - \eta(0)|^2}{2\varepsilon(1-\varepsilon)^2(1+2\varepsilon)(1-e^{-K_4 s})} \right. \\ &\quad \left. + \frac{16K_2^2 s^2 W_\varepsilon(\lambda_p)}{1-4K_1 K_2 s} + \left(K_1^2 r_0 \lambda_p + 2s W_\varepsilon(\lambda_p) \right) \|\xi - \eta\|_\infty^2 \right]. \end{aligned} \quad (4.3)$$

Since $X(t) = Y(t)$ for $t \geq t_0$, it is easy to see from the definition of ϕ_t and (A1), (A3) that

$$\begin{aligned} \int_0^T |\phi_t|^2 dt &\leq \int_0^{t_0} \left\{ \frac{K_1^2(1+\varepsilon)}{\varepsilon} \|X_t - Y_t\|_\infty^2 + \frac{K_2^2(1+\varepsilon)}{\gamma(t)^2} |X(t) - Y(t)|^2 \right\} dt \\ &\quad + K_1^2 r_0 \|X_{t_0} - Y_{t_0}\|_\infty^2. \end{aligned}$$

By this and the Hölder inequality, we obtain

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_p \int_0^T |\phi_t|^2 dt \right] &\leq \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_p K_3^2(1+\varepsilon)^2 \int_0^{t_0} \frac{|X(t) - Y(t)|^2}{\gamma(t)^2} dt \right] \right)^{1/(1+\varepsilon)} \\ &\quad \times \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{2K_1^2(1+\varepsilon)^2 \lambda_p}{\varepsilon^2} \int_0^{t_0} \|X_t - Y_t\|_\infty^2 dt \right] \right)^{\varepsilon/(2+2\varepsilon)} \\ &\quad \times \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{2K_1^2 r_0(1+\varepsilon) \lambda_p}{\varepsilon} \|X_{t_0} - Y_{t_0}\|_\infty^2 \right] \right)^{\varepsilon/(2+2\varepsilon)}. \end{aligned} \quad (4.4)$$

Since $\varepsilon \in \Theta_p$ implies that

$$\lambda_p K_3^2(1+\varepsilon)^2 \leq \frac{(1-\varepsilon)^4}{2(1+\varepsilon)K_2^2},$$

it follows from Lemma 4.2 below that

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_p K_3^2(1+\varepsilon)^2 \int_0^{t_0} \frac{|X(t) - Y(t)|^2}{\gamma(t)^2} dt \right] &\leq \exp \left[\frac{\lambda_p K_3^2(1+\varepsilon)^3 |\xi(0) - \eta(0)|^2}{(1+2\varepsilon)(1-\varepsilon)^2 \gamma(0)} \right] \\ &\quad \times \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{K_1^2 K_2^2 K_3^2 \lambda_p(1+\varepsilon)^3}{8\varepsilon^2(1-\varepsilon)^3} \int_0^{t_0} \|X_t - Y_t\|_\infty^2 dt \right] \right)^{\varepsilon/(1+2\varepsilon)}. \end{aligned} \quad (4.5)$$

Moreover, according to Lemma 4.3 below,

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{2K_1^2 r_0(1+\varepsilon) \lambda_p}{\varepsilon} \|X_{t_0} - Y_{t_0}\|_\infty^2 \right] &\leq \exp \left[1 + \frac{2K_1^2 r_0(1+\varepsilon) \lambda_p}{\varepsilon} \|\xi - \eta\|_\infty^2 \right] \\ &\quad \times \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{8K_1^3 K_2 r_0(1+\varepsilon) \lambda_p(4K_2 K_1 r_0(1+\varepsilon) \lambda_p + \varepsilon)}{\varepsilon^2} \int_0^{t_0} \|X_t - Y_t\|_\infty^2 dt \right] \right)^{1/2}. \end{aligned}$$

Substituting this and (4.5) into (4.4), and using the definition of $W_\varepsilon(\lambda_p)$, we conclude that

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_p \int_0^T |\phi_t|^2 dt \right] &\leq \mathbb{E}_{\mathbb{Q}_T} \exp \left[W_\varepsilon(\lambda_p) \int_0^{t_0} \|X_t - Y_t\|_\infty^2 dt \right] \\ &\times \exp \left[\frac{\lambda_p K_3^2 (1+\varepsilon)^2 |\xi(0) - \eta(0)|^2}{(1+2\varepsilon)(1-\varepsilon)^2 \gamma(0)} + \frac{\varepsilon}{2(1+\varepsilon)} + K_1^2 r_0 \lambda_p \|\xi - \eta\|_\infty^2 \right]. \end{aligned} \quad (4.6)$$

Since $t_0 \leq s_\varepsilon(\lambda_p)$, we have

$$W_\varepsilon(\lambda_p) \leq \frac{1 - 4K_1 K_2 t_0}{8K_2^2 t_0^2}.$$

So, combining (4.6) with Lemma 4.4 below and noting that for $\theta = 2(1 - \varepsilon)$ one has

$$\gamma(0) = \frac{2\varepsilon}{K_4} (1 - e^{-K_4 t_0}),$$

we prove (4.3). \square

Lemma 4.2. For any positive $\lambda \leq \frac{(1-\varepsilon)^4}{2K_2^2(1+\varepsilon)}$ and $s \in [0, t_0]$,

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda \int_0^s \frac{|X(t) - Y(t)|^2}{\gamma(t)^2} dt \right] &\leq \exp \left[\frac{\lambda(1+\varepsilon)|\xi(0) - \eta(0)|^2}{(1+2\varepsilon)(1-\varepsilon)^2 \gamma(0)} \right] \\ &\times \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{K_1^2 K_2^2 (1+\varepsilon) \lambda}{8\varepsilon^2 (1-\varepsilon)^3} \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \right)^{\varepsilon/(1+2\varepsilon)}. \end{aligned}$$

Proof. Since $\theta = 2(1 - \varepsilon)$ and

$$\frac{K_1 K_2}{\gamma(t)} \|X_t - Y_t\|_\infty |X(t) - Y(t)| \leq \frac{K_1^2 K_2^2}{4\theta\varepsilon} \|X_t - Y_t\|_\infty^2 + \theta\varepsilon \frac{|X(t) - Y(t)|^2}{\gamma(t)^2},$$

it follows from (2.10) that

$$0 \leq \tilde{M}(s) + \frac{|\xi(0) - \eta(0)|^2}{\gamma(0)} + \int_0^s \left\{ \frac{K_1^2 K_2^2 \|X_t - Y_t\|_\infty^2}{8\varepsilon(1-\varepsilon)} - \frac{2(1-\varepsilon)^2 |X(t) - Y(t)|^2}{\gamma(t)^2} \right\} dt.$$

Combining this with (A2) and the fact that

$$\mathbb{E}_{\mathbb{Q}_T} e^{N(s)+L} \leq \left(\mathbb{E}_{\mathbb{Q}_T} e^{2(N)(s)+2L} \right)^{1/2} \quad (4.7)$$

holds for a \mathbb{Q}_T -martingale N and a random variable L , we obtain

$$\begin{aligned} \mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda \int_0^s \frac{|X(t) - Y(t)|^2}{\gamma(t)^2} dt - \frac{\lambda |\xi(0) - \eta(0)|^2}{2\gamma(0)(1-\varepsilon)^2} \right] \\ \leq \mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{\lambda}{2(1-\varepsilon)^2} \tilde{M}(s) + \frac{K_1^2 K_2^2 \lambda}{16\varepsilon(1-\varepsilon)^3} \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \\ \leq \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{2K_2^2 \lambda^2}{(1-\varepsilon)^4} \int_0^s \frac{|X(t) - Y(t)|^2}{\gamma(t)^2} dt + \frac{K_1^2 K_2^2 \lambda}{8\varepsilon(1-\varepsilon)^3} \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \right)^{1/2} \end{aligned}$$

$$\leq \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{2K_2^2(1+\varepsilon)\lambda^2}{(1-\varepsilon)^4} \int_0^s \frac{|X(t) - Y(t)|^2}{\gamma(t)^2} dt \right] \right)^{1/(2+2\varepsilon)} \\ \times \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\frac{K_1^2 K_2^2(1+\varepsilon)\lambda}{8\varepsilon^2(1-\varepsilon)^3} \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \right)^{\varepsilon/(2+2\varepsilon)}.$$

Since

$$\frac{2K_2^2(1+\varepsilon)\lambda^2}{(1-\varepsilon)^4} \leq \lambda$$

and up to an approximation argument as in [18, Proof of Lemma 2.2] we may assume that

$$\mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda \int_0^s \frac{|X(t) - Y(t)|^2}{\gamma(t)^2} dt \right] < \infty,$$

this implies the desired inequality. \square

Lemma 4.3. For any $\lambda > 0$ and $s \in [0, t_0]$,

$$\mathbb{E}_{\mathbb{Q}_T} e^{\lambda \|X_s - Y_s\|_\infty^2} \leq e^{1+\lambda \|\xi - \eta\|_\infty^2} \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[4\lambda K_2(2\lambda K_2 + K_1) \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \right)^{1/2}.$$

Proof. Let

$$N(t) = 2 \int_0^t \langle X(r) - Y(r), (\sigma(r, X(r)) - \sigma(r, Y(r))) d\tilde{B}(r) \rangle, \quad r \leq s,$$

which is a \mathbb{Q}_T -martingale. By (2.7) and noting that $K_4 \leq \frac{2}{\gamma(r)}$, we obtain

$$\|X_t - Y_t\|_\infty^2 \leq \left\{ \sup_{r \in [0, t]} |X(r) - Y(r)|^2 \right\} \vee \|\xi - \eta\|_\infty^2 \\ \leq \|\xi - \eta\|_\infty^2 + \sup_{r \in [0, t]} \left\{ N(r) + 2K_1 K_2 \int_0^r \|X_u - Y_u\|_\infty^2 du \right\}.$$

Combining this with (4.7) and noting that the Doob inequality implies

$$\mathbb{E}_{\mathbb{Q}_T} \sup_{r \in [0, t]} e^{M(r)} = \lim_{p \rightarrow \infty} \mathbb{E}_{\mathbb{Q}_T} \left(\sup_{r \in [0, t]} e^{M(r)/p} \right)^p \\ \leq \lim_{p \rightarrow \infty} \left(\frac{p}{p-1} \right)^p \mathbb{E}_{\mathbb{Q}_T} \left(e^{M(t)/p} \right)^p = e \mathbb{E}_{\mathbb{Q}_T} e^{M(t)}$$

for a \mathbb{Q}_T -submartingale $M(r)$, we arrive at

$$\mathbb{E}_{\mathbb{Q}_T} e^{\lambda \|X_s - Y_s\|_\infty^2 - \lambda \|\xi - \eta\|_\infty^2} \leq \mathbb{E}_{\mathbb{Q}_T} \sup_{t \in [0, s]} \exp \left[\lambda N(t) + 2\lambda K_1 K_2 \int_0^t \|X_r - Y_r\|_\infty^2 dr \right] \\ \leq e \mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda N(s) + 2\lambda K_1 K_2 \int_0^s \|X_t - Y_t\|_\infty^2 dt \right]$$

$$\begin{aligned}
&\leq e \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[2\lambda^2 \langle N \rangle(s) + 4\lambda K_1 K_2 \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \right)^{1/2} \\
&\leq e \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[(8K_2^2 \lambda^2 + 4\lambda K_1 K_2) \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \right)^{1/2}. \quad \square
\end{aligned}$$

Lemma 4.4. For any $s \in (0, t_0]$ and positive $\lambda \leq \frac{1-4K_1K_2s}{8K_2^2s^2}$,

$$\mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \leq \exp \left[\frac{16K_2^2s^2\lambda}{1-4K_1K_2s} + 2s\lambda\|\xi - \eta\|_\infty^2 \right].$$

Proof. Let

$$\lambda_0 = \frac{1-4K_1K_2s}{8K_2^2s^2},$$

which is positive since $s \in (0, s_\varepsilon(\lambda_p)]$. It is easy to see that

$$4K_2s\lambda_0(2K_2s\lambda_0 + K_1) = \lambda_0.$$

So, it follows from the Jensen inequality and Lemma 4.3 that

$$\begin{aligned}
\mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_0 \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] &\leq \frac{1}{s} \int_0^s \mathbb{E}_{\mathbb{Q}_T} e^{\lambda_0 s \|X_t - Y_t\|_\infty^2} ds \\
&\leq e^{1+\lambda_0 s \|\xi - \eta\|_\infty^2} \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[4\lambda_0 K_2 s (2\lambda_0 K_2 s + K_1) \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \right)^{1/2} \\
&= e^{1+\lambda_0 s \|\xi - \eta\|_\infty^2} \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_0 \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \right)^{1/2}.
\end{aligned}$$

Up to an approximation argument as in [18, Proof of Lemma 2.2], we may assume that

$$\mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_0 \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] < \infty,$$

so that this implies

$$\mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_0 \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \leq e^{2+2\lambda_0 s \|\xi - \eta\|_\infty^2}.$$

Therefore, by the Jensen inequality, for any $\lambda \in [0, \lambda_0]$

$$\begin{aligned}
\mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] &\leq \left(\mathbb{E}_{\mathbb{Q}_T} \exp \left[\lambda_0 \int_0^s \|X_t - Y_t\|_\infty^2 dt \right] \right)^{\lambda/\lambda_0} \\
&\leq \exp \left[\frac{2\lambda}{\lambda_0} + 2\lambda s \|\xi - \eta\|_\infty^2 \right]. \quad \square
\end{aligned}$$

Acknowledgment

The authors would like to thank the referee for useful comments and corrections.

References

- [1] S. Aida, T. Zhang, On the small time asymptotics of diffusion processes on path groups, *Pot. Anal.* 16 (2002) 67–78.
- [2] M. Arnaudon, A. Thalmaier, F.-Y. Wang, Harnack inequality and heat kernel estimates on manifolds with curvature unbounded below, *Bull. Sci. Math.* 130 (2006) 223–233.
- [3] M. Arnaudon, A. Thalmaier, F.-Y. Wang, Gradient estimates and Harnack inequalities on non-compact Riemannian manifolds, *Stoch. Proc. Appl.* 119 (2009) 3653–3670. 130 (2006) 223–233.
- [4] G. Da Prato, M. Röckner, F.-Y. Wang, Singular stochastic equations on Hilbert spaces: Harnack inequalities for their transition semigroups, *J. Funct. Anal.* 257 (2009) 992–1017.
- [5] G. Da Prato, J. Zabczyk, *Ergodicity for Infinite Dimensional Systems*, Cambridge University Press, 1996.
- [6] A. Es-Sarhir, O.V. Gaans, M. Scheutzow, Invariant measure for stochastic functional differential equations with superlinear drift term, [arXiv:0903.1959](https://arxiv.org/abs/0903.1959).
- [7] A. Es-Sarhir, M.-K. von Renesse, M. Scheutzow, Harnack inequality for functional SDEs with bounded memory, *Electron. Commun. Probab.* 14 (2009) 560–565.
- [8] M. Hairer, J. Mattingly, M. Scheutzow, Asymptotic coupling and a weak form of Harris’ theorem with applications to stochastic delay equations, [arXiv:0902.4495v2](https://arxiv.org/abs/0902.4495v2).
- [9] H. Kawabi, The parabolic Harnack inequality for the time dependent Ginzburg–Landau type SPDE and its application, *Pot. Anal.* 22 (2005) 61–84.
- [10] W. Liu, Harnack inequality and applications for stochastic evolution equations with monotone drifts, *J. Evol. Equ.* 9 (2009) 747–770.
- [11] W. Liu, F.-Y. Wang, Harnack inequality and strong Feller property for stochastic fast diffusion equations, *J. Math. Anal. Appl.* 342 (2008) 651–662.
- [12] M. Röckner, F.-Y. Wang, Log-Harnack inequality for stochastic differential equations in Hilbert spaces and its consequences, *Infin. Dimens. Anal. Quant. Probab. Relat. Topics* 13 (2010) 27–37.
- [13] M.K.R. Scheutzow, M.K. von Renesse, Existence and uniqueness of solutions of stochastic functional differential equations, *Random Oper. Stoch. Equ.* 18 (3) (2010) 267–284.
- [14] F.-Y. Wang, Logarithmic Sobolev inequalities on noncompact Riemannian manifolds, *Probab. Theory Relat. Fields* 109 (1997) 417–424.
- [15] F.-Y. Wang, Harnack inequalities for log-Sobolev functions and estimates of log-Sobolev constants, *Ann. Probab.* 27 (1999) 653–663.
- [16] F.-Y. Wang, Harnack inequality and applications for stochastic generalized porous media equations, *Ann. Probab.* 35 (2007) 1333–1350.
- [17] F.-Y. Wang, F.-Y. Wang, Harnack inequalities on manifolds with boundary and applications, *J. Math. Pures Appl.* 94 (2010) 304–321.
- [18] F.-Y. Wang, Harnack inequality for SDE with multiplicative noise and extension to Neumann semigroup on non-convex manifolds, in press in *Ann. Probab.* Available online [arXiv:0911.1644](https://arxiv.org/abs/0911.1644).
- [19] F.-Y. Wang, L. Xu, Derivative formula and applications for hyperdissipative stochastic Navier–Stokes/Burgers equations. Available online [arXiv:1009.1464](https://arxiv.org/abs/1009.1464).
- [20] T.-S. Zhang, White noise driven SPDEs with reflection: strong Feller properties and Harnack inequalities, *Potential Anal.* 33 (2010) 137–151.