



ELSEVIER

Available online at www.sciencedirect.com

ScienceDirect

Stochastic Processes and their Applications xx (xxxx) xxx–xxx

stochastic
processes
and their
applicationswww.elsevier.com/locate/spa

Optimal stopping under nonlinear expectation

Ibrahim Ekren^a, Nizar Touzi^b, Jianfeng Zhang^{a,*}

^a University of Southern California, Department of Mathematics, United States

^b CMAP, Ecole Polytechnique Paris, France

Received 9 February 2013; received in revised form 7 January 2014; accepted 12 April 2014

Abstract

Let $X : [0, T] \times \Omega \rightarrow \mathbb{R}$ be a bounded càdlàg process with positive jumps defined on the canonical space of continuous paths Ω . We consider the problem of optimal stopping the process X under a nonlinear expectation operator \mathcal{E} defined as the supremum of expectations over a weakly compact but nondominated family of probability measures. We introduce the corresponding nonlinear Snell envelope. Our main objective is to extend the Snell envelope characterization to the present context. Namely, we prove that the nonlinear Snell envelope is an \mathcal{E} -supermartingale, and an \mathcal{E} -martingale up to its first hitting time of the obstacle X . This result is obtained under an additional uniform continuity property of X . We also extend the result in the context of a random horizon optimal stopping problem.

This result is crucial for the newly developed theory of viscosity solutions of path-dependent PDEs as introduced in Ekren et al. (2014), in the semilinear case, and extended to the fully nonlinear case in the accompanying papers (Ekren et al., 0000, 0000).

© 2014 Published by Elsevier B.V.

MSC: 35D40; 35K10; 60H10; 60H30

Keywords: Nonlinear expectation; Optimal stopping; Snell envelope

1. Introduction

On the canonical space of continuous paths Ω , we consider a bounded càdlàg process $X : [0, T] \times \Omega \rightarrow \mathbb{R}$, with positive jumps, and satisfying some uniform continuity condition.

* Corresponding author. Tel.: +1 213 740 9805.

E-mail addresses: ekren@usc.edu (I. Ekren), nizar.touzi@polytechnique.edu (N. Touzi), jianfenz@usc.edu (J. Zhang).

<http://dx.doi.org/10.1016/j.spa.2014.04.006>

0304-4149/© 2014 Published by Elsevier B.V.

Let H_0 be the first exit time of the canonical process from some convex domain, and $H := H_0 \wedge t_0$ for some $t_0 > 0$. This paper focuses on the problem

$$\sup_{\tau \in \mathcal{T}} \mathcal{E}[X_{\tau \wedge H}], \quad \text{where } \mathcal{E}[\cdot] := \sup_{\mathbb{P} \in \mathcal{P}} \mathbb{E}^{\mathbb{P}}[\cdot],$$

\mathcal{T} is the collection of all stopping times, relative to the natural filtration of the canonical process, and \mathcal{P} is a weakly compact non-dominated family of probability measures.

Our main result is the following. Similar to the standard theory of optimal stopping, we introduce the corresponding nonlinear Snell envelope Y , and we show that the classical Snell envelope characterization holds true in the present context. More precisely, we prove that the Snell envelope Y is an \mathcal{E} -supermartingale, and an \mathcal{E} -martingale up to its first hitting time τ^* of the obstacle. Consequently, τ^* is an optimal stopping time for our problem of optimal stopping under nonlinear expectation.

This result is proved by adapting the classical arguments available in the context of the standard optimal stopping problem under linear expectation. However, such an extension turns out to be highly technical. The first step is to derive the dynamic programming principle in the present context, implying the \mathcal{E} -supermartingale property of the Snell envelope Y . To establish the \mathcal{E} -martingale property on $[0, \tau^*]$, we need to use some limiting argument for a sequence Y_{τ_n} , where τ_n 's are stopping times increasing to τ^* . However, we face one major difficulty related to the fact that in a nonlinear expectation framework the dominated convergence theorem fails in general. It was observed in Denis, Hu and Peng [3] that the monotone convergence theorem holds in this framework if the decreasing sequence of random variables are quasi-continuous. Therefore, one main contribution of this paper is to construct convenient quasi-continuous approximations of the sequence Y_{τ_n} . This allows us to apply the arguments in [3] on Y_{τ_n} , which is decreasing under expectation (but not pointwise!) due to the supermartingale property. The weak compactness of the class \mathcal{P} is crucial for the limiting arguments.

We note that in an one dimensional Markov model with uniformly non-degenerate diffusion, Krylov [10] studied a similar optimal stopping problem in the language of stochastic control (instead of nonlinear expectation). However, his approach relies heavily on the smoothness of the (deterministic) value function, which we do not have here. Indeed, one of the main technical difficulties in our situation is to obtain the locally uniform regularity of the value process.

Our interest in this problem is motivated from the recent notion of viscosity solutions of path-dependent partial differential equations, as developed in [5] and the accompanying papers [6,7]. Our definition is in the spirit of Crandall, Ishii and Lions [2], see also Fleming and Soner [9], but avoids the difficulties related to the fact that our canonical space fails to be locally compact. The key point is that the pointwise maximality condition, in the standard theory of viscosity solution, is replaced by a problem of optimal stopping under nonlinear expectation.

Our previous paper [5] was restricted to the context of semilinear path-dependent partial differential equations. In this special case, our definition of viscosity solutions can be restricted to the context where \mathcal{P} consists of equivalent measures on the canonical space (and hence \mathcal{P} has dominating measures). Consequently, the Snell envelope characterization of the optimal stopping problem under nonlinear expectation is available in the existing literature on reflected backward stochastic differential equations, see e.g. El Karoui et al. [8], Bayraktar, Karatzas and Yao [1]. However, the extension of our definition to the fully nonlinear case requires to consider a nondominated family of measures.

The paper is organized as follows. Section 2 introduces the probabilistic framework. Section 3 formulates the problem of optimal stopping under nonlinear expectation, and contains the

statement of our main results. The proof of the Snell envelope characterization in the deterministic maturity case is reported in Section 4. The more involved case of a random maturity is addressed in Section 5. Finally, in the Appendix we present some additional results.

2. Nondominated family of measures on the canonical space

2.1. The canonical spaces

Let $\Omega := \{\omega \in C([0, T], \mathbb{R}^d) : \omega_0 = \mathbf{0}\}$, the set of continuous paths starting from the origin, B the canonical process, $\mathbb{F} = \{\mathcal{F}_t\}_{0 \leq t \leq T}$ the natural filtration generated by B , \mathbb{P}_0 the Wiener measure, \mathcal{T} the set of \mathbb{F} -stopping times, and $\Lambda := [0, T] \times \Omega$. Moreover, for any sub- σ -field $\mathcal{G} \subset \mathcal{F}_T$, let $\mathbb{L}^0(\mathcal{G})$ denote the set of \mathcal{G} -measurable random variables, and $\mathbb{H}^0(\mathbb{F})$ the set of \mathbb{F} -progressively measurable processes. Here and in the sequel, for notational simplicity, we use $\mathbf{0}$ to denote vectors or matrices with appropriate dimensions whose components are all equal to 0. We define a seminorm on Ω and a pseudometric on Λ as follows: for any $(t, \omega), (t', \omega') \in \Lambda$,

$$\|\omega\|_t := \sup_{0 \leq s \leq t} |\omega_s|, \quad \mathbf{d}_\infty((t, \omega), (t', \omega')) := |t - t'| + \|\omega_{\cdot \wedge t} - \omega'_{\cdot \wedge t'}\|_T. \quad (2.1)$$

Then $(\Omega, \|\cdot\|_T)$ is a Banach space and $(\Lambda, \mathbf{d}_\infty)$ is a complete pseudometric space. In fact, the subspace $\{(t, \omega_{\cdot \wedge t}) : (t, \omega) \in \Lambda\}$ is a complete metric space under \mathbf{d}_∞ .

We next introduce the shifted spaces. Let $0 \leq s \leq t \leq T$.

- Let $\Omega^t := \{\omega \in C([t, T], \mathbb{R}^d) : \omega_t = \mathbf{0}\}$ be the shifted canonical space; B^t the shifted canonical process on Ω^t ; \mathbb{F}^t the shifted filtration generated by B^t , \mathbb{P}_0^t the Wiener measure on Ω^t , \mathcal{T}^t the set of \mathbb{F}^t -stopping times, and $\Lambda^t := [t, T] \times \Omega^t$. Moreover, for any $\mathcal{G} \subset \mathcal{F}_T^t$, $\mathbb{L}^0(\mathcal{G})$ and $\mathbb{H}^0(\mathbb{F}^t)$ are the corresponding sets of measurable random variables and processes, respectively.
- For $\omega \in \Omega^s$ and $\omega' \in \Omega^t$, define the concatenation path $\omega \otimes_t \omega' \in \Omega^s$ by:

$$(\omega \otimes_t \omega')(r) := \omega_r \mathbf{1}_{[s, t)}(r) + (\omega_t + \omega'_r) \mathbf{1}_{[t, T]}(r), \quad \text{for all } r \in [s, T].$$

- Let $0 \leq s < t \leq T$ and $\omega \in \Omega^s$. For any $\xi \in \mathbb{L}^0(\mathcal{F}_t^s)$ and $X \in \mathbb{H}^0(\mathbb{F}^s)$ on Ω^s , define the shifted $\xi^{t, \omega} \in \mathbb{L}^0(\mathcal{F}_t^t)$ and $X^{t, \omega} \in \mathbb{H}^0(\mathbb{F}^t)$ on Ω^t by:

$$\xi^{t, \omega}(\omega') := \xi(\omega \otimes_t \omega'), \quad X^{t, \omega}(\omega') := X(\omega \otimes_t \omega'), \quad \text{for all } \omega' \in \Omega^t.$$

2.2. Capacity and nonlinear expectation

A probability measure \mathbb{P} on Ω is called a semimartingale measure if the canonical process B is a semimartingale under \mathbb{P} . For every constant $L > 0$, we denote by \mathcal{P}^L the collection of all continuous semimartingale measures \mathbb{P} on Ω whose drift and diffusion characteristics are bounded by L and $\sqrt{2L}$, respectively. To be precise, let $\tilde{\Omega} := \Omega^3$ be an enlarged canonical space, $\tilde{B} := (B, A, M)$ be the canonical processes, and $\tilde{\omega} = (\omega, a, m) \in \tilde{\Omega}$ be the paths. For any $\mathbb{P} \in \mathcal{P}^L$, there exists an extension \mathbb{Q} on $\tilde{\Omega}$ such that:

$$\begin{aligned} B &= A + M, \quad A \text{ is absolutely continuous, } M \text{ is a martingale,} \\ |\alpha^\mathbb{P}| &\leq L, \quad \frac{1}{2} \text{tr}((\beta^\mathbb{P})^2) \leq L, \quad \text{where } \alpha_t^\mathbb{P} := \frac{dA_t}{dt}, \quad \mathbb{Q}\text{-a.s.} \\ \beta_t^\mathbb{P} &:= \sqrt{\frac{d\langle M \rangle_t}{dt}}, \end{aligned} \quad (2.2)$$

Similarly, for any $t \in [0, T)$, we may define \mathcal{P}_t^L on Ω^t .

Remark 2.1. Let \mathbb{S}_+^d denote the set of $d \times d$ nonnegative definite matrices.

(i) In \mathbb{Q} -a.s. sense, clearly $\beta^{\mathbb{P}} \in \mathbb{L}^0(\mathbb{F}^B)$ and then $\alpha^{\mathbb{P}} \in \mathbb{L}^0(\mathbb{F}^{B,M})$.

(ii) We may also have the following equivalent characterization of \mathcal{P}^L . Consider the canonical space $\Omega' := \Omega^2$ with canonical processes (B, B') . For any $\mathbb{P} \in \mathcal{P}^L$, there exist a probability measure \mathbb{Q}' and $\alpha^{\mathbb{P}} \in \mathbb{L}^0(\mathbb{F}^{B,B'}, \mathbb{R}^d)$, $\beta^{\mathbb{P}} \in \mathbb{L}^0(\mathbb{F}^B, \mathbb{S}_+^d)$ such that

$$\begin{aligned} |\alpha^{\mathbb{P}}| \leq L, \quad \frac{1}{2} \text{tr}((\beta^{\mathbb{P}})^2) \leq L, \quad \mathbb{Q}'|_{\mathcal{F}_T^B} = \mathbb{P}, \quad \mathbb{Q}'|_{\mathcal{F}_T^{B'}} = \text{Wiener measure}; \\ dB_t = \alpha_t^{\mathbb{P}}(B, B')dt + \beta_t^{\mathbb{P}}(B)dB'_t, \quad \mathbb{Q}'\text{-a.s.} \end{aligned} \quad (2.3)$$

(iii) For any deterministic measurable functions $\alpha : [0, T] \rightarrow \mathbb{R}^d$ and $\beta : [0, T] \rightarrow \mathbb{S}_+^d$ satisfying $|\alpha| \leq L$, $\frac{1}{2} \text{tr}(\beta^2) \leq L$, there exists unique $\mathbb{P} \in \mathcal{P}^L$ such that $\alpha^{\mathbb{P}} = \alpha$, $\beta^{\mathbb{P}} = \beta$, \mathbb{P} -a.s., where $\alpha^{\mathbb{P}}$, $\beta^{\mathbb{P}}$ can be understood in the sense of either (2.2) or (2.3). ■

Throughout this paper, we shall consider a family $\{\mathcal{P}_t, t \in [0, T]\}$ of semimartingale measures on Ω^t satisfying:

(P1) *there exists some L_0 such that, for all t , \mathcal{P}_t is a weakly compact subset of $\mathcal{P}_t^{L_0}$.*

(P2) *For any $0 \leq t \leq T$, $\tau \in \mathcal{T}^t$, and $\mathbb{P} \in \mathcal{P}_t$, the regular conditional probability distribution $\mathbb{P}^{\tau, \omega} \in \mathcal{P}_{\tau(\omega)}$ for \mathbb{P} -a.e. $\omega \in \Omega^t$.*

(P3) *For any $0 \leq s \leq t \leq T$, $\mathbb{P} \in \mathcal{P}_s$, $\{E_i, i \geq 1\} \subset \mathcal{F}_t^s$ disjoint, and $\mathbb{P}^i \in \mathcal{P}_t$, the following $\hat{\mathbb{P}}$ is also in \mathcal{P}_s :*

$$\hat{\mathbb{P}} := \mathbb{P} \otimes_t \left[\sum_{i=1}^{\infty} \mathbb{P}^i \mathbf{1}_{E_i} + \mathbb{P} \mathbf{1}_{\cap_{i=1}^{\infty} E_i^c} \right]. \quad (2.4)$$

Here (2.4) means, for any event $E \in \mathcal{F}_T^s$ and denoting $E^{t, \omega} := \{\omega' \in \Omega^t : \omega \otimes_t \omega' \in E\}$:

$$\hat{\mathbb{P}}[E] := \mathbb{E}^{\mathbb{P}} \left[\sum_{i=1}^{\infty} \mathbb{P}^i[E^{t, B}] \mathbf{1}_{E_i}(B) \right] + \mathbb{P}[E \cap (\cap_{i=1}^{\infty} E_i^c)].$$

We refer to the seminal work of Stroock and Varadhan [18] for the introduction of regular conditional probability distribution (r.c.p.d. for short). See also Appendix A.1, in particular (A.2) below for the precise meaning of $\mathbb{P}^{\tau, \omega}$.

Remark 2.2. (i) The weak compactness of (P1) is crucial for the existence of the optimal stopping time. As explained in Introduction, the major technical difficulty we face is the failure of the dominated convergence theorem in our nonlinear expectation framework. To overcome this, we shall use the regularity of the processes and the weak compactness of the classes \mathcal{P}_t . See e.g. Step 2 in Section 4.4.

(ii) The regular conditional probability distribution is a convenient tool for proving the dynamic programming principle, see e.g. Soner, Touzi, and Zhang [16]. In particular, (P2) is used to prove one inequality in the dynamic programming principle, see e.g. Step 1 in the proof of Lemma 4.1.

(iii) The concatenation Property (P3) is used to prove the opposite inequality in the dynamic programming principle, see e.g. Step 2 in the proof of Lemma 4.1. We remark that this condition can be weakened by using the more abstract framework in Nutz and van Handel [11]. ■

We first observe that

Lemma 2.3. For all $L > 0$, the family $\{\mathcal{P}_t^L, t \in [0, T]\}$ satisfies conditions (P1)–(P3).

The proof is quite straightforward, by using the definition of r.c.p.d. We nevertheless provide a proof in the [Appendix](#).

The following are some other typical examples of such a family $\{\mathcal{P}_t, t \in [0, T]\}$. Their Properties (P1)–(P3) can be checked similarly.

Example 2.4. Let $L, L_1, L_2 > 0$ be some constants.

Wiener measure $\mathcal{P}_t^0 := \{\mathbb{P}_t^0\} = \{\mathbb{P} : \alpha^\mathbb{P} = 0, \beta^\mathbb{P} = I_d\}$.

Finite variation $\mathcal{P}_t^{\text{FV}}(L) := \{\mathbb{P} : |\alpha^\mathbb{P}| \leq L, \beta^\mathbb{P} = 0\}$.

Drifted Wiener measure $\mathcal{P}_t^{\text{ac}}(L) := \{\mathbb{P} : |\alpha^\mathbb{P}| \leq L, \beta^\mathbb{P} = I_d\}$.

Relaxed bounds $\mathcal{P}_t(L_1, L_2) := \{\mathbb{P} : |\alpha^\mathbb{P}| \leq L_1, 0 \leq \beta^\mathbb{P} \leq L_2 I_d\}$.

Relaxed bounds, Uniformly elliptic $\mathcal{P}_t^{\text{UE}}(L_1, L_2, L) := \{\mathbb{P} : |\alpha^\mathbb{P}| \leq L_1, L I_d \leq \beta^\mathbb{P} \leq L_2 I_d\}$.

Equivalent martingale measures $\mathcal{P}_t^e(L_1, L_2, L) := \{\mathbb{P} \in \mathcal{P}_t(L_1, L_2) : \exists |\gamma^\mathbb{P}| \leq L, \alpha^\mathbb{P} = \beta^\mathbb{P} \gamma^\mathbb{P}\}$.

We denote by $\mathbb{L}^1(\mathcal{F}_T^t, \mathcal{P}_t)$ the set of all $\xi \in \mathbb{L}^0(\mathcal{F}_T^t)$ with $\sup_{\mathbb{P} \in \mathcal{P}_t} \mathbb{E}^\mathbb{P}[|\xi|] < \infty$. The set \mathcal{P}_t induces the following capacity and nonlinear expectation:

$$C_t[A] := \sup_{\mathbb{P} \in \mathcal{P}_t} \mathbb{P}[A] \quad \text{for } A \in \mathcal{F}_T^t, \quad \text{and} \quad \mathcal{E}_t[\xi] := \sup_{\mathbb{P} \in \mathcal{P}_t} \mathbb{E}^\mathbb{P}[\xi]$$

$$\text{for } \xi \in \mathbb{L}^1(\mathcal{F}_T^t, \mathcal{P}_t). \quad (2.5)$$

When $t = 0$, we shall omit t and abbreviate them as $\mathcal{P}, \mathcal{C}, \mathcal{E}$. Clearly \mathcal{E} is a G -expectation, in the sense of Peng [13]. We remark that, when ξ satisfies certain regularity condition, then $\mathcal{E}_t[\xi^{t, \omega}]$ can be viewed as the conditional G -expectation of ξ , and as a process it is the solution of a Second Order BSDE, as introduced by Soner, Touzi and Zhang [15].

We remark that the last three families of measures in [Example 2.4](#) are non-dominated, which are most interesting to us. In particular, in these cases the dominated convergence theorem fails under the corresponding nonlinear expectation as we see in the following simple example.

Example 2.5. Consider the relaxed bounds $\mathcal{P}_t(L_1, L_2)$ in [Example 2.4](#) with $d = 1$. Let

$$\xi_n := \mathbf{1}_{\{0 < \langle B \rangle_T \leq \frac{1}{n}\}},$$

where $\langle B \rangle$ is the pathwise quadratic variation. Then $\xi_n \downarrow 0$ for all ω as $n \rightarrow \infty$, but $\mathcal{E}_0[\xi_n] = 1$ for all $n \geq \frac{1}{2L_2}$. ■

Given a family of probability measures \mathcal{P} on Ω , abusing the terminology of Denis and Martini [4], we say that a property holds \mathcal{P} -q.s. (quasi-surely) if it holds \mathbb{P} -a.s. for all $\mathbb{P} \in \mathcal{P}$. Moreover, a random variable $\xi : \Omega \rightarrow \mathbb{R}$ is

- \mathcal{P} -quasi-continuous if for any $\varepsilon > 0$, there exists a closed set $\Omega_\varepsilon \subset \Omega$ such that $\mathcal{C}(\Omega_\varepsilon^c) < \varepsilon$ and ξ is continuous in Ω_ε ,
- \mathcal{P} -uniformly integrable if $\mathcal{E}[|\xi|] \mathbf{1}_{\{|\xi| \geq n\}} \rightarrow 0$, as $n \rightarrow \infty$.

Since \mathcal{P} is weakly compact, by Denis, Hu and Peng [3, Lemma 4 and Theorems 22, 28], we have:

Proposition 2.6. (i) Let $(\Omega_n)_{n \geq 1}$ be a sequence of open sets with $\Omega_n \uparrow \Omega$. Then $\mathcal{C}(\Omega_n^c) \downarrow 0$.
(ii) Let $(\xi_n)_{n \geq 1}$ be a sequence of \mathcal{P} -quasi-continuous and \mathcal{P} -uniformly integrable maps from Ω to \mathbb{R} . If $\xi_n \downarrow \xi$, \mathcal{P} -q.s. then $\mathcal{E}[\xi_n] \downarrow \mathcal{E}[\xi]$.

We finally recall the notion of martingales under nonlinear expectation.

Definition 2.7. Let $X \in \mathbb{H}^0(\mathbb{F})$ such that $X_\tau \in \mathbb{L}^1(\mathcal{F}_\tau, \mathcal{P})$ for all $\tau \in \mathcal{T}$. We say that X is a \mathcal{E} -supermartingale (resp. submartingale, martingale) if, for any $(t, \omega) \in \Lambda$ and any $\tau \in \mathcal{T}^t$, $\mathcal{E}_t[X_\tau^{t, \omega}] \leq$ (resp. $\geq, =$) $X_t(\omega)$ for \mathcal{P} -q.s. $\omega \in \Omega$.

We remark that we require the \mathcal{E} -supermartingale property holds for stopping times. Under linear expectation $\mathbb{E}^\mathbb{P}$, this is equivalent to the \mathbb{P} -supermartingale property for deterministic times, due to the Doob's optional sampling theorem. However, under nonlinear expectation, they are in general not equivalent.

3. Optimal stopping under nonlinear expectations

We now fix a process $X \in \mathbb{H}^0(\mathbb{F})$.

Assumption 3.1. X is a bounded càdlàg process with positive jumps, and there exists a modulus of continuity function ρ_0 such that for any $(t, \omega), (t', \omega') \in \Lambda$:

$$X(t, \omega) - X(t', \omega') \leq \rho_0(\mathbf{d}_\infty((t, \omega), (t', \omega'))) \quad \text{whenever } t \leq t'. \quad (3.1)$$

Remark 3.2. There is some redundancy in the above assumption. Indeed, it is shown in the [Appendix](#) that (3.1) implies that X has left-limits and $X_{t-} \leq X_t$ for all $t \in (0, T]$. Moreover, the fact that X has only positive jumps is important to ensure that the random times τ^* in (3.2), $\hat{\tau}^*$ in (3.5), and τ_n in (4.7) and (5.15) are \mathbb{F} -stopping times. ■

We define the nonlinear Snell envelope and the corresponding obstacle first hitting time:

$$Y_t(\omega) := \sup_{\tau \in \mathcal{T}^t} \mathcal{E}_t[X_\tau^{t, \omega}], \quad \text{and} \quad \tau^* := \inf\{t \geq 0 : Y_t = X_t\}. \quad (3.2)$$

Our first result is the following *nonlinear Snell envelope* characterization of the deterministic maturity optimal stopping problem Y_0 .

Theorem 3.3 (Deterministic Maturity). Under [Assumption 3.1](#), the process Y is an \mathcal{E} -supermartingale on $[0, T]$, $Y_{\tau^*} = X_{\tau^*}$, and $Y_{\cdot \wedge \tau^*}$ is an \mathcal{E} -martingale. Consequently, τ^* is an optimal stopping time for the problem Y_0 .

To prove the partial comparison principle for viscosity solutions of path-dependent partial differential equations in our accompanying paper [7], we need to consider optimal stopping problems with random maturity time $H \in \mathcal{T}$ of the form

$$H := \inf\{t \geq 0 : B_t \in O^c\} \wedge t_0, \quad (3.3)$$

for some $t_0 \in (0, T]$ and some open convex set $O \subset \mathbb{R}^d$ containing the origin. We shall extend the previous result to the following stopped process:

$$\hat{X}_s^H := X_s \mathbf{1}_{\{s < H\}} + X_H \mathbf{1}_{\{s \geq H\}} \quad \text{for } s \in [0, T]. \quad (3.4)$$

The corresponding Snell envelope and obstacle first hitting time are denoted:

$$\hat{Y}_t^H(\omega) := \sup_{\tau \in \mathcal{T}^t} \mathcal{E}_t[(\hat{X}_\tau^H)^{t, \omega}], \quad \text{and} \quad \hat{\tau}^* := \inf\{t \geq 0 : \hat{Y}_t^H = \hat{X}_t^H\}. \quad (3.5)$$

Our second main result requires the following additional assumption.

- Assumption 3.4.** (i) For some $L > 0$, $\mathcal{P}_t^{\text{FV}}(L) \subset \mathcal{P}_t$ for all $t \in [0, T]$, where $\mathcal{P}_t^{\text{FV}}(L)$ is defined in [Example 2.4](#).
 (ii) For any $0 \leq t < t + \delta \leq T$, $\mathcal{P}_t \subset \mathcal{P}_{t+\delta}$ in the following sense: for any $\mathbb{P} \in \mathcal{P}_t$ we have $\tilde{\mathbb{P}} \in \mathcal{P}_{t+\delta}$, where $\tilde{\mathbb{P}}$ is the probability measure on $\Omega^{t+\delta}$ such that the $\tilde{\mathbb{P}}$ -distribution of $B^{t+\delta}$ is equal to the \mathbb{P} -distribution of $\{B_s^t, t \leq s \leq T - \delta\}$.

Remark 3.5. The above assumption is a technical condition used to prove the dynamic programming principle in [Section 5.1](#) below.

- (i) All sets in [Example 2.4](#) satisfy [Assumption 3.4\(ii\)](#), and the relaxed bounds $\mathcal{P}_t(L_1, L_2)$ satisfies [Assumption 3.4\(i\)](#). We remark that, for the viscosity theory of path-dependent partial differential equations in our accompanying papers [\[6,7\]](#), we shall use $\mathcal{P}_t(L, \sqrt{2}L)$ which satisfies both (i) and (ii) of [Assumption 3.4](#).
 (ii) By a little more involved arguments, we may prove the results in [Section 5.1](#) by replacing [Assumption 3.4](#) (i) with: for $\mathcal{P}_t^{\text{UE}}$ defined in [Example 2.4](#),

$$\text{for some constants } L, L_1, L_2, \quad \mathcal{P}_t^{\text{UE}}(L_1, L_2, L) \subset \mathcal{P}_t \quad \text{for all } t \in [0, T], \quad (3.6)$$

- (iii) If \mathcal{P}_t is uniformly nondegenerate, namely

$$\text{there exists } c > 0 \text{ such that } \beta^{\mathbb{P}} \geq cI_d \quad \text{for all } t \text{ and } \mathbb{P} \in \mathcal{P}_t, \quad (3.7)$$

then we shall use [\(3.6\)](#) instead of [Assumption 3.4](#) (i). In this case, under the additional condition that X is uniformly continuous in (t, ω) , \hat{Y}^H is left continuous at H and the arguments for our main result [Theorem 3.6](#) below can be simplified significantly, see [Lemma A.1](#) and [Remark 5.10](#) below. ■

Theorem 3.6 (*Random Maturity*). Under [Assumptions 3.1](#) and [3.4](#), the process \hat{Y}^H is an \mathcal{E} -supermartingale on $[0, H]$, $\hat{Y}_{\hat{\tau}^*}^H = \hat{X}_{\hat{\tau}^*}^H$, and $\hat{Y}_{\cdot \wedge \hat{\tau}^*}^H$ is an \mathcal{E} -martingale. In particular, $\hat{\tau}^*$ is an optimal stopping time for the problem \hat{Y}_0^H .

Remark 3.7. The main idea for proving [Theorem 3.6](#) is to show that $\mathcal{E}[\hat{Y}_{\tau_n}^H]$ converges to $\mathcal{E}[\hat{Y}_{\hat{\tau}^*}^H]$, where τ_n is defined by [\(5.15\)](#) below and increases to $\hat{\tau}^*$. However, we face a major difficulty that the dominated convergence theorem fails in our nonlinear expectation framework. Notice that Y is an \mathcal{E} -supermartingale and thus Y_{τ_n} are decreasing under expectation (but not pointwise!). We shall extend the arguments of [\[3\]](#) for the monotone convergence theorem, [Proposition 2.6](#), to our case. For this purpose, we need to construct certain continuous approximations of the stopping times τ_n , and the requirement that the random maturity H is of the form [\(3.3\)](#) is crucial. We remark that, in his Markov model, Krylov [\[10\]](#) also considers this type of hitting times. We also remark that, in a special case, Song [\[17\]](#) proved that H is quasi-continuous. ■

4. Deterministic maturity optimal stopping

We now prove [Theorem 3.3](#). Throughout this section, [Assumption 3.1](#) is always in force, and we consider the nonlinear Snell envelope Y together with the first obstacle hitting time τ^* , as defined in [\(3.2\)](#). Assume $|X| \leq C_0$, and without loss of generality that $\rho_0 \leq 2C_0$. It is obvious that

$$|Y| \leq C_0, \quad Y \geq X, \quad \text{and} \quad Y_T = X_T. \quad (4.1)$$

Throughout this section, we shall use the following modulus of continuity function:

$$\bar{\rho}_0(\delta) := \rho_0(\delta) \vee \left[\rho_0(\delta^{\frac{1}{3}}) + \delta^{\frac{1}{3}} \right], \quad (4.2)$$

and we shall use a generic constant C which depends only on C_0, T, d , and the L_0 in Property (P1), and it may vary from line to line.

4.1. Dynamic programming principle

Similar to the standard Snell envelope characterization under linear expectation, our first step is to establish the dynamic programming principle. We start by the case of deterministic times.

Lemma 4.1. *For each t , the random variable Y_t is uniformly continuous in ω , with the modulus of continuity function ρ_0 , and satisfies*

$$Y_{t_1}(\omega) = \sup_{\tau \in T^1} \mathcal{E}_{t_1} \left[X_{\tau}^{t_1, \omega} \mathbf{1}_{\{\tau < t_2\}} + Y_{t_2}^{t_1, \omega} \mathbf{1}_{\{\tau \geq t_2\}} \right] \quad \text{for all } 0 \leq t_1 \leq t_2 \leq T, \omega \in \Omega. \quad (4.3)$$

Proof. (i) First, for any t , any $\omega, \omega' \in \Omega$, and any $\tau \in T^t$, by (3.1) we have

$$\begin{aligned} |X_{\tau}^{t, \omega} - X_{\tau}^{t, \omega'}| &= |X(\tau(B^t), \omega \otimes_t B^t) - X(\tau(B^t), \omega' \otimes_t B^t)| \\ &\leq \rho_0 \left(d_{\infty}((\tau(B^t), \omega \otimes_t B^t), (\tau(B^t), \omega' \otimes_t B^t)) \right) = \rho_0(\|\omega - \omega'\|_t). \end{aligned}$$

Since τ is arbitrary, this proves uniform continuity of Y_t in ω .

(ii) When $t_2 = T$, since $Y_T = X_T$ (4.3) coincides with the definition of Y . Without loss of generality we assume $(t_1, \omega) = (0, \mathbf{0})$ and $t := t_2 < T$. Recall that we omit the subscript 0.

Step 1. We first prove “ \leq ”. For any $\tau \in T$ and $\mathbb{P} \in \mathcal{P}$:

$$\mathbb{E}^{\mathbb{P}}[X_{\tau}] = \mathbb{E}^{\mathbb{P}}[X_{\tau} \mathbf{1}_{\{\tau < t\}} + \mathbb{E}_t^{\mathbb{P}}[X_{\tau}] \mathbf{1}_{\{\tau \geq t\}}].$$

By the definition of the regular conditional probability distribution, we have $\mathbb{E}_t^{\mathbb{P}}[X_{\tau}](\omega) = \mathbb{E}^{\mathbb{P}^{t, \omega}}[X_{\tau}^{t, \omega}] \leq Y_t(\omega)$ for \mathbb{P} -a.e. $\omega \in \{\tau \geq t\}$, where the inequality follows from Property (P2) of the family $\{\mathcal{P}_t\}$ that $\mathbb{P}^{t, \omega} \in \mathcal{P}_t$. Then:

$$\mathbb{E}^{\mathbb{P}}[X_{\tau}] \leq \mathbb{E}^{\mathbb{P}}[X_{\tau} \mathbf{1}_{\{\tau < t\}} + Y_t \mathbf{1}_{\{\tau \geq t\}}].$$

By taking the sup over τ and \mathbb{P} , it follows that:

$$Y_0 = \sup_{\tau \in T} \mathcal{E}[X_{\tau}] \leq \sup_{\tau \in T} \mathcal{E}[X_{\tau} \mathbf{1}_{\{\tau < t\}} + Y_t \mathbf{1}_{\{\tau \geq t\}}].$$

Step 2. We next prove “ \geq ”. Fix arbitrary $\tau \in T$ and $\mathbb{P} \in \mathcal{P}$, we shall prove

$$\mathbb{E}^{\mathbb{P}}[X_{\tau} \mathbf{1}_{\{\tau < t\}} + Y_t \mathbf{1}_{\{\tau \geq t\}}] \leq Y_0. \quad (4.4)$$

Let $\varepsilon > 0$, and $\{E_i\}_{i \geq 1}$ be an \mathcal{F}_t -measurable partition of the event $\{\tau \geq t\} \in \mathcal{F}_t$ such that $\|\omega - \tilde{\omega}\|_t \leq \varepsilon$ for all $\omega, \tilde{\omega} \in E_i$. For each i , fix an $\omega^i \in E_i$, and by the definition of Y we have

$$Y_t(\omega^i) \leq \mathbb{E}^{\mathbb{P}^i}[X_{\tau}^{t, \omega^i}] + \varepsilon \quad \text{for some } (\tau^i, \mathbb{P}^i) \in T^t \times \mathcal{P}_t.$$

By (3.1) and the uniform continuity of Y , proved in (i), we have

$$|Y_t(\omega) - Y_t(\omega^i)| \leq \rho_0(\varepsilon), \quad |X_{\tau}^{t, \omega} - X_{\tau}^{t, \omega^i}| \leq \rho_0(\varepsilon), \quad \text{for all } \omega \in E_i.$$

Thus, for $\omega \in E_i$,

$$Y_t(\omega) \leq Y_t(\omega^i) + \rho_0(\varepsilon) \leq \mathbb{E}^{\mathbb{P}^i}[X_{\tau^i}^{t, \omega^i}] + \varepsilon + \rho_0(\varepsilon) \leq \mathbb{E}^{\mathbb{P}^i}[X_{\tau^i}^{t, \omega}] + \varepsilon + 2\rho_0(\varepsilon). \quad (4.5)$$

Thanks to Property (P3) of the family $\{\mathcal{P}_t\}$, we may define the following pair $(\tilde{\tau}, \tilde{\mathbb{P}}) \in \mathcal{T} \times \mathcal{P}$:

$$\tilde{\tau} := \mathbf{1}_{\{\tau < t\}}\tau + \mathbf{1}_{\{\tau \geq t\}} \sum_{i \geq 1} \mathbf{1}_{E_i} \tau^i(B^i); \quad \tilde{\mathbb{P}} := \mathbb{P} \otimes_t \left[\sum_{i \geq 1} \mathbf{1}_{E_i} \mathbb{P}^i + \mathbf{1}_{\{\tau < t\}} \mathbb{P} \right].$$

It is obvious that $\{\tau < t\} = \{\tilde{\tau} < t\}$. Then, by (4.5),

$$\begin{aligned} \mathbb{E}^{\mathbb{P}}[X_{\tau} \mathbf{1}_{\{\tau < t\}} + Y_t \mathbf{1}_{\{\tau \geq t\}}] &= \mathbb{E}^{\mathbb{P}} \left[X_{\tau} \mathbf{1}_{\{\tau < t\}} + \sum_{i \geq 1} Y_t \mathbf{1}_{E_i} \right] \\ &\leq \mathbb{E}^{\mathbb{P}} \left[X_{\tau} \mathbf{1}_{\{\tau < t\}} + \sum_{i \geq 1} \mathbb{E}^{\mathbb{P}^i}[X_{\tau^i}^{t, \cdot}] \mathbf{1}_{E_i} \right] + \varepsilon + 2\rho_0(\varepsilon) \\ &= \mathbb{E}^{\tilde{\mathbb{P}}} \left[X_{\tilde{\tau}} \mathbf{1}_{\{\tilde{\tau} < t\}} + \sum_{i \geq 1} X_{\tilde{\tau}} \mathbf{1}_{E_i} \right] + \varepsilon + 2\rho_0(\varepsilon) \\ &= \mathbb{E}^{\tilde{\mathbb{P}}}[X_{\tilde{\tau}}] + \varepsilon + 2\rho_0(\varepsilon) \leq Y_0 + \varepsilon + 2\rho_0(\varepsilon), \end{aligned}$$

which provides (4.4) by sending $\varepsilon \rightarrow 0$. ■

We now derive the regularity of Y in t .

Lemma 4.2. For each $\omega \in \Omega$ and $0 \leq t_1 < t_2 \leq T$,

$$|Y_{t_1}(\omega) - Y_{t_2}(\omega)| \leq C \bar{\rho}_0(\mathbf{d}_{\infty}((t_1, \omega), (t_2, \omega))).$$

Proof. Denote $\delta := \mathbf{d}_{\infty}((t_1, \omega), (t_2, \omega))$. If $\delta \geq \frac{1}{8}$, then clearly $|Y_{t_1}(\omega) - Y_{t_2}(\omega)| \leq 2C_0 \leq C \bar{\rho}_0(\delta)$. So we continue the proof assuming $\delta \leq \frac{1}{8}$. First, by setting $\tau = t_2$ in Lemma 4.1,

$$\begin{aligned} \delta Y &:= Y_{t_2}(\omega) - Y_{t_1}(\omega) \leq Y_{t_2}(\omega) - \mathcal{E}_{t_1}[Y_{t_2}^{t_1, \omega}] \\ &\leq \mathcal{E}_{t_1}[Y_{t_2}(\omega) - Y_{t_2}(\omega \otimes_{t_1} B^{t_1})] \\ &\leq \mathcal{E}_{t_1}[\rho_0(\mathbf{d}_{\infty}((t_2, \omega), (t_2, \omega \otimes_{t_1} B^{t_1})))] \\ &\leq \mathcal{E}_{t_1}[\rho_0(\delta + \|B^{t_1}\|_{t_1+\delta})]. \end{aligned}$$

On the other hand, by the inequality $X \leq Y$, Lemma 4.1, and (3.1), we have

$$\begin{aligned} -\delta Y &\leq \sup_{\tau \in \mathcal{T}^{t_1}} \mathcal{E}_{t_1} \left[[X_{t_2}^{t_1, \omega} + \rho_0(\mathbf{d}_{\infty}((\tau, \omega \otimes_{t_1} B^{t_1}), (t_2, \omega \otimes_{t_1} B^{t_1})))] \mathbf{1}_{\{\tau < t_2\}} \right. \\ &\quad \left. + Y_{t_2}^{t_1, \omega} \mathbf{1}_{\{\tau \geq t_2\}} \right] - Y_{t_2}(\omega) \\ &\leq \mathcal{E}_{t_1} [Y_{t_2}^{t_1, \omega} - Y_{t_2}(\omega) + \rho_0(\mathbf{d}_{\infty}((t_1, \omega), (t_2, \omega \otimes_{t_1} B^{t_1})))] \end{aligned}$$

$$\begin{aligned} &\leq \mathcal{E}_{t_1} \left[\rho_0(\mathbf{d}_\infty((t_2, \omega), (t_2, \omega \otimes_{t_1} B^{t_1}))) + \rho_0(\mathbf{d}_\infty((t_1, \omega), (t_2, \omega \otimes_{t_1} B^{t_1}))) \right] \\ &\leq 2\mathcal{E}_{t_1} [\rho_0(\delta + \|B^{t_1}\|_{t_1+\delta})]. \end{aligned}$$

Hence

$$|\delta Y| \leq 2\mathcal{E}_{t_1} [\rho_0(\delta + \|B^{t_1}\|_{t_1+\delta})] \leq \mathcal{E}_{t_1} \left[\rho_0 \left(\delta + \frac{3}{4}\delta^{\frac{1}{3}} \right) + 2C_0 \mathbf{1}_{\{\|B^{t_1}\|_{t_1+\delta} \geq \frac{3}{4}\delta^{\frac{1}{3}}\}} \right].$$

Since $\delta + \frac{3}{4}\delta^{\frac{1}{3}} \leq \delta^{\frac{1}{3}}$ for $\delta \leq \frac{1}{8}$, this provides:

$$|\delta Y| \leq \rho_0(\delta^{\frac{1}{3}}) + C\delta^{-\frac{2}{3}} \mathcal{E}_{t_1} [\|B^{t_1}\|_{t_1+\delta}^2] \leq \rho_0(\delta^{\frac{1}{3}}) + C\delta^{-\frac{2}{3}}\delta \leq C\bar{\rho}_0(\delta). \quad \blacksquare \quad (4.6)$$

We are now ready to prove the dynamic programming principle for stopping times.

Theorem 4.3. For any $(t, \omega) \in \Lambda$ and $\tau \in T^t$, we have

$$Y_t(\omega) = \sup_{\tilde{\tau} \in T^t} \mathcal{E}_t \left[X_{\tilde{\tau}}^{t,\omega} \mathbf{1}_{\{\tilde{\tau} < \tau\}} + Y_{\tilde{\tau}}^{t,\omega} \mathbf{1}_{\{\tilde{\tau} \geq \tau\}} \right].$$

Consequently, Y is an \mathcal{E} -supermartingale on $[0, T]$.

Proof. First, follow the arguments in Lemma 4.1(ii) Step 1 and note that Property (P2) of the family $\{\mathcal{P}_t\}$ holds for stopping times, one can prove straightforwardly that

$$Y_t(\omega) \leq \sup_{\tilde{\tau} \in T^t} \mathcal{E}_t \left[X_{\tilde{\tau}}^{t,\omega} \mathbf{1}_{\{\tilde{\tau} < \tau\}} + Y_{\tilde{\tau}}^{t,\omega} \mathbf{1}_{\{\tilde{\tau} \geq \tau\}} \right].$$

On the other hand, let $\tau_k \downarrow \tau$ such that τ_k takes only finitely many values. By Lemma 4.1 one can easily show that Theorem 4.3 holds for τ_k . Then for any $\mathbb{P} \in \mathcal{P}_t$ and $\tilde{\tau} \in T^t$, by denoting $\tilde{\tau}_m := [\tilde{\tau} + \frac{1}{m}] \wedge T$ we have

$$\mathbb{E}^{\mathbb{P}} \left[X_{\tilde{\tau}_m}^{t,\omega} \mathbf{1}_{\{\tilde{\tau}_m < \tau_k\}} + Y_{\tau_k}^{t,\omega} \mathbf{1}_{\{\tilde{\tau}_m \geq \tau_k\}} \right] \leq Y_t(\omega).$$

Sending $k \rightarrow \infty$, by Lemma 4.2 and the dominated convergence theorem (under \mathbb{P}):

$$\mathbb{E}^{\mathbb{P}} \left[X_{\tilde{\tau}_m}^{t,\omega} \mathbf{1}_{\{\tilde{\tau}_m \leq \tau\}} + Y_{\tau}^{t,\omega} \mathbf{1}_{\{\tilde{\tau}_m > \tau\}} \right] \leq Y_t(\omega).$$

Since the process X is right continuous in t , we obtain by sending $m \rightarrow \infty$:

$$Y_t(\omega) \geq \mathbb{E}^{\mathbb{P}} \left[X_{\tilde{\tau}}^{t,\omega} \mathbf{1}_{\{\tilde{\tau} < \tau\}} + Y_{\tau}^{t,\omega} \mathbf{1}_{\{\tilde{\tau} \geq \tau\}} \right],$$

which provides the required result by the arbitrariness of \mathbb{P} and $\tilde{\tau}$. \blacksquare

4.2. Preparation for the \mathcal{E} -martingale property

If $Y_0 = X_0$, then $\tau^* = 0$ and obviously all the statements of Theorem 3.3 hold true. Therefore, we focus on the non-trivial case $Y_0 > X_0$.

We continue following the proof of the Snell envelope characterization in the standard linear expectation context. Let

$$\tau_n := \inf \left\{ t \geq 0 : Y_t - X_t \leq \frac{1}{n} \right\} \wedge T, \quad \text{for } n > (Y_0 - X_0)^{-1}. \quad (4.7)$$

Lemma 4.4. *The process Y is an \mathcal{E} -martingale on $[0, \tau_n]$.*

Proof. By the dynamic programming principle of [Theorem 4.3](#),

$$Y_0 = \sup_{\tau \in \mathcal{T}} \mathcal{E} \left[X_\tau \mathbf{1}_{\{\tau < \tau_n\}} + Y_{\tau_n} \mathbf{1}_{\{\tau \geq \tau_n\}} \right].$$

For any $\varepsilon > 0$, there exist $\tau_\varepsilon \in \mathcal{T}$ and $\mathbb{P}_\varepsilon \in \mathcal{P}$ such that

$$Y_0 \leq \mathbb{E}^{\mathbb{P}_\varepsilon} \left[X_{\tau_\varepsilon} \mathbf{1}_{\{\tau_\varepsilon < \tau_n\}} + Y_{\tau_n} \mathbf{1}_{\{\tau_\varepsilon \geq \tau_n\}} \right] + \varepsilon \leq \mathbb{E}^{\mathbb{P}_\varepsilon} \left[Y_{\tau_\varepsilon \wedge \tau_n} - \frac{1}{n} \mathbf{1}_{\{\tau_\varepsilon < \tau_n\}} \right] + \varepsilon, \quad (4.8)$$

where we used the fact that $Y_t - X_t > \frac{1}{n}$ for $t < \tau_n$, by the definition of τ_n . On the other hand, it follows from the \mathcal{E} -supermartingale property of Y in [Theorem 4.3](#) that $\mathbb{E}^{\mathbb{P}_\varepsilon} \left[Y_{\tau_\varepsilon \wedge \tau_n} \right] \leq \mathcal{E}[Y_{\tau_\varepsilon \wedge \tau_n}] \leq Y_0$, which implies by (4.8) that $\mathbb{P}_\varepsilon[\tau_\varepsilon < \tau_n] \leq n\varepsilon$. We then get from (4.8) that:

$$\begin{aligned} Y_0 &\leq \mathbb{E}^{\mathbb{P}_\varepsilon} \left[(X_{\tau_\varepsilon} - Y_{\tau_n}) \mathbf{1}_{\{\tau_\varepsilon < \tau_n\}} + Y_{\tau_n} \right] + \varepsilon \leq C\mathbb{P}_\varepsilon[\tau_\varepsilon < \tau_n] + \mathbb{E}^{\mathbb{P}_\varepsilon}[Y_{\tau_n}] + \varepsilon \\ &\leq \mathcal{E}[Y_{\tau_n}] + (Cn + 1)\varepsilon. \end{aligned}$$

Since ε is arbitrary, we obtain $Y_0 \leq \mathcal{E}[Y_{\tau_n}]$. Similarly one can prove Y is an \mathcal{E} -submartingale on $[0, \tau_n]$. By the \mathcal{E} -supermartingale property of Y established in [Theorem 4.3](#), this implies that Y is an \mathcal{E} -martingale on $[0, \tau_n]$. ■

By [Lemma 4.2](#) we have

$$Y_0 - \mathcal{E}[Y_{\tau^*}] = \mathcal{E}[Y_{\tau_n}] - \mathcal{E}[Y_{\tau^*}] \leq C\mathcal{E} \left[\bar{\rho}_0 \left(\mathbf{d}_\infty((\tau_n, \omega), (\tau^*, \omega)) \right) \right]. \quad (4.9)$$

Clearly, $\tau_n \nearrow \tau^*$, and $\bar{\rho}_0 \left(\mathbf{d}_\infty((\tau_n, \omega), (\tau^*, \omega)) \right) \searrow 0$. However, in general the stopping times τ_n, τ^* are not \mathcal{P} -quasi-continuous, so we cannot apply [Proposition 2.6\(ii\)](#) to conclude $Y_0 \leq \mathcal{E}[Y_{\tau^*}]$. To overcome this difficulty, we need to approximate τ_n by continuous random variables.

4.3. Continuous approximation

The following lemma is crucial for us.

Lemma 4.5. *Let $\underline{\theta} \leq \theta \leq \bar{\theta}$ be random variables on Ω , with values in a compact interval $I \subset \mathbb{R}$, such that for some $\Omega_0 \subset \Omega$ and $\delta > 0$:*

$$\underline{\theta}(\omega) \leq \theta(\omega') \leq \bar{\theta}(\omega) \quad \text{for all } \omega \in \Omega_0 \quad \text{and} \quad \|\omega - \omega'\| \leq \delta.$$

Then for any $\varepsilon > 0$, there exists a uniformly continuous function $\hat{\theta} : \Omega \rightarrow I$ and an open subset $\Omega_\varepsilon \subset \Omega$ such that

$$C[\Omega_\varepsilon^c] \leq \varepsilon \quad \text{and} \quad \underline{\theta} - \varepsilon \leq \hat{\theta} \leq \bar{\theta} + \varepsilon \quad \text{in } \Omega_\varepsilon \cap \Omega_0.$$

Proof. If I is a single point set, then θ is a constant and the result is obviously true. Thus at below we assume the length $|I| > 0$. Let $\{\omega_j\}_{j \geq 1}$ be a dense sequence in Ω . Denote $O_j := \{\omega \in \Omega : \|\omega - \omega_j\| < \frac{\delta}{2}\}$ and $\Omega_n := \cup_{j=1}^n O_j$. It is clear that Ω_n is open and $\Omega_n \uparrow \Omega$ as $n \rightarrow \infty$. Let $f_n : [0, \infty) \rightarrow [0, 1]$ be defined as follows: $f_n(x) = 1$ for $x \in [0, \frac{\delta}{2}]$, $f_n(x) = \frac{1}{n^2|x|}$

for $x \geq \delta$, and f_n is linear in $[\frac{\delta}{2}, \delta]$. Define

$$\theta_n(\omega) := \phi_n(\omega) \sum_{j=1}^n \theta(\omega_j) \varphi_{n,j}(\omega) \quad \text{where } \varphi_{n,j}(\omega) := f_n(\|\omega - \omega_j\|)$$

$$\text{and } \phi_n := \left(\sum_{j=1}^n \varphi_{n,j} \right)^{-1}.$$

Then clearly θ_n is uniformly continuous and takes values in I . For each $\omega \in \Omega_n \cap \Omega_0$, the set $J_n(\omega) := \{1 \leq j \leq n : \|\omega - \omega_j\| \leq \delta\} \neq \emptyset$ and $\phi_n(\omega) \leq 1$. Then, by our assumption,

$$\begin{aligned} \theta_n(\omega) - \bar{\theta}(\omega) &= \phi_n(\omega) \left(\sum_{j \in J_n(\omega)} [\theta(\omega_j) - \bar{\theta}(\omega)] \varphi_{n,j}(\omega) + \sum_{j \notin J_n(\omega)} [\theta(\omega_j) - \bar{\theta}(\omega)] \varphi_{n,j}(\omega) \right) \\ &\leq \phi_n(\omega) \sum_{j \notin J_n(\omega)} |\theta(\omega_j) - \bar{\theta}(\omega)| \varphi_{n,j}(\omega) \leq \phi_n(\omega) \sum_{j \notin J_n(\omega)} \frac{1}{n^2} \leq \frac{1}{n}. \end{aligned}$$

Similarly one can show that $\underline{\theta} - \frac{1}{n} \leq \theta_n$ in $\Omega_n \cap \Omega_0$. Finally, since $\Omega_n \uparrow \Omega$ as $n \rightarrow \infty$, it follows from Proposition 2.6(i) that $\lim_{n \rightarrow \infty} \mathcal{C}[\Omega_n^c] = 0$. ■

4.4. Proof of Theorem 3.3

We proceed in two steps.

Step 1. For each n , let $\delta_n > 0$ be such that $3C\bar{\rho}_0(\delta_n) \leq \frac{1}{n(n+1)}$ for the constant C in Lemma 4.2. Now for any ω and ω' such that $\|\omega - \omega'\|_T \leq \delta_n$, by (3.1), the uniform continuity of Y in Lemma 4.1, and the fact that $\rho_0 \leq \bar{\rho}_0$, we have

$$(Y - X)_{\tau_{n+1}(\omega)}(\omega') \leq (Y - X)_{\tau_{n+1}(\omega)}(\omega) + 3C\bar{\rho}_0(\delta_n) \leq \frac{1}{n+1} + \frac{1}{n(n+1)} = \frac{1}{n}.$$

Then $\tau_n(\omega') \leq \tau_{n+1}(\omega)$. Since $3C\bar{\rho}_0(\delta_n) \leq \frac{1}{n(n+1)} \leq \frac{1}{n(n-1)}$, similarly we have $\tau_{n-1}(\omega) \leq \tau_n(\omega')$. We may then apply Lemma 4.5 with $\underline{\theta} = \tau_{n-1}$, $\theta = \tau_n$, $\bar{\theta} = \tau_{n+1}$, and $\Omega_0 = \Omega$. Thus, there exist an open set $\Omega_n \subset \Omega$ and a continuous random variable $\tilde{\tau}_n$ valued in $[0, T]$ such that

$$\mathcal{C}[\Omega_n^c] \leq 2^{-n} \quad \text{and} \quad \tau_{n-1} - 2^{-n} \leq \tilde{\tau}_n \leq \tau_{n+1} + 2^{-n} \quad \text{in } \Omega_n.$$

Step 2. By Lemma 4.4, for each n large, there exists $\mathbb{P}_n \in \mathcal{P}$ such that

$$Y_0 = \mathcal{E}[Y_{\tau_n}] \leq \mathbb{E}^{\mathbb{P}_n}[Y_{\tau_n}] + 2^{-n}.$$

By Property (P1), \mathcal{P} is weakly compact. Then, there exists a subsequence $\{n_j\}$ and $\mathbb{P}^* \in \mathcal{P}$ such that \mathbb{P}_{n_j} converges weakly to \mathbb{P}^* . Now for any n large and any $n_j \geq n$, note that $\tau_{n_j} \geq \tau_n$. Since Y is an \mathcal{E} -supermartingale and thus a \mathbb{P}_{n_j} -supermartingale, we have

$$Y_0 - 2^{-n_j} \leq \mathbb{E}^{\mathbb{P}_{n_j}}[Y_{\tau_{n_j}}] \leq \mathbb{E}^{\mathbb{P}_{n_j}}[Y_{\tau_n}] \leq \mathbb{E}^{\mathbb{P}_{n_j}}[Y_{\tilde{\tau}_n}] + \mathbb{E}^{\mathbb{P}_{n_j}}[|Y_{\tilde{\tau}_n} - Y_{\tau_n}|]. \quad (4.10)$$

By the boundedness of Y in (4.1) and the uniform continuity of Y in Lemma 4.2, we have

$$\begin{aligned} |Y_{\tilde{\tau}_n} - Y_{\tau_n}| &\leq C\bar{\rho}_0(\mathbf{d}_\infty((\tilde{\tau}_n, \omega), (\tau_n, \omega))) \\ &\leq C\bar{\rho}_0(\mathbf{d}_\infty((\tilde{\tau}_n, \omega), (\tau_n, \omega))) \mathbf{1}_{\Omega_{n-1} \cap \Omega_{n+1}} + C\mathbf{1}_{\Omega_{n-1}^c \cup \Omega_{n+1}^c}. \end{aligned}$$

Notice that $\tilde{\tau}_{n-1} - 2^{1-n} \leq \tau_n \leq \tilde{\tau}_{n+1} + 2^{-1-n}$ on $\Omega_{n-1} \cap \Omega_{n+1}$. Then

$$\begin{aligned} |Y_{\tilde{\tau}_n} - Y_{\tau_n}| &\leq C\bar{\rho}_0\left(\mathbf{d}_\infty((\tilde{\tau}_n, \omega), (\tilde{\tau}_{n-1} - 2^{1-n}, \omega))\right)\mathbf{1}_{\Omega_{n-1} \cap \Omega_{n+1}} \\ &\quad + C\bar{\rho}_0\left(\mathbf{d}_\infty((\tilde{\tau}_n, \omega), (\tilde{\tau}_{n+1} + 2^{-1-n}, \omega))\right)\mathbf{1}_{\Omega_{n-1} \cap \Omega_{n+1}} + C\mathbf{1}_{\Omega_{n-1}^c \cup \Omega_{n+1}^c} \\ &\leq C\bar{\rho}_0\left(\mathbf{d}_\infty((\tilde{\tau}_n, \omega), (\tilde{\tau}_{n-1} - 2^{1-n}, \omega))\right) \\ &\quad + C\bar{\rho}_0\left(\mathbf{d}_\infty((\tilde{\tau}_n, \omega), (\tilde{\tau}_{n+1} + 2^{-1-n}, \omega))\right) + C\mathbf{1}_{\Omega_{n-1}^c \cup \Omega_{n+1}^c}. \end{aligned}$$

Then (4.10) together with the estimate $\mathcal{C}[\Omega_n^c] \leq 2^{-n}$ lead to

$$\begin{aligned} Y_0 - 2^{-n_j} &\leq \mathbb{E}^{\mathbb{P}^{n_j}}[Y_{\tilde{\tau}_n}] + C\mathbb{E}^{\mathbb{P}^{n_j}}\left[\bar{\rho}_0\left(\mathbf{d}_\infty((\tilde{\tau}_n, \omega), (\tilde{\tau}_{n-1} - 2^{1-n}, \omega))\right)\right] \\ &\quad + C\mathbb{E}^{\mathbb{P}^{n_j}}\left[\bar{\rho}_0\left(\mathbf{d}_\infty((\tilde{\tau}_n, \omega), (\tilde{\tau}_{n+1} + 2^{-1-n}, \omega))\right)\right] + C2^{-n}. \end{aligned}$$

Notice that Y and $\tilde{\tau}_{n-1}, \tilde{\tau}_n, \tilde{\tau}_{n+1}$ are continuous. Send $j \rightarrow \infty$, we obtain

$$\begin{aligned} Y_0 &\leq \mathbb{E}^{\mathbb{P}^*}[Y_{\tilde{\tau}_n}] + C\mathbb{E}^{\mathbb{P}^*}\left[\bar{\rho}_0\left(\mathbf{d}_\infty((\tilde{\tau}_n, \omega), (\tilde{\tau}_{n-1} - 2^{1-n}, \omega))\right)\right] \\ &\quad + C\mathbb{E}^{\mathbb{P}^*}\left[\bar{\rho}_0\left(\mathbf{d}_\infty((\tilde{\tau}_n, \omega), (\tilde{\tau}_{n+1} - 2^{-1-n}, \omega))\right)\right] + C2^{-n}. \end{aligned} \quad (4.11)$$

Since $\sum_n \mathbb{P}^*[|\tilde{\tau}_n - \tau_n| \geq 2^{-n}] \leq \sum_n \mathcal{C}[|\tilde{\tau}_n - \tau_n| \geq 2^{-n}] \leq \sum_n 2^{-n} < \infty$ and $\tau_n \uparrow \tau^*$, by the Borel–Cantelli lemma under \mathbb{P}^* we see that $\tilde{\tau}_n \rightarrow \tau^*$, \mathbb{P}^* -a.s. Send $n \rightarrow \infty$ in (4.11) and apply the dominated convergence theorem under \mathbb{P}^* , we obtain

$$Y_0 \leq \mathbb{E}^{\mathbb{P}^*}[Y_{\tau^*}] \leq \mathcal{E}[Y_{\tau^*}].$$

Similarly $Y_t(\omega) \leq \mathcal{E}_t[Y_{\tau^*}^{\tau, \omega}]$ for $t < \tau^*(\omega)$. By the \mathcal{E} -supermartingale property of Y established in Theorem 4.3, this implies that Y is an \mathcal{E} -martingale on $[0, \tau^*]$. ■

5. Random maturity optimal stopping

In this section, we prove Theorem 3.6. The main idea follows that of Theorem 3.3. However, since \hat{X}_H is not continuous in ω , the estimates become much more involved.

Throughout this section, let $X, H, O, t_0, \hat{X} := \hat{X}^H, \hat{Y} := \hat{Y}^H$, and $\hat{\tau}^*$ be as in Theorem 3.6. Assumptions 3.1 and 3.4 will always be in force. We shall emphasize when the additional Assumption 3.4 is needed, and we fix the constant L as in Assumption 3.4 (i). Assume $|X| \leq C_0$, and without loss of generality that $\rho_0 \leq 2C_0$ and $L \leq 1$. It is clear that

$$|\hat{Y}| \leq C_0, \hat{X} \leq \hat{Y}, \quad \text{and} \quad \hat{Y}_H = \hat{X}_H = X_{H-}. \quad (5.1)$$

By (3.1) and the fact that X has positive jumps, one can check straightforwardly that,

$$\begin{aligned} \hat{X}(t, \omega) - \hat{X}(t', \omega') &\leq \rho_0(\mathbf{d}_\infty((t, \omega), (t', \omega'))) \quad \text{for } t \leq t', t \leq H(\omega), t' \leq H(\omega') \\ &\text{except the case } t = t' = H(\omega') < H(\omega) \leq t_0. \end{aligned} \quad (5.2)$$

In particular,

$$\hat{X}(t, \omega) - \hat{X}(t', \omega) \leq \rho_0(\mathbf{d}_\infty((t, \omega), (t', \omega))) \quad \text{whenever } t \leq t' \leq H(\omega). \quad (5.3)$$

Moreover, we define

$$\rho_1(\delta) := \rho_0(\delta) \vee \left[\rho_0((L^{-1}\delta)^{\frac{1}{3}}) + \delta^{\frac{1}{3}} \right], \quad \rho_2(\delta) := [\rho_1(\delta) + \delta] \vee [\rho_1(\delta^{\frac{1}{3}}) + \delta^{\frac{1}{3}}], \quad (5.4)$$

and in this section, the generic constant C may depend on L as well.

5.1. Dynamic programming principle

We start with the regularity in ω .

Lemma 5.1. *For any $t < H(\omega) \wedge H(\omega')$ we have:*

$$|\hat{Y}_t(\omega) - \hat{Y}_t(\omega')| \leq C\rho_1(\|\omega - \omega'\|_t).$$

To motivate our proof, we first follow the arguments in Lemma 4.1(i) and see why it does not work here. Indeed, note that

$$\hat{Y}_t(\omega) - \hat{Y}_t(\omega') \leq \sup_{\tau \in \mathcal{T}^t} \sup_{\mathbb{P} \in \mathcal{P}_t} \mathbb{E}^{\mathbb{P}} \left[\hat{X}_{\tau \wedge H^t, \omega}^{t, \omega} - \hat{X}_{\tau \wedge H^t, \omega'}^{t, \omega'} \right].$$

Since we do not have $H^t, \omega \leq H^t, \omega'$, we cannot apply (5.2) to obtain the required estimate.

Proof. Let $\tau \in \mathcal{T}^t$ and $\mathbb{P} \in \mathcal{P}_t$. Denote $\delta := \frac{1}{L}\|\omega - \omega'\|_t$, $t_\delta := [t + \delta] \wedge t_0$ and $\tilde{B}_s^{t_\delta} := B_{s+\delta}^t - B_{t_\delta}^t$ for $s \geq t$. Set $\tau'(B^t) := [\tau(\tilde{B}^{t_\delta}) + \delta] \wedge t_0$, then $\tau' \in \mathcal{T}^t$. Moreover, by Assumption 3.4 and Property (P3), we may choose $\mathbb{P}' \in \mathcal{P}_t$ defined as follows: $\alpha^{\mathbb{P}'} := \frac{1}{\delta}(\omega_t - \omega'_t)$, $\beta^{\mathbb{P}'} := \mathbf{0}$ on $[t, t_\delta]$, and the \mathbb{P}' -distribution of \tilde{B}^{t_δ} is equal to the \mathbb{P} -distribution of B^t . We claim that

$$I := \mathbb{E}^{\mathbb{P}}[\hat{X}_{\tau \wedge H^t, \omega}^{t, \omega}] - \mathbb{E}^{\mathbb{P}'}[\hat{X}_{\tau' \wedge H^t, \omega'}^{t, \omega'}] \leq C\rho_1(L\delta). \quad (5.5)$$

Then $\mathbb{E}^{\mathbb{P}}[\hat{X}_{\tau \wedge H^t, \omega}^{t, \omega}] - \hat{Y}_t(\omega') \leq \mathbb{E}^{\mathbb{P}}[\hat{X}_{\tau \wedge H^t, \omega}^{t, \omega}] - \mathbb{E}^{\mathbb{P}'}[\hat{X}_{\tau' \wedge H^t, \omega'}^{t, \omega'}] \leq C\rho_1(L\delta)$, and it follows from the arbitrariness of $\mathbb{P} \in \mathcal{P}_t$ and $\tau \in \mathcal{T}^t$ that $\hat{Y}_t(\omega) - \hat{Y}_t(\omega') \leq C\rho_1(L\delta)$. By exchanging the roles of ω and ω' , we obtain the required estimate.

It remains to prove (5.5). Denote

$$\tilde{\omega}'_s := \omega'_s \mathbf{1}_{[0, t)}(s) + [\omega'_t + \alpha^{\mathbb{P}'}(s - t)] \mathbf{1}_{[t, T]}(s).$$

Since $t < H(\omega) \wedge H(\omega')$, we have $\omega_t, \omega'_t \in O$. By the convexity of O , this implies that $\tilde{\omega}'_s \in O$ for $s \in [t, t_\delta]$, and thus $H^{t, \omega'}(B^t) = (H^{t, \omega}(\tilde{B}^{t_\delta}) + \delta) \wedge t_0$, \mathbb{P}' -a.s. Therefore,

$$\begin{aligned} \mathbb{E}^{\mathbb{P}'}[\hat{X}_{\tau' \wedge H^t, \omega'}^{t, \omega'}] &= \mathbb{E}^{\mathbb{P}'}\left[\hat{X}\left(\tau'(B^t) \wedge H^{t, \omega'}(B^t), \omega' \otimes_t B^t\right)\right] \\ &= \mathbb{E}^{\mathbb{P}'}\left[\hat{X}\left([\tau(\tilde{B}^{t_\delta}) + \delta] \wedge [H^{t, \omega}(\tilde{B}^{t_\delta}) + \delta] \wedge t_0, \tilde{\omega}' \otimes_{t_\delta} \tilde{B}_{\cdot - \delta}^{t_\delta}\right)\right] \\ &= \mathbb{E}^{\mathbb{P}}\left[\hat{X}\left([\tau(B^t) + \delta] \wedge [H^{t, \omega}(B^t) + \delta] \wedge t_0, \tilde{\omega}' \otimes_{t_\delta} B_{\cdot - \delta}^t\right)\right], \end{aligned} \quad (5.6)$$

while

$$\mathbb{E}^{\mathbb{P}}[\hat{X}_{\tau \wedge H^t, \omega}^{t, \omega}] = \mathbb{E}^{\mathbb{P}}\left[\hat{X}\left(\tau(B^t) \wedge H^{t, \omega}(B^t), \omega \otimes_t B^t\right)\right].$$

Notice that, whenever $\tau(B^t) \wedge H^{t, \omega}(B^t) = [\tau(B^t) + \delta] \wedge [H^{t, \omega}(B^t) + \delta] \wedge t_0$, we have $\tau(B^t) \wedge H^{t, \omega}(B^t) = t_0$. This excludes the exceptional case in (5.2). Then it follows from (5.6)

and (5.2) that

$$I \leq \mathbb{E}^{\mathbb{P}} \left[\rho_0 \left(\delta + \|(\omega \otimes_t B^t) \cdot \wedge \tau(B^t) \wedge H^{t,\omega}(B^t) - (\tilde{\omega}' \otimes_{t_\delta} B^t_{-\delta}) \cdot \wedge [\tau(B^t) + \delta] \wedge [H^{t,\omega}(B^t) + \delta] \wedge_{t_0} \|_{t_0} \right) \right].$$

Note that, denoting $\theta := \tau(B^t) \wedge H^{t,\omega}(B^t)$,

$$\begin{aligned} & \|(\omega \otimes_t B^t) \cdot \wedge \tau(B^t) \wedge H^{t,\omega}(B^t) - (\tilde{\omega}' \otimes_{t_\delta} B^t_{-\delta}) \cdot \wedge [\tau(B^t) + \delta] \wedge [H^{t,\omega}(B^t) + \delta] \wedge_{t_0} \|_{t_0} \\ & \leq \| \omega \otimes_t B^t - \tilde{\omega}' \otimes_{t_\delta} B^t_{-\delta} \|_{t_0} + \sup_{0 \leq r \leq \delta} |(\omega \otimes_t B^t)_{\theta+r} - (\omega \otimes_t B^t)_\theta| \\ & \leq \left[\|\omega - \omega'\|_t \right] \vee \left[\sup_{t \leq s \leq t_\delta} |\omega_t + B_s^t - \tilde{\omega}'_s| \right] \vee \left[\sup_{t_\delta \leq s \leq t_0} |\omega_t + B_s^t - \tilde{\omega}'_{t_\delta} - B_{s-\delta}^t| \right] \\ & \quad + \sup_{0 \leq r \leq \delta} |(\omega \otimes_t B^t)_{\theta+r} - (\omega \otimes_t B^t)_\theta| \\ & \leq 2L\delta + \|B^t\|_{t_\delta} + \sup_{t_\delta \leq s \leq t_0} |B_s^t - B_{s-\delta}^t| + \sup_{0 \leq r \leq \delta} |B_{\theta+r}^t - B_\theta^t|. \end{aligned}$$

Since $L \leq 1$, we have

$$I \leq \mathbb{E}^{\mathbb{P}} \left[\rho_0 \left(3\delta + \|B^t\|_{t_\delta} + \sup_{t_\delta \leq s \leq t_0} |B_s^t - B_{s-\delta}^t| + \sup_{0 \leq r \leq \delta} |B_{\theta+r}^t - B_\theta^t| \right) \right].$$

If $\delta \geq \frac{1}{8}$, then $I \leq 2C_0 \leq C\rho_1(L\delta)$. We then continue assuming $\delta \leq \frac{1}{8}$, and thus $3\delta + \frac{1}{4}\delta^{\frac{1}{3}} \leq \delta^{\frac{1}{3}}$. Therefore,

$$\begin{aligned} I & \leq \rho_0(\delta^{\frac{1}{3}}) + C\mathbb{P} \left(\|B^t\|_{t_\delta} + \sup_{t_\delta \leq s \leq t_0} |B_s^t - B_{s-\delta}^t| + \sup_{0 \leq r \leq \delta} |B_{\theta+r}^t - B_\theta^t| \geq \frac{1}{4}\delta^{\frac{1}{3}} \right) \\ & \leq \rho_0(\delta^{\frac{1}{3}}) + C\delta^{-\frac{8}{3}} \mathbb{E}^{\mathbb{P}} \left[\|B^t\|_{t_\delta}^8 + \sup_{t_\delta \leq s \leq t_0} |B_s^t - B_{s-\delta}^t|^8 + \sup_{0 \leq r \leq \delta} |B_{\theta+r}^t - B_\theta^t|^8 \right] \\ & \leq \rho_0(\delta^{\frac{1}{3}}) + C\delta^{\frac{4}{3}} + C\delta^{-\frac{8}{3}} \mathbb{E}^{\mathbb{P}} \left[\sup_{t_\delta \leq s \leq t_0} |B_s^t - B_{s-\delta}^t|^8 \right]. \end{aligned}$$

Set $t_\delta = s_0 < \dots < s_n = t_0$ such that $\delta \leq s_{i+1} - s_i \leq 2\delta$, $i = 0, \dots, n-1$. Then

$$\begin{aligned} \mathbb{E}^{\mathbb{P}} \left[\sup_{t_\delta \leq s \leq t_0} |B_s^t - B_{s-\delta}^t|^8 \right] & = \mathbb{E}^{\mathbb{P}} \left[\max_{0 \leq i \leq n-1} \sup_{s_i \leq s \leq s_{i+1}} |B_s^t - B_{s-\delta}^t|^8 \right] \\ & \leq \sum_{i=0}^{n-1} \mathbb{E}^{\mathbb{P}} \left[\sup_{s_i \leq s \leq s_{i+1}} [|B_s^t - B_{s_i-\delta}^t| + |B_{s_i-\delta}^t - B_{s_i-\delta}^t|]^8 \right] \\ & \leq C \sum_{i=0}^{n-1} (s_{i+1} - s_i + \delta)^4 \leq C\delta^{-1}\delta^4 = C\delta^3. \end{aligned}$$

Thus $I \leq \rho_0(\delta^{\frac{1}{3}}) + C\delta^{\frac{4}{3}} + C\delta^{-\frac{8}{3}}\delta^3 \leq \rho_0(\delta^{\frac{1}{3}}) + C\delta^{\frac{1}{3}} \leq C\rho_1(L\delta)$, proving (5.5) and hence the lemma. ■

We next show that the dynamic programming principle holds along deterministic times.

Lemma 5.2. *Let $t_1 < H(\omega)$ and $t_2 \in [t_1, t_0]$. We have:*

$$\widehat{Y}_{t_1}(\omega) = \sup_{\tau \in \mathcal{T}^{t_1}} \mathcal{E}_{t_1} \left[\widehat{X}_{\tau \wedge H^{t_1, \omega}}^{t_1, \omega} \mathbf{1}_{\{\tau \wedge H^{t_1, \omega} < t_2\}} + \widehat{Y}_{t_2}^{t_1, \omega} \mathbf{1}_{\{\tau \wedge H^{t_1, \omega} \geq t_2\}} \right].$$

Proof. When $t_2 = t_0$, the lemma coincides with the definition of \hat{Y} . Without loss of generality we assume $(t_1, \omega) = (0, \mathbf{0})$ and $t := t_2 < t_0$. First, follow the arguments in Lemma 4.1(ii) Step 1, one can easily prove

$$\hat{Y}_0 \leq \sup_{\tau \in \mathcal{T}} \mathcal{E} \left[\hat{X}_{\tau \wedge H} \mathbf{1}_{\{\tau \wedge H < t\}} + \hat{Y}_t \mathbf{1}_{\{\tau \wedge H \geq t\}} \right]. \quad (5.7)$$

To show that equality holds in the above inequality, fix arbitrary $\mathbb{P} \in \mathcal{P}$ and $\tau \in \mathcal{T}$ satisfying $\tau \leq H$ (otherwise reset τ as $\tau \wedge H$), we shall prove

$$\mathbb{E}^{\mathbb{P}} \left[\hat{X}_{\tau} \mathbf{1}_{\{\tau < t\}} + \hat{Y}_t \mathbf{1}_{\{\tau \geq t\}} \right] \leq \hat{Y}_0.$$

Since $\hat{Y}_H = \hat{X}_H$, this amounts to show that:

$$\mathbb{E}^{\mathbb{P}} \left[\hat{X}_{\tau} \mathbf{1}_{\{\tau < t\} \cup \{H \leq t\}} + \hat{Y}_t \mathbf{1}_{\{\tau \geq t, H > t\}} \right] \leq \hat{Y}_0. \quad (5.8)$$

We adapt the arguments in Lemma 4.1(ii) Step 2 to the present situation. Fix $0 < \delta \leq t_0 - t$. Let $\{E_i\}_{i \geq 1}$ be an \mathcal{F}_t measurable partition of the event $\{\tau \geq t, H > t\} \in \mathcal{F}_t$ such that $\|\omega - \tilde{\omega}\| \leq L\delta$ for all $\omega, \tilde{\omega} \in E_i$. Fix an $\omega^i \in E_i$ for each i . By the definition of \hat{Y} we have

$$\hat{Y}_t(\omega^i) \leq \mathbb{E}^{\mathbb{P}^i} \left[\hat{X}_{\tau^i \wedge H^t, \omega^i}^{t, \omega^i} \right] + \delta \quad \text{for some } (\tau^i, \mathbb{P}^i) \in \mathcal{T}^t \times \mathcal{P}_t. \quad (5.9)$$

As in Lemma 5.1, we set $t_\delta := t + \delta < t_0$, $\tilde{B}_s^{t_\delta} := B_{s+\delta}^t - B_{t_\delta}^t$ for $s \geq t$, and $\tilde{\tau}^i(B^t) := [\tau^i(\tilde{B}^{t_\delta}) + \delta] \wedge t_0$. Then $\tilde{\tau}^i \in \mathcal{T}^t$. Moreover by Assumption 3.4 and Property (P3), for each $\omega \in E_i$, we may define $\mathbb{P}^{i, \omega} \in \mathcal{P}_t$ as follows: $\alpha^{\mathbb{P}^{i, \omega}} := \frac{1}{\delta}(\omega_t^i - \omega_t)$, $\beta^{\mathbb{P}^{i, \omega}} := \mathbf{0}$ on $[t, t_\delta]$, and the $\mathbb{P}^{i, \omega}$ -distribution of \tilde{B}^{t_δ} is equal to the \mathbb{P}^i -distribution of B^t . By (5.5), we have

$$\mathbb{E}^{\mathbb{P}^i} \left[\hat{X}_{\tau^i \wedge H^t, \omega^i}^{t, \omega^i} \right] - \mathbb{E}^{\mathbb{P}^{i, \omega}} \left[\hat{X}_{\tilde{\tau}^i \wedge H^t, \omega}^{t, \omega} \right] \leq C\rho_1(L\delta). \quad (5.10)$$

Then by Lemma 5.1 and (5.9), (5.10) we have

$$\hat{Y}_t(\omega) \leq \hat{Y}_t(\omega^i) + C\rho_1(L\delta) \leq \mathbb{E}^{\mathbb{P}^{i, \omega}} \left[\hat{X}_{\tilde{\tau}^i \wedge H^t, \omega}^{t, \omega} \right] + \delta + C\rho_1(L\delta), \quad \text{for all } \omega \in E_i. \quad (5.11)$$

We next define:

$$\tilde{\tau} := \mathbf{1}_{\{\tau < t\} \cup \{H \leq t\}} \tau + \sum_{i \geq 1} \mathbf{1}_{E_i} \tilde{\tau}^i(B^t), \quad \text{and then } \{\tau < t\} \cup \{H \leq t\} = \{\tilde{\tau} < t\} \cup \{H \leq t\}.$$

Since $\tau \leq H$, we see that $\{\tau < t\} \cup \{H \leq t\} = \{\tau < t\} \cup \{\tau = H = t\}$, and thus it is clear that $\tilde{\tau} \in \mathcal{T}$. Moreover, we claim that there exists $\tilde{\mathbb{P}} \in \mathcal{P}$ such that

$$\begin{aligned} \tilde{\mathbb{P}} &= \mathbb{P} \text{ on } \mathcal{F}_t \text{ and the regular conditional probability distribution} \\ (\tilde{\mathbb{P}})^{t, \omega} &= \mathbb{P}^{i, \omega} \quad \text{for } \mathbb{P}\text{-a.e. } \omega \in E_i, i \geq 1, \\ (\tilde{\mathbb{P}})^{t, \omega} &= \mathbb{P}^t, \omega \quad \text{for } \mathbb{P}\text{-a.e. } \omega \in \{\tau < t\} \cup \{H \leq t\}. \end{aligned} \quad (5.12)$$

Then, by (5.11) we have

$$\hat{Y}_t(\omega) \leq \mathbb{E}^{(\tilde{\mathbb{P}})^{t, \omega}} \left[\hat{X}_{\tilde{\tau} \wedge H^t, \omega}^{t, \omega} \right] + \delta + C\rho_1(L\delta), \quad \mathbb{P}\text{-a.e. } \omega \in \{\tau \geq t, H > t\}, \quad (5.13)$$

and therefore:

$$\begin{aligned}\mathbb{E}^{\mathbb{P}}\left[\widehat{X}_{\tau}\mathbf{1}_{\{\tau < t\} \cup \{H \leq t\}} + \widehat{Y}_t\mathbf{1}_{\{\tau \geq t, H > t\}}\right] &\leq \mathbb{E}^{\tilde{\mathbb{P}}}\left[\widehat{X}_{\tilde{\tau} \wedge H}\mathbf{1}_{\{\tau < t\} \cup \{H \leq t\}} + \widehat{X}_{\tilde{\tau} \wedge H}\mathbf{1}_{\{\tau \geq t, H > t\}}\right] \\ &\quad + \delta + C\rho_1(L\delta) \\ &= \mathbb{E}^{\tilde{\mathbb{P}}}\left[\widehat{X}_{\tilde{\tau} \wedge H}\right] + \delta + C\rho_1(L\delta) \leq \widehat{Y}_0 + \delta + C\rho_1(L\delta),\end{aligned}$$

which implies (5.8) by sending $\delta \rightarrow 0$. Then the reverse inequality of (5.7) follows from the arbitrariness of \mathbb{P} and τ .

It remains to prove (5.12). For any $\varepsilon > 0$ and each $i \geq 1$, there exists a partition $\{E_j^i, j \geq 1\}$ of E_i such that $\|\omega - \omega'\|_t \leq \varepsilon$ for any $\omega, \omega' \in E_j^i$. Fix an $\omega^{ij} \in E_j^i$ for each (i, j) . By Property (P3) we may define $\tilde{\mathbb{P}}^\varepsilon \in \mathcal{P}$ by:

$$\tilde{\mathbb{P}}^\varepsilon := \mathbb{P} \otimes_t \left[\sum_{i \geq 1} \sum_{j \geq 1} \mathbb{P}^{i, \omega^{ij}} \mathbf{1}_{E_j^i} + \mathbb{P} \mathbf{1}_{\{\tau < t\} \cup \{H \leq t\}} \right].$$

By Property (P1), \mathcal{P} is weakly compact. Then $\tilde{\mathbb{P}}^\varepsilon$ has a weak limit $\tilde{\mathbb{P}} \in \mathcal{P}$ as $\varepsilon \rightarrow 0$. We now show that $\tilde{\mathbb{P}}$ satisfies all the requirements in (5.12). Indeed, for any partition $0 = s_0 < \dots < s_m = t < s_{m+1} < \dots < s_M = t_\delta < s_{M+1} < \dots < s_N = T$ and any bounded and uniformly continuous function $\varphi : \mathbb{R}^{N \times d} \rightarrow \mathbb{R}$, let $\xi := \varphi(B_{s_1} - B_{s_0}, \dots, B_{s_N} - B_{s_{N-1}})$. Then, denoting $\Delta s_k := s_{k+1} - s_k$, $\Delta \omega_k := \omega_{s_k} - \omega_{s_{k-1}}$, we see that

$$\mathbb{E}^{\mathbb{P}^{i, \omega}}[\xi^{t, \omega}] = \eta_t^i(\omega), \quad \mathbb{E}^{\mathbb{P}^{i, \omega^{ij}}}[\xi^{t, \omega}] = \eta_t^{i, j}(\omega),$$

where:

$$\begin{aligned}\eta_t^i(\omega) &:= \mathbb{E}^{\mathbb{P}^i} \left[\varphi \left((\Delta \omega_k)_{1 \leq k \leq m}, \frac{\omega_t^i - \omega_t}{\delta} (\Delta s_k)_{m+1 \leq k \leq M}, (B_{s_k - \delta} - B_{s_{k-1} - \delta})_{M+1 \leq k \leq N} \right) \right]; \\ \eta_t^{i, j}(\omega) &:= \mathbb{E}^{\mathbb{P}^i} \left[\varphi \left((\Delta \omega_k)_{1 \leq k \leq m}, \frac{\omega_t^i - \omega_t^{ij}}{\delta} (\Delta s_k)_{m+1 \leq k \leq M}, (B_{s_k - \delta} - B_{s_{k-1} - \delta})_{M+1 \leq k \leq N} \right) \right].\end{aligned}$$

Let ρ denote the modulus of continuity function of φ . Then

$$\left| \mathbb{E}^{\mathbb{P}^{i, \omega^{ij}}}[\xi^{t, \omega}] - \mathbb{E}^{\mathbb{P}^{i, \omega}}[\xi^{t, \omega}] \right| \leq \rho(\varepsilon) \quad \text{for all } \omega \in E_j^i,$$

and thus

$$\begin{aligned}&\left| \mathbb{E}^{\tilde{\mathbb{P}}^\varepsilon}[\xi] - \mathbb{E}^{\mathbb{P}} \left[\xi \mathbf{1}_{\{\tau < t\} \cup \{H \leq t\}} + \sum_{i \geq 1} \eta_t^i \mathbf{1}_{E_i} \right] \right| \\ &= \left| \mathbb{E}^{\mathbb{P}} \left[\sum_{i, j \geq 1} \mathbb{E}^{\mathbb{P}^{i, \omega^{ij}}}[\xi^{t, \cdot}] \mathbf{1}_{E_j^i} \right] - \mathbb{E}^{\mathbb{P}} \left[\sum_{i, j \geq 1} \eta_t^i \mathbf{1}_{E_j^i} \right] \right| \\ &\leq \mathbb{E}^{\mathbb{P}} \left[\sum_{i, j \geq 1} \left| \mathbb{E}^{\mathbb{P}^{i, \omega^{ij}}}[\xi^{t, \cdot}] - \mathbb{E}^{\mathbb{P}^{i, \cdot}}[\xi^{t, \cdot}] \right| \mathbf{1}_{E_j^i} \right] \\ &\leq \mathbb{E}^{\mathbb{P}} \left[\sum_{i, j \geq 1} \rho(\varepsilon) \mathbf{1}_{E_j^i} \right] \leq \rho(\varepsilon).\end{aligned}$$

By sending $\varepsilon \rightarrow 0$, we obtain $\mathbb{E}^{\tilde{\mathbb{P}}}[\xi] = \mathbb{E}^{\mathbb{P}}[\xi \mathbf{1}_{\{\tau < t\} \cup \{H \leq t\}} + \sum_{i \geq 1} \eta_i^t \mathbf{1}_{E_i}]$, which proves (5.12) by the arbitrariness of ξ . ■

We now prove the regularity in the t -variable. Recall the ρ_2 defined in (5.4).

Lemma 5.3. *Let $0 \leq t_1 < H(\omega^1)$, $0 \leq t_2 < H(\omega^2)$, and $t_1 \leq t_2$. Then we have:*

$$|\widehat{Y}_{t_1}(\omega^1) - \widehat{Y}_{t_2}(\omega^2)| \leq C \left[1 + \frac{1}{d(\omega_{t_1}^1, O^c)} \right] \rho_2(\mathbf{d}_{\infty}((t_1, \omega^1), (t_2, \omega^2))).$$

Proof. Without loss of generality we assume $t_1 < t_2$. Also, in view of the uniform continuity in ω of Lemma 5.1, it suffices to prove the lemma in the case $\omega^1 = \omega^2 = \omega$.

Denote $\delta := \mathbf{d}_{\infty}((t_1, \omega), (t_2, \omega))$ and $\varepsilon := d(\omega_{t_1}, O^c)$. For $\delta \geq \frac{1}{8}$, we have $|\widehat{Y}_{t_1}(\omega) - \widehat{Y}_{t_2}(\omega)| \leq 2C_0 \leq C\varepsilon^{-1}\rho_2(\delta)$. So we assume in the rest of this proof that $\delta < \frac{1}{8}$.

First, by Assumption 3.4, we may consider the measure $\mathbb{P} \in \mathcal{P}_{t_1}$ such that $\alpha_t^{\mathbb{P}} := 0$, $\beta_t^{\mathbb{P}} := 0$, $t \in [t_1, t_2]$. Then, by setting $\tau := t_0$ in Lemma 5.2, we see that $\widehat{Y}_{t_1}(\omega) \geq \mathcal{E}_{t_1}[\widehat{Y}_{t_2}^{t_1, \omega}] \geq \mathbb{E}^{\mathbb{P}}[\widehat{Y}_{t_2}^{t_1, \omega}] = \widehat{Y}_{t_2}(\omega_{\cdot \wedge t_1})$. Note that $H(\omega_{\cdot \wedge t_1}) = t_0 > t_2$. Thus, by Lemma 5.1,

$$\widehat{Y}_{t_2}(\omega) - \widehat{Y}_{t_1}(\omega) \leq C\rho_1(\mathbf{d}_{\infty}((t_2, \omega_{\cdot \wedge t_1}), (t_2, \omega))) \leq C\rho_1(\delta) \leq C\rho_2(\delta). \quad (5.14)$$

Next, for arbitrary $\tau \in \mathcal{T}^{t_1}$, noting that $\widehat{X} \leq \widehat{Y}$ we have

$$\begin{aligned} I(\tau) &:= \mathcal{E}_{t_1} \left[\widehat{X}_{\tau \wedge H^{t_1, \omega}}^{t_1, \omega} \mathbf{1}_{\{\tau \wedge H^{t_1, \omega} < t_2\}} + \widehat{Y}_{t_2}^{t_1, \omega} \mathbf{1}_{\{\tau \wedge H^{t_1, \omega} \geq t_2\}} \right] - \widehat{Y}_{t_2}(\omega) \\ &= \mathcal{E}_{t_1} \left[\widehat{X}_{\tau}^{t_1, \omega} \mathbf{1}_{\{\tau < H^{t_1, \omega} \wedge t_2\}} + \widehat{X}_{H^{t_1, \omega}}^{t_1, \omega} \mathbf{1}_{\{H^{t_1, \omega} < t_2, H^{t_1, \omega} \leq \tau\}} + \widehat{Y}_{t_2}^{t_1, \omega} \mathbf{1}_{\{\tau \wedge H^{t_1, \omega} \geq t_2\}} \right] - \widehat{Y}_{t_2}(\omega) \\ &\leq \mathcal{E}_{t_1} \left[(\widehat{X}_{\tau}^{t_1, \omega} - \widehat{X}_{H^{t_1, \omega} \wedge t_2}^{t_1, \omega}) \mathbf{1}_{\{\tau < H^{t_1, \omega} \wedge t_2\}} + \widehat{Y}_{H^{t_1, \omega} \wedge t_2}^{t_1, \omega} \right] - \widehat{Y}_{t_2}(\omega) \\ &\leq \mathcal{E}_{t_1} \left[(\widehat{X}_{\tau}^{t_1, \omega} - \widehat{X}_{H^{t_1, \omega} \wedge t_2}^{t_1, \omega}) \mathbf{1}_{\{\tau < H^{t_1, \omega} \wedge t_2\}} \right] + \mathcal{E}_{t_1} \left[|\widehat{Y}_{t_2}^{t_1, \omega} - \widehat{Y}_{t_2}(\omega)| \mathbf{1}_{\{H^{t_1, \omega} > t_2\}} \right] \\ &\quad + C C_{t_1} [H^{t_1, \omega} \leq t_2]. \end{aligned}$$

By (5.3) and Lemma 5.1 we have

$$\begin{aligned} I(\tau) &\leq \mathcal{E}_{t_1} \left[\rho_0(\mathbf{d}_{\infty}((t_1, \omega), (t_2, \omega \otimes_{t_1} B^{t_1}))) \right] + C\mathcal{E}_{t_1} \left[\rho_1(\|\omega - \omega \otimes_{t_1} B^{t_1}\|_{t_2}) \right] \\ &\quad + CC_{t_1} [\|B^{t_1}\|_{t_2} \geq \varepsilon] \\ &\leq \mathcal{E}_{t_1} \left[\rho_0(\delta + \|B^{t_1}\|_{t_2}) \right] + C\mathcal{E}_{t_1} \left[\rho_1(\delta + \|B^{t_1}\|_{t_2}) \right] + C\varepsilon^{-1}\mathcal{E}_{t_1} [\|B^{t_1}\|_{t_2}] \\ &\leq C[1 + \varepsilon^{-1}]\mathcal{E}_{t_1} [\rho_1(\delta + \|B^{t_1}\|_{t_2})]. \end{aligned}$$

Since $\delta \leq \frac{1}{8}$, following the proof of (4.6) we have

$$I(\tau) \leq C[1 + \varepsilon^{-1}] \left[\rho_1(\delta^{\frac{1}{3}}) + \delta^{\frac{1}{3}} \right] \leq C[1 + \varepsilon^{-1}]\rho_2(\delta).$$

By the arbitrariness of τ and the dynamic programming principle of Theorem 5.4, we obtain $\widehat{Y}_{t_1}(\omega) - \widehat{Y}_{t_2}(\omega) \leq C\varepsilon^{-1}\rho_2(\delta)$, and the proof is complete by (5.14). ■

Applying Lemmas 5.1–5.3, and following the same arguments as those of Theorem 4.3, we establish the dynamic programming principle in the present context.

Theorem 5.4. Let $t < H(\omega)$ and $\tau \in \mathcal{T}^t$. Then

$$\widehat{Y}_t(\omega) = \sup_{\tilde{\tau} \in \mathcal{T}^t} \mathcal{E}_t \left[\widehat{X}_{\tilde{\tau} \wedge H^t, \omega}^t \mathbf{1}_{\{\tilde{\tau} \wedge H^t, \omega < \tau\}} + \widehat{Y}_{\tilde{\tau}}^t \mathbf{1}_{\{\tilde{\tau} \wedge H^t, \omega \geq \tau\}} \right].$$

Consequently, \widehat{Y} is a \mathcal{E} -supermartingale on $[0, H]$.

By Lemma 5.3, \widehat{Y} is continuous for $t \in [0, H)$. Moreover, since \widehat{Y} is an \mathcal{E} -supermartingale, we see that \widehat{Y}_{H-} exists. However, Example A.2 below shows that in general \widehat{Y} may be discontinuous at H . This issue is crucial for our purpose, and we will discuss more in Section 5.4 below.

5.2. Continuous approximation of the hitting times

Similar to the proof of Theorem 3.3, we need to apply some limiting arguments. We therefore assume without loss of generality that $\widehat{Y}_0 > \widehat{X}_0$ and introduce the stopping times: for any $m \geq 1$ and $n > (\widehat{Y}_0 - \widehat{X}_0)^{-1}$,

$$\begin{aligned} H_m &:= \inf \left\{ t \geq 0 : \mathbf{d}(\omega_t, O^c) \leq \frac{1}{m} \right\} \wedge \left(t_0 - \frac{1}{m} \right), \\ \tau_n &:= \inf \left\{ t \geq 0 : \widehat{Y}_t - \widehat{X}_t \leq \frac{1}{n} \right\}. \end{aligned} \quad (5.15)$$

Here we abuse the notation slightly by using the same notation τ_n as in (4.7). Our main task in this subsection is to build an approximation of H_m and τ_n by continuous random variables. This will be obtained by a repeated use of Lemma 4.5.

We start by a continuous approximation of the sequence $(H_m)_{m \geq 1}$ defined in (5.15).

Lemma 5.5. For all $m \geq 2$:

- (i) $H_{m-1}(\omega) \leq H_m(\omega') \leq H_{m+1}(\omega)$, whenever $\|\omega - \omega'\|_{t_0} \leq \frac{1}{m(m+1)}$,
- (ii) there exists an open subset $\Omega_0^m \subset \Omega$, and a uniformly continuous \widehat{H}_m such that
$$\mathcal{C}[(\Omega_0^m)^c] < 2^{-m} \quad \text{and} \quad H_{m-1} - 2^{-m} \leq \widehat{H}_m \leq H_{m+1} + 2^{-m} \quad \text{on } \Omega_0^m,$$
- (iii) there exist $\delta_m > 0$ such that $|\widehat{H}_m(\omega) - \widehat{H}_m(\omega')| \leq 2^{-m}$ whenever $\|\omega - \omega'\|_{t_0} \leq \delta_m$, and:
$$\mathcal{C}[(\widehat{\Omega}_0^m)^c] \leq 2^{-m} \quad \text{where } \widehat{\Omega}_0^m := \{\omega \in \Omega_0^m : \mathbf{d}(\omega, [\Omega_0^m]^c) > \delta_m\}.$$

Proof. Notice that (ii) is a direct consequence of (i) obtained by applying Lemma 4.5 with $\varepsilon = 2^{-m}$. To prove (i), we observe that for $\|\omega - \omega'\|_{t_0} \leq \frac{1}{m(m+1)}$ and $t < H_m(\omega')$, we have

$$\mathbf{d}(\omega_t, O^c) \geq \mathbf{d}(\omega'_t, O^c) - \frac{1}{m(m+1)} > \frac{1}{m} - \frac{1}{m(m+1)} = \frac{1}{m+1}.$$

This shows that $H_m(\omega') \leq H_{m+1}(\omega)$ whenever $\|\omega - \omega'\|_{t_0} \leq \frac{1}{m(m+1)}$. Similarly, $H_{m-1}(\omega) \leq H_m(\omega')$ whenever $\|\omega - \omega'\|_{t_0} \leq \frac{1}{m(m-1)}$, and the inequality (i) follows.

It remains to prove (iii). The first claim follows from the uniform continuity of \widehat{H}_m . For each $\delta > 0$, define $h_\delta : [0, \infty) \rightarrow [0, 1]$ as follows:

$$h_\delta(x) := 1 \quad \text{for } x \leq \delta, \quad h_\delta(x) = 0 \quad \text{for } x \geq 2\delta, \quad \text{and } h_\delta \text{ is linear on } [\delta, 2\delta]. \quad (5.16)$$

Then the map $\omega \mapsto \psi_\delta(\omega) := h_\delta(d(\omega, [\Omega_0^m]^c))$ is continuous, and $\psi_\delta \downarrow \mathbf{1}_{[\Omega_0^m]^c}$ as $\delta \downarrow 0$. Applying Proposition 2.6(ii) we have

$$\lim_{\delta \rightarrow 0} \mathcal{E}[\psi_\delta] = \mathcal{E}[\mathbf{1}_{(\Omega_0^m)^c}] = \mathcal{C}[(\Omega_0^m)^c] < 2^{-m}.$$

By definition of $\hat{\Omega}_0^m$, notice that $\mathbf{1}_{(\hat{\Omega}_0^m)^c} \leq \psi_{\delta_m}$. Then $\mathcal{C}[(\hat{\Omega}_0^m)^c] \leq \mathcal{E}[\psi_{\delta_m}]$, and (iii) holds true for sufficiently small δ_m . ■

We next derive a continuous approximation of the sequences

$$\tau_n^m := \tau_n \wedge \hat{H}_m, \quad (5.17)$$

where τ_n and \hat{H}_m are defined in (5.15) and Lemma 5.5(ii), respectively.

Lemma 5.6. For all $m \geq 2, n > (\hat{Y}_0 - \hat{X}_0)^{-1}$, there exists an open subset $\Omega_n^m \subset \Omega$ and a uniformly continuous map $\hat{\tau}_n^m$ such that

$$\tau_{n-1}^m - 2^{1-m} - 2^{-n} \leq \hat{\tau}_n^m \leq \tau_{n+1}^m + 2^{1-m} + 2^{-n}$$

$$\text{on } \hat{\Omega}_0^m \cap \Omega_n^m, \quad \text{and} \quad \mathcal{C}[(\Omega_n^m)^c] \leq 2^{-n}.$$

Proof. Fix m , and recall the modulus of continuity ρ_1 introduced in (5.4). For each n , let $0 < \delta_n^m < \delta^m$ such that $(\rho_0 + C\rho_1)(\delta_n^m) \leq \frac{1}{n(n+1)}$, where C is the constant in Lemma 5.1. We shall prove

$$(\tau_{n-1} \wedge \hat{H}_m)(\omega) - 2^{1-m} \leq (\tau_n \wedge \hat{H}_m)(\omega') \leq (\tau_{n+1} \wedge \hat{H}_m)(\omega) + 2^{1-m}$$

$$\text{whenever } \omega \in \hat{\Omega}_0^m, \quad \|\omega - \omega'\|_{t_0} \leq \delta_n^m. \quad (5.18)$$

Then the required statement follows from Lemma 4.5 with $\varepsilon = 2^{-n}$.

We shall prove only the right inequality of (5.18). The left one can be proved similarly. Let ω, ω' be as in (5.18). First, by Lemma 5.5(iii) we have

$$\omega' \in \Omega_0^m \quad \text{and} \quad \hat{H}_m(\omega') \leq \hat{H}_m(\omega) + 2^{-m}. \quad (5.19)$$

We now prove the right inequality of (5.18) in three cases.

Case 1. if $\tau_{n+1}(\omega) \geq \hat{H}_m(\omega') - 2^{-m}$, then $\hat{H}_m(\omega') \leq (\tau_{n+1} \wedge \hat{H}_m)(\omega) + 2^{-m}$ and thus the result is true.

Case 2. If $\tau_{n+1}(\omega) = H(\omega)$, then by Lemma 5.5(ii) we have $\hat{H}_m(\omega) \leq H_{m+1}(\omega) + 2^{-m} \leq \tau_{n+1}(\omega) + 2^{-m}$, and thus $\hat{H}_m(\omega') \leq \hat{H}_m(\omega) + 2^{-m} \leq \tau_{n+1}(\omega) + 2^{1-m}$. This, together with (5.19), proves the desired inequality.

Case 3. We now assume $\tau_{n+1}(\omega) < \hat{H}_m(\omega') - 2^{-m}$ and $\tau_{n+1}(\omega) < H(\omega)$. By Lemma 5.5(ii) we have $\tau_{n+1}(\omega) < H_{m+1}(\omega')$, and thus $\tau_{n+1}(\omega) < H(\omega')$. Then it follows from Lemma 5.1 that

$$(Y - X)_{\tau_{n+1}(\omega)}(\omega') \leq (Y - X)_{\tau_{n+1}(\omega)}(\omega) + (\rho_0 + C\rho_1)(\delta_n^m) \leq \frac{1}{n+1} + \frac{1}{n(n+1)} = \frac{1}{n}.$$

That is, $\tau_n(\omega') \leq \tau_{n+1}(\omega)$. This, together with (5.19), proves the desired inequality. ■

For our final approximation result, we introduce the notations:

$$\bar{\tau}_n := \tau_n \wedge H_n, \quad \underline{\theta}_n^* := \hat{\tau}_{n-1}^{n-1} - 2^{3-n}, \quad \bar{\theta}_n^* := \hat{\tau}_{n+1}^{n+1} + 2^{1-n}, \quad (5.20)$$

and

$$\Omega_n^* := \hat{\Omega}_0^{n-1} \cap \Omega_{n-1}^{n-1} \cap \hat{\Omega}_0^{n+1} \cap \Omega_{n+1}^{n+1}. \quad (5.21)$$

Lemma 5.7. For all $n \geq (\hat{Y}_0 - \hat{X}_0)^{-1} \vee 2$, θ_n^* , $\bar{\theta}_n^*$ are uniformly continuous, and $\theta_n^* \leq \bar{\tau}_n \leq \bar{\theta}_n^*$ on Ω_n^* .

Proof. This is a direct combination of Lemmas 5.5 and 5.6.

5.3. Proof of Theorem 3.6

We first prove the \mathcal{E} -martingale property under an additional condition.

Lemma 5.8. Let $\tau \in \mathcal{T}$ such that $\tau \leq \tau^*$ and $\mathcal{E}[Y_{\tau-}] = \mathcal{E}[Y_{\tau}]$ (in particular if $\tau < H$). Then \hat{Y} is an \mathcal{E} -martingale on $[0, \tau]$.

Proof. If $\hat{Y}_0 = \hat{X}_0$, then $\hat{\tau}^* = 0$ and obviously the statement is true. We then assume $\hat{Y}_0 > \hat{X}_0$, and prove the lemma in several steps.

Step 1. Let n be sufficiently large so that $\frac{1}{n} < \hat{Y}_0 - \hat{X}_0$. Follow the same arguments as that of Lemma 4.4, one can easily prove:

$$\hat{Y} \text{ is an } \mathcal{E}\text{-martingale on } [0, \tau_n]. \quad (5.22)$$

Step 2. Recall the sequence of stopping times $(\bar{\tau}_n)_{n \geq 1}$ introduced in (5.20). By Step 1 we have $\hat{Y}_0 = \mathcal{E}[\hat{Y}_{\bar{\tau}_n}]$. Then for any $\varepsilon > 0$, there exists $\mathbb{P}_n \in \mathcal{P}$ such that $\hat{Y}_0 - \varepsilon < \mathbb{E}^{\mathbb{P}_n}[\hat{Y}_{\bar{\tau}_n}]$. Since \mathcal{P} is weakly compact, there exists subsequence $\{n_j\}$ and $\mathbb{P}^* \in \mathcal{P}$ such that \mathbb{P}_{n_j} converges weakly to \mathbb{P}^* . Now for any n and $n_j \geq n$, since Y is a supermartingale under each \mathbb{P}_{n_j} and $(\bar{\tau}_n)_{n \geq 1}$ is increasing, we have

$$\hat{Y}_0 - \varepsilon < \mathbb{E}^{\mathbb{P}_{n_j}}[\hat{Y}_{\bar{\tau}_{n_j}}] \leq \mathbb{E}^{\mathbb{P}_{n_j}}[\hat{Y}_{\bar{\tau}_n}]. \quad (5.23)$$

Our next objective is to send $j \nearrow \infty$, for fixed n , and use the weak convergence of \mathbb{P}_{n_j} towards \mathbb{P}^* . To do this, we need to approximate $\hat{Y}_{\bar{\tau}_n}$ with continuous random variables. Denote

$$\psi_n(\omega) := h_n \left(\inf_{0 \leq t \leq \bar{\theta}_n^*(\omega)} \mathbf{d}(\omega_t, O^c) \right)$$

$$\text{with } h_n(x) := 1 \wedge [(n+3)(n+4)x - (n+3)]^+. \quad (5.24)$$

Then ψ_n is continuous in ω , and

$$\{\psi_n > 0\} \subset \left\{ \inf_{0 \leq t \leq \bar{\theta}_n^*(\omega)} \mathbf{d}(\omega_t, O^c) > \frac{1}{n+4} \right\} \subset \{\bar{\theta}_n^* < H_{n+4}\}. \quad (5.25)$$

In particular, this implies that $\hat{Y}_{\bar{\theta}_n^*} \psi_n$ and $\hat{Y}_{\bar{\theta}_n^*} \psi_n$ are continuous in ω . We now decompose the right hand-side term of (5.23) into:

$$\hat{Y}_0 - \varepsilon \leq \mathbb{E}^{\mathbb{P}_{n_j}} \left[[\hat{Y}_{\bar{\theta}_n^*} + (\hat{Y}_{\bar{\tau}_n} - \hat{Y}_{\bar{\theta}_n^*}) \mathbf{1}_{\Omega_n^*}] (\psi_n + (1 - \psi_n)) + (\hat{Y}_{\bar{\tau}_n} - \hat{Y}_{\bar{\theta}_n^*}) \mathbf{1}_{(\Omega_n^*)^c} \right].$$

Note that $\theta_n^* \leq \bar{\tau}_n \leq \bar{\theta}_n^*$ on Ω_n^* . Then

$$\hat{Y}_0 - \varepsilon \leq \mathbb{E}^{\mathbb{P}_{n_j}} \left[\left(\hat{Y}_{\bar{\theta}_n^*} + \sup_{\theta_n^* \leq t \leq \bar{\theta}_n^*} (\hat{Y}_t - \hat{Y}_{\bar{\theta}_n^*}) \right) \psi_n \right] + CC[\psi_n < 1] + CC[(\Omega_n^*)^c].$$

Send $j \rightarrow \infty$, we obtain

$$\widehat{Y}_0 - \varepsilon \leq \mathbb{E}^{\mathbb{P}^*} \left[\psi_n \widehat{Y}_{\underline{\theta}_n^*} \right] + \mathbb{E}^{\mathbb{P}^*} \left[\psi_n \sup_{\underline{\theta}_n^* \leq t \leq \bar{\theta}_n^*} (\widehat{Y}_t - \widehat{Y}_{\underline{\theta}_n^*}) \right] + C\mathcal{C}[\psi_n < 1] + C\mathcal{C}[(\Omega_n^*)^c]. \quad (5.26)$$

Step 3. In this step we show that

$$\lim_{n \rightarrow \infty} \mathbb{E}^{\mathbb{P}^*} \left[\psi_n \sup_{\underline{\theta}_n^* \leq t \leq \bar{\theta}_n^*} (\widehat{Y}_t - \widehat{Y}_{\underline{\theta}_n^*}) \right] = \lim_{n \rightarrow \infty} C[\psi_n < 1] = \lim_{n \rightarrow \infty} C[(\Omega_n^*)^c] = 0. \quad (5.27)$$

(i) First, by the definition of Ω_n^* in (5.21) together with Lemmas 5.5(iii) and 5.6, it follows that $C[(\Omega_n^*)^c] \leq C2^{-n} \rightarrow 0$ as $n \rightarrow \infty$.

(ii) Next, notice that

$$\{\psi_n < 1\} = \left\{ \inf_{0 \leq t \leq \bar{\theta}_n^*(\omega)} \mathbf{d}(\omega_t, O^c) < \frac{1}{n+3} \right\} \subset \{\bar{\theta}_n^* > H_{n+3}\}.$$

Moreover, by (5.20) and Lemma 5.7,

$$\bar{\theta}_n^* = \hat{\tau}_{n+1}^{n+1} + 2^{1-n} = \underline{\theta}_{n+2}^* + 2^{2-n} \leq \bar{\tau}_{n+2} + 2^{2-n} \leq H_{n+2} + 2^{2-n}, \quad \text{on } \Omega_{n+2}^*.$$

Then

$$\begin{aligned} \{\psi_n < 1\} &\subset (\Omega_{n+2}^*)^c \cup \{H_{n+3} < H_{n+2} + 2^{2-n}\} \\ &\subset (\Omega_{n+2}^*)^c \cup \left\{ \sup_{H_{n+2} \leq t \leq H_{n+2} + 2^{2-n}} |B_t - B_{H_{n+2}}| \geq \frac{1}{(n+2)(n+3)} \right\}. \end{aligned}$$

Then one can easily see that $C[\psi_n < 1] \rightarrow 0$, as $n \rightarrow \infty$.

(iii) Finally, it is clear that $\underline{\theta}_n^* \rightarrow \hat{\tau}^*$, $\bar{\theta}_n^* \rightarrow \hat{\tau}^*$. Recall that $\widehat{Y}_{\hat{\tau}^*-}$ exists. By (5.25), we see that $\psi_n \sup_{\underline{\theta}_n^* \leq t \leq \bar{\theta}_n^*} (\widehat{Y}_t - \widehat{Y}_{\underline{\theta}_n^*}) \rightarrow 0$, \mathbb{P}^* -a.s. as $n \rightarrow \infty$. Then by applying the dominated convergence theorem under \mathbb{P}^* we obtain the first convergence in (5.27).

Step 4. By the dominated convergence theorem under \mathbb{P}^* we obtain $\lim_{n \rightarrow \infty} \mathbb{E}^{\mathbb{P}^*} [\psi_n \widehat{Y}_{\underline{\theta}_n^*}] = \mathbb{E}^{\mathbb{P}^*} [\widehat{Y}_{\hat{\tau}^*-}]$. This, together with (5.26) and (5.27), implies that

$$\widehat{Y}_0 \leq \mathbb{E}^{\mathbb{P}^*} [\widehat{Y}_{\hat{\tau}^*-}] + \varepsilon.$$

Note that \widehat{Y} is an \mathbb{P}^* -supermartingale and $\tau \leq \hat{\tau}^*$, then

$$\widehat{Y}_0 \leq \mathbb{E}^{\mathbb{P}^*} [\widehat{Y}_{\tau-}] + \varepsilon.$$

Since ε is arbitrary, we obtain $\widehat{Y}_0 \leq \mathcal{E}[\widehat{Y}_{\tau-}]$, and thus by the assumption $\mathcal{E}[\widehat{Y}_{\tau-}] = \mathcal{E}[\widehat{Y}_{\tau}]$ we have $\widehat{Y}_0 \leq \mathcal{E}[\widehat{Y}_{\tau}]$. This, together with the fact that \widehat{Y} is a \mathcal{E} -supermartingale, implies that

$$\widehat{Y}_0 = \mathcal{E}[\widehat{Y}_{\tau}]. \quad (5.28)$$

Similarly, one can prove $\widehat{Y}_t(\omega) = \mathcal{E}_t[\widehat{Y}_{\tau}^{t,\omega}]$ for $t < \tau(\omega)$, and thus $\widehat{Y}_{\cdot \wedge \tau}$ is a \mathcal{E} -martingale. ■

In light of Lemma 5.8, the following result is obviously important for us.

Proposition 5.9. *It holds that $\mathcal{E}[\widehat{Y}_{\hat{\tau}^*-}] = \mathcal{E}[\widehat{Y}_{\hat{\tau}^*}]$.*

We recall again that $\widehat{Y}_{\hat{\tau}^*-} = \widehat{Y}_{\hat{\tau}^*}$ whenever $\hat{\tau}^* < H$. So the only possible discontinuity is at H . The proof of Proposition 5.9 is reported in Section 5.4 below. Let us first show how it allows to complete the

Proof of Theorem 3.6. By Lemma 5.8 and Proposition 5.9, \widehat{Y} is an \mathcal{E} -martingale on $[0, \widehat{\tau}^*]$. Moreover, since $\widehat{X}_{\widehat{\tau}^*} = \widehat{Y}_{\widehat{\tau}^*}$, then $\widehat{Y}_0 = \mathcal{E}[\widehat{X}_{\widehat{\tau}^*}]$ and thus $\widehat{\tau}^*$ is an optimal stopping time. ■

Remark 5.10. Assume Assumption 3.4(ii) and the conditions of Lemma A.1 below hold, by Remark 3.5(iii) and Lemma A.1 we see that Proposition 5.9 and hence Theorem 3.6 hold. That is, in this case the Section 5.4 below is not needed. ■

5.4. \mathcal{E} -continuity of \widehat{Y} at the random maturity

This subsection is dedicated to the proof of Proposition 5.9. We first reformulate some path-wise properties established in previous subsections. For that purpose, we introduce the following additional notation: for any $\mathbb{P} \in \mathcal{P}$, $\tau \in \mathcal{T}$, and $E \in \mathcal{F}_\tau$

$$\mathcal{P}(\mathbb{P}, \tau, E) := \left\{ \mathbb{P}' \in \mathcal{P} : \mathbb{P}' = \mathbb{P} \otimes_\tau [\mathbb{P}' \mathbf{1}_E + \mathbb{P} \mathbf{1}_{E^c}] \right\}, \quad \mathcal{P}(\mathbb{P}, \tau) := \mathcal{P}(\mathbb{P}, \tau, \Omega). \quad (5.29)$$

That is, $\mathbb{P}' \in \mathcal{P}(\mathbb{P}, \tau, E)$ means $\mathbb{P}' = \mathbb{P}$ on \mathcal{F}_τ and $(\mathbb{P}')^{\tau, \omega} = \mathbb{P}^{\tau, \omega}$ for \mathbb{P} -a.e. $\omega \in E^c$.

The first result corresponds to Theorem 5.4.

Lemma 5.11. Let $\mathbb{P} \in \mathcal{P}$, $\tau_1, \tau_2 \in \mathcal{T}$, and $E \in \mathcal{F}_{\tau_1}$. Assume $\tau_1 \leq \tau_2 \leq H$, and $\tau_1 < H$ on E . Then for any $\varepsilon > 0$, there exist $\mathbb{P}_\varepsilon \in \mathcal{P}(\mathbb{P}, \tau_1, E)$ and $\tau_\varepsilon \in \mathcal{T}$ with values in $[\tau_1, \tau_2]$, s.t.

$$\mathbb{E}^\mathbb{P}[\widehat{Y}_{\tau_1} \mathbf{1}_E] \leq \mathbb{E}^{\mathbb{P}_\varepsilon}[\widehat{X}_{\tau_\varepsilon} \mathbf{1}_{\{\tau_\varepsilon < \tau_2\}} + \widehat{Y}_{\tau_2} \mathbf{1}_{\{\tau_\varepsilon = \tau_2\}}] \mathbf{1}_E + \varepsilon.$$

Proof. Let τ_1^n be a sequence of stopping times such that $\tau_1^n \downarrow \tau$ and each τ_1^n takes only finitely many values. Applying Lemma 5.3 together with the dominated convergence theorem under \mathbb{P} , we see that $\lim_{n \rightarrow \infty} \mathbb{E}^\mathbb{P}[|\widehat{Y}_{\tau_1^n \wedge \tau_2} - \widehat{Y}_{\tau_1}|] = 0$. Fix n such that

$$\mathbb{E}^\mathbb{P}[|\widehat{Y}_{\tau_1^n \wedge \tau_2} - \widehat{Y}_{\tau_1}|] \leq \frac{\varepsilon}{2}. \quad (5.30)$$

Assume τ_1^n takes values $\{t_i, i = 1, \dots, m\}$, and for each i , denote $E_i := E \cap \{\tau_1^n = t_i < \tau_2\} \in \mathcal{F}_{t_i}$. By (5.13), there exists $\tilde{\tau}_i \in \mathcal{T}$ and $\tilde{\mathbb{P}}_i \in \mathcal{P}(\mathbb{P}, t_i)$ such that $\tilde{\tau}_i \geq t_i$ on E_i and

$$\widehat{Y}_{t_i} \leq \mathbb{E}^{\tilde{\mathbb{P}}_i}[\widehat{X}_{\tilde{\tau}_i \wedge H}] + \frac{\varepsilon}{2}, \quad \mathbb{P}\text{-a.s. on } E_i. \quad (5.31)$$

Here $\mathbb{E}^{\tilde{\mathbb{P}}_i}[\cdot] := \mathbb{E}^{\tilde{\mathbb{P}}_i}[\cdot | \mathcal{F}_{t_i}]$ denotes the conditional expectation. Define

$$\tilde{\tau} := \tau_2 \mathbf{1}_{E^c \cup \{\tau_2 \leq \tau_1^n\}} + \sum_{i=1}^m \tilde{\tau}_i \mathbf{1}_{E_i}, \quad \tilde{\mathbb{P}} := \mathbb{P} \mathbf{1}_{E^c \cup \{\tau_2 \leq \tau_1^n\}} + \sum_{i=1}^m \tilde{\mathbb{P}}_i \mathbf{1}_{E_i}. \quad (5.32)$$

Then one can check straightforwardly that

$$\tilde{\tau} \in \mathcal{T} \quad \text{and} \quad \tilde{\tau} \geq \tau_2 \wedge \tau_1^n; \quad (5.33)$$

and $\tilde{\mathbb{P}} \in \mathcal{P}(\mathbb{P}, \tau_2 \wedge \tau_1^n, E) \subset \mathcal{P}(\mathbb{P}, \tau_1, E)$. Moreover, by (5.31) and (5.32),

$$\begin{aligned} \mathbb{E}^{\tilde{\mathbb{P}}}[\widehat{Y}_{\tau_2 \wedge \tau_1^n} \mathbf{1}_E] &= \mathbb{E}^{\tilde{\mathbb{P}}} \left[\left[\widehat{Y}_{\tau_2} \mathbf{1}_{\{\tau_2 \leq \tau_1^n\}} + \sum_{i=1}^m \widehat{Y}_{t_i} \mathbf{1}_{E_i} \right] \mathbf{1}_E \right] \\ &\leq \mathbb{E}^{\tilde{\mathbb{P}}} \left[\left[\widehat{Y}_{\tau_2} \mathbf{1}_{\{\tau_2 \leq \tau_1^n\}} + \left(\widehat{X}_{\tilde{\tau} \wedge H} + \frac{\varepsilon}{2} \right) \mathbf{1}_{\{\tau_1^n < \tau_2\}} \right] \mathbf{1}_E \right]. \end{aligned}$$

This, together with (5.30) and (5.33), leads to

$$\begin{aligned} & \mathbb{E}^{\tilde{\mathbb{P}}} \left[\left(\widehat{Y}_{\tau_1} - \widehat{X}_{\tilde{\tau}} \mathbf{1}_{\{\tilde{\tau} < \tau_2\}} - \widehat{Y}_{\tau_2} \mathbf{1}_{\{\tilde{\tau} \geq \tau_2\}} \right) \mathbf{1}_E \right] \\ & \leq \varepsilon + \mathbb{E}^{\tilde{\mathbb{P}}} \left[\left(\widehat{Y}_{\tau_2} \mathbf{1}_{\{\tau_2 \leq \tau_1^n\}} + \widehat{X}_{\tilde{\tau} \wedge H} \mathbf{1}_{\{\tau_1^n < \tau_2\}} - \widehat{X}_{\tilde{\tau}} \mathbf{1}_{\{\tilde{\tau} < \tau_2\}} - \widehat{Y}_{\tau_2} \mathbf{1}_{\{\tilde{\tau} \geq \tau_2\}} \right) \mathbf{1}_E \right] \\ & = \varepsilon + \mathbb{E}^{\tilde{\mathbb{P}}} \left[\left(\widehat{X}_{\tilde{\tau} \wedge H} - \widehat{Y}_{\tau_2} \right) \mathbf{1}_{\{\tau_1^n < \tau_2 \leq \tilde{\tau}\}} \mathbf{1}_E \right] \\ & = \varepsilon + \mathbb{E}^{\tilde{\mathbb{P}}} \left[\left(\mathbb{E}_{\tau_2}^{\tilde{\mathbb{P}}} [\widehat{X}_{\tilde{\tau} \wedge H}] - \widehat{Y}_{\tau_2} \right) \mathbf{1}_{\{\tau_1^n < \tau_2 \leq \tilde{\tau}\}} \mathbf{1}_E \right] \leq \varepsilon, \end{aligned}$$

where the last inequality follows from the definition of \widehat{Y} . Then, by setting $\tau_\varepsilon := \tilde{\tau} \wedge \tau_2$ we prove the result. ■

Next result corresponds to Lemma 5.8.

Lemma 5.12. *Let $\mathbb{P} \in \mathcal{P}$, $\tau \in \mathcal{T}$, and $E \in \mathcal{F}_\tau$ such that $\tau \leq \widehat{\tau}^*$ on E . Then for all $\varepsilon > 0$:*

$$\mathbb{E}^{\mathbb{P}} [\mathbf{1}_E \widehat{Y}_\tau] \leq \mathbb{E}^{\mathbb{P}_\varepsilon} [\mathbf{1}_E \widehat{Y}_{\widehat{\tau}^* -}] + \varepsilon \quad \text{for some } \mathbb{P}_\varepsilon \in \mathcal{P}(\mathbb{P}, \tau, E).$$

Proof. We proceed in three steps.

Step 1. We first assume $\tau = t < \widehat{\tau}^*$ on E . We shall prove the result following the arguments in Lemma 5.8. Recall the notations in Section 5.2 and the ψ_n defined in (5.24), and let ρ_n denote the modulus of continuity functions of $\underline{\theta}_n^*$, $\bar{\theta}_n^*$, and ψ_n .

Denote $\bar{\tau}_n := 0$ for $n \leq (\widehat{Y}_0 - \widehat{X}_0)^{-1}$. For any n and $\delta > 0$, let $\{E_i^{n,\delta}, i \geq 1\} \subset \mathcal{F}_t$ be a partition of $E \cap \{\bar{\tau}_{n-1} \leq t < \bar{\tau}_n\}$ such that $\|\omega - \omega'\|_t \leq \delta$ for any $\omega, \omega' \in E_i^{n,\delta}$. For each (n, i) , fix $\omega^{n,i} := \omega^{n,\delta,i} \in E_i^{n,\delta}$. By Lemma 5.8, $\widehat{Y}_{E_i^{n,\delta}}$ is an \mathcal{E} -martingale on $[t, \bar{\tau}_n]$. Then $\widehat{Y}_t(\omega^{n,i}) = \mathcal{E}_t[\widehat{Y}_{\bar{\tau}_n}^{t,\omega^{n,i}}]$, and thus there exists $\mathbb{P}_i^{n,\delta} \in \mathcal{P}_t$ such that

$$\widehat{Y}_t(\omega^{n,i}) \leq \mathbb{E}^{\mathbb{P}_i^{n,\delta}} \left[\widehat{Y}_{\bar{\tau}_n}^{t,\omega^{n,i}} \right] + \varepsilon. \quad (5.34)$$

Note that $\bigcup_{m=1}^n \bigcup_{i \geq 1} E_i^{m,\delta} = E \cap \{t < \bar{\tau}_n\}$. Set

$$\mathbb{P}^{n,\delta} := \mathbb{P} \otimes_t \left[\sum_{m=1}^n \sum_{i \geq 1} \mathbb{P}_i^{m,\delta} \mathbf{1}_{E_i^{m,\delta}} + \mathbb{P} \mathbf{1}_{E^c \cup \{t \geq \bar{\tau}_n\}} \right] \in \mathcal{P}(\mathbb{P}, t, E). \quad (5.35)$$

Recall the h_δ defined by (5.16). We claim that, for any $N \geq n$,

$$\begin{aligned} & \mathbb{E}^{\mathbb{P}} [\widehat{Y}_t \mathbf{1}_E] - \mathbb{E}^{\mathbb{P}^{N,\delta}} [\widehat{Y}_{t \vee \underline{\theta}_n^*} \psi_n \mathbf{1}_E] \\ & \leq Cn\varepsilon \left[\rho_2(\delta + \rho_n(\delta) + 2\eta_n(\delta)) \right] + C\rho_n(\delta) + \varepsilon + C2^{-n} + C\mathcal{C}(\psi_n < 1) \\ & \quad + 2\mathbb{E}^{\mathbb{P}^{N,\delta}} \left[\sup_{\underline{\theta}_n^* \leq s \leq \bar{\theta}_n^*} |\widehat{Y}_s - \widehat{Y}_{\underline{\theta}_n^*}| \psi_n \mathbf{1}_E \right] + C\varepsilon \left[h_\delta(d(\omega, (\Omega_n^*)^c)) \right], \end{aligned} \quad (5.36)$$

$$\text{where } \eta_n(\delta) := \sup_{t \leq s_1 < s_2 \leq t_0, s_2 - s_1 \leq \rho_n(\delta)} |B_{s_1}^t - B_{s_2}^t|.$$

Moreover, one can easily find \mathcal{F}_t -measurable continuous random variables φ_k such that $|\varphi_k| \leq 1$ and $\lim_{k \rightarrow \infty} \mathbb{E}^\mathbb{P}[\mathbf{1}_E - \varphi_k] = 0$. Then

$$\begin{aligned} & \mathbb{E}^\mathbb{P}[\widehat{Y}_t \mathbf{1}_E] - \mathbb{E}^{\mathbb{P}^{N,\delta}}[\widehat{Y}_{t \vee \underline{\theta}_n^*} \psi_n \varphi_k] \\ & \leq Cn\mathcal{E}\left[\rho_2\left(\delta + \rho_n(\delta) + 2\eta_n(\delta)\right)\right] + C\rho_n(\delta) + \varepsilon + C2^{-n} + CC(\psi_n < 1) \\ & \quad + C\mathbb{E}^{\mathbb{P}^{N,\delta}}\left[\sup_{\underline{\theta}_n^* \leq s \leq \bar{\theta}_n^*} |\widehat{Y}_s - \widehat{Y}_{\underline{\theta}_n^*}| \psi_n \varphi_k\right] + C\mathcal{E}\left[h_\delta(d(\omega, (\Omega_n^*)^c))\right] + C\mathbb{E}^\mathbb{P}[\mathbf{1}_E - \varphi_k]. \end{aligned}$$

Send $\delta \rightarrow 0$. First note that $[\delta + \rho_n(\delta) + 2\eta_n(\delta)] \downarrow 0$ and $h_\delta \downarrow \mathbf{1}_{\{0\}}$, then by Proposition 2.6(ii) we have

$$\begin{aligned} & \lim_{\delta \rightarrow 0} \mathcal{E}\left[\rho_2\left(\delta + \rho_n(\delta) + 2\eta_n(\delta)\right)\right] = 0; \\ & \lim_{\delta \rightarrow 0} \mathcal{E}\left[h_\delta(d(\omega, (\Omega_n^*)^c))\right] = C\left[d(\omega, (\Omega_n^*)^c) = 0\right] = C[(\Omega_n^*)^c] \leq C2^{-n}. \end{aligned}$$

Moreover, for each N , by the weak compactness assumption (P1) we see that $\mathbb{P}^{N,\delta}$ has a weak limit $\mathbb{P}^N \in \mathcal{P}$. It is straightforward to check that $\mathbb{P}^N \in \mathcal{P}(\mathbb{P}, t, E)$. Note that the random variables $\widehat{Y}_{t \vee \underline{\theta}_n^*} \psi_n \varphi_k$ and $\sup_{\underline{\theta}_n^* \leq s \leq \bar{\theta}_n^*} |\widehat{Y}_s - \widehat{Y}_{\underline{\theta}_n^*}| \psi_n \varphi_k$ are continuous. Then

$$\begin{aligned} & \mathbb{E}^\mathbb{P}[\widehat{Y}_t \mathbf{1}_E] - \mathbb{E}^{\mathbb{P}^N}[\widehat{Y}_{t \vee \underline{\theta}_n^*} \psi_n \varphi_k] \\ & \leq \varepsilon + C2^{-n} + CC(\psi_n < 1) + C\mathbb{E}^{\mathbb{P}^N}\left[\sup_{\underline{\theta}_n^* \leq s \leq \bar{\theta}_n^*} |\widehat{Y}_s - \widehat{Y}_{\underline{\theta}_n^*}| \psi_n \varphi_k\right] + C\mathbb{E}^\mathbb{P}[\mathbf{1}_E - \varphi_k]. \end{aligned}$$

Again by the weak compactness assumption (P1), \mathbb{P}^N has a weak limit $\mathbb{P}^* \in \mathcal{P}(\mathbb{P}, t, E)$ as $N \rightarrow \infty$. Now send $N \rightarrow \infty$, by the continuity of the random variables we obtain

$$\begin{aligned} & \mathbb{E}^\mathbb{P}[\widehat{Y}_t \mathbf{1}_E] - \mathbb{E}^{\mathbb{P}^*}[\widehat{Y}_{t \vee \underline{\theta}_n^*} \psi_n \varphi_k] \\ & \leq \varepsilon + C2^{-n} + CC(\psi_n < 1) + C\mathbb{E}^{\mathbb{P}^*}\left[\sup_{\underline{\theta}_n^* \leq s \leq \bar{\theta}_n^*} |\widehat{Y}_s - \widehat{Y}_{\underline{\theta}_n^*}| \psi_n \varphi_k\right] + C\mathbb{E}^\mathbb{P}[\mathbf{1}_E - \varphi_k]. \end{aligned}$$

Send $k \rightarrow \infty$ and recall that $\mathbb{P}^* = \mathbb{P}$ on \mathcal{F}_t , we have

$$\begin{aligned} & \mathbb{E}^\mathbb{P}[\widehat{Y}_t \mathbf{1}_E] - \mathbb{E}^{\mathbb{P}^*}[\widehat{Y}_{t \vee \underline{\theta}_n^*} \psi_n \mathbf{1}_E] \\ & \leq \varepsilon + C2^{-n} + CC(\psi_n < 1) + 2\mathbb{E}^{\mathbb{P}^*}\left[\sup_{\underline{\theta}_n^* \leq s \leq \bar{\theta}_n^*} |\widehat{Y}_s - \widehat{Y}_{\underline{\theta}_n^*}| \psi_n \mathbf{1}_E\right]. \end{aligned}$$

Finally send $n \rightarrow \infty$, by (5.27) and applying the dominated convergence theorem under \mathbb{P} and \mathbb{P}^* we have

$$\mathbb{E}^\mathbb{P}[\widehat{Y}_t \mathbf{1}_E] - \mathbb{E}^{\mathbb{P}^*}[\widehat{Y}_{\widehat{\tau}^*} \mathbf{1}_E] \leq \varepsilon.$$

That is, $\mathbb{P}_\varepsilon := \mathbb{P}^*$ satisfies the requirement in the case $\tau = t < \widehat{\tau}^*$ on E .

Step 2. We now prove Claim (5.36). Indeed, for any $m \leq n$ and any $\omega \in E_i^{m,\delta}$, by Lemma 5.1 we have

$$\begin{aligned} & \widehat{Y}_t(\omega) - \mathbb{E}^{\mathbb{P}_i^{m,\delta}}\left[\widehat{Y}_{\widehat{\tau}_n^{t,\omega}}^{t,\omega}\right] \\ & = \widehat{Y}_t(\omega) - \widehat{Y}_t(\omega^{m,i}) + \widehat{Y}_t(\omega^{m,i}) - \mathbb{E}^{\mathbb{P}_i^{m,\delta}}\left[\widehat{Y}_{\widehat{\tau}_n^{t,\omega^{m,i}}}^{t,\omega^{m,i}}\right] + \mathbb{E}^{\mathbb{P}_i^{m,\delta}}\left[\widehat{Y}_{\widehat{\tau}_n^{t,\omega^{m,i}}}^{t,\omega^{m,i}} - \widehat{Y}_{\widehat{\tau}_n^{t,\omega}}^{t,\omega}\right] \end{aligned}$$

$$\begin{aligned}
&\leq C\rho_1(\delta) + \varepsilon + \mathbb{E}^{\mathbb{P}_i^{m,\delta}} \left[\left| \widehat{Y}_{\tau_n^{t,\omega^{m,i}}}^{t,\omega^{m,i}} - \widehat{Y}_{\tau_n^{t,\omega}}^{t,\omega} \right| \mathbf{1}_{(\Omega_n^*)^{t,\omega^{m,i}} \cap (\Omega_n^*)^{t,\omega}} \psi_n^{t,\omega^{m,i}} \psi_n^{t,\omega} \right] \\
&\quad + C\mathbb{P}_i^{m,\delta} \left[[(\Omega_n^*)^{t,\omega^{m,i}}]^c \cup [(\Omega_n^*)^{t,\omega}]^c \right] + C\mathbb{E}^{\mathbb{P}_i^{m,\delta}} \left[1 - \psi_n^{t,\omega^{m,i}} + 1 - \psi_n^{t,\omega} \right]. \quad (5.37)
\end{aligned}$$

Note that

$$\begin{aligned}
\mathbb{E}^{\mathbb{P}_i^{m,\delta}} \left[1 - \psi_n^{t,\omega^{m,i}} + 1 - \psi_n^{t,\omega} \right] &\leq 2\mathbb{E}^{\mathbb{P}_i^{m,\delta}} \left[1 - \psi_n^{t,\omega} \right] + \rho_n(\delta); \\
\mathbb{P}_i^{m,\delta} \left[[(\Omega_n^*)^{t,\omega^{m,i}}]^c \cup [(\Omega_n^*)^{t,\omega}]^c \right] &\leq 2\mathbb{P}_i^{m,\delta} \left[[(\Omega_n^*)^{t,\omega}]^c \right] \\
&\quad + \mathbb{P}_i^{m,\delta} \left[[(\Omega_n^*)^{t,\omega^{m,i}}]^c \cap (\Omega_n^*)^{t,\omega} \right] \\
&\leq 2\mathbb{P}_i^{m,\delta} \left[[(\Omega_n^*)^{t,\omega}]^c \right] \\
&\quad + \mathbb{P}_i^{m,\delta} \left[0 < d(\omega \otimes_t B^t, (\Omega_n^*)^c) < \delta \right] \\
&\leq 2\mathbb{P}_i^{m,\delta} \left[[(\Omega_n^*)^{t,\omega}]^c \right] \\
&\quad + \mathbb{E}^{\mathbb{P}_i^{m,\delta}} \left[h_\delta(d(\omega \otimes_t B^t, (\Omega_n^*)^c)) \right]. \quad (5.38)
\end{aligned}$$

Moreover, on $(\Omega_n^*)^{t,\omega^{m,i}} \cap (\Omega_n^*)^{t,\omega} \cap \{\psi_n^{t,\omega^{m,i}} > 0\} \cap \{\psi_n^{t,\omega} > 0\}$, by Lemma 5.7 and (5.25) we have

$$(\underline{\theta}_n^*)^{t,\omega^{m,i}} \leq \bar{\tau}_n^{t,\omega^{m,i}} \leq (\bar{\theta}_n^*)^{t,\omega^{m,i}} < H_{n+4}^{t,\omega^{m,i}}; \quad (\underline{\theta}_n^*)^{t,\omega} \leq \bar{\tau}_n^{t,\omega} \leq (\bar{\theta}_n^*)^{t,\omega} < H_{n+4}^{t,\omega}.$$

Then

$$\begin{aligned}
&\left| \widehat{Y}_{\tau_n^{t,\omega^{m,i}}}^{t,\omega^{m,i}} - \widehat{Y}_{\tau_n^{t,\omega}}^{t,\omega} \right| \leq \left| \widehat{Y}_{(\underline{\theta}_n^*)^{t,\omega^{m,i}}}^{t,\omega^{m,i}} - \widehat{Y}_{(\underline{\theta}_n^*)^{t,\omega}}^{t,\omega} \right| \\
&\quad + \sup_{(\underline{\theta}_n^*)^{t,\omega^{m,i}} \leq s \leq (\bar{\theta}_n^*)^{t,\omega^{m,i}}} \left| \widehat{Y}_s^{t,\omega^{m,i}} - \widehat{Y}_{(\underline{\theta}_n^*)^{t,\omega^{m,i}}}^{t,\omega^{m,i}} \right| + \sup_{(\underline{\theta}_n^*)^{t,\omega} \leq s \leq (\bar{\theta}_n^*)^{t,\omega}} \left| \widehat{Y}_s^{t,\omega} - \widehat{Y}_{(\underline{\theta}_n^*)^{t,\omega}}^{t,\omega} \right| \\
&= \left| \widehat{Y}_{(\underline{\theta}_n^*)^{t,\omega^{m,i}}}^{t,\omega^{m,i}} - \widehat{Y}_{(\underline{\theta}_n^*)^{t,\omega}}^{t,\omega} \right| + 2 \sup_{(\underline{\theta}_n^*)^{t,\omega} \leq s \leq (\bar{\theta}_n^*)^{t,\omega}} \left| \widehat{Y}_s^{t,\omega} - \widehat{Y}_{(\underline{\theta}_n^*)^{t,\omega}}^{t,\omega} \right| \\
&\quad + \sup_{(\underline{\theta}_n^*)^{t,\omega^{m,i}} \leq s \leq (\bar{\theta}_n^*)^{t,\omega^{m,i}}} \left| \widehat{Y}_s^{t,\omega^{m,i}} - \widehat{Y}_{(\underline{\theta}_n^*)^{t,\omega^{m,i}}}^{t,\omega^{m,i}} \right| - \sup_{(\underline{\theta}_n^*)^{t,\omega} \leq s \leq (\bar{\theta}_n^*)^{t,\omega}} \left| \widehat{Y}_s^{t,\omega} - \widehat{Y}_{(\underline{\theta}_n^*)^{t,\omega}}^{t,\omega} \right|.
\end{aligned}$$

Applying Lemma 5.3 we get

$$\begin{aligned}
&\left| \widehat{Y}_{(\underline{\theta}_n^*)^{t,\omega^{m,i}}}^{t,\omega^{m,i}} - \widehat{Y}_{(\underline{\theta}_n^*)^{t,\omega}}^{t,\omega} \right| \leq Cn\rho_2 \left(d_\infty((\underline{\theta}_n^*)^{t,\omega^{m,i}}, \omega^{m,i} \otimes_t B^t), ((\underline{\theta}_n^*)^{t,\omega}, \omega \otimes_t B^t) \right) \\
&\leq Cn\rho_2 \left(\delta + \rho_n(\delta) + 2 \sup_{(\underline{\theta}_n^*)^{t,\omega} - \rho_n(\delta) \leq s \leq (\underline{\theta}_n^*)^{t,\omega} + \rho_n(\delta)} |B_s^t - B_{(\underline{\theta}_n^*)^{t,\omega}}^t| \right) \\
&\leq Cn\rho_2 \left(\delta + \rho_n(\delta) + 2\eta_n(\delta) \right),
\end{aligned}$$

and, similarly,

$$\begin{aligned}
 & \sup_{(\theta_n^*)^{t,\omega^{m,i}} \leq s \leq (\bar{\theta}_n^*)^{t,\omega^{m,i}}} |\widehat{Y}_s^{t,\omega^{m,i}} - \widehat{Y}_{(\theta_n^*)^{t,\omega^{m,i}}}^{t,\omega^{m,i}}| - \sup_{(\theta_n^*)^{t,\omega} \leq s \leq (\bar{\theta}_n^*)^{t,\omega}} |\widehat{Y}_s^{t,\omega} - \widehat{Y}_{(\theta_n^*)^{t,\omega}}^{t,\omega}| \\
 & \leq \sup_{(\theta_n^*)^{t,\omega^{m,i}} \leq s \leq (\bar{\theta}_n^*)^{t,\omega^{m,i}} \vee (\theta_n^*)^{t,\omega}} |\widehat{Y}_s^{t,\omega^{m,i}} - \widehat{Y}_{(\theta_n^*)^{t,\omega^{m,i}}}^{t,\omega^{m,i}}| \\
 & \quad + \sup_{(\theta_n^*)^{t,\omega^{m,i}} \vee (\theta_n^*)^{t,\omega} \leq s \leq (\bar{\theta}_n^*)^{t,\omega^{m,i}} \wedge (\bar{\theta}_n^*)^{t,\omega}} |\widehat{Y}_s^{t,\omega^{m,i}} - \widehat{Y}_s^{t,\omega}| + |\widehat{Y}_{(\theta_n^*)^{t,\omega^{m,i}}}^{t,\omega^{m,i}} - \widehat{Y}_{(\theta_n^*)^{t,\omega}}^{t,\omega}| \\
 & \quad + \sup_{(\bar{\theta}_n^*)^{t,\omega^{m,i}} \wedge (\bar{\theta}_n^*)^{t,\omega} \leq s \leq (\bar{\theta}_n^*)^{t,\omega^{m,i}}} |\widehat{Y}_s^{t,\omega^{m,i}} - \widehat{Y}_{(\bar{\theta}_n^*)^{t,\omega^{m,i}}}^{t,\omega^{m,i}}| + |\widehat{Y}_{(\bar{\theta}_n^*)^{t,\omega^{m,i}}}^{t,\omega^{m,i}} - \widehat{Y}_{(\bar{\theta}_n^*)^{t,\omega}}^{t,\omega}| \\
 & \leq Cn\rho_2(\delta + \rho_n(\delta) + 2\eta_n(\delta)) + C\rho_1(\delta) \leq Cn\rho_2(\delta + \rho_n(\delta) + 2\eta_n(\delta)).
 \end{aligned}$$

Then

$$|\widehat{Y}_{\bar{\tau}_n^{t,\omega^{m,i}}}^{t,\omega^{m,i}} - \widehat{Y}_{\bar{\tau}_n^{t,\omega}}^{t,\omega}| \leq Cn\rho_2(\delta + \rho_n(\delta) + 2\eta_n(\delta)) + 2 \sup_{(\theta_n^*)^{t,\omega} \leq s \leq (\bar{\theta}_n^*)^{t,\omega}} |\widehat{Y}_s^{t,\omega} - \widehat{Y}_{(\theta_n^*)^{t,\omega}}^{t,\omega}|.$$

Plug this and (5.38) into (5.37), for $\omega \in E_i^{m,\delta}$ we obtain

$$\begin{aligned}
 \widehat{Y}_t(\omega) - \mathbb{E}^{\mathbb{P}_i^{m,\delta}}[\widehat{Y}_{\bar{\tau}_n^{t,\omega}}^{t,\omega}] & \leq Cn\mathbb{E}^{\mathbb{P}_i^{m,\delta}}\left[\rho_2(\delta + \rho_n(\delta) + 2\eta_n(\delta))\right] + C\rho_n(\delta) + \varepsilon \\
 & \quad + 2\mathbb{E}^{\mathbb{P}_i^{m,\delta}}\left[\sup_{(\theta_n^*)^{t,\omega} \leq s \leq (\bar{\theta}_n^*)^{t,\omega}} |\widehat{Y}_s^{t,\omega} - \widehat{Y}_{(\theta_n^*)^{t,\omega}}^{t,\omega}| \psi_n^{t,\omega}\right] \\
 & \quad + C\mathbb{P}_i^{m,\delta}\left[(\Omega_n^*)^{t,\omega}\right]^c \\
 & \quad + C\mathbb{E}^{\mathbb{P}_i^{m,\delta}}\left[1 - \psi_n^{t,\omega}\right] + C\mathbb{E}^{\mathbb{P}_i^{m,\delta}}\left[h_\delta(d(\omega \otimes_t B^t, (\Omega_n^*)^c))\right].
 \end{aligned}$$

Then by (5.35) we have, for any $N \geq n$,

$$\begin{aligned}
 \mathbb{E}^{\mathbb{P}}[\widehat{Y}_t \mathbf{1}_E] - \mathbb{E}^{\mathbb{P}^{N,\delta}}[\widehat{Y}_{\bar{\tau}_n} \mathbf{1}_E] & = \mathbb{E}^{\mathbb{P}^{N,\delta}}\left[(\widehat{Y}_t - \widehat{Y}_{\bar{\tau}_n}) \mathbf{1}_{E \cap \{t < \bar{\tau}_n\}}\right] \\
 & \leq Cn\mathbb{E}^{\mathbb{P}^{N,\delta}}\left[\rho_2(\delta + \rho_n(\delta) + 2\eta_n(\delta))\right] + C\rho_n(\delta) + \varepsilon + C\mathbb{P}^{N,\delta}\left[(\Omega_n^*)^c\right] \\
 & \quad + C\mathbb{E}^{\mathbb{P}^{N,\delta}}\left[1 - \psi_n\right] \\
 & \quad + 2\mathbb{E}^{\mathbb{P}^{N,\delta}}\left[\sup_{\bar{\theta}_n^* \leq s \leq \bar{\theta}_n^*} |\widehat{Y}_s - \widehat{Y}_{\bar{\theta}_n^*}| \psi_n \mathbf{1}_E\right] + C\mathbb{E}^{\mathbb{P}^{N,\delta}}\left[h_\delta(d(\omega, (\Omega_n^*)^c))\right] \\
 & \leq Cn\mathcal{E}\left[\rho_2(\delta + \rho_n(\delta) + 2\eta_n(\delta))\right] + C\rho_n(\delta) + \varepsilon + C2^{-n} + C\mathcal{C}(\psi_n < 1) \\
 & \quad + 2\mathbb{E}^{\mathbb{P}^{N,\delta}}\left[\sup_{\bar{\theta}_n^* \leq s \leq \bar{\theta}_n^*} |\widehat{Y}_s - \widehat{Y}_{\bar{\theta}_n^*}| \psi_n \mathbf{1}_E\right] + C\mathcal{E}\left[h_\delta(d(\omega, (\Omega_n^*)^c))\right]. \tag{5.39}
 \end{aligned}$$

Similarly we have

$$\begin{aligned} & \mathbb{E}^{\mathbb{P}^{N,\delta}} \left[[\widehat{Y}_{t \vee \bar{\tau}_n} - \widehat{Y}_{t \vee \theta_n^*} \psi_n] \mathbf{1}_E \right] \\ & \leq C2^{-n} + CC(\psi_n < 1) + \mathbb{E}^{\mathbb{P}^{N,\delta}} \left[[\widehat{Y}_{t \vee \bar{\tau}_n} - \widehat{Y}_{t \vee \theta_n^*} \mathbf{1}_{E \cap \Omega_n^*} \psi_n] \right] \\ & \leq C2^{-n} + CC(\psi_n < 1) + 2\mathbb{E}^{\mathbb{P}^{N,\delta}} \left[\sup_{\theta_n^* \leq s \leq \bar{\theta}_n^*} |\widehat{Y}_s - \widehat{Y}_{\theta_n^*}| \psi_n \mathbf{1}_E \right]. \end{aligned}$$

This, together with (5.39), implies (5.36).

Step 3. Finally we prove the lemma for general stopping time τ . We follow the arguments in Lemma 5.11. Let τ^n be a sequence of stopping times such that $\tau^n \downarrow \tau$ and each τ^n takes only finitely many values. By applying the dominated convergence theorem under \mathbb{P} , we may fix n such that

$$\mathbb{E}^{\mathbb{P}} \left[|\widehat{Y}_{\tau^n \wedge \widehat{\tau}^*} - \widehat{Y}_{\tau} \mathbf{1}_E| \right] \leq \frac{\varepsilon}{2}.$$

Assume τ^n takes values $\{t_i, i = 1, \dots, m\}$, and for each i , denote $E_i := E \cap \{\tau^n = t_i < \widehat{\tau}^*\} \in \mathcal{F}_{t_i}$. Then $\{E_i, 1 \leq i \leq m\}$ form a partition of $\tilde{E} := E \cap \{\tau^n < \widehat{\tau}^*\}$. For each i , by Step 1 there exists $\mathbb{P}^i \in \mathcal{P}(\mathbb{P}, t_i, E_i)$ such that

$$\mathbb{E}^{\mathbb{P}} [\widehat{Y}_{t_i} \mathbf{1}_{E_i}] \leq \mathbb{E}^{\mathbb{P}^i} [\widehat{Y}_{\widehat{\tau}^*} \mathbf{1}_{E_i}] + \frac{\varepsilon}{2m}.$$

Now define $\mathbb{P}_\varepsilon := \sum_{i=1}^m \mathbb{P}^i \mathbf{1}_{E_i} + \mathbb{P} \mathbf{1}_{\tilde{E}^c} \in \mathcal{P}(\mathbb{P}, \tau^n, \tilde{E}) \subset \mathcal{P}(\mathbb{P}, \tau, E)$. Recall that $\tilde{E} \in \mathcal{F}_{\tau^n}$ and note that $\widehat{Y}_{\widehat{\tau}^*} \leq \widehat{Y}_{\widehat{\tau}^*}$, thanks to the supermartingale property of \widehat{Y} . Then

$$\begin{aligned} \mathbb{E}^{\mathbb{P}} [\widehat{Y}_{\tau} \mathbf{1}_E] - \mathbb{E}^{\mathbb{P}_\varepsilon} [\widehat{Y}_{\widehat{\tau}^*} \mathbf{1}_E] & \leq \frac{\varepsilon}{2} + \mathbb{E}^{\mathbb{P}} [\widehat{Y}_{\tau^n \wedge \widehat{\tau}^*} \mathbf{1}_E] - \mathbb{E}^{\mathbb{P}_\varepsilon} [\widehat{Y}_{\widehat{\tau}^*} \mathbf{1}_E] \\ & \leq \frac{\varepsilon}{2} + \mathbb{E}^{\mathbb{P}} [\widehat{Y}_{\tau^n} \mathbf{1}_{\tilde{E}}] - \mathbb{E}^{\mathbb{P}_\varepsilon} [\widehat{Y}_{\widehat{\tau}^*} \mathbf{1}_{\tilde{E}}] \\ & = \frac{\varepsilon}{2} + \sum_{i=1}^m \left(\mathbb{E}^{\mathbb{P}} [\widehat{Y}_{t_i} \mathbf{1}_{E_i}] - \mathbb{E}^{\mathbb{P}_\varepsilon} [\widehat{Y}_{\widehat{\tau}^*} \mathbf{1}_{E_i}] \right) \\ & \leq \frac{\varepsilon}{2} + \sum_{i=1}^m \frac{\varepsilon}{2m} = \varepsilon. \end{aligned}$$

The proof is complete now. ■

We need one more lemma.

Lemma 5.13. *Let $\mathbb{P} \in \mathcal{P}$, $\tau \in \mathcal{T}$, and $E \in \mathcal{F}_\tau$ such that $\tau \leq H$ on E . For any $\varepsilon > 0$, there exists $\mathbb{P}_\varepsilon \in \mathcal{P}(\mathbb{P}, \tau, E)$ such that*

$$H \leq \tau + \frac{1}{L} d(\omega_\tau, O^c) + 3\varepsilon + \sup_{\tau \leq t \leq \tau + \varepsilon} |\omega_t - \omega_\tau|, \quad \mathbb{P}_\varepsilon\text{-a.s. on } E.$$

Proof. First, there exists $\tilde{\tau} \in \mathcal{T}$ such that $\tau \leq \tilde{\tau} \leq \tau + \varepsilon$ and $\tilde{\tau}$ takes only finitely many values $0 \leq t_1 < \dots < t_n = t_0$. Denote $E_i := E \cap \{\tilde{\tau} = t_i < H\} \in \mathcal{F}_{t_i}$. Then $\{E_i, 1 \leq i \leq n\}$ is a partition of $E \cap \{\tilde{\tau} < H\}$ and

$$H \leq \tilde{\tau} \leq \tau + \varepsilon \quad \text{on } E \cap \{\tilde{\tau} \geq H\}. \quad (5.40)$$

For any i , there exists a partition $(E_j^i)_{j \geq 1}$ of E_i such that $|\omega_{t_i} - \omega'_{t_i}| \leq L\varepsilon$ for any $\omega, \omega' \in E_j^i$. For each (i, j) , fix an $\omega^{ij} \in E_j^i$ and a unit vector α^{ij} pointing to the direction from $\omega_{t_i}^{ij}$ to O^c . Now for any $\omega \in E_j^i$, define $\mathbb{P}^{i,j,\omega} \in \mathcal{P}_{t_i}$ as follows:

$$\beta = \mathbf{0}, \quad \alpha_t = \frac{1}{\varepsilon}[\omega_{t_i}^{ij} - \omega_{t_i}]\mathbf{1}_{[t_i, t_i + \varepsilon)}(t) + L\alpha^{ij}\mathbf{1}_{[t_i + \varepsilon, T]}(t).$$

We see that

$$H^{i,\omega} = \left[t_i + \varepsilon + \frac{1}{L}d(\omega_{t_i}^{ij}, O^c) \right] \wedge t_0, \quad \mathbb{P}^{i,j,\omega}\text{-a.s. on } E_j^i.$$

Similar to the proof of (5.12), there exists $\mathbb{P}_\varepsilon \in \mathcal{P}(\mathbb{P}, \tilde{\tau}, E) \subset \mathcal{P}(\mathbb{P}, \tau, E)$ such that the regular conditional probability distribution $\mathbb{P}_\varepsilon^{i,\omega} = \mathbb{P}^{i,j,\omega}$ for \mathbb{P} -a.e. $\omega \in E_j^i$. Then

$$\begin{aligned} H &\leq \tau + 2\varepsilon + \frac{1}{L}[d(\omega_{t_i}, O^c) + L\varepsilon] \leq \tau + 3\varepsilon + \frac{1}{L}[d(\omega_\tau, O^c) + |\omega_\tau - \omega_{t_i}|] \\ &\leq \tau + 3\varepsilon + \frac{1}{L}\left[d(\omega_\tau, O^c) + \sup_{\tau \leq t \leq \tau + \varepsilon} |\omega_t - \omega_\tau|\right], \quad \mathbb{P}_\varepsilon\text{-a.s. on } E_j^i. \end{aligned}$$

This, together with (5.40), proves the lemma. ■

We are now ready to complete the

Proof of Proposition 5.9. The inequality $\mathcal{E}[\widehat{Y}_{\tilde{\tau}^*}] \leq \mathcal{E}[\widehat{Y}_{\tilde{\tau}^* -}]$ is a direct consequence of the \mathcal{E} -supermartingale property of \widehat{Y} established in Theorem 5.4. As for the reverse inequality, since \widehat{Y} is continuous on $[0, H)$ and $H_n \uparrow H$ with $H_n < H$, it suffices to show that, for any $\mathbb{P} \in \mathcal{P}$ and any $\varepsilon > 0$

$$I_n := \mathbb{E}^\mathbb{P}[\widehat{Y}_{\tilde{\tau}^* \wedge H_n}] - \mathcal{E}[\widehat{Y}_{\tilde{\tau}^*}] \leq 5\varepsilon \quad \text{for sufficiently large } n. \quad (5.41)$$

Let $\delta > 0, n > \frac{1}{L\delta}$. Set $t_n := t_0 - \frac{1}{n}$, $\tau^0 := \tilde{\tau}^* \wedge H_n$, and $\mathbb{P}^0 := \mathbb{P}$. We proceed in two steps.

Step 1. Apply Lemma 5.11 with $\mathbb{P}^0, \tau^0, \tilde{\tau}^*$, and Ω , there exist $\mathbb{P}^{1,1} \in \mathcal{P}(\mathbb{P}^0, \tau^0, \Omega)$ and a stopping time $\tilde{\tau}^1$ taking values in $[\tau^0, \tilde{\tau}^*]$, such that

$$\mathbb{E}^{\mathbb{P}^0}[\widehat{Y}_{\tau^0}] \leq \mathbb{E}^{\mathbb{P}^{1,1}}\left[\widehat{X}_{\tilde{\tau}^1}\mathbf{1}_{\{\tilde{\tau}^1 < \tilde{\tau}^*\}} + \widehat{Y}_{\tilde{\tau}^*}\mathbf{1}_{\{\tilde{\tau}^1 = \tilde{\tau}^*\}}\right] + \varepsilon.$$

Denote $E_1 := \{\tilde{\tau}^1 < t_n\} \in \mathcal{F}_{\tilde{\tau}^1}$. By (5.3) and following the same argument as for the estimate in (4.6), we have: $\mathbb{P}^{1,1}$ -a.s. on $E_1^c \cap \{\tilde{\tau}^1 < \tilde{\tau}^*\}$,

$$\begin{aligned} \widehat{X}_{\tilde{\tau}^1} &\leq \widehat{X}_{\tilde{\tau}^1} - \mathbb{E}^{\mathbb{P}^{1,1}}[\widehat{X}_{\tilde{\tau}^*}] + \mathbb{E}^{\mathbb{P}^{1,1}}[\widehat{Y}_{\tilde{\tau}^*}] \\ &\leq \mathbb{E}^{\mathbb{P}^{1,1}}\left[\rho_0\left(\frac{1}{n} + \|B^{\tilde{\tau}^1}\|_{\tilde{\tau}^1 + \frac{1}{n}}\right)\right] + \mathbb{E}^{\mathbb{P}^{1,1}}[\widehat{Y}_{\tilde{\tau}^*}] \leq C\bar{\rho}_0(n^{-1}) + \mathbb{E}^{\mathbb{P}^{1,1}}[\widehat{Y}_{\tilde{\tau}^*}]. \end{aligned}$$

Then, denoting $E_2 := E_1 \cap \{\tilde{\tau}^1 < \tilde{\tau}^*\} \in \mathcal{F}_{\tilde{\tau}^1}$, we get:

$$\begin{aligned} \mathbb{E}^{\mathbb{P}^0}[\widehat{Y}_{\tau^0}] &\leq \mathbb{E}^{\mathbb{P}^{1,1}}\left[\widehat{X}_{\tilde{\tau}^1}\mathbf{1}_{E_2} + \widehat{X}_{\tilde{\tau}^1}\mathbf{1}_{E_1^c \cap \{\tilde{\tau}^1 < \tilde{\tau}^*\}} + \widehat{Y}_{\tilde{\tau}^*}\mathbf{1}_{\{\tilde{\tau}^1 = \tilde{\tau}^*\}}\right] + \varepsilon \\ &\leq \mathbb{E}^{\mathbb{P}^{1,1}}\left[\widehat{X}_{\tilde{\tau}^1}\mathbf{1}_{E_2} + \widehat{Y}_{\tilde{\tau}^*}\mathbf{1}_{E_2^c}\right] + C\bar{\rho}_0(n^{-1})\mathbb{P}^0[E_1^c] + \varepsilon. \end{aligned} \quad (5.42)$$

Next, set $\tilde{\delta} := [\delta^2 \bar{\rho}_0(3\delta)] \wedge \frac{\delta}{3}$. Apply Lemma 5.13 on $\mathbb{P}^{1,1}$, $\tilde{\tau}^1$, E_2 , and $\tilde{\delta}$, there exists $\mathbb{P}^{1,2} \in \mathcal{P}(\mathbb{P}^{1,1}, \tilde{\tau}^1, E_2)$ such that

$$H \leq \tilde{\tau}^1 + \frac{1}{L} d(\omega_{\tilde{\tau}^1}, O^c) + \delta + \|\omega_{\tilde{\tau}^1}^{\tilde{\tau}^1}\|_{\tilde{\tau}^1 + \tilde{\delta}}, \quad \mathbb{P}^{1,2}\text{-a.s. on } E_2.$$

Since $\tilde{\tau}^1 \leq \hat{\tau}^* \leq H$, we have

$$\hat{\tau}^* - \tilde{\tau}^1 \leq 3\delta, \quad \mathbb{P}^{1,2}\text{-a.s. on } E_2 \cap \{d(\omega_{\tilde{\tau}^1}, O^c) \leq L\delta\} \cap \{\|\omega_{\tilde{\tau}^1}^{\tilde{\tau}^1}\|_{\tilde{\tau}^1 + \tilde{\delta}} \leq \delta\}.$$

Then, by (5.3) and (4.6) again we have: $\mathbb{P}^{1,2}$ -a.s. on $E_2 \cap \{d(\omega_{\tilde{\tau}^1}, O^c) \leq L\delta\} \in \mathcal{F}_{\tilde{\tau}^1}$,

$$\begin{aligned} \hat{X}_{\tilde{\tau}^1} &\leq \mathbb{E}_{\tilde{\tau}^1}^{\mathbb{P}^{1,2}}[\hat{X}_{\hat{\tau}^*}] + \mathbb{E}_{\tilde{\tau}^1}^{\mathbb{P}^{1,2}}\left[\rho_0\left(\mathbf{d}_{\infty}((\tilde{\tau}^1, B), (\hat{\tau}^*, B))\right)\right] \\ &= \mathbb{E}_{\tilde{\tau}^1}^{\mathbb{P}^{1,2}}[\hat{X}_{\hat{\tau}^*}] + \mathbb{E}_{\tilde{\tau}^1}^{\mathbb{P}^{1,2}}\left[\rho_0\left(\mathbf{d}_{\infty}((\tilde{\tau}^1, B), (\hat{\tau}^*, B))\right)\left[\mathbf{1}_{\{\|B^{\tilde{\tau}^1}\|_{\tilde{\tau}^1 + \tilde{\delta}} \leq \delta\}} + \mathbf{1}_{\{\|B^{\tilde{\tau}^1}\|_{\tilde{\tau}^1 + \tilde{\delta}} > \delta\}}\right]\right] \\ &\leq \mathbb{E}_{\tilde{\tau}^1}^{\mathbb{P}^{1,2}}[\hat{X}_{\hat{\tau}^*}] + \mathbb{E}_{\tilde{\tau}^1}^{\mathbb{P}^{1,2}}\left[\rho_0\left(3\delta + \|B^{\tilde{\tau}^1}\|_{\tilde{\tau}^1 + 3\delta}\right)\right] + C\delta^{-2}\mathbb{E}_{\tilde{\tau}^1}^{\mathbb{P}^{1,2}}[\|B^{\tilde{\tau}^1}\|_{\tilde{\tau}^1 + \tilde{\delta}}^2] \\ &\leq \mathbb{E}_{\tilde{\tau}^1}^{\mathbb{P}^{1,2}}[\hat{X}_{\hat{\tau}^*}] + C\bar{\rho}_0(3\delta) + \frac{C\tilde{\delta}}{\delta^2} \leq \mathbb{E}_{\tilde{\tau}^1}^{\mathbb{P}^{1,2}}[\hat{X}_{\hat{\tau}^*}] + C\bar{\rho}_0(3\delta). \end{aligned}$$

Note that $n^{-1} \leq L\delta \leq 3\delta$. Thus, denoting $E_3 := E_2 \cap \{d(\omega_{\tilde{\tau}^1}, O^c) > L\delta\} \in \mathcal{F}_{\tilde{\tau}^1}$, (5.42) leads to:

$$\mathbb{E}^{\mathbb{P}^0}[\hat{Y}_{\tau^0}] \leq \mathbb{E}^{\mathbb{P}^{1,2}}[\hat{X}_{\tilde{\tau}^1} \mathbf{1}_{E_3} + \hat{Y}_{\tilde{\tau}^1} \mathbf{1}_{E_3^c}] + C\bar{\rho}_0(3\delta)\mathbb{P}^{1,2}(E_3^c) + \varepsilon. \quad (5.43)$$

Moreover, apply Lemma 5.12 with $\mathbb{P}^{1,2}$, $\tilde{\tau}^1$, E_3 , and ε , there exists $\mathbb{P}^{1,3} \in \mathcal{P}(\mathbb{P}^{1,2}, \tilde{\tau}^1, E_3)$ such that

$$\mathbb{E}^{\mathbb{P}^{1,2}}[\hat{X}_{\tilde{\tau}^1} \mathbf{1}_{E_3}] \leq \mathbb{E}^{\mathbb{P}^{1,2}}[\hat{Y}_{\tilde{\tau}^1} \mathbf{1}_{E_3}] \leq \mathbb{E}^{\mathbb{P}^{1,3}}[\hat{Y}_{\tilde{\tau}^1} \mathbf{1}_{E_3}] + \varepsilon.$$

Define $\tau^1 := \inf\{t \geq \tilde{\tau}^1 : d(\omega_t, O^c) \leq \frac{1}{n}\} \wedge \hat{\tau}^*$. Note that $\tau^1 < H$ on E_3 and \hat{Y} is a $\mathbb{P}^{1,3}$ -supermartingale. Then

$$\mathbb{E}^{\mathbb{P}^{1,3}}[\hat{Y}_{\tilde{\tau}^1} \mathbf{1}_{E_3}] \leq \mathbb{E}^{\mathbb{P}^{1,3}}[\hat{Y}_{\tau^1} \mathbf{1}_{E_3}].$$

Thus

$$\mathbb{E}^{\mathbb{P}^{1,2}}[\hat{X}_{\tilde{\tau}^1} \mathbf{1}_{E_3}] \leq \mathbb{E}^{\mathbb{P}^{1,3}}[\hat{Y}_{\tau^1} \mathbf{1}_{E_3}] + \varepsilon.$$

Plug this into (5.43), we obtain

$$\mathbb{E}^{\mathbb{P}^0}[\hat{Y}_{\tau^0}] \leq \mathbb{E}^{\mathbb{P}^{1,3}}[\hat{Y}_{\tau^1} \mathbf{1}_{E_3} + \hat{Y}_{\tilde{\tau}^1} \mathbf{1}_{E_3^c}] + C\bar{\rho}_0(3\delta)\mathbb{P}^{1,3}(E_3^c) + 2\varepsilon.$$

We now denote $\mathbb{P}^1 := \mathbb{P}^{1,3} \in \mathcal{P}(\mathbb{P}^0, \tau^0, \Omega)$, and

$$D_1 := E_3 \cap \{\tau^1 < \hat{\tau}^*\} = \{\tau^1 < t_n \wedge \tilde{\tau}^*\} \cap \{d(\omega_{\tilde{\tau}^1}, O^c) > L\delta\} \cap \{\tau^1 < \hat{\tau}^*\} \in \mathcal{F}_{\tau^1}. \quad (5.44)$$

Then

$$\mathbb{E}^{\mathbb{P}^0}[\hat{Y}_{\tau^0}] \leq \mathbb{E}^{\mathbb{P}^1}[\hat{Y}_{\tau^1} \mathbf{1}_{D_1} + \hat{Y}_{\tilde{\tau}^1} \mathbf{1}_{D_1^c}] + C\bar{\rho}_0(3\delta)\mathbb{P}^1(D_1^c) + 2\varepsilon. \quad (5.45)$$

Step 3: Iterating the arguments of Step 1, we may define $(\tilde{\tau}^m, \tau^m, \mathbb{P}^m, D_m)_{m \geq 1}$ such that:

$$\mathbb{P}^{m+1} \in \mathcal{P}(\mathbb{P}^m, \tau^m, D_m), \quad \tau^m \leq \tilde{\tau}^{m+1} \leq \hat{\tau}^*;$$

$$\tau^{m+1} := \inf \left\{ t \geq \tilde{\tau}^{m+1} : d(\omega_t, O^c) \leq \frac{1}{n} \right\} \wedge \hat{\tau}^*$$

$$D_{m+1} := D_m \cap \{\tilde{\tau}^{m+1} < t_n \wedge \hat{\tau}^*\} \cap \{d(\omega_{\tilde{\tau}^{m+1}}, O^c) > L\delta\} \cap \{\tau^{m+1} < \hat{\tau}^*\};$$

and

$$\begin{aligned} \mathbb{E}^{\mathbb{P}^m} [\hat{Y}_{\tau^m} \mathbf{1}_{D_m}] &\leq \mathbb{E}^{\mathbb{P}^{m+1}} [\hat{Y}_{\tau^{m+1}} \mathbf{1}_{D_{m+1}} + \hat{Y}_{\tilde{\tau}^*} \mathbf{1}_{D_m \cap D_{m+1}^c}] + C \bar{\rho}_0(3\delta) \mathbb{P}^{m+1}(D_m \cap D_{m+1}^c) \\ &\quad + 2^{1-m} \varepsilon. \end{aligned}$$

By induction, for any $m \geq 1$ we have

$$\begin{aligned} \mathbb{E}^{\mathbb{P}^0} [\hat{Y}_{\tau^0}] &\leq \mathbb{E}^{\mathbb{P}^m} [\hat{Y}_{\tau^m} \mathbf{1}_{D_m} + \hat{Y}_{\tilde{\tau}^*} \mathbf{1}_{D_m^c}] + C \bar{\rho}_0(3\delta) \mathbb{P}^m(D_m^c) + 4\varepsilon \\ &\leq \mathbb{E}^{\mathbb{P}^m} [\hat{Y}_{\tilde{\tau}^*}] + 2C_0 \mathbb{P}^m[D_m] + C \bar{\rho}_0(3\delta) + 4\varepsilon. \end{aligned} \quad (5.46)$$

Note that

$$\begin{aligned} \mathbb{P}^m[D_m] &\leq \mathbb{P}^m \left[\bigcap_{i=1}^m \left\{ |B_{\tilde{\tau}^i} - B_{\tau^{i-1}}| \geq L\delta - \frac{1}{n} \right\} \cap \left\{ |B_{\tau^i} - B_{\tilde{\tau}^i}| \geq L\delta - \frac{1}{n} \right\} \right] \\ &\leq \mathbb{P}^m \left[\sum_{i=1}^m [|B_{\tilde{\tau}^i} - B_{\tau^{i-1}}|^2 + |B_{\tau^i} - B_{\tilde{\tau}^i}|^2] \geq 2m \left(L\delta - \frac{1}{n} \right)^2 \right] \\ &\leq \frac{1}{2m \left(L\delta - \frac{1}{n} \right)^2} \mathbb{E}^{\mathbb{P}^m} \left[\sum_{i=1}^m [|B_{\tilde{\tau}^i} - B_{\tau^{i-1}}|^2 + |B_{\tau^i} - B_{\tilde{\tau}^i}|^2] \right] \leq \frac{C}{2m \left(L\delta - \frac{1}{n} \right)^2}. \end{aligned}$$

Then, (5.46) leads to

$$I_n \leq \frac{C}{2m \left(L\delta - \frac{1}{n} \right)^2} + C \bar{\rho}_0(3\delta) + 4\varepsilon,$$

which implies, by sending $m \rightarrow \infty$ that

$$I_n \leq C \bar{\rho}_0(3\delta) + 4\varepsilon.$$

Hence, by choosing δ small enough such that $\bar{\rho}_0(3\delta) \leq \varepsilon$, we see that (5.41) holds true for $n > \frac{1}{L\delta}$. ■

Uncited references

[12] and [14].

Acknowledgements

The second author's research was supported by the Chair *Financial Risks* of the *Risk Foundation* sponsored by Société Générale, and the Chair *Finance and Sustainable Development* sponsored by EDF and Calyon. The third author's research was supported in part by NSF grant DMS 10-08873.

Appendix

A.1. Regular conditional probability distribution

We first recall the definition of r.c.p.d. from Stroock–Varadhan [18]. Let $(\Omega, \mathcal{F}, \mathbb{P})$ be a probability space, $\mathcal{G} \subset \mathcal{F}$ is a sub- σ -field. An r.c.p.d. $\{\mathbb{P}_\omega^\mathcal{G}, \omega \in \Omega\}$ is a family of probability measures on \mathcal{F} satisfying the following requirements:

- For all $E \in \mathcal{F}$, the mapping $\omega \rightarrow \mathbb{P}_\omega^\mathcal{G}(E)$ is \mathcal{G} -measurable;
- For all $\xi \in \mathbb{L}^\infty(\mathcal{F})$, the conditional expectation $\mathbb{E}^{\mathbb{P}_\omega^\mathcal{G}}[\xi] = \mathbb{E}[\xi|\mathcal{G}](\omega)$, for \mathbb{P} -a.e. ω ;
- For any $\omega \in \Omega$, $\mathbb{P}_\omega^\mathcal{G}(\Omega_\mathcal{G}^\omega) = 1$, where $\Omega_\mathcal{G}^\omega := \cap \{E \in \mathcal{G} : \omega \in E\}$.

We note that an r.c.p.d. exists whenever \mathcal{G} is countably generated.

In the special case that $\Omega := \{\omega \in C([0, T], \mathbb{R}^d) : \omega_0 = \mathbf{0}\}$ is our canonical space and $\mathcal{G} = \mathcal{F}_\tau$ for some $\tau \in \mathcal{T}$, it holds that

$$\Omega_{\mathcal{F}_\tau}^\omega := \{\omega' \in \Omega : \tau(\omega') = \tau(\omega) \text{ and } \omega'_{\wedge \tau} = \omega_{\wedge \tau}\} = \{\omega \otimes_{\tau(\omega)} \omega' : \omega' \in \Omega^{\tau(\omega)}\}. \quad (\text{A.1})$$

Then, as in [16], we define for all $\omega \in \Omega$ a probability measure $\mathbb{P}^{\tau, \omega}$ on $\mathcal{F}_T^{\tau(\omega)}$ by:

$$\mathbb{P}^{\tau, \omega}(E) := \mathbb{P}_\omega^\omega(\{\omega \otimes_{\tau(\omega)} \omega' : \omega' \in E\}), \quad \forall E \in \mathcal{F}_T^{\tau(\omega)}, \quad (\text{A.2})$$

and still call it an r.c.p.d. of \mathbb{P} conditional on \mathcal{F}_τ . One may easily check that $\omega \mapsto \mathbb{E}^{\mathbb{P}^{\tau, \omega}}[\xi^{\tau, \omega}]$ is \mathcal{F}_τ -measurable, for all $\xi \in \mathbb{L}^\infty(\mathcal{F}_T)$, and $\mathbb{E}^{\mathbb{P}^{\tau, \omega}}[\xi^{\tau, \omega}] = \mathbb{E}[\xi|\mathcal{F}_\tau](\omega)$, for \mathbb{P} -a.e. ω and for all $\xi \in \mathbb{L}^\infty(\mathcal{F})$.

A.2. Proof of Lemma 2.3

Recall the notations in the beginning of Section 2.2. Let $\mathbb{F} := \mathbb{F}^B$ and $\tilde{\mathbb{F}} := \tilde{\mathbb{F}}^{\tilde{B}}$ be the natural filtrations on Ω and $\tilde{\Omega}$, respectively. Moreover, we may identify \mathbb{F} with the filtration $\tilde{\mathbb{F}}^B$ on $\tilde{\Omega}$ generated by B : $\tilde{\mathcal{F}}_t^B = \{E \times \Omega^2 : E \in \mathcal{F}_t^B\}$.

- First, it follows from standard arguments, see e.g. Zheng [19, Theorem 3], that \mathcal{P}_t^L is weakly compact. Then Property (P1) holds.
- We next check without loss of generality Property (P2) only at $t = 0$. Let $\tau \in \mathcal{T}$ and $\mathbb{P} \in \mathcal{P}^L$ with corresponding \mathbb{Q} as in (2.2). Define $\tilde{\tau}(\tilde{\omega}) := \tau(\omega)$ for $\tilde{\omega} := (\omega, a, m) \in \tilde{\Omega}$, then clearly $\tilde{\tau}$ is an $\tilde{\mathbb{F}}^B$ -stopping time, hence also an $\tilde{\mathbb{F}}$ -stopping time. By Stroock–Varadhan [18], the r.c.p.d. $\mathbb{Q}_{\tilde{\mathcal{F}}_{\tilde{\tau}}^B}^{\tilde{\omega}}$ exists. Note that $\tilde{\omega} \mapsto \mathbb{Q}_{\tilde{\mathcal{F}}_{\tilde{\tau}}^B}^{\tilde{\omega}}(E)$ is $\tilde{\mathcal{F}}_{\tilde{\tau}}^B$ -measurable for any $E \in \tilde{\mathcal{F}}_T$, it follows that $\mathbb{Q}_{\tilde{\mathcal{F}}_{\tilde{\tau}}^B}^{\tilde{\omega}}$ depends only on ω and thus we may denote it as $\mathbb{Q}_{\tilde{\mathcal{F}}_{\tilde{\tau}}^B}^\omega$.

Recall the shifted spaces Ω^t , $\tilde{\Omega}^t$, \mathbb{F}^t , and $\tilde{\mathbb{F}}^t$. We now define the following probability measure on the shifted space $\Omega^{\tau(\omega)}$:

$$\begin{aligned} \mathbb{Q}^{\tau, \omega}[\tilde{E}] &:= \mathbb{Q}_{\tilde{\mathcal{F}}_{\tilde{\tau}}^B}^\omega(\tilde{\omega}^1 \otimes_{\tau(\omega)} \tilde{\omega}^2 : \tilde{\omega}^1 \in \tilde{\Omega}, \tilde{\omega}^2 \in \tilde{E}), \quad \forall \tilde{E} \in \tilde{\mathcal{F}}_T^{\tau(\omega)}; \\ \mathbb{P}^{\tau, \omega}[E] &:= \mathbb{Q}^{\tau, \omega}[E \times (\Omega^{\tau(\omega)})^2], \quad \forall E \in \mathcal{F}_T^{\tau(\omega)}. \end{aligned} \quad (\text{A.3})$$

It is straightforward to check that $\mathbb{P}^{\tau, \omega}$ is an r.c.p.d. of \mathbb{P} conditional on \mathcal{F}_τ , and $\mathbb{Q}^{\tau, \omega}$ is the required extension on $\tilde{\Omega}^{\tau(\omega)}$ satisfying (2.2) for \mathbb{P} -a.e. ω . This verifies (P2).

(iii) It remains to check Property (P3). Assume \mathbb{Q} and \mathbb{Q}^i are the corresponding extensions of \mathbb{P} and \mathbb{P}^i . Define

$$\hat{\mathbb{Q}} := \mathbb{Q} \otimes_t \left[\sum_{i=1}^{\infty} \mathbb{Q}^i \mathbf{1}_{E_i \times (\Omega^s)^2} + \mathbb{Q} \mathbf{1}_{\cap_{i=1}^{\infty} (E_i^c \times (\Omega^s)^2)} \right].$$

Following similar arguments as in (ii) one can show that $\hat{\mathbb{Q}}$ satisfies (2.2). It is clear that $\hat{\mathbb{P}}(E) = \hat{\mathbb{Q}}(E \times (\Omega^s)^2)$ for all $E \in \mathcal{F}_T^s$. Then $\hat{\mathbb{P}} \in \mathcal{P}_s^L$ and thus (P3) holds. ■

A.3. Some additional results

In this subsection we provide some results which are interesting for our discussion, although they are technically not used in the paper.

Proof of Remark 3.2. Fix $\omega \in \Omega$, and let $\{t_n\}$ and $\{s_n\}$ be two sequences such that $t_n \uparrow t$, $s_n \uparrow t$, and $X_{t_n}(\omega) \rightarrow \lim_{s \uparrow t} X_s(\omega)$, $X_{s_n}(\omega) \rightarrow \lim_{s \uparrow t} X_s(\omega)$. Here and in the sequel, in $\lim_{s \uparrow t}$ we take the notational convention that $s < t$. Without loss of generality, we may assume $t_n < s_n < t_{n+1}$ for $n = 1, 2, \dots$. Then for the ρ_0 defined in (3.1) we have

$$\begin{aligned} 0 &\leq \overline{\lim}_{s \uparrow t} X_s(\omega) - \underline{\lim}_{s \uparrow t} X_s(\omega) = \lim_{n \rightarrow \infty} X_{t_n}(\omega) - \lim_{n \rightarrow \infty} X_{s_n}(\omega) \\ &\leq \underline{\lim}_{n \rightarrow \infty} \rho_0(\mathbf{d}_{\infty}((t_n, \omega), (s_n, \omega))) = 0. \end{aligned}$$

This implies the existence of $X_{t-}(\omega)$. Moreover,

$$X_{t-}(\omega) - X_t(\omega) = \lim_{s \uparrow t} X_s(\omega) - X_t(\omega) \leq \underline{\lim}_{s \uparrow t} \rho(\mathbf{d}_{\infty}((s, \omega), (t, \omega))) = 0,$$

completing the proof. ■

Lemma A.1. Let the nondegeneracy condition (3.7) hold and X be bounded and uniformly continuous in (t, ω) under d_{∞} . Then \hat{Y}^H defined in (3.5) is left continuous at H .

Proof. We first claim that, for any $\omega \in \Omega$ and $\varepsilon > 0$

$$\lim_{t \uparrow H(\omega)} \mathbb{C}_t[H^{t, \omega} \geq t + \varepsilon] = 0. \quad (\text{A.4})$$

Indeed, let H correspond to O and t_0 as in (3.3). If $H(\omega) = t_0$, since $H^{t, \omega} \leq t_0$, (A.4) is obvious. We now assume $t_1 := H(\omega) < t_0$ and thus $\omega_{t_1} \in O^c$. Note that $t < H(\omega)$ implies $\omega_t \in O$. Denote $\delta := d(\omega_t, O^c)$, then $0 < \delta \leq |\omega_t - \omega_{t_1}|$. Let η be a unit vector pointing to the direction from ω_t to O^c . Since O is convex, we see that

$$\text{for any } x \in \mathbb{R}^d, x \cdot \eta \geq \delta \text{ implies } x + \omega_t \in O^c. \quad (\text{A.5})$$

Since we will send $t \uparrow t_1$, we may assume $\delta \leq \varepsilon$. Then, for any $\mathbb{P} \in \mathcal{P}_t$ with corresponding α, β , and W , we have

$$\mathbb{P}(H^{t, \omega} \geq t + \varepsilon) \leq \mathbb{P}(H^{t, \omega} \geq t + \delta) \leq \mathbb{P}\left(\sup_{t \leq s \leq t + \delta} (B_s^t \cdot \eta) < \delta\right) \leq \mathbb{P}\left(\sup_{t \leq s \leq t + \delta} M_s \leq C\delta\right)$$

where $M_s := \int_t^s \beta_r dW_r \cdot \eta$ is a scalar \mathbb{P} -martingale. Denote $A_s := \int_t^s |\beta_r \eta|^2 dr$ and introduce the time change: $\tau_r := \inf\{s \geq t : A_s \geq r - t\}$ and $N_r := M_{\tau_r}$. Then N is a \mathbb{P} -Brownian motion.

Since $\beta \geq cI_d$, then $c^2(\tau_r - t) \leq r - t$, and thus

$$\mathbb{P}(H^{t,\omega} \geq t + \varepsilon) \leq \mathbb{P}\left(\sup_{t \leq s \leq t+\delta} M_s \leq C\delta\right) \leq \mathbb{P}\left(\sup_{t \leq r \leq t+c^2\delta} N_r \leq C\delta\right) = C\sqrt{\delta},$$

where C is independent of \mathbb{P} . Then $C_t[H^{t,\omega} \geq t + \varepsilon] \leq C\sqrt{\delta}$ for $\delta \leq \varepsilon$. Now send $t \uparrow H(\omega)$, we have $\delta \rightarrow 0$ and thus (A.4) holds.

We now prove the lemma. Let ρ denote the modulus of continuity function of X . Note that in this case $\hat{X}^H = X_{H \wedge \cdot}$. Fix $\omega \in \Omega$. For $t < t_1 := H(\omega)$ and $\varepsilon > 0$, denoting $E := \{H^{t,\omega} \leq t + \varepsilon\} \cap \{\|B^t\|_{t+\varepsilon} \leq \varepsilon^{\frac{1}{3}}\}$, we have

$$\begin{aligned} |\hat{Y}_t^H(\omega) - \hat{Y}_H^H(\omega)| &\leq \sup_{\tau \in T^t} \mathcal{E}_t \left[|X_{\tau \wedge H^{t,\omega}}^{t,\omega} - X_H(\omega)| \right] \\ &\leq CC_t[E^c] + \sup_{\tau \in T^t} \mathcal{E}_t \left[\rho(d_\infty((\tau \wedge H^{t,\omega}, \omega \otimes_t B^t), (t_1, \omega))) \mathbf{1}_E \right] \\ &\leq CC_t[H^{t,\omega} \geq t + \varepsilon] + CC_t[\|B^t\|_{t+\varepsilon} \geq \varepsilon^{\frac{1}{3}}] \\ &\quad + \mathcal{E}_t \left[\rho(d_\infty((H^{t,\omega}, \omega \otimes_t B^t), (t, \omega)) + d_\infty((t, \omega), (t_1, \omega))) \mathbf{1}_E \right] \\ &\leq CC_t[H^{t,\omega} \geq t + \varepsilon] + C\varepsilon^{\frac{1}{3}} + \rho\left(\varepsilon + \varepsilon^{\frac{1}{3}} + d_\infty((t, \omega), (t_1, \omega))\right). \end{aligned}$$

Then, by (A.4) we have

$$\lim_{t \uparrow H(\omega)} |\hat{Y}_t^H(\omega) - \hat{Y}_H^H(\omega)| \leq C\varepsilon^{\frac{1}{3}} + \rho\left(\varepsilon + \varepsilon^{\frac{1}{3}}\right).$$

Since ε is arbitrary, we prove the result. ■

However, in the degenerate case in general \hat{Y}^H may be discontinuous at H .

Example A.2. Set $X_t(\omega) := t$ and let H correspond to O and t_0 . Clearly $\hat{X}^H = X$, $\hat{Y}_H^H = H$ and $\hat{Y}_t^H(\omega) \leq t_0$. However, for any $t < H(\omega)$, set $\tau := t_0$ and $\mathbb{P} \in \mathcal{P}_t$ such that $\alpha^\mathbb{P} = 0$, $\beta^\mathbb{P} = 0$, we see that $\hat{Y}_t^H(\omega) \geq \mathbb{E}^\mathbb{P} \left[X(H(\omega \otimes_t B^t), \omega \otimes_t B^t) \right] = X(H(\omega_{\cdot \wedge t}), \omega_{\cdot \wedge t}) = H(\omega_{\cdot \wedge t}) = t_0$. That is, $\hat{Y}_t^H(\omega) = t_0$. Thus \hat{Y}^H is discontinuous at H whenever $H(\omega) < t_0$. ■

References

- [1] E. Bayraktar, I. Karatzas, S. Yao, Optimal stopping for dynamic convex risk measures, *Illinois J. Math.* 54 (3) (2010) 1025–1067.
- [2] M.G. Crandall, H. Ishii, P.-L. Lions, User's guide to viscosity solutions of second order partial differential equations, *Bull. Amer. Math. Soc. (N.S.)* 27 (1992) 1–67.
- [3] L. Denis, M. Hu, S. Peng, Function spaces and capacity related to a sublinear expectation: application to G -Brownian motion paths, *Potential Anal.* 34 (2) (2011) 139–161.
- [4] L. Denis, C. Martini, A theoretical framework for the pricing of contingent claims in the presence of model uncertainty, *Ann. Appl. Probab.* 16 (2) (2006) 827–852.
- [5] I. Ekren, C. Keller, N. Touzi, J. Zhang, On viscosity solutions of path dependent PDEs, *Ann. Probab.* 42 (2014) 204–236.
- [6] I. Ekren, N. Touzi, J. Zhang, Viscosity solutions of fully nonlinear parabolic path dependent PDEs: Part I, Preprint, [arXiv:1210.0006](https://arxiv.org/abs/1210.0006).
- [7] I. Ekren, N. Touzi, J. Zhang, Viscosity solutions of fully nonlinear parabolic path dependent PDEs: Part II, Preprint [arXiv:1210.0007](https://arxiv.org/abs/1210.0007).
- [8] N. El Karoui, C. Kapoudjian, E. Pardoux, S. Peng, M.C. Quenez, Reflected solutions of backward SDE's, and related obstacle problems for PDE's, *Ann. Probab.* 25 (2) (1997) 702–737.

- [9] W. Fleming, H.M. Soner, Controlled Markov Processes and Viscosity Solutions, second ed., Springer, New York, 2006. 1
- [10] N.V. Krylov, Controlled Diffusion Processes, Springer-Verlag, 1980. 2
- [11] M. Nutz, R. van Handel, Constructing sublinear expectations on path space, Stochastic Process. Appl. 123 (2013) 3100–3121. 3
- [12] E. Pardoux, S. Peng, Adapted solutions of backward stochastic differential equations, Systems Control Lett. 14 (1990) 55–61. 4
- [13] S. Peng, G -Brownian motion and dynamic risk measure under volatility uncertainty, 2007, Preprint, [arXiv:0711.2834](https://arxiv.org/abs/0711.2834). 5
- [14] S. Peng, Backward stochastic differential equation, nonlinear expectation and their applications, in: Proceedings of the International Congress of Mathematicians, Hyderabad, India, 2010. 6
- [15] M. Soner, N. Touzi, J. Zhang, Wellposedness of second order BSDEs, Probab. Theory Related Fields 153 (2012) 149–190. 7
- [16] M. Soner, N. Touzi, J. Zhang, Dual formulation of second order target problems, Ann. Appl. Probab. 23 (2013) 308–347. 8
- [17] Y. Song, Properties of hitting times for G -martingales and their applications, Stochastic Process. Appl. 121 (2011) 1770–1784. 9
- [18] D.W. Stroock, S.R.S. Varadhan, Multidimensional Diffusion Processes, Springer-Verlag, Berlin, Heidelberg, New York, 1979. 10
- [19] W. Zheng, Tightness results for laws of diffusion processes application to stochastic mechanics, Ann. Inst. Henri Poincaré Probab. Stat. 21 (2) (1985) 103–124. 11