



Transforming spatial point processes into Poisson processes

Frederic Schoenberg ^{*,1}

Department of Statistics, University of California at Los Angeles, Los Angeles, CA 90095-1554, USA

Received 23 March 1998; received in revised form 16 November 1998; accepted 27 November 1998

Abstract

In 1986, Merzbach and Nualart demonstrated a method of transforming a two-parameter point process into a planar Poisson process of unit rate, using random stopping sets. Merzbach and Nualart's theorem applies only to a special class of point processes, since it requires two restrictive conditions: (F4) condition of conditional independence and the convexity of the 1-compensator. (F4) condition was removed in 1990 by Nair, but the convexity condition remained. Here both (F4) condition and the convexity condition are removed by making use of predictable sets rather than stopping sets. As with Nair's theorem, the result extends to point processes in higher dimensions. © 1999 Elsevier Science B.V. All rights reserved.

Keywords: Compensator; Intensity; Point process; Poisson process; Predictable set; Random space change; Spatial process; Stopping time

1. Introduction

Suppose N is a point process. Is it possible to rescale the domain in such a way that N is transformed into a Poisson process with rate 1?

When N is a simple point process on the line, the question is answered by Papangelou (1972) and by Brémaud (1972), using the characterization of the Poisson process of Watanabe (1964). Provided its compensator is continuous, any such N can be transformed into a unit rate Poisson process via random stopping times.

Now suppose that N is a multivariate point process, i.e. a countable sequence of point processes on the line. Meyer (1971) shows that provided an orthogonality condition is satisfied, N can be transformed into a sequence of independent unit rate Poisson processes on the line. This result is also proven by Aalen and Hoem (1978) for the self-exciting case, and an elegant proof by Brown and Nair (1988) generalizes Meyer's result to include a wide class of multivariate point processes.

When N is a point process on the plane, the situation is more complex. Merzbach and Nualart (1986) show that N can be transformed into a unit rate Poisson process using

^{*} E-mail address: frederic@stat.ucla.edu (F. Schoenberg)

¹ Research supported by Michel and Line Loève fellowship and by the University of California Campus-Laboratory Collaboration program "Advanced Earthquake Hazard Research".

random stopping sets, provided several conditions are met. For instance, N must satisfy the conditional independence condition (F4) of Cairoli and Walsh (1975). The need for this condition is removed by Nair (1990). However, both Merzbach and Nualart (1986) and Nair (1990) assume the convexity of the 1-compensator. This convexity condition is rather stringent; for example self-exciting point processes generally do not satisfy this condition.

The current paper investigates transformations based on \mathcal{F}^1 -predictable sets rather than stopping sets and eliminates the need for the conditions mentioned above. However, the existence of the \mathcal{F}^1 - and \mathcal{F}^2 -intensities is assumed, which is also required by Merzbach and Nualart (1986) but not by Nair (1990). As with the result of Nair (1990), the result extends to the case where N is a point process in \mathbb{R}^k , for $k > 2$.

The next section introduces planar point processes, predictable sets, stopping sets, and some related concepts. In Section 3 the result on transforming planar point processes to Poisson processes is presented, and a brief comparison of the use of stopping sets and predictable sets is given. An example is provided in Section 4. In Section 5, the result is extended to point processes in higher dimensions.

2. Preliminaries

First, some notation. In what follows, z , z' , and z'' represent elements of \mathbb{R}_+^2 , the positive quadrant of the plane, while s, s', t, t', u, x and y denote elements of \mathbb{R}_+ .

Before defining planar point processes, some ordering of points in the plane is required. For $z = (s, t)$ and $z' = (s', t') \in \mathbb{R}_+^2$, say $z < z'$ if $s < s'$ and $t < t'$. Similarly, $z \leq z'$ if $s \leq s'$ and $t \leq t'$. Let $(z, z']$ denote the rectangular region in \mathbb{R}_+^2 consisting of all points greater than z and less than or equal to z' ; i.e. $(z, z'] = \{z'' : z < z'' \leq z'\}$.

Let (Ω, \mathcal{F}, P) be a complete probability space. A *filtration* $\mathcal{F}(z)$ is a collection of sub- σ -fields of \mathcal{F} which is increasing (i.e. $\mathcal{F}(z) \subseteq \mathcal{F}(z')$ for $z \leq z'$), right-continuous (i.e. $\mathcal{F}(z) = \bigcap_{z' > z} \mathcal{F}(z')$) and complete (i.e. each $\mathcal{F}(z)$ contains the null sets of \mathcal{F}).

Let \mathcal{B} denote the Borel subsets of \mathbb{R}_+^2 , and let μ denote Lebesgue measure on \mathcal{B} . Let $\hat{\mathcal{F}} = \mathcal{F} \otimes \mathcal{B}$, and let π denote the product measure $P \otimes \mu$ on $\hat{\mathcal{F}}$.

For $z = (s, t)$, define $\mathcal{F}^1(z)$ as $\bigvee_{t' \geq 0} \mathcal{F}(s, t')$. Similarly, $\mathcal{F}^2(z) = \bigvee_{s' \geq 0} \mathcal{F}(s', t)$. The focus of most of this paper is on properties related to the filtration \mathcal{F}^1 ; the definitions and results related to \mathcal{F}^2 are analogous.

The σ -field \mathcal{P}^1 generated by sets of the form $F \times (z, z']$, for $z \leq z'$ and $F \in \mathcal{F}^1(z)$, is called the \mathcal{F}^1 -predictable σ -field, and an element of \mathcal{P}^1 is an \mathcal{F}^1 -predictable set. A process X on \mathbb{R}_+^2 is called \mathcal{F}^1 -predictable if it is \mathcal{P}^1 -measurable; i.e. if $\{(\omega, z) : \omega \in \Omega, z \in \mathbb{R}_+^2, X(\omega, z) \in B\} \in \mathcal{P}^1$, for any Borel set $B \in \mathbb{R}$.

A mapping D from Ω to the closed subsets of \mathbb{R}_+^2 is called an \mathcal{F}^1 -stopping set provided $z' \in D(\omega)$ implies $z \in D(\omega)$ for all $z < z'$, and $\{\omega : z \in D(\omega)\} \in \mathcal{F}^1(z)$ for all $z \in \mathbb{R}_+^2$.

For X a process on \mathbb{R}_+^2 and B a Borel set in \mathbb{R}_+^2 , let $X(B)$ denote $\int 1_B(z) dX(z)$, provided the integral exists. In particular, if $z = (s, t) < (s', t') = z'$, then $X(z, z']$ can be written as

$$X(z') - X(s, t') - X(s', t) + X(z). \quad (1)$$

Following Nair (1990), a process $N(z)$ on \mathbb{R}_+^2 is *increasing* if $N(z, z') \geq 0$ for every $z < z'$, and $N(s, 0) = N(0, t) = 0$ for every $s, t \geq 0$. If an adapted process $N(z)$ on \mathbb{R}_+^2 taking values in $\mathbb{Z}^+ \cup \{\infty\}$ is right-continuous and increasing, then N is a *point process*.

A point process is called *simple* if all its jumps are of size 1, i.e. if $\lim_{\Delta \downarrow 0} N((s - \Delta, t - \Delta), (s, t]) = 0$ or 1, for each $s, t \in \mathbb{R}_+$. A *Poisson process* on \mathbb{R}_+^2 is a simple point process N where for any disjoint Borel sets B_1, \dots, B_n in \mathbb{R}_+^2 , $N(B_1), \dots, N(B_n)$ are independent Poisson random variables. If the mean of N satisfies $EN(B) = \mu(B)$ for any Borel set $B \subset \mathbb{R}_+^2$, then N is said to have *unit rate*.

A *1-martingale* is an integrable \mathcal{F} -adapted process X where for each $z \leq z'$, $E[X(z, z') | \mathcal{F}^1(z)] = 0$. If N is an \mathcal{F}^1 -adapted point process, then a *1-compensator* A of N is an increasing \mathcal{F}^1 -predictable process so that $N - A$ is a 1-martingale. The existence and uniqueness of A for simple, integrable N are proven by Jacod (1975).

Suppose that A is the 1-compensator of N and that there exists an integrable, non-negative, real-valued, \mathcal{F}^1 -predictable process λ such that with probability 1, for each $z \in \mathbb{R}_+^2$,

$$\int_{z' < z} \lambda(z') d\mu(z') = A(z). \quad (2)$$

Then λ is called an \mathcal{F}^1 -intensity of N .

3. Transformation of planar processes

This section contains a result on transformations changing point processes on \mathbb{R}_+^2 into Poisson processes. First, recall the following lemma of Nair (1990), which generalizes a similar result in Brown et al. (1986) for point processes on $[0, 1]^2$.

Lemma 3.1. *If N is an \mathcal{F}^1 -adapted point process on \mathbb{R}_+^2 with deterministic, continuous 1-compensator A and at most one point on any vertical line, then N is a Poisson process whose mean corresponds to A .*

Theorem 3.2. *Suppose N is a simple \mathcal{F} -adapted point process on \mathbb{R}_+^2 with \mathcal{F}^1 -intensity λ_1 and \mathcal{F}^2 -intensity λ_2 . If with probability one, $\int_0^\infty \lambda_1(\omega, s, t) dt = \infty$ for all $s \in \mathbb{R}_+$, then there is a family of \mathcal{F}^1 -predictable sets $\{D_z\}$ such that $M(z) := \int \mathbf{1}_{D_z} dN$ is a Poisson process on \mathbb{R}_+^2 with unit rate.*

Proof. Define a process τ_z on \mathbb{R}_+ as follows. Fix $z = (s, t) \in \mathbb{R}_+^2$. For $s' \leq s$, let

$$\tau_z(\omega, s') = \inf \left\{ t' : \int_0^{t'} \lambda_1(\omega, s', u) du > t \right\} \quad (3)$$

with the convention that $\inf \{\emptyset\} = \infty$. For $s' > s$, let $\tau_z(\omega, s') = 0$.

By assumption, for any $s' \in \mathbb{R}_+$, $\int_0^\infty \lambda_1(\omega, s', u) du = \infty$ a.s., so $\tau_z(\omega, s') < \infty$ a.s. In addition, observe that by definition of τ_z , for almost all $\omega \in \Omega$,

$$\int_0^{\tau_z(\omega, s')} \lambda_1(\omega, s', t') dt' \geq t$$

and since $\lambda_1 < \infty$, in fact, equality holds, i.e.

$$\int_0^{\tau_z(\omega, s')} \lambda_1(\omega, s', t') dt' = t. \quad (4)$$

Let $M(z) = \int \mathbf{1}_{D_z} dN$, where D_z is the random closed region bounded by τ_z and the axes. That is,

$$D_z(\omega) = \left\{ (x, y) \in \mathbb{R}_+^2 : x \leq s, \int_0^y \lambda_1(x, u) du \leq t \right\}. \quad (5)$$

Note that since λ_1 is an \mathcal{F}^1 -predictable process, so is $\int_0^y \lambda_1(x, u) du$; thus D_z is an \mathcal{F}^1 -predictable set.

In order to show that M is a Poisson process, it is first necessary to verify that M is a well-defined \mathcal{F}^1 -adapted point process. For each $\omega \in \Omega$, since $\mathbf{1}_{D_z}$ is nonnegative, the ordinary Lebesgue integral $\int \mathbf{1}_{D_z}(z') dN(z')$ is clearly well-defined. To see that M is \mathcal{F}^1 -adapted, note that since D_z is an \mathcal{F}^1 -predictable set, the indicator $\mathbf{1}_{D_z}$ is an \mathcal{F}^1 -predictable (and hence \mathcal{F}^1 -adapted) process. Since $\mathbf{1}_{D_z}$ and N are both \mathcal{F}^1 -adapted, the integral $\int \mathbf{1}_{D_z} dN \in \mathcal{F}^1$.

To be a point process, M must furthermore take values in $\mathbb{Z}^+ \cup \{\infty\}$, be increasing, and be right-continuous. It is clear from the definition of M that M inherits from N the property of taking values in $\mathbb{Z}^+ \cup \{\infty\}$.

To show that M is increasing, recall from Eq. (1) that for $(s, t) \leq (s', t') \in \mathbb{R}_+^2$,

$$\begin{aligned} M((s, t), (s', t')) &= M(s', t') - M(s, t') - M(s', t) + M(s, t) \\ &= \int \mathbf{1}_{D(s', t')} dN - \int \mathbf{1}_{D(s, t')} dN - \int \mathbf{1}_{D(s', t)} dN + \int \mathbf{1}_{D(s, t)} dN \\ &= \int \mathbf{1}_{D(s', t') \setminus D(s, t')} dN - \int \mathbf{1}_{D(s', t) \setminus D(s, t)} dN, \end{aligned} \quad (6)$$

the last equation following from the fact that $D_{(s, t')} \subseteq D_{(s', t')}$ and $D_{(s, t)} \subseteq D_{(s', t)}$, as is evident from the definition of D_z in Eq. (5). Further, since

$$D_{(s', t')} \setminus D_{(s, t')} = \left\{ (x, y) : s < x \leq s'; \int_0^y \lambda_1(x, u) du \leq t' \right\} \quad (7)$$

and

$$D_{(s', t)} \setminus D_{(s, t)} = \left\{ (x, y) : s < x \leq s'; \int_0^y \lambda_1(x, u) du \leq t \right\}, \quad (8)$$

one sees that $D_{(s', t')} \setminus D_{(s, t')} \supseteq D_{(s', t)} \setminus D_{(s, t)}$. From the fact that the point process N is nonnegative, it follows that

$$\int \mathbf{1}_{D(x', y') \setminus D(x, y')} dN \geq \int \mathbf{1}_{D(x', y) \setminus D(x, y)} dN$$

which with Eq. (6) establishes that $M((s, t), (s', t')) \geq 0$. In order for M to be increasing, M must also satisfy

$$M(s, 0) = M(0, t) = 0 \quad (9)$$

for every $s, t \geq 0$. N is a point process, so Eq. (9) holds with M replaced by N . Since $M = \int \mathbf{1}_{D_z} dN$, clearly Eq. (9) holds for M as well.

In order to establish that M is a point process, only the right-continuity of M remains to be verified. Fix any $\omega \in \Omega$. From the definition of τ in Eq. (3) and the fact that $\lambda_1 < \infty$, it follows that for any $x \in \mathbb{R}_+$, $\tau_{z'}(\omega, x) \downarrow \tau_z(\omega, x)$ as $z' \downarrow z$. Consequently, for any $(x, y) \in \mathbb{R}_+^2$, $\mathbf{1}_{D_{z'}}(\omega, x, y) \downarrow \mathbf{1}_{D_z}(\omega, x, y)$ as $z' \downarrow z$, and the right-continuity of M follows by monotone convergence. Thus M is a well-defined, \mathcal{F}^1 -adapted point process.

Using Lemma 2.1 of Nair (1990) and the assumption that the \mathcal{F}^2 -intensity of N exists, N contains at most one point on any vertical line a.s. This is true also of M , since the definition of M implies that N has a point at (s, t) if and only if M has a point at $(s, \int_0^t \lambda_1(s, u) du)$.

For $z = (s, t)$, Let $C(z) = \int \mathbf{1}_{D_z} dA$, where A is the 1-compensator of N , i.e. $A(z) = \int_0^s \int_0^t \lambda_1(s', t') dt' ds'$. In other words, $C(z) = \int_0^s \int_0^{\tau_z(s)} \lambda_1(s', t') dt' ds'$. Recall from Eq. (4) that for all $s' \leq s$, $\int_0^{\tau_z(s')} \lambda_1(s', t') dt' = t$, so $C(z) = \int_0^s t ds' = st$. Thus, C is the 1-compensator of the unit rate Poisson process.

In light of Lemma 3.1, all that remains is to show that C is the 1-compensator of M , i.e. that $M - C$ is a 1-martingale.

Choose any two points $z = (s, t)$ and $z' = (s', t')$ in \mathbb{R}_+^2 such that $z \leq z'$, and let F be any set in \mathcal{F}_z . From Eq. (6),

$$\begin{aligned} M(z, z') &= \int [\mathbf{1}_{D(s', t') \setminus D(s, t)} - \mathbf{1}_{D(s', t) \setminus D(s, t)}] dN \\ &= \int \mathbf{1}_{D(z, z')} dN, \end{aligned}$$

where $D(z, z') := \{D(s', t') \setminus D(s, t')\} \setminus \{D(s', t) \setminus D(s, t)\}$.

Note that from Eqs. (7) and (8), $D(z, z')$ is a random subset of $(s, s'] \times \mathbb{R}_+$. Thus its intersection with $F \times \mathbb{R}_+^2$ is given by

$$D(z, z') \cap \{F \times (s, s'] \times \mathbb{R}_+\}.$$

Further, note that $F \in \mathcal{F}^1(s, 0)$, so that $\{F \times (s, s'] \times \mathbb{R}_+\}$ is a predictable set.

Therefore,

$$\begin{aligned} E[M(z, z') | F] &= E \left[\int \mathbf{1}_{F \times \mathbb{R}_+^2} \mathbf{1}_{D(z, z')} dN \right] \\ &= E \left[\int \mathbf{1}_{F \times (s, s'] \times \mathbb{R}_+} \mathbf{1}_{D(z, z')} dN \right] \\ &= E \left[\int \mathbf{1}_{F \times (s, s'] \times \mathbb{R}_+} \mathbf{1}_{D(z, z')} dA \right] \\ &= E[C(z, z') | F] \end{aligned} \tag{10}$$

with relation (10) following from the martingale property (see e.g. Eq. (1) of Nair, 1990), using the fact that $\mathbf{1}_{F \times (s, s'] \times \mathbb{R}_+}$ and $\mathbf{1}_{D(z, z')}$ are bounded, nonnegative, predictable processes. From Lemma 3.1, the proof is complete. \square

Remark 3.3. The relation of Theorem 3.2 to the results of Nair (1990) and Merzbach and Nualart (1986) is of interest. The results of the previous authors involve transforming the point process N via a sequence of stopping sets. Notice that the sets D_z

defined in Theorem 3.2 are generally not stopping sets, since they may fail to meet the requirement that if a stopping set contains a point z' , then it must also contain all points less than z' .

Theorem 3.3 of Nair (1990) assumes that A is 1-convex, i.e. that $A(s + \Delta s, t) - A(s, t) \leq A(s + 2\Delta s, t) - A(s + \Delta s, t)$, for all $s, \Delta s$, and $t \in \mathbb{R}_+$. When λ_1 exists, the 1-convexity of A is equivalent to the assumption in Theorem 4 of Merzbach and Nualart (1986) that for any t , $\int_0^t \lambda_1(s, t') dt'$ is a nondecreasing function of s . This means that each function $\tau_z(s')$ is decreasing in s' . This condition ensures that D_z is an \mathcal{F}^1 -stopping set.

Note that Theorem 5 of Merzbach and Nualart (1986) and Theorem 3.4 of Nair (1990) relax the convexity condition slightly. For example, in Theorem 5 of Merzbach and Nualart (1986), $\lambda_1(s, t)/\beta(s)$ is assumed nondecreasing in s , for all $t \geq 0$, where β is some positive decreasing function. Though a bit weaker than the convexity condition, this condition also holds only in rather special cases. The situation is the same in Nair (1990).

Remark 3.4. In applications, one often observes a point process N on a finite subregion $S \subset \mathbb{R}_+^2$, and typically the requirement in Theorem 3.2 that $\int_0^\infty \lambda_1(\omega, s, t) dt = \infty$ is not met. One can nevertheless apply Theorem 3.2 to the process $\tilde{N} := N + N'$, where N' is an \mathcal{F} -adapted Poisson process independent of N with \mathcal{F}^1 -intensity 1 on S^c and \mathcal{F}^1 -intensity 0 on S . Let $\tilde{\lambda}_1$ be the \mathcal{F}^1 -intensity of the process \tilde{N} . In transforming \tilde{N} as in Theorem 3.2, a point $z = (s, t)$ is moved to $(s, \int_0^t \tilde{\lambda}_1(s, t') dt')$. Similarly, the region S corresponds in the transformed plane to a region $T := \{(s, \int_0^t \tilde{\lambda}_1(s, t') dt') : (s, t) \in S\} \subset \mathbb{R}_+^2$. Note that the shape of T is random. If λ_1 is a left-continuous version of the \mathcal{F}^1 -intensity of N , and S is the set of points under a left-continuous function $f(x)$, i.e. $S = \{(x, y) \in \mathbb{R}_+^2 : y \leq f(x)\}$, then from the definitions of T and λ_1 it follows that T is the set of points under some left-continuous function $g(x)$. Further properties of T may be a subject for future research.

Corollary 3.5. Suppose that N satisfies the conditions of Theorem 3.2, and let λ_1^* be any integrable, nonnegative, \mathcal{F}^1 -predictable process. Define D_z^* and M^* by

$$D_z^* := \left\{ (\omega, s', t') \in \Omega \times \mathbb{R}_+^2 : s' \leq s, \int_0^{t'} \lambda_1^*(\omega, s', u) du \leq t \right\},$$

$$M^*(z) := \int \mathbf{1}_{D_z^*} dN.$$

Then M^* is a Poisson process on \mathbb{R}_+^2 with unit rate if and only if $\lambda_1^* = \lambda_1$, π -a.e.

Proof. If $\lambda_1^* = \lambda_1$ π -a.e., then λ_1^* is a version of the \mathcal{F}^1 -intensity of N , so M^* is a unit-rate Poisson process by Theorem 3.2.

For the other direction, suppose that M^* is a unit-rate Poisson process, and that $\pi(\{\lambda_1^* \neq \lambda_1\}) > 0$. We must show that this leads to a contradiction.

Either $\pi(\{\lambda_1 < \lambda_1^*\}) > 0$ or $\pi(\{\lambda_1 > \lambda_1^*\}) > 0$; suppose that $\pi(\{\lambda_1 < \lambda_1^*\}) > 0$. (The case where $\pi(\{\lambda_1 > \lambda_1^*\}) > 0$ can be proven equivalently.)

Some additional definitions are required. For each $z \in \mathbb{R}_+^2$, define random sets L_z , K_z , and K_z^* as follows:

$$L_z(\omega) := \{z' \leq z; \lambda_1(\omega, z') < \lambda_1^*(\omega, z')\},$$

$$K_z(\omega) := \left\{ \left(s, \int_0^t \lambda_1(\omega, s, u) du \right); (s, t) \in L_z \right\},$$

$$K_z^*(\omega) := \left\{ \left(s, \int_0^t \lambda_1^*(\omega, s, u) du \right); (s, t) \in L_z \right\}.$$

Let M be defined as in Theorem 3.2. Recall that N has a point at (s, t) if and only if M contains a point at $(s, \int_0^t \lambda_1(s, u) du)$. Similarly, the definition of M^* implies that N has a point at (s, t) if and only if M^* has a point at $(s, \int_0^t \lambda_1^*(s, u) du)$. It follows that for any $\omega \in \Omega$,

$$\int \mathbf{1}_{L_z} dN = \int \mathbf{1}_{K_z} dM = \int \mathbf{1}_{K_z^*} dM^*. \quad (11)$$

By Theorem 3.2, M is a Poisson process with unit rate, and by assumption so is M^* . Thus, since K_z and K_z^* are predictable sets,

$$E \left[\int \mathbf{1}_{K_z} dM \right] = E \left[\int \mathbf{1}_{K_z} d\mu \right] \quad (12)$$

and

$$E \left[\int \mathbf{1}_{K_z^*} dM^* \right] = E \left[\int \mathbf{1}_{K_z^*} d\mu \right]. \quad (13)$$

Combining Eqs. (11)–(13) yields

$$E \left[\int \mathbf{1}_{K_z} d\mu \right] = E \left[\int \mathbf{1}_{K_z^*} d\mu \right]. \quad (14)$$

Note that for all $\omega \in \Omega$,

$$\int \mathbf{1}_{K_z} d\mu \leq \int \mathbf{1}_{K_z^*} d\mu. \quad (15)$$

Relation (15) follows directly from the definitions of K_z and K_z^* , in light of the fact that $\lambda_1 < \lambda_1^*$ on L_z . Further, $\pi(\{\lambda_1 < \lambda_1^*\}) > 0$ by assumption, which implies that for some $z \in \mathbb{R}_+^2$, the inequality in Eq. (15) is strict on a subset of Ω with positive probability. Therefore,

$$E \left[\int \mathbf{1}_{K_z} d\mu \right] < E \left[\int \mathbf{1}_{K_z^*} d\mu \right],$$

contradicting Eq. (14). \square

Remark 3.6. Corollary 3.5 may be useful for the evaluation of point process models. Given a point process N and a model specifying the \mathcal{F}^1 -conditional intensity, λ_1^* , by Corollary 3.5 the problem of assessing the fit of the model boils down to examining whether M^* is a planar Poisson process of unit rate. Much has been written on the latter problem; see e.g. Ripley (1979), Diggle (1983), Heinrich (1991), or Cressie (1993).

4. Example

Rathbun (1995) describes a planar version of the self-exciting point process analyzed by Hawkes (1971). Here the \mathcal{F}^1 -intensity may be given by

$$\lambda_1(s, t) = f(s, t) + \int \mathbf{1}_{\{s' < s\}} g(s - s', |t - t'|) dN(s', t'),$$

where f and g are deterministic, nonnegative functions from \mathbb{R}^2 to \mathbb{R} . Similarly, the \mathcal{F}^1 -intensity of a k -dimensional version of a Hawkes process can be given by

$$\begin{aligned} \lambda_1(t_1, \dots, t_k) &= f(t_1, \dots, t_k) \\ &+ \int \mathbf{1}_{\{t'_1 < t_1\}} g(t_1 - t'_1, |t_2 - t'_2|, \dots, |t_k - t'_k|) dN(t'_1, \dots, t'_k), \end{aligned}$$

where f and g are now deterministic nonnegative functions from \mathbb{R}^k to \mathbb{R} .

Such processes generally do not satisfy the convexity condition of Merzbach and Nualart (1986) and Nair (1990). For instance, suppose N is a planar Hawkes process and f and g are decreasing functions. Then for any s, s' , and $t \in \mathbb{R}_+$ such that $N(s, t) = N(s', t)$, $\lambda_1(x, y)$ is decreasing in x for $s < x < s'$ and $y \leq t$. Thus $\int_0^t \lambda_1(x, y) dy$ is decreasing in x for $s < x < s'$, which violates the convexity condition.

5. Extension to higher dimensions

In this section, Theorem 3.2 is generalized to include the case where N is a point process in \mathbb{R}_+^k , for $k \geq 2$. First, a few of the previous definitions must be extended.

For $z = (t_1, \dots, t_k) \in \mathbb{R}_+^k$ and $z' = (t'_1, \dots, t'_k) \in \mathbb{R}_+^k$, say $z < z'$ if $t_i < t'_i$ for each i , and say $z \leq z'$ if $t_i \leq t'_i$ for each i . A filtration \mathcal{F} on \mathbb{R}_+^k may be defined exactly as in Section 2. Let $\mathcal{F}^1(z)$ denote $\bigvee_{u_2, u_3, \dots, u_k} \mathcal{F}(t_1, u_2, u_3, \dots, u_k)$, where $z = (t_1, \dots, t_k)$. The \mathcal{F}^1 -predictable σ -field \mathcal{P}^1 , the 1-compensator A and the \mathcal{F}^1 -intensity λ of N can also be defined exactly as in Section 2.

The following lemma is an extension of Lemma 3.3 of Brown et al. (1986). A slightly stronger version is given in Proposition 4.2 of Nair (1990) and proven for the three-dimensional case.

Lemma 5.1. *If N is an \mathcal{F}^1 -adapted point process on \mathbb{R}_+^k with deterministic, continuous 1-compensator A and at most one point on every hyperplane $\{t_1 = t\}$, then N is a Poisson process whose mean corresponds to A .*

Theorem 5.2. *Suppose that N is a simple \mathcal{F} -adapted point process on \mathbb{R}_+^k with \mathcal{F}^1 -intensity λ_1 , \mathcal{F}^2 -intensity λ_2, \dots , and \mathcal{F}^k -intensity λ_k . If with probability one, for all $t_1, t_2, \dots, t_{k-1} \in \mathbb{R}_+$,*

$$\int_0^\infty \lambda_1(\omega, t_1, t_2, t_3, \dots, t_k) dt_k = \infty,$$

then there is a sequence of \mathcal{F}^1 -predictable sets D_z , such that $M(z) := \int \mathbf{1}_{D_z} dN$ is a Poisson process on \mathbb{R}_+^k with unit rate.

Proof. Fix $z = (t_1, \dots, t_k)$, and for $s_1 \leq t_1, s_2 \leq t_2, \dots, s_{k-1} \leq t_{k-1}$ define $\tau_z(\omega, s_1, s_2, \dots, s_{k-1})$ as $\inf\{u: \int_0^u \lambda_1(\omega, s_1, s_2, \dots, s_{k-1}, s_k) ds_k > t_k\}$, letting $\tau_z(\omega, s_1, \dots, s_{k-1}) = 0$ otherwise.

Define D_z and M by

$$D_z = \left\{ (\omega, s_1, \dots, s_k) \in \Omega \times \mathbb{R}_+^k : s_1 \leq t_1, \dots, s_{k-1} \leq t_{k-1}, \int_0^{s_k} \lambda_1(\omega, s_1, \dots, s_{k-1}, u) du \leq t_k \right\},$$

$$M(z) = \int \mathbf{1}_{D_z} dN.$$

It follows by the same argument as in Theorem 3.2 that D_z is \mathcal{F}^1 -predictable and that M is a well-defined \mathcal{F}^1 -adapted point process. Further, from Proposition 4.1 of Nair (1990) it follows that with probability 1, N has at most one point on any hyperplane perpendicular to the t_1 -axis, and therefore the same is true for M .

Let $C(z) = \int \mathbf{1}_{D_z} dA$, where A is the 1-compensator of N . It follows from exactly the same argument as in Theorem 3.2 that C is the 1-compensator of the unit rate Poisson process in \mathbb{R}_+^k and that C is also the 1-compensator of M . This along with Lemma 5.1 completes the proof. \square

Remark 5.3. Note that the transformation in Theorem 5.2 involves rescaling the k th coordinate. This choice is arbitrary; one could similarly rescale the l th coordinate (for $l > 1$) by defining τ_z as $\inf\{u: \int_0^u \lambda_1(\omega, s_1, \dots, s_{l-1}, u, s_{l+1}, \dots, s_k) du > t_l\}$ and end the proof of Theorem 5.2 similarly. Thus in dimension $k > 2$, the transformation described here is not unique.

Acknowledgements

I am deeply indebted to David Brillinger for introducing me to this problem and providing many helpful ideas and comments, as well as motivation. I am also grateful to David Aldous, Tim Brown, Steve Evans, Ed Ionides, Lucien LeCam, Gopalan Nair, Jean Paik, and Jim Pitman for their helpful comments and encouragement. An anonymous referee also provided numerous valuable improvements. Any mistakes are my own though. This problem was motivated by research on multidimensional models for earthquake processes.

References

- Aalen, O., Hoem, J., 1978. Random time changes for multivariate counting processes. *Scand. Actuar. J.* 81–101.
- Brémaud, P.M., 1972. A martingale approach to point processes. Memorandum ERL-M345, Electronics Research Laboratory, University of California, Berkeley, CA.
- Brown, T., Ivanoff, B.G., Weber, N.C., 1986. Poisson convergence in two dimensions with applications to row exchangeable arrays. *Stochastic Process. Appl.* 23, 307–318.
- Brown, T., Nair, M., 1988. A simple proof of the multivariate random time change theorem for point processes. *J. Appl. Probab.* 25, 210–214.

- Cairolì, R., Walsh, J.B., 1975. Stochastic integrals in the plane. *Acta Math.* 134, 37–57.
- Cressie, N.A., 1993. *Statistics for Spatial Data*, Revised ed. Wiley, New Jersey.
- Diggle, P.J., 1983. *Statistical Analysis of Spatial Point Patterns*. Academic Press, London.
- Hawkes, A.G., 1971. Point spectra of some mutually exciting point processes. *J. Roy. Statist. Soc. B* 33, 438–443.
- Heinrich, L., 1991. Goodness-of-fit tests for the second moment function of a stationary multidimensional Poisson process. *Statistics* 22, 245–278.
- Jacod, J., 1975. Multivariate point processes: predictable projections, Radon–Nikodym derivatives, representation of martingales. *Z. Wahrsch. Verw. Gebiete* 31, 235–253.
- Merzbach, E., Nualart, D., 1986. A characterization of the spatial Poisson process and changing time. *Ann. Probab.* 14, 1380–1390.
- Meyer, P.A., 1971. Démonstration simplifiée d'un théorème de Knight. *Lecture Notes in Mathematics*, vol. 191. Springer, Berlin, pp. 191–195.
- Nair, M., 1990. Random space change for multiparameter point processes. *Ann. Probab.* 18, 1222–1231.
- Papangelou, F., 1972. Integrability of expected increments of point processes and a related random change of scale. *Trans. Amer. Math. Soc.* 165, 483–506.
- Rathbun, S., 1995. Modeling marked spatio-temporal point patterns. *Bull. Int. Statist. Inst.* 55, 379–396.
- Ripley, B., 1979. Tests of 'randomness' for spatial point patterns. *J. Roy. Statist. Soc. B* 41, 368–374.
- Watanabe, S., 1964. On discontinuous additive functionals and Lévy measures of a Markov process. *Jpn. J. Math.* 34, 53–70.