



Non-uniqueness of stationary measures for self-stabilizing processes

S. Herrmann*, J. Tugaut

*Institut de Mathématiques Elie Cartan - UMR 7502, Nancy-Université, CNRS, INRIA, B.P. 239,
54506 Vandoeuvre-lès-Nancy Cedex, France*

Received 18 March 2009; received in revised form 16 March 2010; accepted 18 March 2010
Available online 27 March 2010

Abstract

We investigate the existence of invariant measures for self-stabilizing diffusions. These stochastic processes represent roughly the behavior of some Brownian particle moving in a double-well landscape and attracted by its own law. This specific self-interaction leads to nonlinear stochastic differential equations and permits pointing out singular phenomena like non-uniqueness of associated stationary measures. The existence of several invariant measures is essentially based on the non-convex environment and requires generalized Laplace's method approximations.

© 2010 Elsevier B.V. All rights reserved.

MSC: primary 60H10; secondary 60J60; 60G10; 41A60

Keywords: Self-interacting diffusion; Stationary measures; Double-well potential; Perturbed dynamical system; Laplace's method; Fixed point theorem; McKean–Vlasov stochastic differential equations

1. Introduction

The aim of this paper is to present some new and surprising results concerning the existence of invariant probability measures for one-dimensional self-stabilizing diffusions. The specificity of such diffusion is the attraction of its paths by the own law of the stochastic process. The dynamical system solved by self-stabilizing diffusions can be characterized by three essential elements: first the system is governed by a double-well potential V which represents roughly

* Corresponding address: Nancy-Université, CNRS, INRIA, Institut de Mathématiques Elie Cartan - UMR 7502, F-54500 Vandoeuvre lès Nancy Cedex, France. Tel.: +33 3 83 68 46 07; fax: +33 3 83 68 45 34.

E-mail address: Samuel.Herrmann@iecn.u-nancy.fr (S. Herrmann).

the environment of the process, secondly some interaction potential F describes how strong the attraction between the process and its own law is, and finally the system is perturbed by some Brownian motion with small amplitude ($\sqrt{\epsilon}B_t, t \geq 0$).

Let us denote by $u_t^\epsilon(dx)$ the law of the self-stabilizing diffusion ($X_t^\epsilon, t \geq 0$), then the so-called McKean–Vlasov SDE satisfied by (X_t^ϵ) is given by:

$$X_t^\epsilon = X_0 + \sqrt{\epsilon}B_t - \int_0^t V'(X_s^\epsilon)ds - \int_0^t \int_{\mathbb{R}} F'(X_s^\epsilon - x)du_s^\epsilon(x)ds, \quad \epsilon > 0. \quad (E^{\epsilon, X_0})$$

Introducing the notation of the convolution product, (E^{ϵ, X_0}) can be written as follows:

$$X_t^\epsilon = X_0 + \sqrt{\epsilon}B_t - \int_0^t (V' + F' * u_s^\epsilon) (X_s^\epsilon) ds. \quad (1.1)$$

Let us just note that the interaction part of the drift term is related to the diffusion in some simple way: $F' * u_t^\epsilon(x) = \mathbb{E}[F'(x - X_t^\epsilon)]$. This way of characterizing the drift term essentially points out the structure of the attraction between the paths of the diffusion and its law. Self-interaction corresponds obviously to *mean fields* stabilization.

Such interacting particle systems have been studied from various points of view. A survey about the general setting for interaction (under global Lipschitz and boundedness assumptions) may be found in [11].

The aim of this paper is to consider both the existence and the uniqueness of stationary measures for the self-stabilizing diffusion (E^{ϵ, X_0}). In [6] Herrmann, Imkeller and Peithmann proved the existence of some unique strong solution to Eq. (1.1) generalizing previous results obtained by Benachour, Roynette, Talay and Vallois [2] in the context of constant environment potential V ($V'(x) = 0$ for all $x \in \mathbb{R}$). We choose their work as basis for developing our study. Nevertheless there exist several different papers dealing with the existence problem for self-stabilizing diffusion, each of them concerning other families of interaction functions. Let us cite McKean who studied in some earlier work a class of Markov processes that contains the solution of the limiting equation under restrictive global Lipschitz assumptions for the interaction [7], Stroock and Varadhan who considered some local form of interaction [10], Oelschläger who studied the particular case where interaction is represented by the derivative of the Dirac measure at zero [9] and finally Funaki who addressed existence and uniqueness for the martingale problem associated with self-stabilizing diffusions [4].

Let us now focus our attention to the stationary measures. In [2] the authors emphasize that the invariant measure, corresponding to some given average, is unique in this particular constant potential V situation. This feature is essential for further developments. The natural convergence question between the law of the process and the invariant measure, as time elapses, can then be analyzed, see [3]. This kind of convergence was also considered by Tamura under different assumptions on the structure of the interaction, see [12,13].

The presence of some potential gradient which describes the environment of the self-stabilizing diffusion is essential for the question of existence and uniqueness of invariant measures. In particular, if the landscape is represented by some symmetric double-well potential then surprising effects appear due to the lack of convexity: we shall prove that, under suitable conditions, there exist at least three invariant measures of which one is symmetric (Theorem 4.5) and two are asymmetric or so-called *outlying* (Theorem 4.6). In the particular linear interaction case ($F'(x) = \alpha x$ with $\alpha > 0$), these three measures constitute the whole set of invariant measures (Theorem 3.2) provided that V'' is a convex function.

The material of this paper is organized as follows: first we list several assumptions concerning both the interaction function F and the environment potential V which permit in particular to assure the existence of the self-stabilizing diffusion (E^ϵ, X_0) . In Section 2 preliminary results concerning the structure of the invariant measure (if it exists!) are developed. These results are essential for the construction of such measures. The question of existence starts to be addressed in Section 3 in the particular linear interaction context. After pointing out some symmetric and asymmetric invariant measures, we point out some nice context for which the whole set of stationary measures can be described. This study is finally extended to the general interaction case in the last section. We postpone different tools concerning asymptotic analysis based on Laplace’s method to the [Appendix](#).

1.1. Main assumptions

In order to study invariant measures for self-stabilizing diffusions, we especially need that (1.1) admits some unique strong solution. For this reason, we assume that both the potential landscape V and the interaction function F satisfy some growth conditions and some regularity properties. Moreover we add some technical assumptions which permit to simplify the statements.

We assume the following properties for the function V :

- (V-1) *Regularity*: $V \in C^\infty(\mathbb{R}, \mathbb{R})$. C^∞ denotes the Banach space of infinitely bounded continuously differentiable function.
- (V-2) *Symmetry*: V is an even function.
- (V-3) V is a double-well potential. The equation $V'(x) = 0$ admits exactly three solutions: a , $-a$ and 0 with $a > 0$; $V''(a) > 0$ and $V''(0) < 0$. The bottoms of the wells are reached for $x = a$ and $x = -a$.
- (V-4) There exist two constants $C_4, C_2 > 0$ such that $\forall x \in \mathbb{R}, V(x) \geq C_4x^4 - C_2x^2$.
- (V-5) $\lim_{x \rightarrow \pm\infty} V''(x) = +\infty$ and $\forall x \geq a, V''(x) > 0$.
- (V-6) The growth of the potential V is at most polynomial: there exist $q \in \mathbb{N}^*$ and $C_q > 0$ such that $|V'(x)| \leq C_q (1 + x^{2q})$.
- (V-7) *Initialization*: $V(0) = 0$.

Typically, V is a double-well polynomial function (Fig. 1). Nevertheless the results presented in the following are applied to more general functions: regular double-well functions with polynomial growth as $|x|$ becomes large. Moreover, let us note that the hypothesis (V-5) implies V' not to be globally Lipschitz. We introduce the parameter ϑ which plays some important role in the following:

$$\vartheta = \sup_{x \in \mathbb{R}} -V''(x). \tag{1.2}$$

Let us note that the simplest example (most famous in the literature) is $V(x) = \frac{x^4}{4} - \frac{x^2}{2}$ which bottoms are localized in -1 and 1 and with parameter $\vartheta = 1$.

Let us now present the assumptions concerning the attraction function F .

- (F-1) F is an even polynomial function. Indeed we consider some classical situation: the attraction between two points x and y only depends on the distance $F(x - y) = F(y - x)$.
- (F-2) F is a convex function.
- (F-3) F' is a convex function on \mathbb{R}_+ therefore for any $x \geq 0$ and $y \geq 0$ such that $x \geq y$ we get $F'(x) - F'(y) \geq F''(0)(x - y)$.




Fig. 1. Potential V .

(F-4) The polynomial growth of the attraction function F is related to the growth condition (V-6):
 $|F'(x) - F'(y)| \leq C_q|x - y|(1 + |x|^{2q-2} + |y|^{2q-2})$.

The function F' is globally Lipschitz if and only if the degree of the polynomial function F is equal to 2. Let us define the parameter $\alpha \geq 0$ which shall play an essential role in the following:

$$F'(x) = \alpha x + F'_0(x) \quad \text{with } \alpha = F''(0) \geq 0. \tag{1.3}$$

In [6], Herrmann, Imkeller and Peithmann presented sufficient conditions for the SDE (1.1) to admit a unique strong solution. In particular, if $\mathbb{E}[X_0^{8q^2}] < +\infty$, where q is defined by (V-6), and if all main assumptions just defined are satisfied, the existence of a unique solution is proved and $\mathbb{E}[X_t^{8q^2}] < +\infty$ for all $t \geq 0$. In the following we shall always assume that the $(8q^2)$ -th moment of the initial value X_0 is finite. This permits to study further the self-stabilizing diffusions and exhibit invariant measures. Also, we will use the same notation for the measures absolutely continuous with respect to the Lebesgue measure and their density.

2. General structure of the invariant measures

This section deals with different preliminary results describing the main structure of the invariant measures of (E^{ϵ, X_0}) . First of all, there is some classical link between the stochastic differential equation and the associated parabolic partial differential equation which permits to characterize stationary measures.

Lemma 2.1. *We recall that $\int_{\mathbb{R}} x^{8q^2} u_0^\epsilon(dx) < \infty$. Let du_t^ϵ denote the probability measure of $(X_t^\epsilon; t \geq 0)$. Then, for all $t > 0$, du_t^ϵ admits a C^∞ -continuous density u_t^ϵ with respect to the Lebesgue measure. Moreover, u^ϵ is solution of the following PDE:*

$$\frac{\partial}{\partial t} u_t^\epsilon(x) = \frac{\epsilon}{2} \frac{\partial^2}{\partial x^2} u_t^\epsilon(x) + \frac{\partial}{\partial x} [u_t^\epsilon(x) (V'(x) + (F' * u^\epsilon)(t, x))] \tag{2.1}$$

for all $t > 0$, $x \in \mathbb{R}$ and $du_0^\epsilon(x) = \mathbb{P}(X_0 \in dx)$.

Proof. In [8] (Step 3), McKean proves, using Weyl’s lemma that the unique strong solution of (1.1) admits a regular density function solution to the PDE (2.1) provided that the drift term of the SDE is a C^∞ -continuous function. In fact the hypothesis (F-1) implies that $x \mapsto F' * u_t^\epsilon(x)$ is a polynomial function whose parameters are the moments of orders 1 to $2n - 1$. We assume

that the $(8q^2)$ -th moment of the initial value X_0 is finite so $\int_{\mathbb{R}} x^{8q^2} du_t^\epsilon(x) < \infty$ for all $t \geq 0$ consequently the drift $x \mapsto V'(x) + F' * u_t^\epsilon(x)$ is regular. \square

Let us note that Tamura (Theorem 2.1 in [13]) extended the initial result of McKean to unbounded drift terms. This extension was essential to prove the existence of self-stabilizing processes but was not necessary in the study of the density regularity. Since the existence of some unique solution of (1.1) was already presented in [6], we do not really need the extension of Tamura.

The density of $(X_t^\epsilon, t > 0)$ with respect to the Lebesgue measure is solution to the parabolic PDE (2.1) (non-linear Kolmogorov equation): this implies in particular that any stationary measure whose $(8q^2)$ -moment is finite (if such a measure exists!) satisfies an elliptic differential equation. This link between non-linear differential equations and self-stabilizing diffusions permits to express the invariant measure in some exponential form.

Lemma 2.2. *If there exists an invariant measure u_ϵ to (E^ϵ, X_0) whose $(8q^2)$ -moment is finite, then:*

$$\begin{aligned} u_\epsilon(x) &= \frac{1}{\lambda(u_\epsilon)} \exp \left[-\frac{2}{\epsilon} \left(\int_0^x F' * u_\epsilon(y) dy + V(x) \right) \right] \\ &= \frac{1}{\lambda(u_\epsilon)} \exp \left[-\frac{2}{\epsilon} (F * u_\epsilon(x) - F * u_\epsilon(0) + V(x)) \right], \end{aligned} \tag{2.2}$$

where $\lambda(u_\epsilon)$ denotes the normalization factor: $\int_{\mathbb{R}} u_\epsilon(x) dx = 1$. Conversely any measure whose density satisfies (2.2) is invariant for (E^ϵ, X_0) and admits a $8q^2$ finite moment.

Proof. 1. First we shall prove that any measure u_ϵ satisfying (2.2) is an invariant measure for (1.1). Let X_0 be some random variable with distribution u_ϵ . We consider the diffusion $(Y_t^\epsilon)_{t \geq 0}$ solution of the SDE:

$$Y_t^\epsilon = X_0 + \sqrt{\epsilon} B_t - \int_0^t W_\epsilon'(Y_s^\epsilon) ds \tag{2.3}$$

where $W_\epsilon(x) = V(x) + F * u_\epsilon(x) - F * u_\epsilon(0)$. Since $(Y_t^\epsilon)_{t \geq 0}$ is a Kolmogorov diffusion process, it admits a unique stationary measure of probability v_ϵ given by $v_\epsilon(x) = \frac{e^{-\frac{2}{\epsilon} W_\epsilon(x)}}{\int_{\mathbb{R}} e^{-\frac{2}{\epsilon} W_\epsilon(y)} dy} = u_\epsilon(x)$. Consequently the law of Y_t^ϵ corresponds with u_ϵ for all $t \geq 0$. Moreover (2.3) becomes (1.1). Hence, u_ϵ is an invariant measure for (1.1).

Finally the hypotheses (V-4) and (F-3) imply that there exists $R_\epsilon > 0$ such that $V(x) + F * u_\epsilon(x) - F * u_\epsilon(0) \geq x^2$ for all $x \geq R_\epsilon$. Therefore the $(8q^2)$ -moment of u_ϵ is finite.

2. Let us prove now that any invariant measure satisfies to some exponential implicit structure. By (2.1), any stationary measure u_ϵ whose $(8q^2)$ -moment is finite satisfies

$$\frac{\epsilon}{2} u_\epsilon''(x) + (u_\epsilon(x) (V'(x) + F' * u_\epsilon(x)))' = 0, \quad \text{for all } x \in \mathbb{R}. \tag{2.4}$$

By integrating (2.4), we obtain the existence of some constant $C_\epsilon \in \mathbb{R}$ such that

$$\frac{\epsilon}{2} u_\epsilon'(x) + u_\epsilon(x) (V'(x) + F' * u_\epsilon(x)) = C_\epsilon, \quad \text{for all } x \in \mathbb{R}.$$

Using the method of variation of parameters, u_ϵ takes the following form

$$u_\epsilon(x) = \Lambda_\epsilon(x) \exp \left[-\frac{2}{\epsilon} \left(\int_0^x F' * u_\epsilon(y) dy + V(x) \right) \right],$$

with

$$\Lambda'_\epsilon(x) = \frac{2}{\epsilon} C_\epsilon \exp \left[\frac{2}{\epsilon} \left(\int_0^x F' * u_\epsilon(y) dy + V(x) \right) \right].$$

Hence

$$\begin{aligned} u_\epsilon(x) &= \Lambda_\epsilon(0) \exp \left[-\frac{2}{\epsilon} \left(\int_0^x F' * u_\epsilon(y) dy + V(x) \right) \right] \\ &\quad + \frac{2}{\epsilon} C_\epsilon \int_0^x \exp \left[\frac{2}{\epsilon} \left(\int_0^y F' * u_\epsilon(z) dz + V(y) \right) \right] dy \\ &\quad \times \exp \left[-\frac{2}{\epsilon} \left(\int_0^x F' * u_\epsilon(y) dy + V(x) \right) \right]. \end{aligned}$$

Let us assume that $C_\epsilon \neq 0$. Applying Lemma A.1 to the function $U(x) = \int_0^x F' * u_\epsilon(y) dy + V(x)$, whose second derivative is positive for $|x|$ large enough (using hypotheses (V-5) and (F-2)), permits to exhibit the equivalent of $\Lambda_\epsilon(x)$:

$$\Lambda_\epsilon(x) \approx \frac{2}{\epsilon} C_\epsilon \frac{\exp \left[\frac{2}{\epsilon} \left(\int_0^x F' * u_\epsilon(y) dy + V(x) \right) \right]}{\frac{2}{\epsilon} (V'(x) + F' * u_\epsilon(x))} \quad \text{as } x \rightarrow \pm\infty.$$

Hence $u_\epsilon(x) \approx \frac{C_\epsilon}{V'(x) + F' * u_\epsilon(x)}$. We can note that $V'(x) + F' * u_\epsilon(x)$ is positive in a neighborhood of $+\infty$ and negative for $-\infty$. Since u_ϵ is a probability distribution and consequently needs to be nonnegative, we get $C_\epsilon = 0$ and obtain (2.2) after normalization. \square

Lemma 2.2 presents the essential structure of any invariant measure. The global exponential form will play a crucial role in next sections: to prove the existence of some stationary measure, it is necessary and sufficient to solve Eq. (2.2).

3. The linear interaction case

First we shall analyze the existence problem for stationary measures in the simple linear case. In this case $F'(x) = \alpha x$ with $\alpha > 0$, the interaction gradient function is quadratic: $F(x) = \frac{\alpha}{2} x^2$ and the stochastic differential equation takes an interesting simple form. The non-linearity of the drift term is limited to the average of the density $u_t^\epsilon(x)$:

$$X_t^\epsilon = X_0 + \sqrt{\epsilon} B_t - \int_0^t V'(X_s^\epsilon) ds - \alpha \int_0^t \left(X_s^\epsilon - \int_{\mathbb{R}} x du_s^\epsilon(x) \right) ds, \quad \epsilon > 0.$$

The study of this particular case emphasizes the existence of several invariant measures. The interesting problem is then to determine in which situations the number of such measures is perfectly known.

3.1. Existence of invariant measures

The existence question is really simplified in the linear interaction case, it is just reduced in fine to the following parametrization problem. Let us denote the first moment of an invariant measure u_ϵ by

$$m_1(\epsilon) = \int_{\mathbb{R}} x u_\epsilon(x) dx, \tag{3.1}$$

then (2.2) becomes

$$u_\epsilon(x) = \frac{\exp\left[-\frac{2}{\epsilon}\left(V(x) + \alpha\frac{x^2}{2} - \alpha m_1(\epsilon)x\right)\right]}{\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon}\left(V(y) + \alpha\frac{y^2}{2} - \alpha m_1(\epsilon)y\right)\right] dy}. \tag{3.2}$$

Consequently: u_ϵ is an invariant measure if and only if (3.1) and (3.2) are satisfied. It suffices then to point out the convenient parameters $m_1(\epsilon)$ since there is a one to one correspondence between these parameters and the invariant measures. In other words, we shall find the solution of the equation

$$m = \Psi_\epsilon(m) \quad \text{with} \quad \Psi_\epsilon(m) = \frac{\int_{\mathbb{R}} x \exp\left[-\frac{2}{\epsilon}\left(V(x) + \alpha\frac{x^2}{2} - \alpha mx\right)\right] dx}{\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon}\left(V(x) + \alpha\frac{x^2}{2} - \alpha mx\right)\right] dx}. \tag{3.3}$$

Obviously, $m_1^0(\epsilon) = 0$ is a candidate. The corresponding measure u_ϵ^0 is invariant and symmetric:

$$u_\epsilon^0(x) = \exp\left[-\frac{2}{\epsilon}\left(V(x) + \alpha\frac{x^2}{2}\right)\right] \left(\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon}\left(V(y) + \alpha\frac{y^2}{2}\right)\right] dy\right)^{-1}.$$

In fact u_ϵ^0 is the unique symmetric stationary measure. Indeed, on one hand the average of such measure vanishes and on the other hand any invariant measure satisfies (2.2); finally we obtain the previous expression for u_ϵ^0 .

Of course the natural question concerns the existence of other real solutions $m_1(\epsilon)$ of (3.2). In fact the basic dynamical system associated to self-stabilizing diffusions is symmetric since F and V are assumed to be even functions. The consequence is immediate: if the initial law of the diffusion ($X_t^\epsilon, t \geq 0$) is symmetric so will be the law of X_t^ϵ for all $t > 0$. In [2], the authors consider self-stabilizing diffusions without the environment potential V . They proved the existence of some unique symmetric invariant measure and describe the behavior of the diffusion: for any initial law satisfying the moment condition of order $8q^2$ the law of $X_t - \mathbb{E}[X_0]$ converges to the invariant symmetric law as time elapses.

Adding some double-well potential V in the main structure of the stochastic differential equation drastically changes the situation. In particular we prove the existence of several invariant measures, one of them being symmetric.

Proposition 3.1. *Let a be the unique positive real which minimizes V (see (V-3)). For all $\delta \in]0, 1[$, there exists $\epsilon_0 > 0$ such that for all $\epsilon \leq \epsilon_0$, the Eq. (3.3) admits a solution satisfying the estimates:*

$$\left| m_1(\epsilon) - a + \frac{V^{(3)}(a)}{4V''(a)(\alpha + V''(a))} \epsilon \right| \leq \delta \epsilon. \tag{3.4}$$

Moreover $-m_1(\epsilon)$ satisfies (3.3) too.

Let us note that, for ϵ small enough, the preceding proposition implies the existence of at least three invariant measures corresponding to the averages: $0, m_1(\epsilon)$ and $-m_1(\epsilon)$.

Proof. Set $\tau > 0$. Let's proceed to the first order asymptotic development of the expression $\Psi_\epsilon(a - \tau\epsilon)$.

$$\begin{aligned} \Psi_\epsilon(a - \tau\epsilon) &= \frac{\int_{\mathbb{R}} x \exp\left[-\frac{2}{\epsilon}\left(V(x) + \alpha\frac{x^2}{2} - \alpha(a - \tau\epsilon)x\right)\right] dx}{\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon}\left(V(x) + \alpha\frac{x^2}{2} - \alpha(a - \tau\epsilon)x\right)\right] dx} \\ &= \frac{\int_{\mathbb{R}} x e^{-2\alpha\tau x} \exp\left[-\frac{2}{\epsilon}\left(V(x) + \alpha\frac{x^2}{2} - \alpha ax\right)\right] dx}{\int_{\mathbb{R}} e^{-2\alpha\tau x} \exp\left[-\frac{2}{\epsilon}\left(V(x) + \alpha\frac{x^2}{2} - \alpha ax\right)\right] dx}. \end{aligned}$$

By Lemma A.3 applied to the context: $f(x) = -2\alpha\tau x$, $n = 1$, $U(x) = V(x) + \frac{\alpha}{2}x^2 - \alpha ax$ and $\mu = 0$, we get:

$$\begin{aligned} \Psi_\epsilon(a - \tau\epsilon) &= a - \frac{1}{4\alpha(\alpha + V''(a))^2} \left[aV^{(3)}(a) + 4\alpha\alpha\tau(\alpha + V''(a)) \right] \epsilon + o(\epsilon) \\ &= a - \tau\epsilon + \frac{V''(a)}{\alpha + V''(a)} \left[\tau - \frac{V^{(3)}(a)}{4V''(a)(\alpha + V''(a))} \right] \epsilon + o(\epsilon). \end{aligned}$$

Set $\tau^0 = \frac{V^{(3)}(a)}{4V''(a)(\alpha + V''(a))}$. Then $a - \tau^0\epsilon$ is the first order approximation of the fixed point. Indeed for $\delta \in]0; 1[$ we can define

$$d_\pm := \Psi_\epsilon\left(a - (\tau^0 \pm \delta)\epsilon\right) - \left(a - \tau^0(1 \pm \delta)\epsilon\right) = \pm\delta \frac{V''(a)}{\alpha + V''(a)}\epsilon + o(\epsilon).$$

For ϵ small enough, $d_+ > 0$ and $d_- < 0$. Since the function Ψ_ϵ is C^0 continuous, there exists $m_1(\epsilon) \in [a - (\tau^0 + \delta)\epsilon; a - (\tau^0 - \delta)\epsilon]$ which satisfies $\Psi_\epsilon(m_1(\epsilon)) = m_1(\epsilon)$. Finally, by the change of variable $x := -x$ in the integral expression (3.3), we obtain $\Psi_\epsilon(-m_1(\epsilon)) = -\Psi_\epsilon(m_1(\epsilon)) = -m_1(\epsilon)$. \square

3.2. Description of the set of invariant measures

According to Proposition 3.1, we know there are at least three invariants measures. One of them is symmetric corresponding to the average 0 and two others will be called *outlying measures*, one wrapped around a and the other one around $-a$. The aim of this section is to study if there are exactly three invariants measures or more.

For this purpose, we study the asymptotic behavior of the function Ψ_ϵ defined by (3.3) in the small noise limit.

Theorem 3.2. *If V'' is a convex function then, in the small noise limit, there exist exactly three stationary measures.*

Proof. Let $m > 0$. Let us recall that the interaction function is linear: $F'(x) = \alpha x$ with $\alpha > 0$. In order to study the invariant measures, we have to consider the fixed points of the application $\Psi_\epsilon(m)$ defined by (3.3). We introduce the following potential function:

$$W_m(x) = V(x) + \frac{\alpha}{2}x^2 - \alpha mx.$$

Since $V'(0) = 0$, we have $W'_m(0) < 0$. Moreover $\lim_{x \rightarrow +\infty} W'_m(x) = +\infty$. So we denote by x_m the positive real for which the potential W_m admits its global minimum. It is uniquely determined

since V'' (and so W_m'') is a convex function. Let us prove by using *reductio ad absurdum* that the global minimum is uniquely determined: we assume the existence of two positive reals $x_m^{(1)} < x_m^{(2)}$ such that W_m reaches its minimum on \mathbb{R} for $x = x_m^{(1)}$ and $x = x_m^{(2)}$. Hence $W_m'(x_m^{(1)}) = W_m'(x_m^{(2)}) = 0$ and $W_m''(x_m^{(1)}) \geq 0$. Since W_m'' is non-decreasing, we deduce that $W_m''(x) = 0$ for all $x \in [x_m^{(1)}; x_m^{(2)}]$. The convexity of W_m'' implies that $W_m^{(3)}$ is a non-decreasing function. Moreover $W_m^{(3)}(0) = V^{(3)}(0) = 0$ which leads to $W_m^{(3)}(x) = 0$ and so $W_m''(x) = 0$ for all $x \in [0; x_m^{(2)}]$. The following contradiction holds: $W_m'(0) = -\alpha m \neq 0 = W_m'(x_m^{(2)})$. Consequently, x_m is unique.

Moreover, x_m satisfies

$$V'(x_m) + \alpha(x_m - m) = 0 \quad \text{and} \quad V''(x_m) + \alpha > 0. \tag{3.5}$$

Indeed, since x_m is a global minimum, the equality $V''(x_m) + \alpha = 0$ implies that $V^{(3)}(x_m) = 0$ that is $x_m = 0$ which contradicts the assumption concerning the positivity of x_m .

We define

$$\chi_\epsilon(m) = \Psi_\epsilon(m) - m \quad \text{and} \quad \chi_0(m) = x_m - m.$$

We obtain the expression:

$$\chi_\epsilon(m) = x_m - m + \frac{\int_{\mathbb{R}} (x - x_m) \exp\left[-\frac{2}{\epsilon} \left(V(x) + \frac{\alpha x^2}{2} - \alpha m x\right)\right] dx}{\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon} \left(V(x) + \frac{\alpha x^2}{2} - \alpha m x\right)\right] dx}. \tag{3.6}$$

It suffices to prove that χ_ϵ has just one zero in \mathbb{R}_+^* .

Step 1. For all $\epsilon > 0$ and $m > 0$, we observe that $\chi_\epsilon(m) \leq \chi_0(m) = x_m - m$.

We apply the change of variable $x := y + x_m$ to the integrals in (3.6) and obtain

$$\begin{aligned} \chi_\epsilon(m) &= \chi_0(m) + \frac{\int_{\mathbb{R}} y \exp\left[-\frac{2}{\epsilon} \left(V(y + x_m) + \frac{\alpha y^2}{2} + \alpha(x_m - m)y\right)\right] dy}{\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon} \left(V(y + x_m) + \frac{\alpha y^2}{2} + \alpha(x_m - m)y\right)\right] dy} \\ &= \chi_0(m) + \frac{\int_0^\infty y \exp\left[-\frac{\alpha}{\epsilon} y^2\right] \Omega_{\epsilon,m}(y) dy}{\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon} \left(V(y + x_m) + \frac{\alpha y^2}{2} + \alpha(x_m - m)y\right)\right] dy}, \end{aligned}$$

with

$$\begin{aligned} \Omega_{\epsilon,m}(y) &= \exp\left[-\frac{2}{\epsilon} \left(V(y + x_m) + \alpha(x_m - m)y\right)\right] \\ &\quad - \exp\left[-\frac{2}{\epsilon} \left(V(y - x_m) - \alpha(x_m - m)y\right)\right]. \end{aligned}$$

We introduce the function

$$A_m(y) = V(y + x_m) - V(y - x_m) + 2\alpha(x_m - m)y.$$

Since V is an even function, $A_m(0) = 0$ and $A_m''(0) = 0$. According to the definition of x_m , $A_m'(0) = 0$. V'' is a convex function therefore $V^{(3)}$ is increasing. So $A_m^{(3)}(y) = V^{(3)}(y + x_m) - V^{(3)}(y - x_m) \geq 0$ for all y . We deduce that A_m'' is increasing. Hence A_m'' is nonnegative on \mathbb{R}_+^* so does $A_m(y)$ for $y > 0$. Finally we get $\Omega_{\epsilon,m}(y) \leq 0$ for all $y > 0$. We obtain the announced result: $\chi_\epsilon(m) \leq \chi_0(m)$ for $m > 0$.

Step 2. χ_0 has a unique zero on \mathbb{R}_+^* .

Let us compute $\chi_0(a)$ with a defined in (V-3). We know that a is solution of $V'(x) + \alpha(x - a) = 0$ with $V''(x) + \alpha > 0$. Hence $\chi_0(a) = 0$.

Let us focus our attention on the variations of the function χ_0 on the interval $]0, +\infty[$. Since $V'(x_m) + \alpha x_m = \alpha m$, and $\alpha + V''(x_m) > 0$ we deduce that $m \rightarrow x_m$ is derivable, we obtain

$$\chi_0'(m) = \frac{d}{dm} x_m - 1$$

and

$$\frac{d}{dm} x_m = \frac{\alpha}{\alpha + V''(x_m)} > 0 \quad \text{which implies } \chi_0'(m) = -\frac{V''(x_m)}{\alpha + V''(x_m)}. \tag{3.7}$$

The denominator is positive due to the definition of x_m . According to (V-5), $V''(x) > 0$ for all $x > a$. Hence $\chi_0'(m) < 0$ for all $m > a$. Since $\chi_0(a) = 0$ we deduce that, for all $m > a$, $\chi_0(m)$ is negative and therefore the function χ_0 has no zero on $]a; +\infty[$.

It remains to study χ_0 on the interval $]0, a]$. Since V'' is a convex function, we deduce that the derivative of χ_0 is non positive for $x_m \geq c$ with $c > 0$ satisfying $V''(c) = 0$. We know that $c > 0$ is unique since $V''(0) < 0$ and V'' is a convex function. Moreover $c < a$. Since the function $m \rightarrow x_m$ is increasing for $m > 0$, we deduce that χ_0' is negative for $m \in]\max(0, m_c), a]$ where m_c is such that c is the solution of (3.5). By computation, $m_c = c + \frac{V'(c)}{\alpha}$.

We observe then two different cases:

- If $m_c \leq 0$ i.e. $\alpha < \frac{|V'(c)|}{c}$: χ_0 is decreasing on \mathbb{R}_+^* with $\chi_0(a) = 0$. The unique zero of χ_0 on \mathbb{R}_+^* is a .
- If $m_c > 0$ then χ_0 , which is a continuous function on \mathbb{R}_+^* , is increasing on $]0, m_c[$ and decreasing on $]m_c, +\infty[$ with $\chi_0(a) = 0$. It suffices to prove that $\lim_{m \rightarrow 0+} \chi_0(m) \geq 0$ in order to conclude that a is the unique zero of χ_0 on \mathbb{R}_+^* . Due to the definition of x_m we get: $\lim_{m \rightarrow 0+} \chi_0(m) = \lim_{m \rightarrow 0+} x_m \geq 0$. Indeed $m \rightarrow x_m$ is increasing on $]0; m_c[$ and $x_m > 0$ for all $m \in]0; m_c[$ so the extension to $m = 0$ is non negative.

In these two cases, there is a unique zero of χ_0 on \mathbb{R}_+^* .

Step 3. The family of functions $(\chi_\epsilon)_\epsilon$ (respectively $(\chi'_\epsilon)_\epsilon$) converges uniformly towards χ_0 (resp. χ'_0) on each compact subset of \mathbb{R}_+^* .

First we prove the convergence of $\Psi_\epsilon(m)$ for $m > 0$. Recall that

$$\Psi_\epsilon(m) = \frac{\int_{\mathbb{R}} x \exp \left[-\frac{2}{\epsilon} \left(V(x) + \frac{\alpha x^2}{2} - \alpha m x \right) \right] dx}{\int_{\mathbb{R}} \exp \left[-\frac{2}{\epsilon} \left(V(x) + \frac{\alpha x^2}{2} - \alpha m x \right) \right] dx}.$$

By Lemma A.3 with $U(x) = V(x) + \frac{\alpha x^2}{2}$, $n = 1$, $\mu = m$ and $G = -\alpha x$ we obtain the announced convergence result:

$$\chi_\epsilon(m) - \chi_0(m) = \Psi_\epsilon(m) - x_m = -\frac{V^{(3)}(x_m)}{4(\alpha + V''(x_m))^2} \epsilon + o(\epsilon).$$

Moreover this convergence is uniform with respect to the variable m on compact subsets of \mathbb{R}_+^* .

We estimate now the asymptotics of $\chi'_\epsilon(m)$ as ϵ becomes small. Taking the derivative of Ψ_ϵ , we obtain

$$\Psi'_\epsilon(m) = \frac{2\alpha}{\epsilon} \left\{ \frac{\int_{\mathbb{R}} x^2 \exp\left[-\frac{2}{\epsilon} W_m(x)\right] dx}{\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon} W_m(x)\right] dx} - \left(\frac{\int_{\mathbb{R}} x \exp\left[-\frac{2}{\epsilon} W_m(x)\right] dx}{\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon} W_m(x)\right] dx} \right)^2 \right\}.$$

We recognize the variance of the measure $u_\epsilon^{(m)}$ which is the measure associated to the average m by (3.2). Hence

$$\chi'_\epsilon(m) = \frac{2\alpha}{\epsilon} \text{Var}(u_\epsilon^{(m)}) - 1. \tag{3.8}$$

Applying again Lemma A.3 with $U = V(x) + \frac{\alpha x^2}{2}$, $G = -\alpha x$, $\mu = m$ and $n = 2$, we obtain

$$\frac{\int_{\mathbb{R}} x^2 \exp\left[-\frac{2}{\epsilon} W_m(x)\right] dx}{\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon} W_m(x)\right] dx} = x_m^2 - \frac{(x_m V^{(3)}(x_m) - (\alpha + V''(x_m)))}{2(\alpha + V''(x_m))^2} \epsilon + o(\epsilon).$$

Applying the same lemma with $n = 1$ permits to compute the first moment:

$$\frac{\int_{\mathbb{R}} x \exp\left[-\frac{2}{\epsilon} W_m(x)\right] dx}{\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon} W_m(x)\right] dx} = x_m - \frac{x_m V^{(3)}(x_m)}{4x_m(\alpha + V''(x_m))^2} \epsilon + o(\epsilon).$$

By (3.8) and the computations of the two first moments, we get

$$\chi'_\epsilon(m) = \frac{-V''(x_m)}{\alpha + V''(x_m)} + o(1) = \chi'_0(m) + o(1). \tag{3.9}$$

Furthermore this convergence is uniform with respect to the variable m on compact subsets of \mathbb{R}_+^* .

Step 4. For any $\delta > 0$ small enough, there exists $\epsilon_0 > 0$ such that χ_ϵ has a unique zero on $[\delta, \infty[$ for all $\epsilon \leq \epsilon_0$.

Since there is no zero of χ_ϵ on the interval $]a, +\infty[$ (Step 1 and 2), we focus our attention on the interval $]0, a[$. On each compact subset of this interval, χ_ϵ converges uniformly towards the limit function χ_0 (Step 3). Hence the zeros of χ_ϵ are in a small neighborhood of the unique zero of χ_0 namely a (Step 2). Let us study the derivative of χ_ϵ in a neighborhood of a . Since χ'_ϵ converges uniformly towards χ'_0 (Step 3) and $\chi'_0(m) < 0$ in a neighborhood of a (Step 2), we obtain that $\chi'_\epsilon(m) < 0$ in a neighborhood of a for ϵ small enough. Finally we proved that, as soon as ϵ is small enough, the function χ_ϵ cannot admit two zeros or more on \mathbb{R}_+^* .

Step 5. There exists $\delta > 0$ and $\epsilon_0 > 0$ such that χ_ϵ doesn't vanish on $]0, \delta[$ for all $\epsilon \leq \epsilon_0$.

In this last step, we have to distinguish three different cases depending on the values $\vartheta (= -V''(0)$ since V'' is convex) and α defined by (1.2) and (1.3).

Step 5.1. We assume $\alpha < \vartheta$. In this particular case $W_0(x) = V(x) + \alpha x^2/2$ reaches a unique global minimum on \mathbb{R}_+ for $x = x_0 > 0$.

Let us fix some small $\delta > 0$ (depending on x_0 : we shall precise it in the following). We prove that, for ϵ small enough, $\chi_\epsilon(m) = \Psi_\epsilon(m) - m > 0$ on $]0, \delta[$. By the definition of Ψ_ϵ , see (3.3), it suffices to prove that $N_\epsilon(m) > 0$ for $m \in]0, \delta[$ where

$$N_\epsilon(m) = \int_{\mathbb{R}} x \exp\left[-\frac{2}{\epsilon} W_m(x)\right] dx - m \int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon} W_m(x)\right] dx. \tag{3.10}$$

Obviously $N_\epsilon(0) = 0$. Let us prove that N_ϵ is non decreasing. Taking the derivative, we get

$$N'_\epsilon(m) = \frac{2\alpha}{\epsilon} \int_{\mathbb{R}} \left(x^2 - mx - \frac{\epsilon}{2\alpha}\right) \exp\left[-\frac{2}{\epsilon}W_m(x)\right] dx.$$

This expression is in fact non negative. Indeed, using the symmetry property of $W_0(x)$ and the upper bound $m \leq \delta$, we obtain

$$\begin{aligned} N'_\epsilon(m) &= \frac{2\alpha}{\epsilon} \int_0^\infty \left\{ \left(x^2 - \frac{\epsilon}{2\alpha}\right) \cosh\left(\frac{2\alpha mx}{\epsilon}\right) - mx \sinh\left(\frac{2\alpha mx}{\epsilon}\right) \right\} e^{-\frac{2}{\epsilon}W_0(x)} dx \\ &\geq \frac{\alpha}{\epsilon} \int_0^\infty P_\delta(x) e^{\frac{2\alpha mx}{\epsilon}} e^{-\frac{2}{\epsilon}W_0(x)} dx \quad \text{with } P_\delta(x) = x^2 - \delta x - \frac{\epsilon}{\alpha}. \end{aligned}$$

We split the preceding integral into two parts: the first integral I_0 concerns the support $[0, 2\delta]$ and the second integral $I_{2\delta}$ the complementary support $[2\delta, \infty[$. We get $N'_\epsilon(m) \geq \frac{\alpha}{\epsilon}(I_0 + I_{2\delta})$.

Since the roots of the polynomial function P_δ satisfy

$$x_\pm = \frac{1}{2} \left(\delta \pm \sqrt{\delta^2 + \frac{4\epsilon}{\alpha}} \right) < 2\delta,$$

the polynomial is positive on the interval $[2\delta, \infty[$ and can be lower bounded by $P_\delta(2\delta) = 2\delta^2 - \epsilon/\alpha$. Lemma A.3 implies the existence of some constant $C > 0$ leading to the following estimate as $\epsilon \rightarrow 0$:

$$I_{2\delta} \geq (2\delta^2 - \epsilon/\alpha) \int_{2\delta}^{x_0+1} e^{-\frac{2}{\epsilon}W_0(x)} dx \geq C\delta^2\sqrt{\epsilon}e^{-\frac{2}{\epsilon}(V(x_0)+\alpha x_0^2/2)} \tag{3.11}$$

provided that $x_0 > 2\delta$ (it suffices then to chose δ small enough).

Let us finally focus our attention to the lower bound of the integral term I_0 . Since the minimum value of P_δ is $-(\delta^2/4 + \epsilon/\alpha)$ and since $W''(0) < 0$, we have

$$I_0 \geq -\left(\frac{\delta^2}{4} + \frac{\epsilon}{\alpha}\right) \int_0^{2\delta} e^{\frac{2\alpha mx}{\epsilon}} e^{-\frac{2}{\epsilon}W_0(x)} dx \geq -2\delta \left(\frac{\delta^2}{4} + \frac{\epsilon}{\alpha}\right) e^{-\frac{V(2\delta)}{\epsilon}}. \tag{3.12}$$

For $\delta > 0$ small enough, $V(2\delta) > V(x_0) + \alpha x_0^2/2$ (since the minimum of $V(x) + \alpha x^2/2$ is only reached for $x = x_0$). Consequently the negative lower bound of I_0 (3.12) is negligible with respect to the positive lower bound of $I_{2\delta}$ as ϵ becomes small. We deduce that there exists ϵ_0 such that $N'_\epsilon(m) > 0$ for all $m \in [0, \delta]$ and $\epsilon \leq \epsilon_0$. Since $N_\epsilon(0) = 0$ we conclude that $N_\epsilon(m) > 0$ on $]0, \delta]$ and so is χ_ϵ .

Step 5.2. We assume $\alpha > \vartheta$. In this case $W_0(x)$ admits a unique minimum reached for $x = 0$ and x_m converges continuously to 0 as $m \rightarrow 0$. Using similar arguments as those presented in Step 3, we claim that χ_ϵ (resp. χ'_ϵ) converges towards χ_0 (resp. χ'_0) uniformly on $[0, a]$ as $\epsilon \rightarrow 0$. Due to the regularity of χ_0 and by the inequality $\chi'_0(0) = -\frac{V''(0)}{\alpha+V''(0)} > 0$ we obtain the existence of $\delta > 0$ and $\epsilon_0 > 0$ such that $\chi'_\epsilon(m) > 0$ for $m \in [0, \delta]$ and $\epsilon \leq \epsilon_0$. χ_ϵ starts in 0 and is increasing on $[0, \delta]$ which implies the announced result.

Step 5.3. We assume that $\alpha = \vartheta$. It suffices then to note that χ_ϵ depends continuously on the parameter α . The following results can be directly deduced from the preceding case (Step 5.2) by continuity: $\chi'_\epsilon(0) > 0$ and $\chi'_\epsilon(m) \geq 0$ for $m \in [0, \delta]$ and $\epsilon \leq \epsilon_0$. In fact χ_ϵ vanishes for $x = 0$ and is increasing on $[0, \delta]$. The inequality $\chi_\epsilon(m) > 0$ for all $m \in]0, \delta]$ and $\epsilon \leq \epsilon_0$ is an obvious consequence.

Conclusion: Step 4 and 5 lead to the existence of $\epsilon_0 > 0$ such that for all $\epsilon < \epsilon_0$, χ_ϵ has exactly three zeros: 0 and two other reals, one in the neighborhood of a , the other one near $-a$. To each of these averages corresponds a unique invariant measure obtained by (3.2). \square

4. The general interaction case

We assumed for this study that the self-attraction phenomenon is represented by a polynomial function F' , see (F-1). In previous section, we analyzed the particular linear situation: $F'(x) = \alpha x$ and proved under suitable conditions that there exist exactly three invariant measures in the small noise limit. In this section we shall focus our attention to the general case: the polynomial function F is of degree $n \geq 2$. First we shall present results, concerning symmetric invariant measures, derived from the application of a fixed point theorem to some suitable function space. Secondly, we deal with the uniqueness of stationary symmetric measures in particular situations using similar methods as those developed in Proposition 3.1. Thirdly, we discuss the existence of asymmetric measures based on the application of a fixed point theorem to some finite-dimensional space.

4.1. Symmetric invariant measures

In the linear case we proved the existence of a unique symmetric invariant measure. The result is obvious since it suffices to solve the Eq. (3.3) with $m_1(\epsilon) = 0$. In the general case in order to find the symmetric measure we have to solve some equation like (3.2) but depending on much more parameters than just the mean $m_1(\epsilon)$. The total number of parameters depends in fact on the degree of F . Instead of trying to solve such a system, we choose some other kind of proof based on a fixed point theorem which permits to prove the existence of symmetric invariant measures in even more general cases: the interaction function does not need to be polynomial. In [2], Benachour, Roynette, Talay and Vallois introduced this method of proof for a self-stabilizing diffusion in the constant environment case ($V'(x) = 0$). This proof can be adapted to our situation and is based on the following Schauder’s theorem (see for instance [5] Corollary 11.2 p. 280):

Proposition 4.1. *Let \mathbb{B} a Banach space, \mathbb{C} a closed convex subset and \mathbb{A} a continuous application $\mathbb{C} \rightarrow \mathbb{C}$ such that $\overline{\mathbb{A}(\mathbb{C})}$ is compact. Then \mathbb{A} admits a fixed point in \mathbb{C} .*

In order to use this proposition we introduce some definitions and notations:

1. Let us choose $p > 4q$ where q is defined in (V-6).
2. $\mathbb{B} = \{f : \mathbb{R} \rightarrow \mathbb{R}; \sup_{x \in \mathbb{R}} (1 + |x|^p) |f(x)| < \infty\}$. \mathbb{B} is equipped with the norm $|\cdot|_\infty$ where $|f|_\infty = \sup_{x \in \mathbb{R}} (1 + |x|^p) |f(x)|$.
3. For all $M > 0$ we define the function space \mathbb{C}_M as the subset of all non negative and even function belonging to \mathbb{B} which satisfy:

$$\int_{\mathbb{R}} f(x)dx = 1 \quad \text{and} \quad \sup_{x \in \mathbb{R}} (1 + |x|^p) f(x) \leq M.$$

4. For any function $f \in \mathbb{C}_M$ we define the operator:

$$\begin{aligned} \mathbb{A}^\epsilon(f)(x) &= \frac{\exp\left[-\frac{2}{\epsilon} \left(V(x) + \int_0^x (F' * f)(y)dy\right)\right]}{\int_{z \in \mathbb{R}} \exp\left[-\frac{2}{\epsilon} \left(V(z) + \int_0^z (F' * f)(y)dy\right)\right]} \\ &= \frac{1}{\lambda_\epsilon(f)} \exp\left[-\frac{2}{\epsilon} \left(V(x) + \int_0^x (F' * f)(y)dy\right)\right], \end{aligned} \tag{4.1}$$

where $\lambda_\epsilon(f)$ is the normalization factor.

5. For any function $u \in \mathbb{C}_M$, we define the moments $\gamma_k(u) = \int_{\mathbb{R}} |x|^k u(x) dx$ with $0 \leq k \leq p - 2$.

Let us just point out that \mathbb{C}_M is a closed and convex subset of \mathbb{B} . The aim of this section will consist in proving that the application \mathbb{A}^ϵ is $\mathcal{C}^0(\mathbb{C}_M, \mathbb{C}_M)$ -continuous and that $\overline{\mathbb{A}^\epsilon(\mathbb{C}_M)}$ is compact since M is large enough. Therefore Schauder’s theorem implies the existence of a fixed point and as the matter of fact the existence of an invariant measure in the function space \mathbb{C}_M .

Lemma 4.2. *There exists two constants C_1 and C_2 both independent of M such that for all $u \in \mathbb{C}_M$, we have: $\gamma_k(u) \leq MC_1$ and for all $x \in \mathbb{R}$:*

$$\frac{\alpha}{2} x^2 \leq \int_0^x (F' * u)(y) dy \leq C_2 M x^2 (1 + x^{2q}). \tag{4.2}$$

Proof. 1. Since $u \in \mathbb{C}_M$, we have $\sup_{x \in \mathbb{R}} (1 + |x|^p) u(x) \leq M$ for all $k \leq p - 2$. Therefore

$$\gamma_k(u) = \int_{\mathbb{R}} \frac{|x|^k}{1 + |x|^p} (1 + |x|^p) u(x) dx \leq M \int_{\mathbb{R}} \frac{|x|^k}{1 + |x|^p} dx \leq MC_1.$$

2. Let $x \geq 0$. Since $u \in \mathbb{C}_M$, u is an even function. By (1.3) we have $F'(x) = \alpha x + F'_0(x)$ and (F-3) implies that F' and F'_0 are non negative odd functions, therefore $F'_0 * u$ is also odd and non negative. Using the inequality developed in the statement of Lemma 4.3 in [2] and the assumption (F-3), we have

$$F'_0(x) \leq \frac{1}{2} (F'_0(x - y) + F'_0(x + y)) \quad \text{for } y \in \mathbb{R}, x \geq 0.$$

Therefore, for $x \geq 0$:

$$\begin{aligned} \int_0^x (F'_0 * u)(y) dy &= \int_0^x \int_0^\infty (F'_0(y - z) + F'_0(y + z)) u(z) dz dy \\ &\geq \int_0^x \int_0^\infty 2F'_0(y) u(z) dz dy \geq 0. \end{aligned} \tag{4.3}$$

From the preceding inequality we deduce

$$\int_0^x (F' * u)(y) dy = \int_0^x (F'_0 * u)(y) dy + \frac{\alpha}{2} x^2 \geq \frac{\alpha}{2} x^2 \quad \text{for all } x \geq 0.$$

Since $\int_0^x (F' * u)(y) dy$ is an even function, we get the inequality for all $x \in \mathbb{R}$.

3. Due to the symmetry of $F' * u$ we restrict our study to $x \geq 0$.

$$\int_0^x (F' * u)(y) dy = \frac{1}{2} \int_0^x \int_0^\infty (F'(y - z) + F'(y + z)) u(z) dz dy.$$

According to the assumptions (F-1) and (F-4), F is an even polynomial function of degree smaller than $2q$ with $q \geq 1$. We can therefore write F' as follows

$$F'(x) = \sum_{k=0}^{q-1} \alpha_k x^{2k+1}.$$

Therefore defining $\mathcal{F}(y, z) = F'(y - z) + F'(y + z)$ we get

$$\begin{aligned} \mathcal{F}(y, z) &= y \sum_{k=0}^{q-1} \alpha_k \sum_{j=0}^k C_{2k+1}^{2j+1} y^{2j} z^{2k-2j} \\ &\leq qy \max_{0 \leq k \leq q-1} |\alpha_k| 2^{2q} \max_{0 \leq j \leq q} \sum_{j=0}^k y^{2j} z^{2k-2j} \leq Cy \left(1 + y^{2q}\right) \left(1 + z^{2q}\right). \end{aligned}$$

Finally since $p > 4q$, there exists some constant $C' > 0$ such that:

$$\begin{aligned} \int_0^\infty \mathcal{F}(y, z) u(z) dz &\leq Cy \left(1 + y^{2q}\right) \int_0^\infty \left(1 + z^{2q}\right) u(z) dz \\ &\leq Cy \left(1 + y^{2q}\right) \int_0^\infty \frac{1 + z^{2q}}{1 + z^p} \left((1 + z^p) u(z)\right) dz \\ &\leq C' y M \left(1 + y^{2q}\right). \end{aligned}$$

By integration we obtain $\int_0^x (F' * u)(y) dy \leq C_2 M x^2 (1 + x^{2q})$ for all $x \in \mathbb{R}_+$. \square

Lemma 4.3. *There exists $M_0 > 0$ such that for any $M \geq M_0$, $\mathbb{A}^\epsilon(\mathbb{C}_M) \subset \mathbb{C}_M$. Moreover, there exists a constant $C(\epsilon)$ such that*

$$\frac{1}{\lambda_\epsilon(u)} \leq C(\epsilon) \sqrt{M} \tag{4.4}$$

for all $u \in \mathbb{C}_M$.

Proof. By construction $\mathbb{A}^\epsilon u$ is a non negative even function which satisfies $\int_{\mathbb{R}} \mathbb{A}^\epsilon u(x) dx = 1$. It suffices then to prove that:

$$\sup_{x \in \mathbb{R}} \left(1 + |x|^p\right) \mathbb{A}^\epsilon u(x) \leq M.$$

By (4.1) and according to Lemma 4.2 we obtain some lower bound for the normalization factor:

$$\begin{aligned} \lambda_\epsilon(u) &= \int_{-\infty}^{+\infty} \exp \left[-\frac{2}{\epsilon} \left(V(x) + \int_0^x (F' * u)(y) dy \right) \right] dx \\ &\geq \int_{-\infty}^{+\infty} \exp \left[-\frac{2}{\epsilon} \left(V(x) + C_2 M x^2 (1 + x^{2q}) \right) \right] dx. \end{aligned}$$

According to both (V-3) and (V-7), we know that $V(x) \leq 0$ for all $x \in [-a; a]$. Hence

$$\lambda_\epsilon(u) \geq \int_{-a}^{+a} \exp \left[-\frac{2}{\epsilon} C_2 M x^2 (1 + a^{2q}) \right] dx.$$

By the change of variable $x := \frac{y}{\sqrt{M}}$ and Lemma A.1, the following development holds

$$\begin{aligned} \int_{-a}^{+a} \exp \left[-\frac{2}{\epsilon} C_2 M x^2 (1 + a^{2q}) \right] dx &= \frac{2}{\sqrt{M}} \int_0^{\frac{a}{\sqrt{M}}} \exp \left[-\frac{2}{\epsilon} C_2 x^2 (1 + a^{2q}) \right] dx \\ &= \frac{1}{\sqrt{M}} \{C(\epsilon) + o(1)\} \end{aligned}$$

where $C(\epsilon)$ is independent of M and $o(1) \rightarrow 0$ as $M \rightarrow \infty$. As soon as M is large enough, we have therefore $\frac{1}{\lambda_\epsilon(u)} \leq \frac{2\sqrt{M}}{C(\epsilon)}$ where $C(\epsilon)$ is a positive constant determined by parameters of the global system and ϵ . By (4.1) and the preceding upper bound, we prove that

$$(1 + |x|^p)\mathbb{A}^\epsilon u(x) \leq \frac{2\sqrt{M}}{C(\epsilon)}(1 + |x|^p)e^{-\frac{2}{\epsilon}V(x)} \leq C'(\epsilon)\sqrt{M},$$

where $C'(\epsilon)$ is a positive constant similar to $C(\epsilon)$. In order to conclude, it is sufficient to choose $M \geq C'(\epsilon)^2$: we get immediately $\mathbb{A}^\epsilon u \in \mathbb{C}_M$. \square

Lemma 4.4. \mathbb{A}^ϵ is a continuous operator on \mathbb{C}_M with respect to the uniform norm.

Proof. We shall find some upper bound for the following expression $|\mathbb{A}^\epsilon u - \mathbb{A}^\epsilon v|$.

Step 1. Let $u, v \in \mathbb{C}_M$. We define:

$$\begin{aligned} A^\epsilon(x) &= e^{-\frac{2}{\epsilon}V(x)} \left\{ \exp \left[-\frac{2}{\epsilon} \int_0^x (F' * u)(y)dy \right] - \exp \left[-\frac{2}{\epsilon} \int_0^x (F' * v)(y)dy \right] \right\} \\ &= e^{-\frac{2}{\epsilon}V(x) - \frac{\alpha}{\epsilon}x^2} \left\{ \exp \left[-\frac{2}{\epsilon} \int_0^x (F'_0 * u)(y)dy \right] - \exp \left[-\frac{2}{\epsilon} \int_0^x (F'_0 * v)(y)dy \right] \right\}. \end{aligned}$$

It is well known that $|e^{-a} - e^{-b}| \leq |a - b|$ for $a, b \geq 0$. In order to apply this inequality we have to prove that $\int_0^x (F'_0 * v)(y)dy$ and $\int_0^x (F'_0 * u)(y)dy$ are non negative. For each function $f \in \mathbb{C}_M$ the convolution term $\int_0^x (F'_0 * f)(y)dy$ is non negative due to (4.3). Hence

$$|A^\epsilon(x)| \leq \frac{2}{\epsilon}e^{-\frac{2}{\epsilon}V(x) - \frac{\alpha}{\epsilon}x^2} A_0^\epsilon(x), \tag{4.5}$$

with A_0^ϵ defined by

$$\left| \int_0^x (F'_0 * u)(y)dy - \int_0^x (F'_0 * v)(y)dy \right| = \left| \int_0^x \int_{\mathbb{R}} F'_0(y - z)(u(z) - v(z))dzdy \right|.$$

Since u and v are elements of \mathbb{C}_M , they are even functions and the integral with respect to the variable z becomes

$$\begin{aligned} A_0^\epsilon &= \left| \int_0^x \int_0^\infty (F'_0(z + y) - F'_0(z - y)) (u(z) - v(z)) dzdy \right| \\ &\leq \int_0^x \int_0^\infty |F'_0(z + y) - F'_0(z - y)| |u(z) - v(z)| dzdy. \end{aligned} \tag{4.6}$$

The assumption (F-4) gives information about the increments of the interaction function: there exist two positive constants C'_q and C such that

$$\begin{aligned} |F'_0(z + y) - F'_0(z - y)| &\leq 2|y|C'_q \left(1 + |z + y|^{2q-2} + |z - y|^{2q-2} \right) \\ &\leq 2|y|C'_q \left(1 + 2^{2q-1}|z|^{2q-2} + 2^{2q-1}|y|^{2q-2} \right) \\ &\leq C|y| \left(1 + |y|^{2q-1} + |z|^{2q-1} \right) \\ &\leq C|y| \left(1 + |y|^{2q-1} \right) \left(1 + |z|^{2q-1} \right). \end{aligned} \tag{4.7}$$

We shall now find some upper bound for $|u(z) - v(z)|$ in (4.6). Since $u, v \in \mathbb{C}_M$ then $u(z)(1 + |z|^p) \leq M$ and $v(z)(1 + |z|^p) \leq M, \forall z \in \mathbb{R}$. The obvious upper bound $|u(z) - v(z)|(1 + |z|^p) \leq$

$2M$ permits to obtain $\sqrt{|u(z) - v(z)|} \leq \sqrt{\frac{2M}{1+|z|^p}}$. Consequently, for all z of \mathbb{R} , $|u(z) - v(z)| \leq \sqrt{\|u - v\|_\infty} \sqrt{\frac{2M}{1+|z|^p}}$ where $\|\cdot\|_\infty$ denotes the uniform norm. Using this inequality, (4.7) and (4.6) in order to estimate A_0^ϵ , we get

$$|A_0^\epsilon(x)| \leq C\sqrt{\|u - v\|_\infty} \int_0^x |y| (1 + |y|^{2q-1}) dy \int_0^\infty \sqrt{\frac{2M}{1+|z|^p}} (1 + z^{2q-1}) dz.$$

Since $p > 4q$ the integral with respect to the variable z is finite and can be considered like a constant term. By (4.5) and using the positivity of αx^2 , we obtain directly the existence of some positive constant $C > 0$ such that

$$|A^\epsilon(x)| \leq C \frac{\sqrt{M}}{\epsilon} \sqrt{\|u - v\|_\infty} x^2 (1 + |x|^{2q-1}) e^{-\frac{2}{\epsilon}V(x)}.$$

According to (V-4), the expression $x^2 (1 + |x|^{2q-1}) e^{-\frac{2}{\epsilon}V(x)}$ can be bounded by some constant independent of M . Therefore

$$\|A^\epsilon\|_\infty \leq C(M, \epsilon) \sqrt{\|u - v\|_\infty}. \tag{4.8}$$

Two results can be deduced: firstly $\|A^\epsilon\|_\infty$ is finite and secondly $\|A^\epsilon\|_\infty$ becomes small as $\|u - v\|_\infty$ decreases towards 0.

Step 2. For any $x \in \mathbb{R}$, we introduce:

$$\Omega_\epsilon(x) = \frac{1}{\lambda_\epsilon(u)\lambda_\epsilon(v)} \exp \left[-\frac{2}{\epsilon} \left(\int_0^x (F' * v)(y) dy + V(x) \right) \right]. \tag{4.9}$$

Then the difference $\mathbb{A}^\epsilon u(x) - \mathbb{A}^\epsilon v(x)$ can be decomposed as follows:

$$\mathbb{A}^\epsilon u(x) - \mathbb{A}^\epsilon v(x) = \frac{1}{\lambda_\epsilon(u)} A^\epsilon(x) + (\lambda_\epsilon(v) - \lambda_\epsilon(u)) \Omega_\epsilon(x). \tag{4.10}$$

Taking the uniform norm, we get

$$\|\mathbb{A}^\epsilon u - \mathbb{A}^\epsilon v\|_\infty \leq \frac{1}{\lambda_\epsilon(u)} \|A^\epsilon\|_\infty + |\lambda_\epsilon(v) - \lambda_\epsilon(u)| \|\Omega_\epsilon\|_\infty. \tag{4.11}$$

By (4.4) and by (4.8), we deduce that

$$\frac{1}{\lambda_\epsilon(u)} \|A^\epsilon\|_\infty \leq C'(M, \epsilon) \sqrt{\|u - v\|_\infty}.$$

It is then sufficient to find a similar inequality for the term $|\lambda_\epsilon(v) - \lambda_\epsilon(u)| \|\Omega_\epsilon\|_\infty$ in order to conclude the proof.

$$\begin{aligned} |\lambda_\epsilon(v) - \lambda_\epsilon(u)| &= \left| \int_{\mathbb{R}} A^\epsilon(x) dx \right| \\ &\leq C \frac{\sqrt{M}}{\epsilon} \sqrt{\|u - v\|_\infty} \int_{-\infty}^{+\infty} x^2 (1 + |x|^{2q-1}) e^{-\frac{2}{\epsilon}V(x)} dx. \end{aligned}$$

According to (V-4), the integral with respect to the variable x is finite and does not depend on M . We have immediately

$$|\lambda_\epsilon(v) - \lambda_\epsilon(u)| \leq C(M, \epsilon) \sqrt{\|u - v\|_\infty}.$$

It remains to estimate $\Omega_\epsilon(x)$. By (V-4) and (4.2), we have

$$\int_0^x (F' * v)(y)dy + V(x) \geq C_4x^4 + \left(\frac{\alpha}{2} - C_2\right)x^2$$

for all x positive. Furthermore the symmetry property of V and F permits to extend the bound to all $x \in \mathbb{R}$. The function $\exp\left[-\frac{2}{\epsilon}\left(\int_0^x (F' * v)(y)dy + V(x)\right)\right]$ is then bounded by a constant depending on ϵ . Moreover we have already proved (4.4) that is to say $\frac{1}{\lambda_\epsilon(f)} \leq C(\epsilon)\sqrt{M}$ for all elements f of the function space \mathbb{C}_M . This bound can therefore be applied to u and v . Finally we obtain the existence of some constant $C(\epsilon) > 0$ such that, for all real value x , $|\Omega_\epsilon(x)| \leq C(\epsilon)M$.

By (4.10), we have

$$\|\mathbb{A}^\epsilon u - \mathbb{A}^\epsilon v\|_\infty \leq C'(M, \epsilon)\sqrt{\|u - v\|_\infty} + C(M, \epsilon)\sqrt{\|u - v\|_\infty}C(\epsilon)M.$$

In other words,

$$\|\mathbb{A}^\epsilon u - \mathbb{A}^\epsilon v\|_\infty \leq C''(M, \epsilon)\sqrt{\|u - v\|_\infty}$$

which finishes the proof. \square

We have now all the keys for proving the existence of some symmetric invariant measure. Indeed we have just presented some continuous mapping which stabilizes a convex subset of the Banach space \mathbb{B} .

Theorem 4.5. *There exists a symmetric invariant measure for (1.1).*

Proof. Let M_0 defined by Lemma 4.3. Taking $M \geq M_0$, let us prove that $\overline{\mathbb{A}^\epsilon(\mathbb{C}_M)}$ is a compact set. For this reason we shall estimate the following derivative:

$$(\mathbb{A}^\epsilon u)'(x) = -\frac{2}{\epsilon} \frac{(F' * u)(x) + V'(x)}{\lambda_\epsilon(u)} \exp\left[-\frac{2}{\epsilon}\left(\int_0^x (F' * u)(y)dy + V(x)\right)\right].$$

Let us analyze the different elements of this derivative. We have already seen in the proof of Lemma 4.3 that for any $u \in \mathbb{C}_M$ the normalization factor $\lambda_\epsilon(u)$ satisfies (4.4) that is to say $\frac{1}{\lambda_\epsilon(u)} \leq C(\epsilon)\sqrt{M}$.

By (4.2), we obtain the bound: $0 \leq \int_0^x (F' * u)(y)dy \leq C_2Mx^2(1 + x^{2q})$.

Furthermore by (V-4) and (V-7), we get some estimation of V and its derivative:

$$V(x) \geq C_4x^4 - C_2x^2 \quad \text{and} \quad |V'(x)| \leq C_q(1 + |x|^{2q}) \quad \text{for all } x \in \mathbb{R}. \tag{4.12}$$

It remains to find some upper bound for the convolution term: $|(F' * u)(x)|$ with $x \in \mathbb{R}_+$. By (F-4) and since u is an even function,

$$\begin{aligned} |(F' * u)(x)| &= \left| \int_{\mathbb{R}} F'(x-z)u(z)dz \right| \leq \int_0^\infty |F'(x+z) + F'(x-z)|u(z)dz \\ &\leq C_q \int_0^\infty \left\{ |x+z|(1 + |x+z|^{2q-2}) + |x-z|(1 + |x-z|^{2q-2}) \right\} u(z)dz. \end{aligned}$$

Therefore:

$$\begin{aligned} |(F' * u)(x)| &\leq \int_{\mathbb{R}_+} C_q 2^{2q-1} \left\{ |x|^{2q-1} + |z||x|^{2q-2} + |x|(1 + |z|^{2q-2}) \right. \\ &\quad \left. + |z|(1 + |z|^{2q-2}) \right\} u(z)dz. \end{aligned}$$

By definition of \mathbb{C}_M , we have $u(z) \leq \frac{M}{1+|z|^p}$ for $p > 4q$. Hence the moments of order $1, 2q - 2$ and $2q - 1$ are bounded: there exist some constants C and C' , independent of the function $u \in \mathbb{C}_M$, such that

$$|(F' * u)(x)| \leq C \left(1 + |x| + |x|^{2q-2} + |x|^{2q-1}\right) \leq C' \left(1 + |x|^{2q+1}\right). \tag{4.13}$$

To sum up: using (4.4), (4.12) and (4.13) we obtain

$$\left|(\mathbb{A}^\epsilon u)'(x)\right| \leq \frac{2}{\epsilon} C(\epsilon) \sqrt{M} (1 + |x|^{2q+1}) \exp \left[-\frac{2}{\epsilon} \left(C_4 x^4 - C_2 x^2\right) \right]. \tag{4.14}$$

Finally we deduce that there exists some constant C_ϵ such that $|(\mathbb{A}^\epsilon u)'(x)| \leq C_\epsilon$ for all $x \in \mathbb{R}$.

Let us prove now that $\overline{\mathbb{A}^\epsilon \mathbb{C}_M}$ is compact. To this end, we take some sequence of functions $(u_n)_{n \in \mathbb{N}}$ in \mathbb{C}_M and focus our attention to the sequence $(\mathbb{A}^\epsilon u_n)_{n \in \mathbb{N}}$. We know, by Lemma 4.3 that $(1 + |x|^p) \mathbb{A}^\epsilon u_n(x) \leq M$ for all $x \in \mathbb{R}$. So there exist v and φ such that $(1 + x^p) \mathbb{A}^\epsilon u_{\varphi(n)}(x)$ converges to v , uniformly on compact sets. We extend the uniformity to the whole set of reals. Indeed, (4.14) implies $(1 + |x|^p) |(\mathbb{A}^\epsilon u_n)'(x)| \leq C_{M,\epsilon} e^{-\rho x^2}$ where $C_{M,\epsilon}$ and ρ are positive constants. For any $\delta > 0$, set $R > 0$ such that $C_{M,\epsilon} \int_R^{+\infty} e^{-\rho x^2} dx \leq \delta$. Hence, $|\mathbb{A}^\epsilon u_{\varphi(n)}(x) - \mathbb{A}^\epsilon u_{\varphi(n)}(R)| \leq \delta$ for any $x \geq R$. This proves the convergence of $(1 + x^p) \mathbb{A}^\epsilon u_{\varphi(n)}$ towards v uniformly on \mathbb{R} .

By Lemmas 4.3 and 4.4 we can apply Schauder’s theorem (Proposition 4.1) for the operator \mathbb{A}^ϵ on the function space \mathbb{C}_M with $M \geq M_0$. We deduce the existence of some fixed point which is, by construction, a symmetric stationary measure for the diffusion (1.1). \square

4.2. Example: $F(x) = \frac{\beta}{4}x^4 + \frac{\alpha}{2}x^2$

We have just shown the existence of a symmetric invariant measure for general self-stabilizing diffusions using fixed point arguments. Now let us study the uniqueness for symmetric invariant measure in suitable situations by using the previous result (Theorem 4.5) and a procedure close to that developed in Section 3.2. Let V be a potential satisfying (V-1)–(V-7).

Let u_ϵ be a symmetric invariant measure (Theorem 4.5). We denote by $m_2(\epsilon)$ its second moment. The couple $(m_2(\epsilon), u_\epsilon)$ is solution to some system like (3.1) and (3.2). Indeed

$$\begin{aligned} F * u_\epsilon(x) &= \int_{\mathbb{R}} F(x - z) u_\epsilon(z) dz \\ &= \frac{\alpha}{2}x^2 + \frac{\beta}{4}x^4 + \frac{3\beta m_2(\epsilon)}{2}x^2 + \left(\frac{\alpha}{2}m_2(\epsilon) + \frac{\beta}{4} \int_{\mathbb{R}} z^4 u_\epsilon(z) dz \right), \end{aligned}$$

with $\beta \geq 0$ since F' is a convex function on \mathbb{R}_+ .

The expression delimited by the brackets is just a constant so we obtain the following system of equations for $m_2(\epsilon)$ and u_ϵ : $m_2(\epsilon) = \int_{\mathbb{R}_+} x^2 v(m_2(\epsilon), x) dx$ and $2u_\epsilon(x) = v(m_2(\epsilon), x)$ where

$$v(m, x) = \frac{\exp \left[-\frac{2}{\epsilon} \left(V(x) + F(x) + \frac{3\beta m}{2} x^2 \right) \right]}{\int_0^\infty \exp \left[-\frac{2}{\epsilon} \left(V(z) + F(z) + \frac{3\beta m}{2} z^2 \right) \right] dz}.$$

Therefore we introduce the function $\chi_\epsilon(m) = \int_0^\infty x^2 v(m, x) dx - m$. By Theorem 4.5, we know that χ_ϵ admits at least one zero on \mathbb{R}_+ . Computing the derivative of χ_ϵ , we prove that

the considered function is decreasing:

$$x'_\epsilon(m) = -\frac{3\beta}{\epsilon} \left\{ \int_0^\infty x^4 v(m, x) dx - \left(\int_0^\infty x^2 v(m, x) dx \right)^2 \right\} - 1 < 0.$$

The conclusion is immediate: there is a unique symmetric invariant measure. Obviously this result and the kind of method used to prove it are particular to our simple example. If the degree of the interaction function is larger than 4 then it isn't enough to know the second moment in order to define the invariant measure: we need more moments and the proof of the uniqueness becomes awkward.

4.3. Outlying invariant measures

This section is essentially motivated by the uniqueness question for invariant measures. The existence of some symmetric measure was just proved in Section 4.1. It suffices now to point out asymmetric stationary measures for self-stabilizing diffusions. In the general setting, the interaction function is polynomial: set $F(x) = \sum_{k=1}^n \frac{F^{(2k)}(0)}{(2k)!} x^{2k}$.

Let u be the density of some probability measure with respect to the Lebesgue measure and μ_1, \dots, μ_{2n-1} denote its moments of orders 1 to $2n - 1$ respectively. We assume they are finite. Then the difference $D(x) := F * u(x) - F * u(0)$ satisfies

$$\begin{aligned} D(x) &= F(x - a) - F(-a) + \sum_{p=1}^{2n-1} \frac{(-1)^p}{p!} (\mu_p - a^p) \sum_{j \geq \frac{1+p}{2}}^n \frac{F^{(2j)}(0)}{(2j - p)!} x^{2j-p} \\ &= F(x - a) - F(a) + \sum_{p=1}^{2n-1} \frac{(-1)^p}{p!} (\mu_p - a^p) \left(F^{(p)}(x) - F^{(p)}(0) \right). \end{aligned}$$

Hence $D(x) = Z_\mu(x) - Z_\mu(0)$ where

$$Z_m(x) = F(x - a) + \sum_{p=1}^{2n-1} \frac{(-1)^p}{p!} (m_p - a^p) F^{(p)}(x). \tag{4.15}$$

Since the convolution product can be expressed as a polynomial function which coefficients just depend on the moments of u , then the exponential expression of invariant measure (2.2) can be specified. Indeed Eq. (2.2) can be transformed into some system of equations whose unknown factors are the moments of the measure. In order to introduce this system, let us define, for all $k \in [1; 2n - 1]$, the function

$$\begin{aligned} \varphi_k^{(\epsilon)}(m_1, \dots, m_{2n-1}) &= \frac{\int_{\mathbb{R}} x^k \exp \left[-\frac{2}{\epsilon} (V(x) + Z_m(x) - Z_m(0)) \right] dx}{\int_{\mathbb{R}} \exp \left[-\frac{2}{\epsilon} (V(x) + Z_m(x) - Z_m(0)) \right] dx} \\ &= \frac{\int_{\mathbb{R}} x^k \exp \left[-\frac{2}{\epsilon} W_m(x) \right] dx}{\int_{\mathbb{R}} \exp \left[-\frac{2}{\epsilon} W_m(x) \right] dx} \end{aligned} \tag{4.16}$$

with the potential $W_m(x) = V(x) + Z_m(x)$. We construct the mapping:

$$\Phi^{(\epsilon)} = (\varphi_1^{(\epsilon)}, \dots, \varphi_k^{(\epsilon)}, \dots, \varphi_{2n-1}^{(\epsilon)}). \tag{4.17}$$

The measure associated to the density function u is invariant if and only if its moments vector $(\mu_1, \dots, \mu_{2n-1})$ is a fixed point of the map $\Phi^{(\epsilon)}$.

We are going to show the existence of an asymmetric invariant measure defined by $2n - 1$ parameters close to a, \dots, a^{2n-1} respectively, in other words the outlying measure is close to the Dirac mass in the point a . More precisely, we shall prove that there exists a parallelepiped stable by $\Phi^{(\epsilon)}$, which converges to the point $(a, a^2, \dots, a^{2n-1})$ as ϵ tends to 0. As in the linear case, we shall proceed by applying the mean value theorem in order to obtain asymptotic developments in the small noise limit.

Theorem 4.6. *Let $(\eta_\epsilon)_\epsilon$ some sequence satisfying $\lim_{\epsilon \rightarrow 0} \eta_\epsilon = 0$ and $\lim_{\epsilon \rightarrow 0} \epsilon/\eta_\epsilon = 0$. Under the condition*

$$\sum_{p=0}^{2n-2} \frac{|F^{(p+2)}(a)|}{p!} a^p < \alpha + V''(a), \tag{4.18}$$

for any $\rho > 0$, there are at least two outlying measures u_ϵ^+ and u_ϵ^- satisfying, for ϵ small enough

$$\left| \int_{\mathbb{R}} x^k u_\epsilon^\pm(x) dx - (\pm a)^k \right| \leq \rho \eta_\epsilon. \tag{4.19}$$

Proof. Let $\lambda > 0$. Let us define the parallelepiped

$$C(\epsilon) = \prod_{p=1}^{2n-1} [a^p - pa^{p-1}\lambda\eta_\epsilon, a^p + pa^{p-1}\lambda\eta_\epsilon].$$

Let m be an element of $C(\epsilon)$ then there exist some coordinates $(r_p)_{1 \leq p \leq 2n-1}$ which determine m through the equations $m_p = a^p + r_p \eta_\epsilon$. By (4.15) and (4.16), we get

$$\varphi_k^{(\epsilon)}(m) = \frac{\int_{\mathbb{R}} x^k e^{-\frac{2}{\epsilon}(V(x)+F(x-a))} \exp \left[-\frac{2\eta_\epsilon}{\epsilon} \sum_{p=1}^{2n-1} \frac{(-1)^p r_p}{p!} F^{(p)}(x) \right] dx}{\int_{\mathbb{R}} e^{-\frac{2}{\epsilon}(V(x)+F(x-a))} \exp \left[-\frac{2\eta_\epsilon}{\epsilon} \sum_{p=1}^{2n-1} \frac{(-1)^p r_p}{p!} F^{(p)}(x) \right] dx}.$$

We apply Lemma A.4 and Remark A.5 to the functions $U(x) = V(x) + F(x - a)$, $f(x) = x^k$, $\mu_p = r_p$ and $G_p(x) = \frac{(-1)^p}{p!} F^{(p)}(x)$. We obtain:

$$\varphi_k^{(\epsilon)}(m) = a^k - \eta_\epsilon \frac{ka^{k-1}}{\alpha + V''(a)} \sum_{p=1}^{2n-1} \frac{(-1)^p r_p}{p!} F^{(p+1)}(a) + o(\eta_\epsilon),$$

uniformly with respect to the coordinates $(r_p)_p$. By definition of the parallelepiped $C(\epsilon)$ the coordinates satisfy $|r_p| \leq pa^{p-1}\lambda$. Therefore, under condition (4.18),

$$\begin{aligned} \left| \varphi_k^{(\epsilon)}(m) - a^k \right| &\leq \eta_\epsilon \lambda \frac{ka^{k-1}}{\alpha + V''(a)} \sum_{p=1}^{2n-1} \frac{|F^{(p+1)}(a)|}{p!} pa^{p-1} + o(\eta_\epsilon) \\ &< \eta_\epsilon ka^{k-1}\lambda + o(\eta_\epsilon). \end{aligned}$$

Since this estimate is uniform with respect to the coordinates, as soon as ϵ is small enough, we have $|\varphi_k^{(\epsilon)}(m) - a^k| < ka^{k-1}\lambda\eta_\epsilon$, that means that $\Phi^{(\epsilon)}(m) \in C(\epsilon)$.

Let us note that $C(\epsilon)$ is a convex, closed and bounded subset of \mathbb{R}^{2n-1} . Since the space dimension is finite, the continuity of $\Phi^{(\epsilon)}$ implies that the closure of the parallelepiped's image is a compact set.

We can apply Schauder's Theorem (Proposition 4.1) and obtain that there exists some fixed point in the compact. In other words there exists $m \in C(\epsilon)$ such that the measure associated to the density

$$u_{\epsilon,m}(x) = \frac{\exp\left[-\frac{2}{\epsilon}W_m(x)\right]}{\int_{\mathbb{R}} \exp\left[-\frac{2}{\epsilon}W_m(z)\right] dz} \tag{4.20}$$

is invariant. In a similar way, the measure defined by m^- is also invariant; here $m^-(k) = (-1)^k m_k$. To conclude: we have at least two outlying measures, one around a and the second one around $-a$. \square

We can not prove at this stage the uniqueness of the couple of outlying invariant measures (this question shall be explored in a subsequent work). We can effectively imagine that other outlying measures could exist around a , around $-a$ or even around other areas. Nevertheless we can develop a sharper description of one particular outlying measure: the measure close to δ_a where δ represents the Dirac measure. To do this it suffices to estimate its different moments, that requires the following preliminary result.

Lemma 4.7. *There exists a unique solution $(\tau_1^0, \dots, \tau_{2n-1}^0)$ to the following Cramer's system*

$$\sum_{p=1}^{2n-1} \frac{(-1)^p}{p!} F^{(p+1)}(a)\tau_p + \frac{\alpha + V''(a)}{ka^{k-1}}\tau_k = \frac{V^{(3)}(a)}{4(\alpha + V''(a))} - \frac{k-1}{4a}, \tag{4.21}$$

for $1 \leq k \leq 2n-1$. This solution is given by

$$\tau_k^0 = ka^{k-1} \frac{aV^{(3)}(a) - (k-1)V''(a)}{4aV''(a)(\alpha + V''(a))}, \quad 1 \leq k \leq 2n-1. \tag{4.22}$$

Proof. Let us denote by I_{2n-1} the unit matrix of dimension $2n-1$ and for $A \in \mathbb{R}^{2n-1}$, A^T represents the transpose of the vector A . Moreover we adopt the following notation $(x_k)_{1 \leq k \leq 2n-1} = (x_1, \dots, x_{2n-1})$. The system (4.21) can be written in this way: we define $\mathcal{T} = (\tau_k)_{1 \leq k \leq 2n-1}^T$ then

$$\left[(\alpha + V''(a))I_{2n-1} + C_1C_2^T \right] \mathcal{T} = \left(ka^{k-1} \left(\frac{V^{(3)}(a)}{4\alpha + 4V''(a)} - \frac{k-1}{4a} \right) \right)_{1 \leq k \leq 2n-1}^T$$

with the vectors $C_1^T = (ka^{k-1})_{1 \leq k \leq 2n-1}$ and $C_2^T = \left(\frac{(-1)^k}{k!} F^{(k+1)}(a) \right)_{1 \leq k \leq 2n-1}$. We define therefore

$$A = (\alpha + V''(a))I_{2n-1} + C_1C_2^T. \tag{4.23}$$

Let us note that $C_1C_2^TC_1C_2^T = (C_2^TC_1)C_1C_2^T$ and

$$C_2^TC_1 = \sum_{p=1}^{2n-1} \frac{(-1)^p}{p!} F^{(p+1)}(a)pa^{p-1} = - \sum_{p=0}^{2n-2} \frac{(-1)^p}{p!} F^{(p+2)}(a)a^p = -F''(0).$$

Since $F''(0) = \alpha$, we obtain

$$\begin{aligned} A^2 &= (\alpha + V''(a))^2 I_{2n-1} + \left(2(\alpha + V''(a)) + C_2^T C_1\right) C_1 C_2^T \\ &= (\alpha + V''(a))^2 I_{2n-1} + (2(\alpha + V''(a)) - F''(0)) C_1 C_2^T \\ &= (\alpha + V''(a))^2 I_{2n-1} + (\alpha + 2V''(a)) C_1 C_2^T \\ &= (\alpha + 2V''(a))A - V''(a) (\alpha + V''(a)) I_{2n-1}. \end{aligned}$$

We deduce that A is invertible, that is (4.21) is a Cramer’s system, and using (4.23) we get explicitly the inverse:

$$\begin{aligned} A^{-1} &= \frac{1}{V''(a)(\alpha + V''(a))} ((\alpha + 2V''(a))I_{2n-1} - A) \\ &= \frac{1}{V''(a)(\alpha + V''(a))} \left(V''(a)I_{2n-1} - C_1 C_2^T \right). \end{aligned}$$

Therefore the Cramer’s system (4.21) admits a unique solution given by

$$\begin{aligned} \tau_k^0 &= \frac{1}{V''(a)(\alpha + V''(a))} \left\{ V''(a)ka^{k-1} \frac{aV^{(3)}(a) - (k-1)(\alpha + V''(a))}{4a(\alpha + V''(a))} \right. \\ &\quad \left. - ka^{k-1} \sum_{p=1}^{2n-1} \frac{(-1)^p}{p!} F^{(p+1)}(a) pa^{p-1} \frac{aV^{(3)}(a) - (p-1)(\alpha + V''(a))}{4a(\alpha + V''(a))} \right\} \\ &= \frac{ka^{k-1}}{4aV''(a)(\alpha + V''(a))^2} \left\{ aV^{(3)}(a) \left[V''(a) - \sum_{p=1}^{2n-1} \frac{(-1)^p a^{p-1}}{(p-1)!} F^{(p+1)}(a) \right] \right. \\ &\quad \left. - (\alpha + V''(a)) \left[(k-1)V''(a) - \sum_{p=2}^{2n-1} \frac{(-1)^p}{(p-2)!} F^{(p+1)}(a) a^{p-1} \right] \right\} \\ &= ka^{k-1} \frac{aV^{(3)}(a) - (k-1)V''(a)}{4aV''(a)(\alpha + V''(a))}. \end{aligned}$$

Indeed, we use the two equalities $\sum_{p=1}^{2n-1} \frac{(-1)^p}{(p-1)!} F^{(p+1)}(a) a^{p-1} = -F''(0) = -\alpha$ and $\sum_{p=2}^{2n-1} \frac{(-1)^p}{(p-2)!} F^{(p+1)}(a) a^{p-1} = aF^{(3)}(0) = 0$. \square

Theorem 4.6 points out the existence of two outlying measures, one concentrated around a and an other around $-a$. According to **Lemma 4.7** we get some parallelepiped with sharper edge which contains δ_a and some asymmetric invariant measure.

Theorem 4.8. Under the condition (4.18), for any $\delta > 0$, there exists ϵ_0 such that $\Phi^{(\epsilon)}$ admits two fixed points m^\pm with

$$\left| m_k^\pm(\epsilon) - \left((\pm 1)^k a^k - (\pm 1)^k \tau_k^0 \epsilon \right) \right| \leq \delta \epsilon, \quad 1 \leq k \leq 2n - 1, \quad \epsilon \leq \epsilon_0. \tag{4.24}$$

Proof. It is similar to the proof of **Theorem 4.6**.

Let $\rho > 0$ and $C(\epsilon) = \prod_{p=1}^{2n-1} [a^p - (\tau_p^0 + pa^{p-1}\rho)\epsilon, a^p - (\tau_p^0 - pa^{p-1}\rho)\epsilon]$. We choose an element m in the parallelepiped $C(\epsilon)$. For all $1 \leq p \leq 2n - 1$, there exists a coordinate

$\rho_p \in [-\rho; \rho]$ such that $m_p = a^p - (\tau_p^0 + pa^{p-1}\rho_p)\epsilon$. By (4.15) and (4.16), we obtain

$$\varphi_k^{(\epsilon)}(m) = \frac{\int_{\mathbb{R}} x^k \exp \left[2 \sum_{p=1}^{2n-1} \frac{(-1)^p}{p!} (\tau_p^0 + pa^{p-1}\rho_p) F^{(p)}(x) \right] e^{-\frac{2}{\epsilon}(V(x)+F(x-a))} dx}{\int_{\mathbb{R}} \exp \left[2 \sum_{p=1}^{2n-1} \frac{(-1)^p}{p!} (\tau_p^0 + pa^{p-1}\rho_p) F^{(p)}(x) \right] e^{-\frac{2}{\epsilon}(V(x)+F(x-a))} dx}.$$

We apply Lemma A.3 and Remark A.5 with the following functions: $U(x) = V(x) + F(x - a)$, $\mu_p = \tau_p^0 + pa^{p-1}\rho_p$, $G = 0$ and $f_p(x) = 2\frac{(-1)^p}{p!}F^{(p)}(x)$. Hence

$$\begin{aligned} \varphi_k^{(\epsilon)}(m) &= a^k - \frac{ka^{k-2}}{4(\alpha + V''(a))^2} \left[aV^{(3)}(a) - (\alpha + V''(a)) \right. \\ &\quad \left. \times \left((k-1) + 4a \sum_{p=1}^{2n-1} \frac{(-1)^p}{p!} (\tau_p^0 + pa^{p-1}\rho_p) F^{(p+1)}(a) \right) \right] \epsilon + o(\epsilon) \\ &= a^k - \frac{1}{\alpha + V''(a)} \left[\frac{ka^{k-1}V^{(3)}(a)}{4(\alpha + V''(a))} - ka^{k-1} \sum_{p=1}^{2n-1} \frac{(-1)^p \tau_p^0}{p!} F^{(p+1)}(a) \right. \\ &\quad \left. - \frac{k(k-1)a^{k-2}}{4} - ka^{k-1} \sum_{p=1}^{2n-1} \frac{(-1)^p \rho_p a^{p-1}}{(p-1)!} F^{(p+1)}(a) \right] \epsilon + o(\epsilon). \end{aligned}$$

This estimate is uniform with respect to the variables $(\rho_p)_p$.

We denote by d_k^ϵ the difference $|\varphi_k^{(\epsilon)}(m) - a^k + \tau_k^0 \epsilon|$. We compute this expression:

$$\begin{aligned} d_k^\epsilon &\leq \left| \frac{ka^{k-1}V^{(3)}(a)}{4(\alpha + V''(a))^2} - \frac{ka^{k-1}}{\alpha + V''(a)} \sum_{p=1}^{2n-1} \frac{(-1)^p}{p!} \tau_p^0 F^{(p+1)}(a) - \frac{k(k-1)a^{k-2}}{4(\alpha + V''(a))} \right. \\ &\quad \left. - \tau_k^0 - \frac{ka^{k-1}}{\alpha + V''(a)} \sum_{p=1}^{2n-1} \frac{(-1)^p}{(p-1)!} \rho_p F^{(p+1)}(a) a^{p-1} \right| \epsilon + o(\epsilon). \end{aligned}$$

According to the Lemma 4.7 and using the condition (4.18), we obtain, for ϵ small enough,

$$\begin{aligned} \left| \varphi_k^{(\epsilon)}(m) - a^k + \tau_k^0 \epsilon \right| &\leq \frac{ka^{k-1}}{\alpha + V''(a)} \sum_{p=1}^{2n-1} \frac{a^{p-1}}{(p-1)!} |\rho_p| |F^{(p+1)}(a)| \epsilon + o(\epsilon) \\ &\leq \rho \frac{ka^{k-1}}{\alpha + V''(a)} \sum_{p=0}^{2n-2} \frac{1}{p!} |F^{(p+2)}(a)| a^p \epsilon + o(\epsilon) < ka^{k-1} \rho \epsilon. \end{aligned}$$

In other words, $\Phi^{(\epsilon)}(m) \in C(\epsilon)$ in the small noise limit. The application of Schauder’s theorem (Proposition 4.1) permits to prove the existence of some fixed point in the compact after choosing $\rho < \frac{\delta}{\max_{p \in [1; 2n-1]} pa^{p-1}}$. Therefore there exists $m \in C(\epsilon)$ such that the associated measure $u_{\epsilon, m}(x)$ defined by (4.20) is invariant. In the same way, the measure defined by m^- is invariant with $m^-(k) = (-1)^k m_k$. Finally the continuous map $\Phi^{(\epsilon)}$ admits two fixed points $m^\pm(\epsilon)$ satisfying (4.24). \square

Remark 4.9. 1. In the particular case: $F^{(p)}(a) \geq 0$ for all $p \in \mathbb{N}$, the condition for the existence of outlying measures i.e. (4.18) becomes $V''(a) > F_0''(2a)$ where F_0 is defined by $F(x) = \frac{\alpha}{2}x^2 + F_0(x)$.

2. In the linear interaction case: $F(x) = \frac{\alpha}{2}x^2$, (4.18) is equivalent to the simple condition $V''(a) > 0$ which is in fact always satisfied according to (V-3). In other words we obtain the existence result presented in the linear interaction case.

Appendix. Annex

We shall present here some useful asymptotic results which are close to the classical Laplace’s method. The first lemma is quite classical (see, for instance [1], Corollary 6.4.2., p. 108).

Lemma A.1. *Let $M > 0$. Let us assume that U is $\mathcal{C}^2([M, \infty[)$ -continuous, $U'(x) \neq 0$ and $U''(x) > 0$ for all $x \in [M, \infty[$ and $\lim_{x \rightarrow \infty} \frac{U''(x)}{(U'(x))^2} = 0$. If $x \rightarrow e^{-U(x)}$ is integrable on \mathbb{R} then for any $x \geq M$:*

$$\int_x^{+\infty} e^{-U(t)} dt \approx \frac{e^{-U(x)}}{U'(x)} \quad \text{and} \quad \int_M^x e^{U(t)} dt \approx \frac{e^{U(x)}}{U'(x)} \quad \text{as } x \rightarrow \infty. \tag{A.1}$$

Lemma A.2. *Set $\epsilon > 0$. Let U and G two $\mathcal{C}^\infty(\mathbb{R})$ -continuous functions. We define $U_\mu = U + \mu G$ for μ belonging to some compact interval \mathcal{I} of \mathbb{R} . Let us introduce some interval $[a, b]$ satisfying: $U'_\mu(a) \neq 0$, $U'_\mu(b) \neq 0$ and $U_\mu(x)$ admits some unique global minimum on the interval $[a, b]$ reached at $x_\mu \in]a, b[$ for all $\mu \in \mathcal{I}$. We assume that there exists some exponent k_0 independent of $\mu \in \mathcal{I}$ such that $2k_0 = \min_{r \in \mathbb{N}^*} \{U_\mu^{(r)}(x_\mu) \neq 0\}$. Then taking the limit $\epsilon \rightarrow 0$ we get*

$$I_0 := \int_a^b e^{-\frac{U_\mu(t)}{\epsilon}} dt = \frac{1}{k_0} \left(\frac{\epsilon(2k_0)!}{U_\mu^{2k_0}(x_\mu)} \right)^{\frac{1}{2k_0}} \Gamma\left(\frac{1}{2k_0}\right) e^{-\frac{U_\mu(x_\mu)}{\epsilon}} (1 + o_{\mathcal{I}}(1)), \tag{A.2}$$

where Γ represents the Euler function and $o_{\mathcal{I}}(1)$ converges towards 0 uniformly with respect to $\mu \in \mathcal{I}$.

Proof. Since U_μ is regular and admits some unique global minimum for $x = x_\mu$, there exists $0 < \tau_0 < \min_{\mu \in \mathcal{I}} \min \{x_\mu - a; b - x_\mu\}$ such that the minimum on the interval $[a; x_\mu - \tau] \cup [x_\mu + \tau; b]$ denoted by $\underline{U}_\mu(\tau)$ is reached on the boundary $\{x_\mu - \tau; x_\mu + \tau\}$ for all $\tau < \tau_0$. Consequently

$$\int_a^{x_\mu - \tau} \exp\left[-\frac{U_\mu(t)}{\epsilon}\right] dt + \int_{x_\mu + \tau}^b \exp\left[-\frac{U_\mu(t)}{\epsilon}\right] dt \leq (b - a) \exp\left[-\frac{\underline{U}_\mu(\tau)}{\epsilon}\right].$$

Defining $I_\tau = \int_{x_\mu - \tau}^{x_\mu + \tau} \exp\left[-\frac{U_\mu(t)}{\epsilon}\right] dt$, we obtain the following bound:

$$|I_0 - I_\tau| \leq (b - a) \exp\left[-\frac{\underline{U}_\mu(\tau)}{\epsilon}\right]. \tag{A.3}$$

Let us first estimate I_τ . We define $\eta_\mu = \frac{U_\mu^{(2k_0)}(x_\mu)}{(2k_0)!}$. Let us note that η_μ depends continuously on μ . By the mean value theorem, there exists some constant $C > 0$ independent of $\mu \in \mathcal{I}$ such that,

in a neighborhood of x_μ , the following bound is satisfied: $|U_\mu(t) - U_\mu(x_\mu) - \eta_\mu(t - x_\mu)^{2k_0}| \leq C|t - x_\mu|^{2k_0+1}$. Hence

$$J_\tau \exp\left[-\frac{C\tau^{2k_0+1}}{\epsilon}\right] \leq \frac{I_\tau}{2} \exp\frac{U_\mu(x_\mu)}{\epsilon} \leq J_\tau \exp\left[\frac{C\tau^{2k_0+1}}{\epsilon}\right], \tag{A.4}$$

where

$$J_\tau = \int_0^\tau \exp\left[-\frac{1}{\epsilon}\eta_\mu t^{2k_0}\right] dt = \left(\frac{\epsilon}{\eta_\mu}\right)^{\frac{1}{2k_0}} \frac{1}{2k_0} \int_0^{\tau^{2k_0} \frac{\eta_\mu}{\epsilon}} t^{\frac{1}{2k_0}-1} e^{-t} dt,$$

by the change of variable $t := \left(\frac{\epsilon}{\eta_\mu}\right)^{\frac{1}{2k_0}} (t')^{\frac{1}{2k_0}}$. A simple integration leads to

$$-\tau^{1-2k_0} \left(\frac{\eta_\mu}{\epsilon}\right)^{\frac{1}{2k_0}-1} e^{-\tau^{2k_0} \frac{\eta_\mu}{\epsilon}} \leq \int_0^{\tau^{2k_0} \frac{\eta_\mu}{\epsilon}} t^{\frac{1}{2k_0}-1} e^{-t} dt - \Gamma\left(\frac{1}{2k_0}\right) \leq 0. \tag{A.5}$$

In order to conclude we set $\tau = \exp\left[\frac{\log(\epsilon)}{2k_0+\frac{1}{2}}\right]$. Then we get: for $C \in \mathbb{R}, l > 0$,

$$\lim_{\epsilon \rightarrow 0} e^{C\tau^{2k_0+1}} = 1, \quad \lim_{\epsilon \rightarrow 0} e^{-\eta_\mu \frac{\tau^{2k_0}}{\epsilon}} \frac{\tau^{1-2k_0}}{\epsilon^{\frac{1}{2k_0}-1}} = \lim_{\epsilon \rightarrow 0} \epsilon^{-l} e^{\frac{U_\mu(x_\mu) - U_\mu(\tau)}{\epsilon}} = 0.$$

These convergences are uniform with respect to μ . Applying these asymptotic results to (A.3)–(A.5) permits to prove the statement of the lemma. \square

Lemma A.3. *Let U and G be two $C^\infty(\mathbb{R})$ -continuous functions. We define $U_\mu = U + \mu G$ for the parameter μ belonging to some compact interval \mathcal{I} of \mathbb{R} . We assume that $U_\mu(t) \geq t^2$ for $|t|$ larger than some R independent of μ and that U_μ admits a unique global minimum at x_μ with $U''_\mu(x_\mu) > 0$. Let f_m be a C^3 -continuous function depending on some parameter m which belongs to a compact set \mathcal{M} . Furthermore we assume that there exists some constant $\lambda > 0$ such that $|f_m(t)| \leq \exp[\lambda|U_\mu(t)|]$ for all $t \geq R, \mu \in \mathcal{I}, m \in \mathcal{M}$ and $|f_m^{(i)}|$ is locally bounded uniformly with respect to the parameter $m \in \mathcal{M}$ for $0 \leq i \leq 3$. Let $a, b \in \bar{\mathbb{R}}$ such that $a < x_\mu < b$. Then the following asymptotic result holds as ϵ tends to 0:*

$$\int_a^b f_m(t) e^{\frac{-2U_\mu(t)}{\epsilon}} dt = \sqrt{\frac{\pi\epsilon}{\mathcal{U}_2}} e^{-\frac{2U_\mu(x_\mu)}{\epsilon}} \left\{ f_m(x_\mu) + \gamma_0(\mu)\epsilon + o_{\mathcal{I},\mathcal{M}}^{(1)}(\epsilon) \right\}, \tag{A.6}$$

with

$$\gamma_0(\mu) = f_m(x_\mu) \left(\frac{5\mathcal{U}_3^2}{48\mathcal{U}_2^3} - \frac{\mathcal{U}_4}{16\mathcal{U}_2^2} \right) - f'_m(x_\mu) \frac{\mathcal{U}_3}{4\mathcal{U}_2^2} + \frac{f''_m(x_\mu)}{4\mathcal{U}_2}. \tag{A.7}$$

Here $\mathcal{U}_k = U_\mu^{(k)}(x_\mu)$ and $o_{\mathcal{I},\mathcal{M}}^{(1,2)}(\epsilon)/\epsilon$ converges to 0 as ϵ becomes small uniformly with respect to the parameters m and μ . Moreover, for any $n \geq 1$, we have

$$\frac{\int_{\mathbb{R}} t^n e^{f_m(t)} e^{\frac{-2U_\mu(t)}{\epsilon}} dt}{\int_{\mathbb{R}} e^{f_m(t)} e^{\frac{-2U_\mu(t)}{\epsilon}} dt} - x_\mu^n \approx -\frac{nx_\mu^{n-2}}{4\mathcal{U}_2} \left[x_\mu \frac{\mathcal{U}_3}{\mathcal{U}_2} - n + 1 - 2x_\mu f'_m(x_\mu) \right], \tag{A.8}$$

where the estimate is uniform with respect to the parameters m and μ as $\epsilon \rightarrow 0$.

Proof. First, we consider $[a, b]$ compact and split the integral in (A.6) into two parts:

$$I = \int_{x_\mu - \rho}^{x_\mu + \rho} f_m(t) e^{-\frac{2U_\mu(t)}{\epsilon}} dt + \int_{[x_\mu - \rho; x_\mu + \rho]^c \cap [a; b]} f_m(t) e^{-\frac{2U_\mu(t)}{\epsilon}} dt = I_1 + I_2$$

with some arbitrary $\rho > 0$ which should be specified in the following. Since the hypotheses, there exists $\eta > 0$ such that $|f_m^{(i)}(x)| \leq \eta$ for all $m \in \mathcal{M}$, $x \in [a, b]$ and $0 \leq i \leq 3$. And, x_μ is the unique global minimum on $[a; b]$.

Step 1. We shall prove that the second integral is negligible as $\frac{\rho^2}{\epsilon} \rightarrow \infty$ that means that $I_2 = o_{\mathcal{I}\mathcal{M}}\{\epsilon^{3/2} e^{-\frac{2U(x_\mu)}{\epsilon}}\}$. We get

$$I_2 \leq (b - a) \sup_{z \in [a; b]} |f_m(z)| \exp \left[-2 \frac{\inf_{z \in [x_\mu - \rho; x_\mu + \rho]^c} U_\mu(z)}{\epsilon} \right]. \tag{A.9}$$

Since the global minimum of U_μ is unique and due to the regularity of U_μ with respect to the parameter μ , we deduce that the minimum of the function on the interval $[x_\mu - \rho; x_\mu + \rho]^c \cap [a; b]$ is reached on the boundary provided that ρ is small enough. The development $U_\mu(x_\mu \pm \rho) = U_\mu(x_\mu) + \frac{1}{2} U''_\mu(x_\mu) \rho^2 + o_{\mathcal{I}}(\rho^2)$ implies, as already claimed that $I_2 = o_{\mathcal{I}\mathcal{M}}\left\{\epsilon^{3/2} e^{-\frac{2U(x_\mu)}{\epsilon}}\right\}$ as $\rho^2/\epsilon \rightarrow \infty$.

Step 2. Let us focus our attention to the integral on $[x_\mu - \rho; x_\mu + \rho]$. The function f_m can be developed in the neighborhood of x_μ :

$$f_m(x) = f_m(x_\mu) + f'_m(x_\mu)(x - x_\mu) + \frac{1}{2} f''_m(x_\mu)(x - x_\mu)^2 + \frac{1}{6} f_m^{(3)}(w_{m,\mu}(x))(x - x_\mu)^3$$

with the value $w_{m,\mu}(x)$ between x_μ and x . Taking into account these different terms, the integral I_1 can be split into 4 different integrals respectively $\tilde{I}_0, \dots, \tilde{I}_3$. For each integral we shall analyze the asymptotic behavior.

Step 2.1. Asymptotic behavior of \tilde{I}_3 . By definition $w_{m,\mu}(t) \in [x_\mu - \rho; x_\mu + \rho]$ when $t \in [x_\mu - \rho; x_\mu + \rho]$. Moreover, by assumption $|f_m^{(3)}(w_{m,\mu}(t))|$ is upper bounded by some constant $\eta > 0$ independent of m and μ . By Lemma A.2 applied to $2U_\mu$, for $\rho < 1$ and ϵ small, we obtain the existence of some constant $C > 0$, independent of the parameters m and μ , such that

$$|\tilde{I}_3| \leq \frac{\eta}{6} \rho^3 \int_{(x_\mu - 1) \vee a}^{(x_\mu + 1) \wedge b} e^{-\frac{2U_\mu(t)}{\epsilon}} dt \leq C \sqrt{\pi} \rho^3 \sqrt{\frac{\epsilon}{U''_\mu(x_\mu)}} e^{-\frac{2U_\mu(x_\mu)}{\epsilon}}.$$

Hence, if $\rho^3 = o(\epsilon)$ then the following asymptotic result holds

$$\tilde{I}_3 = o_{\mathcal{I}\mathcal{M}}\left\{\epsilon^{\frac{3}{2}} e^{-\frac{2U(x_0)}{\epsilon}}\right\}. \tag{A.10}$$

Step 2.2. Asymptotic behavior of \tilde{I}_2 . Using the \mathcal{C}^3 -regularity of U_μ that is $U_\mu(t) = U_\mu(x_\mu) + \frac{1}{2} U''_\mu(x_\mu)(t - x_\mu)^2 + \frac{1}{6} U_\mu^{(3)}(y_\mu(t))(t - x_\mu)^3$ with $y_\mu(t)$ belonging to $[x_\mu - \rho; x_\mu + \rho]$, we get

$$\tilde{I}_2 = \frac{f''_m(x_\mu)}{2} e^{-\frac{2U_\mu(x_\mu)}{\epsilon}} \int_{x_\mu - \rho}^{x_\mu + \rho} (t - x_\mu)^2 e^{-\frac{U''_\mu(x_\mu)}{\epsilon}(t - x_\mu)^2 - \frac{U_\mu^{(3)}(y_\mu(t))}{3\epsilon}(t - x_\mu)^3} dt.$$

Since $y_\mu(t)$ belongs to some compact set, the third derivative $U_\mu^{(3)}(y_\mu(t))$ is bounded by some constant independent of μ . Applying the following change of variable $u = (t - x_\mu)^2 U_\mu''(x_\mu)/\epsilon$ yields

$$J_2 e^{-C \frac{\rho^3}{\epsilon}} \left(\frac{\epsilon}{U_\mu''(x_\mu)} \right)^{\frac{3}{2}} \leq \frac{2\tilde{I}_2 e^{\frac{2U_\mu(x_\mu)}{\epsilon}}}{f_m''(x_\mu)} \leq J_2 e^{C \frac{\rho^3}{\epsilon}} \left(\frac{\epsilon}{U_\mu''(x_\mu)} \right)^{\frac{3}{2}},$$

with $J_2 = \int_0^{U_\mu''(x_\mu) \frac{\rho^2}{\epsilon}} \sqrt{u} e^{-u} du$. If $\frac{\rho^3}{\epsilon} \rightarrow 0$ and $\frac{\rho^2}{\epsilon} \rightarrow \infty$ then

$$\tilde{I}_2 = \sqrt{\pi} \frac{f_m''(x_\mu)}{4} e^{-\frac{2U_\mu(x_\mu)}{\epsilon}} \left(\frac{\epsilon}{U_\mu''(x_\mu)} \right)^{\frac{3}{2}} (1 + o_{\mathcal{I}}(1)). \tag{A.11}$$

Step 2.3. Asymptotic behavior of \tilde{I}_1 . We expand the function U_μ in the neighborhood of x_μ : $U_\mu(t + x_\mu) = U_\mu(x_0) + \frac{1}{2}U_\mu''(x_\mu)t^2 + \frac{1}{6}U_\mu^{(3)}(x_\mu)t^3 + \frac{1}{24}U_\mu^{(4)}(y_\mu(t))t^4$ where $y_\mu(t) \in [x_\mu - \rho, x_\mu + \rho]$. The regularity of $U_\mu(x)$ with respect to both x and μ implies the existence of some constant $C > 0$ independent of μ which bounds the fourth derivative of U_μ on the integral support. Therefore we have

$$f'_m(x_\mu) e^{-C \frac{\rho^4}{\epsilon}} J_\rho \leq e^{\frac{2U_\mu(x_\mu)}{\epsilon}} \tilde{I}_1 \leq f'_m(x_\mu) e^{C \frac{\rho^4}{\epsilon}} J_\rho,$$

with $J_\rho = \int_{-\rho}^\rho z e^{-\frac{\mathcal{U}_2}{\epsilon} z^2 - \frac{\mathcal{U}_3}{3\epsilon} z^3} dz$ and $\mathcal{U}_k = U_\mu^{(k)}(x_\mu)$. Since $|e^{-x} - 1 + x - \frac{x^2}{2}| \leq |x|^3 e^{|x|}$, we deduce that, for any $z \in [-\rho; \rho]$:

$$\left| e^{-\frac{\mathcal{U}_3}{3\epsilon} z^3} - 1 + \frac{\mathcal{U}_3 z^3}{3\epsilon} - \frac{\mathcal{U}_3^2 z^6}{18\epsilon^2} \right| \leq \left| \frac{\mathcal{U}_3}{3} \right|^3 \frac{\rho^9}{\epsilon^3} e^{\frac{|\mathcal{U}_3| \rho^3}{3\epsilon}}.$$

We define $m_\rho(l) = \int_{-\rho}^\rho z^l e^{-\frac{\mathcal{U}_2}{\epsilon} z^2} dz$ and $n_\rho(l) = \int_0^\rho |z|^l e^{-\frac{\mathcal{U}_2}{\epsilon} z^2} dz$. Some estimation of the integral J_ρ points out directly:

$$\left| J_\rho - m_\rho(1) + \frac{\mathcal{U}_3}{3\epsilon} m_\rho(4) - \frac{\mathcal{U}_3^2}{18\epsilon^2} m_\rho(7) \right| \leq 2 \left| \frac{\mathcal{U}_3}{3} \right|^3 \frac{\rho^9}{\epsilon^3} e^{\frac{|\mathcal{U}_3| \rho^3}{3\epsilon}} n_\rho(1).$$

Symmetry arguments permit easily to deduce that $m_\rho(1) = m_\rho(7) = 0$. Finally it suffices to compute $m_\rho(4)$ and $n_\rho(1)$. To this end we introduce the change of variable $u := \frac{\mathcal{U}_2}{\epsilon} z^2$ and let ρ^2/ϵ tend to infinity:

$$m_\rho(4) = \frac{3\sqrt{\pi}}{4} \left(\frac{1}{U_\mu''(x_\mu)} \right)^{\frac{5}{2}} \epsilon^{\frac{5}{2}} (1 + o_{\mathcal{I}}(1)) \quad \text{and} \quad n_\rho(1) = \frac{\epsilon}{2U_\mu''(x_\mu)} (1 + o_{\mathcal{I}}(1)).$$

To sum up: if $\frac{\rho^{18}}{\epsilon^7} \rightarrow 0$ (that is $\frac{\rho^9}{\epsilon^2} = o\{\epsilon^{\frac{3}{2}}\}$) then

$$\tilde{I}_1 = -\sqrt{\pi} f'_m(x_\mu) \frac{U_\mu^{(3)}(x_\mu)}{4} \left(\frac{1}{U_\mu''(x_\mu)} \right)^{\frac{5}{2}} \epsilon^{\frac{3}{2}} e^{-\frac{2U_\mu(x_\mu)}{\epsilon}} (1 + o_{\mathcal{I}}(1)). \tag{A.12}$$

Step 2.4. Asymptotic behavior of \tilde{I}_0 . Let us first study the following integral

$$I'_0 = \int_{-\rho}^{\rho} \exp \left[-\frac{U''_{\mu}(x_{\mu})}{\epsilon} z^2 - \frac{U_{\mu}^{(3)}(x_{\mu})}{3\epsilon} z^3 - \frac{U_{\mu}^{(4)}(x_{\mu})}{12\epsilon} z^4 \right] dz.$$

We recall the notations $\mathcal{U}_k = U_{\mu}^{(k)}(x_{\mu})$. The arguments are similar to those used in Step 2.3. Since $|e^{-u} - 1 + u - \frac{u^2}{2}| \leq |u|^3 e^{|u|}$, for any $z \in [-\rho; \rho]$ we get

$$\left| e^{-\frac{\mathcal{U}_3}{3\epsilon} z^3 - \frac{\mathcal{U}_4}{12\epsilon} z^4} - 1 + \frac{\mathcal{U}_3}{3\epsilon} z^3 + \frac{\mathcal{U}_4}{12\epsilon} z^4 - \frac{1}{2} \left(\frac{\mathcal{U}_3}{3\epsilon} z^3 + \frac{\mathcal{U}_4}{12\epsilon} z^4 \right)^2 \right| \leq C \frac{\rho^9}{\epsilon^3}.$$

Adopting the same notations as in Step 2.3 and using symmetry properties, the following bound (uniform with respect to the parameter μ) yields

$$\left| I'_0 - m_{\rho}(0) + \frac{\mathcal{U}_4}{12\epsilon} m_{\rho}(4) - \frac{1}{2} \left(\frac{\mathcal{U}_3}{3\epsilon} \right)^2 m_{\rho}(6) - \frac{1}{2} \left(\frac{\mathcal{U}_4}{12\epsilon} \right)^2 m_{\rho}(8) \right| \leq C \frac{\rho^9}{\epsilon^3} m_{\rho}(0).$$

By the usual change of variable $u := \frac{U''_{\mu}(x_{\mu})}{\epsilon} z^2$ we emphasize some asymptotic estimation of I'_0 as $\rho^2/\epsilon \rightarrow \infty$, $\rho^3/\epsilon \rightarrow 0$ and $\frac{\rho^9}{\epsilon^3} \rightarrow 0$:

$$I'_0 = \sqrt{\frac{\pi\epsilon}{U''_{\mu}(x_{\mu})}} \left\{ 1 - \frac{U_{\mu}^{(4)}(x_{\mu})}{16U''_{\mu}(x_{\mu})^2} \epsilon + \frac{5U_{\mu}^{(3)}(x_{\mu})^2}{48U''_{\mu}(x_{\mu})^3} \epsilon + o_{\mathcal{I}}(\epsilon) \right\}.$$

We apply the mean value theorem to the function U_{μ} :

$$U_{\mu}(x_{\mu} + z) = U_{\mu}(x_{\mu}) + \frac{\mathcal{U}_2}{2} z^2 + \frac{\mathcal{U}_3}{6} z^3 + \frac{\mathcal{U}_4}{24} z^4 + \frac{1}{120} U^{(5)}(y_{\mu}(t)) z^5,$$

with $y_{\mu}(t) \in [x_{\mu} - \rho, x_{\mu} + \rho]$ and $|z| \leq \rho$. From this equality we deduce an estimation of the distance $\mathcal{D} = e^{\frac{2U_{\mu}(x_{\mu})}{\epsilon}} \tilde{I}_0 - f_m(x_{\mu}) I'_0$. Then there exists some constant $C > 0$ independent of μ and m such that

$$\begin{aligned} |\mathcal{D}| &\leq |f_m(x_{\mu})| \int_{-\rho}^{\rho} e^{-\frac{\mathcal{U}_2}{\epsilon} z^2 - \frac{\mathcal{U}_3}{3\epsilon} z^3 - \frac{\mathcal{U}_4}{12\epsilon} z^4} \left| 1 - e^{-\frac{1}{60\epsilon} U^{(5)}(y_{\mu}(z+x_{\mu})) z^5} \right| dz \\ &\leq \frac{|f_m(x_{\mu})| C}{60\epsilon} \rho^5 \int_{-\rho}^{\rho} e^{-\frac{\mathcal{U}_2}{\epsilon} z^2 - \frac{\mathcal{U}_3}{3\epsilon} z^3 - \frac{\mathcal{U}_4}{12\epsilon} z^4 + \frac{1}{60\epsilon} |U^{(5)}(y_{\mu}(z+x_{\mu})) z^5|} dz. \end{aligned}$$

If both conditions $\rho^2/\epsilon \rightarrow \infty$ and $\rho^3/\epsilon \rightarrow 0$ are satisfied then the integral term in the preceding inequality is obviously equivalent to $\sqrt{\frac{\pi\epsilon}{U''_{\mu}(x_{\mu})}}$. The following equivalence holds for the initial integral \tilde{I}_0 : under the assumption that $\frac{\rho^5}{\sqrt{\epsilon}} = o\left(\epsilon^{\frac{3}{2}}\right)$, we get $|\mathcal{D}| = o_{\mathcal{I}\mathcal{M}}\left(\epsilon^{\frac{3}{2}}\right)$ and consequently

$$\tilde{I}_0 = e^{-\frac{2U(x_{\mu})}{\epsilon}} \sqrt{\frac{\pi\epsilon}{\mathcal{U}_2}} \left\{ 1 - \frac{\mathcal{U}_4}{16\mathcal{U}_2^2} \epsilon + \frac{5\mathcal{U}_3^2}{48\mathcal{U}_2^3} \epsilon + o_{\mathcal{I}\mathcal{M}}(\epsilon) \right\}. \tag{A.13}$$

Step 3. To sum up: in Step 1, we proved that it suffices to estimate the integral I_1 which can be split into 4 terms. Each of them has been estimated in Eqs. (A.10)–(A.13). The whole integral has the asymptotic equivalence (A.6) as soon as $\rho^3/\epsilon \rightarrow 0$, $\rho^{18}/\epsilon^7 \rightarrow 0$, $\frac{\rho^9}{\epsilon^3} \rightarrow 0$ and $\rho^5/\epsilon^2 \rightarrow 0$. The particular choice $\rho = \epsilon^{\frac{9}{20}}$ fulfills all these conditions.

Step 4. Now, we will prove (A.6) for $a = -\infty$ and $b = +\infty$. Let $R > \max_{\mu \in \mathcal{I}} |x_\mu|$ such that $U_\mu(t) \geq t^2$ for all $t \geq R$ and for all $\mu \in \mathcal{I}$. The integral on \mathbb{R} can be split into two integrals:

$$L = \int_{-R}^R f_m(t) e^{-\frac{2U_\mu(t)}{\epsilon}} dt + \int_{[-R; R]^c} f_m(t) e^{-\frac{2U_\mu(t)}{\epsilon}} dt = L_1 + L_2.$$

For L_1 it suffices to apply (A.6) with $a := -R$ and $b := R$ in order to get the asymptotic development. It remains then to prove that L_2 is negligible with respect to L_1 that is $L_2 = o_{\mathcal{IM}} \left\{ \epsilon^{\frac{3}{2}} e^{-\frac{2U_\mu(x_\mu)}{\epsilon}} \right\}$. Using the change of variable $t := \left(\frac{2}{\epsilon} - \lambda\right)^{-\frac{1}{2}} s$ the following bound holds:

$$|L_2| \leq 2 \int_R^{+\infty} \exp \left[t^2 \left(\lambda - \frac{2}{\epsilon} \right) \right] dt \leq 2 \sqrt{\frac{\epsilon}{2 - \lambda\epsilon}} \int_{R\sqrt{\frac{2-\lambda\epsilon}{\epsilon}}}^{+\infty} \exp \left[-s^2 \right] ds.$$

Lemma A.1 permits to prove as claimed that L_2 can be neglected.

Step 5. We just apply two times (A.6): the first time to the denominator D^ϵ that is for the function $t \rightarrow e^{f_m(t)}$ and the second time to the numerator N^ϵ for the function $t \rightarrow t^n e^{f_m(t)}$. The following asymptotic result holds

$$D^\epsilon = e^{-\frac{2U_\mu(x_\mu)}{\epsilon}} \sqrt{\frac{\pi\epsilon}{\mathcal{U}_2}} e^{f_m(x_\mu)} \left\{ 1 + \hat{\gamma}_d \epsilon + o_{\mathcal{IM}}(\epsilon) \right\} \tag{A.14}$$

where

$$\hat{\gamma}_d = \left(\frac{5\mathcal{U}_3^2}{48\mathcal{U}_2^3} - \frac{\mathcal{U}_4}{16\mathcal{U}_2^2} \right) - f'_m(x_\mu) \frac{\mathcal{U}_3}{4\mathcal{U}_2^2} + \left(f''_m(x_\mu) + f'_m(x_\mu)^2 \right) \frac{1}{4\mathcal{U}_2}.$$

The numerator normalized by x_μ^n i.e. N^ϵ/x_μ^n satisfies some similar identity as D^ϵ , namely (A.14) with $\hat{\gamma}_d$ replaced by $\hat{\gamma}_d - \frac{n\mathcal{U}_3}{4x_\mu\mathcal{U}_2^2} + \left(\frac{n(n-1)}{x_\mu^2} + 2\frac{n}{x_\mu} f'_m(x_\mu) \right) \frac{1}{4\mathcal{U}_2}$. The estimation of the ratio is then a classical exercise of asymptotic analysis. \square

We will now generalize to functions G depending on the small parameter ϵ .

Lemma A.4. *Let U and G be two $C^\infty(\mathbb{R})$ -continuous functions such that $U(t) \geq t^2$ for $|t|$ large enough and $|G(t)| \leq \lambda|U(t)| + C$ for some constants $\lambda > 0$ and $C > 0$. Moreover we assume that U admits some unique global minimum reached at x_0 with $U''(x_0) > 0$. For any sequence $(\eta_\epsilon)_\epsilon$ satisfying $\lim_{\epsilon \rightarrow 0} \eta_\epsilon = 0$ and $\lim_{\epsilon \rightarrow 0} \epsilon/\eta_\epsilon = 0$ we define $U_{\epsilon,\mu} = U + \eta_\epsilon \mu G$ depending on the parameter μ which belongs to some compact interval \mathcal{I} of \mathbb{R} . Let f a C^3 -continuous function such that $|f(t)| \leq e^{\lambda|U(t)|}$ for all $|t|$ large enough and such that $|f_m^{(i)}|$ is locally bounded uniformly with respect to $m \in \mathcal{M}$ for $0 \leq i \leq 3$. Then, there exists $\epsilon_0 > 0$ such that the potential $U_{\epsilon,\mu}$ admits a unique global minimum reached at $x_{\epsilon,\mu}$ for all $\epsilon \leq \epsilon_0$. Furthermore the following asymptotic results hold*

$$\int_{\mathbb{R}} f(t) e^{-\frac{2U_{\epsilon,\mu}(t)}{\epsilon}} dt = \sqrt{\frac{\pi\epsilon}{U''(x_0)}} e^{-\frac{2U_{\epsilon,\mu}(x_{\epsilon,\mu})}{\epsilon}} \left(f(x_0) + \gamma_\mu \eta_\epsilon + o_{\mathcal{I}}(\eta_\epsilon) \right), \tag{A.15}$$

where

$$x_{\epsilon,\mu} = x_0 - \mu \frac{G'(x_0)}{U''(x_0)} \eta_\epsilon + o_{\mathcal{I}}(\eta_\epsilon), \tag{A.16}$$

$$\gamma_\mu = \frac{\mu}{2U''(x_0)} \left(-2f'(x_0)G'(x_0) - f(x_0)G''(x_0) + f(x_0) \frac{U^{(3)}(x_0)G'(x_0)}{U''(x_0)} \right),$$

and $o_{\mathcal{I}}(\eta_\epsilon)/\eta_\epsilon$ tends to 0 as $\epsilon \rightarrow 0$ uniformly with respect to the parameter μ .

Moreover, as $\epsilon \rightarrow 0$, we have the following estimate

$$\frac{\int_{\mathbb{R}} f(t)e^{-\frac{2U_{\epsilon,\mu}(t)}{\epsilon}} dt}{\int_{\mathbb{R}} e^{-\frac{2U_{\epsilon,\mu}(t)}{\epsilon}} dt} = f(x_0) - \mu \frac{f'(x_0)G'(x_0)}{U''(x_0)} \eta_\epsilon + o_{\mathcal{I}}(\eta_\epsilon) \tag{A.17}$$

where $o_{\mathcal{I}}(\eta_\epsilon)/\eta_\epsilon$ tends to 0 as $\epsilon \rightarrow 0$ uniformly with respect to the parameter μ .

Proof. Let us first prove that the potential $U_{\epsilon,\mu}(x)$ admits a unique minimum for $x = x_{\epsilon,\mu}$ with $\lim_{\epsilon \rightarrow 0} x_{\epsilon,\mu} = x_0$. By the definitions of $(\eta_\epsilon)_\epsilon$ and $U_{\epsilon,\mu}$, the following convergence holds

$$\lim_{\epsilon \rightarrow 0} U_{\epsilon,\mu}(x_0) = U(x_0). \tag{A.18}$$

Since x_0 is the unique global minimum of U , for any small $R > 0$ there exists $\rho_R > 0$ such that $\inf_{x \in [x_0-R, x_0+R]^c} U(x) > U(x_0) + \rho_R$. We deduce the existence of two small constants ρ'_R and ϵ_0 such that

$$U_{\epsilon,\mu}(x) \geq (1 - \mu\lambda\eta_\epsilon)U(x) - \eta_\epsilon\mu C \geq U(x_0) + \rho'_R, \tag{A.19}$$

for all $\epsilon \leq \epsilon_0$ and $x \in [x_0 - R, x_0 + R]^c$. By (A.18) and (A.19) we obtain: for any $R > 0$ the global minimum of the parametrized potential $U_{\epsilon,\mu}$ is reached in the interval $[x_0 - R, x_0 + R]$ provided that ϵ is small enough (uniformly with respect to μ). Moreover this global minimum is unique. Indeed $U''(x_0) > 0$ and the regularity of U implies that $U''(x) > 0$ for all x in some small neighborhood of x_0 . Since $U''_{\epsilon,\mu}$ converges towards U'' as $\epsilon \rightarrow 0$ uniformly on each compact subset of \mathbb{R} , we obtain that $U''_{\epsilon,\mu} > 0$ on $[x_0 - R, x_0 + R]$ provided that R and ϵ are small enough. The minimum is actually unique, we denote its localization $x_{\epsilon,\mu}$ and point out that, for ϵ small, $U''_{\epsilon,\mu}(x_{\epsilon,\mu}) > 0$ uniformly with respect to μ . Let us determine $x_{\epsilon,\mu}$. By applying the mean value theorem to $U_{\epsilon,\mu}$, we get

$$0 = U'_{\epsilon,\mu}(x_{\epsilon,\mu}) = U'(x_0) + \mu\eta_\epsilon G'(x_0) + U''_{\epsilon,\mu}(\tilde{x})(x_{\epsilon,\mu} - x_0),$$

where \tilde{x} is in between x_0 and $x_{\epsilon,\mu}$. Since the second derivative is continuous, $U''_{\epsilon,\mu}(\tilde{x})$ is uniformly bounded. Moreover $U'(x_0) = 0$. Consequently $x_{\epsilon,\mu} - x_0 = \mathcal{O}_{\mathcal{I}}(\eta_\epsilon)$. Using the same argument for the second order asymptotic development of $U'_{\epsilon,\mu}(x_{\epsilon,\mu})$, that is

$$0 = U'(x_0) + \mu\eta_\epsilon G'(x_0) + (U''(x_0) + \mu\eta_\epsilon G''(x_0))(x_{\epsilon,\mu} - x_0) + \frac{U_{\epsilon,\mu}^{(3)}(\tilde{x})}{2}(x_{\epsilon,\mu} - x_0)^2,$$

we obtain the announced estimate (A.16). Finally let us prove the estimate (A.15). The statement of Lemma A.3 can be applied to $U_{\epsilon,\mu}$ since the asymptotic result (A.6) is uniform with respect to the parameter μ . So it suffices to consider the case when μ is replaced by $\mu\eta_\epsilon$. We immediately obtain

$$\int_{\mathbb{R}} f(t)e^{-\frac{U_{\epsilon,\mu}(t)}{\epsilon}} dt = \sqrt{\frac{\pi\epsilon}{U''_{\epsilon,\mu}(x_{\epsilon,\mu})}} f(x_{\epsilon,\mu})e^{-\frac{U_{\epsilon,\mu}(x_{\epsilon,\mu})}{\epsilon}} (1 + o_{\mathcal{I}}(\eta_\epsilon)). \tag{A.20}$$

It remains to approximate $f(x_{\epsilon,\mu})$ and $U''_{\epsilon,\mu}(x_{\epsilon,\mu})$ using (A.16). Due to the regularity of both f and U , the following developments hold

$$f(x_{\epsilon,\mu}) = f(x_0) - \mu\eta_\epsilon f'(x_0) \frac{G'(x_0)}{U''(x_0)} + o_{\mathcal{I}}(\eta_\epsilon),$$

$$U''_{\epsilon,\mu}(x_{\epsilon,\mu}) = U''(x_0) + \mu\eta_\epsilon \left(G''(x_0) - U^{(3)}(x_0) \frac{G'(x_0)}{U''(x_0)} \right) + o_{\mathcal{I}}(\eta_\epsilon).$$

The statement of Lemma A.4 is obtained just by combination of the two preceding asymptotics and (A.20). Then, we apply (A.15) to the numerator and to the denominator. After dividing, we find (A.17). \square

Remark A.5. The statements of Lemmas A.2–A.4 can be easily generalized, replacing the parametrized function $U_\mu = U + \mu G$ by $U_\mu = U + \sum_{i=1}^k \mu_i G_i$ where $\mu = (\mu_1, \dots, \mu_k) \in \mathcal{I}_1 \times \dots \times \mathcal{I}_k$. The convergence results are then uniform with respect to all parameters.

References

- [1] C. Ané, S. Blachère, D. Chafaï, P. Fougères, I. Gentil, F. Malrieu, C. Roberto, G. Scheffer, Sur les inégalités de Sobolev logarithmiques, in: Panoramas et Synthèses (Panoramas and Syntheses), vol. 10, Société Mathématique de France, Paris, 2000.
- [2] S. Benachour, B. Roynette, D. Talay, P. Vallois, Nonlinear self-stabilizing processes. I. Existence, invariant probability, propagation of chaos, Stochastic Process. Appl. 75 (2) (1998) 173–201.
- [3] S. Benachour, B. Roynette, P. Vallois, Nonlinear self-stabilizing processes. II. Convergence to invariant probability, Stochastic Process. Appl. 75 (2) (1998) 203–224.
- [4] T. Funaki, A certain class of diffusion processes associated with nonlinear parabolic equations, Z. Wahrsch. Verw. Gebiete 67 (3) (1984) 331–348.
- [5] D. Gilbarg, N.S. Trudinger, Elliptic partial differential equations of second order, in: Grundlehren der Mathematischen Wissenschaften, vol. 224, Springer-Verlag, Berlin, 1977.
- [6] S. Herrmann, P. Imkeller, D. Peithmann, Large deviations and a Kramers' type law for self-stabilizing diffusions, Ann. Appl. Probab. 18 (4) (2008) 1379–1423.
- [7] H.P. McKean Jr., A class of Markov processes associated with nonlinear parabolic equations, Proc. Natl. Acad. Sci. USA 56 (1966) 1907–1911.
- [8] H.P. McKean Jr., Propagation of chaos for a class of non-linear parabolic equations., in: Stochastic Differential Equations (Lecture Series in Differential Equations, Session 7, Catholic Univ., 1967), Air Force Office Sci. Res., Arlington, VA, 1967, pp. 41–57.
- [9] K. Oelschläger, A law of large numbers for moderately interacting diffusion processes, Z. Wahrsch. Verw. Gebiete 69 (2) (1985) 279–322.
- [10] D.W. Stroock, S.R.S. Varadhan, Multidimensional diffusion processes, in: Grundlehren der Mathematischen Wissenschaften (Fundamental Principles of Mathematical Sciences), vol. 233, Springer-Verlag, Berlin, 1979.
- [11] A.-S. Sznitman, Topics in propagation of chaos, in: École d'Été de Probabilités de Saint-Flour XIX—1989, in: Lecture Notes in Math., vol. 1464, Springer, Berlin, 1991, pp. 165–251.
- [12] Y. Tamura, Free energy and the convergence of distributions of diffusion processes of McKean type, J. Fac. Sci. Univ. Tokyo Sect. IA Math. 34 (2) (1987) 443–484.
- [13] Y. Tamura, On asymptotic behaviors of the solution of a nonlinear diffusion equation, J. Fac. Sci. Univ. Tokyo Sect. IA Math. 31 (1) (1984) 195–221.