Available online at www.sciencedirect.com
ScienceDirect
Neuroscience Letters 417 (2007) 6–9
Neuroscience Letters
www.elsevier.com/locate/neulet

Association of the oxytocin receptor gene (*OXTR*) in Caucasian children and adolescents with autism

Suma Jacob [a,*], Camille W. Brune [a], C.S. Carter [b], Bennett L. Leventhal [a], Catherine Lord [c], Edwin H. Cook Jr. [a]

[a] *Institute for Juvenile Research, Department of Psychiatry, University of Illinois at Chicago, Chicago, IL 60608, USA*
[b] *Brain Body Center, Department of Psychiatry, University of Illinois at Chicago, Chicago, IL 60608, USA*
[c] *University of Michigan Autism and Communication Disorders Center (UMACC), 1111 East Catherine St., Rm. 217, Ann Arbor, MI 48109, USA*

Received 20 October 2006; received in revised form 26 December 2006; accepted 1 February 2007

Abstract

The oxytocin receptor gene (*OXTR*) has been studied in autism because of the role of oxytocin (OT) in social cognition. Linkage has also been demonstrated to the region of *OXTR* in a large sample. Two single nucleotide polymorphisms (SNPs) and a haplotype constructed from them in *OXTR* have been associated with autism in the Chinese Han population. We tested whether these associations replicated in a Caucasian sample with strictly defined autistic disorder. We genotyped the two previously associated SNPs (rs2254298, rs53576) in 57 Caucasian autism trios. Probands met clinical, ADI-R, and ADOS criteria for autistic disorder. Significant association was detected at rs2254298 ($p = 0.03$) but not rs53576. For rs2254298, overtransmission of the G allele to probands with autistic disorder was found which contrasts with the overtransmission of A previously reported in the Chinese Han sample. In both samples, G was more frequent than A. However, in our Caucasian autism trios and the CEU Caucasian HapMap samples the frequency of A was less than that reported in the Chinese Han and Chinese in Bejing HapMap samples. The haplotype test of association did not reveal excess transmission from parents to affected offspring. These findings provide support for association of OXTR with autism in a Caucasian population. Overtransmission of different alleles in different populations may be due to a different pattern of linkage disequilibrium between the marker rs2254298 and an as yet undetermined susceptibility variant in *OXTR*.
© 2007 Elsevier Ireland Ltd. All rights reserved.

Keywords: Oxytocin; OXTR; Autism; Genetics; Linkage disequilibrium; Polymorphisms; Neuropeptide hormone

Autism is a neurodevelopmental disorder that is characterized by impairments in communication and social interaction as well as patterns of restrictive, repetitive interests and behaviors during early childhood (DSM-IV). Deficits in social interaction can include lack of social or emotional reciprocity, absence of shared enjoyment with parents and others, limited nonverbal behavior to regulate social interaction, and difficulty developing friendships. There is a male to female ratio of 4:1 or greater [4]. There is strong evidence for a complex genetic influence of autism with estimates of concordance among monozygotic twins ranging from 64 to 91% and with fraternal twins or siblings from 0 to 9% (reviewed in [24]).

There is growing interest in the role of the neurohypophyseal peptide oxytocin (OT) in the development of autism [9] because of its role in affiliation, social memory and behavior. In animal models, oxytocin has been shown to play critical roles in social processing, recognition and bonding as well as influencing stereotyped behaviors such as exaggerated grooming [9,3,5,25]. Cells mediating centrally-released OT are found distinctly in the paraventricular and supraoptic nuclei of the hypothalamus. In Ferguson et al. [5] OT knockout mice maintained olfaction and cognitive performance, but suffered deficits in social recognition which were recovered by intraventricular OT but not by vasopressin administration. Animal models have shown that altering OT early in life can produce long-lasting and sexually dimorphic changes on brain development and behavior [2].

In human studies, OT administration has been shown to increase trust [12] and amygdala activation compared to placebo in healthy males [11]. Elevated OT has been reported in obsessive compulsive disorder [6,21] and Prader-Willi syndrome [16].

* Corresponding author. Institute for Juvenile Research, Department of Psychiatry (M/C 747), University of Illinois at Chicago, 1747 West Roosevelt Rd, Room 155, Chicago, IL 60608, USA. Tel.: +1 312 413 4623.
E-mail address: sjacob@psych.uic.edu (S. Jacob).

0304-3940/$ – see front matter © 2007 Elsevier Ireland Ltd. All rights reserved.
doi:10.1016/j.neulet.2007.02.001

Children with autism have been shown to have lower blood OT levels [17] and higher precursor OT levels [6] in comparison to typically developing children. Differences in OT peptide processing may result in inactive or less active forms of OT and potentially impact brain development or behavior associated with autism. There are also reports that treatment with OT infusions results in reduced repetitive behaviors [8] and increased retention of affective speech [7] in adults with autism and Asperger disorder.

The OXTR is a G protein-coupled receptor and positively coupled to phospholipase C [13]. In mammals, OXTR are expressed at higher levels in early development [19,23] and are concentrated in brain regions that are involved in social behaviors including the olfactory bulbs, piriform cortex, amygdala, and lateral septum [5]. Compared to wild type, *OXTR* knockout mice emit fewer ultrasonic vocalizations in response to social isolation, experience deficits in social discrimination, and demonstrate more aggressive behavior [22].

Combined linkage analysis of two independent samples of 314 Finnish families demonstrated linkage in the 3p24-25 region containing the OXTR gene [26]. Four polymorphic SNPs were analyzed and two were found to be associated with autism in 195 Chinese Han trios [21]. In order to replicate these findings in a Caucasian population, we genotyped the two SNPs that were significant in the previous sample in 57 Caucasian trios and conducted family-based association analyses.

After complete description of the study to the parents, written informed consent was obtained. Recruitment, assessment, and inclusion criteria were the same as that outlined in a previously described sample in which all subjects met ADI-R criteria for autism and had a best estimate diagnosis of autistic disorder by a clinical psychologist and psychiatrist [10]. Only one sibling was randomly selected from each affected sibling pair in the previous study [10] to avoid confounding linkage and association. For the current study additional inclusion criteria consisted of being Caucasian (because allele frequency differences are reported across populations and there was an insufficient sample of non-Caucasian subjects in the sample), being at least 3 years old at the time the Autism Diagnostic Interview-Revised (ADI-R, [15]) was administered, having sufficient blood or DNA available, meeting Autism Diagnostic Observation Schedule (ADOS) classification for autistic disorder [14] (thereby dropping subjects in the previous sample whose ADOS classification was autistic spectrum disorder), and having complete data for ADI-R algorithm item scores (thereby including subjects recruited from the University of Chicago Developmental Disorders Clinic and excluding subjects recruited from San Diego). There were 57 probands, 45 males and 12 females, with a mean age of 6.4 years (S.D. = 3.5).

All genotyping was performed blind to clinical and demographic data and family relationships. Two SNP markers (rs2254298 [Celera ID: C_15981334_10] and rs53576 [Celera ID: C_3290335_10] were genotyped using TaqMan® SNP Genotyping Assays (Applied Biosystems, Foster City, CA, http://www.appliedbiosystems.com). TaqMan® PCR reactions were done with Universal Master Mix Amperase® UNG, 0.25 μL Taqman probe mix and 2.25 μL of water for a 5 μL total volume. The PCR conditions for the TaqMan® SNP Genotype Assays were: one AmpErase® step at 50.0 °C for 2 min, one enzyme activation step at 95.0 °C for 10 min, and 40 alternating cycles of denaturation at 92.0 °C for 15 s and reannealing and extension at 58.0 °C for one minute. All PCR reactions were performed on a Perkin Elmer 9700 Thermocycler (Applied Biosystems, Foster City, CA). The fluorescence intensity of the final PCR product was measured using an LjL Analyst AD fluorescence microplate reader (LjL Biosystems, Sunnyvale, CA, http://www.moleculardevices.com) using LjL Criterion-Host Software. In addition to the 57 Caucasian trios reported, one trio was excluded from the analyses because of a parent sample genotyping failure. The overall test retest-agreement performed on 16% of the sample was 100%. There were no Mendelian incompatibilities for either marker.

The distributions of rs2254298 and rs53576 genotypes were tested using X^2 for Hardy-Weinberg equilibrium using the HWE program from the LINKUTIL package (http://linkage.rockefeller.edu/ott/linkutil.htm). Transmission disequilibrium tests were calculated using the TDT/S-TDT program (v. 1.1) (http://genomics.med.upenn.edu/spielman/TDT.htm) [20]. To test for association between SNPs, we used Haploview 3.32 (http://www.broad.mit.edu/mpg/haploview/) to calculate two measures of linkage disequilibrium, D' and r^2 [1]. Haplotype association was calculated using the FBAT program version 1.7.2 (http://www.biostat.harvard.edu/~fbat/default.html) under "biallelic" mode [18]. Alpha was set at $p < .05$.

Table 1 reports allele and genotype frequencies of the sample and the TDT for each marker. Genotype distributions for parents and probands were consistent with Hardy-Weinberg equilibrium for rs2254298 (parents: $X^2 = 0.94$, d.f. = 1, $p = 0.34$; probands: $X^2 = 0.53$, $p = 0.47$) and rs53576 (parents: $X^2 = 0.07$, $p = 0.79$; probands: $X^2 = 0.42$, $p = 0.52$).

Table 1
Allele and genotype frequencies of SNPs in pedigrees

SNP	Sample	Allele distribution		Genotype distribution			TDT		
		G	A	GG	AG	AA	*t*(G)/*t*(A)	χ^2	*p*
rs2254298	Parents	196	32	83	30	1	–	–	–
	Autistic probands	104	10	47	10	0	21/9	4.80	0.03
rs53576	Parents	166	62	61	44	9	–	–	
	Autistic probands	84	30	30	24	3	23/21	0.09	0.76

SNP, single nucleotide polymorphism; TDT, transmission disequilibrium test; *t*(G)/*t*(A), number of transmitted G and A alleles from heterozygous parents.

Table 2
Genetic association between haplotypes and autism

Allele rs2254298	Allele rs53576	S	$E(S)$	Z	p^*
G	G	50.250	45.250	1.347	0.178
G	A	30.750	29.750	0.291	0.771
A	G	8.750	12.750	−1.629	0.103
A	A	0			

S, test statistics for the observed number of transmitted alleles; $E(S)$, expected value of S under the null hypothesis (i.e., no linkage or association).
* p, two-tailed.

Preferential transmission of G over A occurred at rs2254298 (see Table 1). No significant association was found for rs53576. The markers showed weak LD with each other ($D' = 0.53$, $r^2 = 0.01$). The haplotype test of association did not reveal excess transmission from parents to affected offspring for the haplotypes ($p > 0.10$) (Table 2).

Since there is growing evidence that OXTR may mediate genetic vulnerability to autism [9,27], we tested two SNPs that were found to be associated with autism in the Chinese Han population [21]. The goal was to replicate these findings in a Caucasian population using standardized diagnostic criteria. The TDT revealed significant transmission disequilibrium for rs2254298, one of the two SNPs that were significant in the Chinese Han sample. In contrast to the Chinese Han population, the G allele was overtransmitted in this population. Allele frequencies differ markedly at this marker in Caucasian and Chinese populations (Table 3). Both the current study and Wu et al. [21] reported allele frequencies representative of their population in the HapMap (http://www.hapmap.org) with the Caucasian samples having a greater frequency of the G allele than the Chinese samples (Table 3).

Although the transmission of different alleles may occur because both studies are false positives or because of phenotypic heterogeneity, the transmission of different marker alleles is also consistent with the alleles being on different haplotypes with an as yet unidentified susceptibility variant in *OXTR*. This is more likely given that the minor allele frequencies at rs2259248 and other SNPs in *OXTR* differ between the 2 populations. Future research should attempt to replicate these preliminary findings across populations using larger samples. Extensive genotyping, resequencing and family-based association testing in larger samples across populations will be necessary to understand the possible role of variants in OXTR in autism susceptibility.

Table 3
Parental allele frequencies for *rs2254298* in autism association studies and the HapMap

Allele	Caucasian		Chinese	
	Current study	HapMap[a]	[21]	HapMap[b]
A	0.140	0.068	0.315	0.289
G	0.860	0.932	0.685	0.711

[a] CEU—CEPH, Utah residents with ancestry from northern and western Europe.
[b] CHB—Han Chinese in Beijing, China.

Acknowledgments

The authors are especially grateful to the families who participated in the study. Kathy Hennessy provided expert technical assistance and Jeremy Veenstra-VanderWeele provided assistance with statistical programs. This work was supported, in part, by NIH U19 HD35482 (E.C., C.L.), R01 MH066496 (C.L.), 5T32MH7631 (S.J.), the Jean Young and Walden W. Shaw Foundation, the Harris Foundation (C.W.B.), and the Children's Brain Research Foundation (E.C.).

References

[1] J. Barrett, B. Fry, J. Maller, M. Daly, Haploview: analysis and visualization of LD and haplotype maps, Bioinformatics 21 (2005) 263–265.
[2] C.S. Carter, Developmental consequences of oxytocin, Physiol. Behav. 79 (2003) 383–397.
[3] C.S. Carter, Neuroendocrine perspectives on social attachment and love, Psychoneuroendocrinology 23 (1998) 779–818.
[4] S. Chakrabarti, E. Fombonne, Pervasive developmental disorders in preschool children: confirmation of high prevalence, Am. J. Psychiatry 162 (2005) 1133–1141.
[5] J.N. Ferguson, L.J. Young, E.F. Hearn, M.M. Matzuk, T.R. Insel, J.T. Winslow, Social amnesia in mice lacking the oxytocin gene, Nat. Genet. 25 (2000) 284–288.
[6] L. Green, D. Fein, C. Modahl, C. Feinstein, L. Waterhouse, M. Morris, Oxytocin and autistic disorder: alterations in peptide forms, Biol. Psychiatry 50 (2001) 609–613.
[7] E. Hollander, J. Bartz, W. Chaplin, A. Phillips, J. Sumner, L. Sooyra, E. Anagnostou, S. Wasserman, Oxytocin increases retention of social cognition in autism, Biol. Psychiatry 61 (4) (2007) 498–503 (Epub. ahead of print).
[8] E. Hollander, S. Novotny, M. Hanratty, R. Yaffe, C.M. DeCaria, B.R. Aronowitz, S. Mosovich, Oxytocin infusion reduces repetitive behaviors in adults with autistic and Asperger's disorders, Neuropsychopharmacology 28 (2003) 193–198.
[9] T.R. Insel, D.J. O'Brien, J.F. Leckman, Oxytocin, vasopressin, and autism: is there a connection? Biol. Psychiatry 45 (1999) 145–157.
[10] S.J. Kim, N. Cox, R. Courchesne, C. Lord, C. Corsello, N. Akshoomoff, S. Guter, B.L. Leventhal, E. Courchesne, E.H. Cook Jr., Transmission disequilibrium mapping at the serotonin transporter gene (SLC6A4) region in autistic disorder, Mol. Psychiatry 7 (2002) 278–288.
[11] P. Kirsch, C. Esslinger, Q. Chen, D. Mier, S. Lis, S. Siddhanti, H. Gruppe, V.S. Mattay, B. Gallhofer, A. Meyer-Lindenberg, Oxytocin modulates neural circuitry for social cognition and fear in humans, J. Neurosci. 25 (2005) 11489–11493.
[12] M. Kosfeld, M. Heinrichs, P.J. Zak, U. Fischbacher, E. Fehr, Oxytocin increases trust in humans, Nature 435 (2005) 673–676.
[13] Y. Kubota, T. Kimura, K. Hashimoto, Y. Tokugawa, K. Nobunaga, C. Azuma, F. Saji, Y. Murata, Structure and expression of the mouse oxytocin receptor gene, Mol. Cell. Endocrinol. 124 (1996) 25–32.
[14] C. Lord, S. Risi, L. Lambrecht, E.H. Cook Jr., B.L. Leventhal, P.C. DiLavore, A. Pickles, M. Rutter, The autism diagnostic observation schedule-generic: a standard measure of social and communication deficits associated with the spectrum of autism, J. Autism Dev. Disorders 30 (2000) 205–223.
[15] C. Lord, M. Rutter, A. Le Couteur, Autism Diagnostic Interview – Revised: a revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders, J. Autism Dev. Disorders 24 (1994) 659–685.
[16] A. Martin, M. State, K. Koenig, R. Schultz, E.M. Dykens, S.B. Cassidy, J.F. Leckman, Prader-Willi syndrome, Am. J. Psychiatry 155 (1998) 1265–1273.
[17] C. Modahl, L. Green, D. Fein, M. Morris, L. Waterhouse, C. Feinstein, H. Levin, Plasma oxytocin levels in autistic children, Biol. Psychiatry 43 (1998) 270–277.

[18] D. Rabinowitz, N. Laird, A unified approach to adjusting association tests for population admixture with arbitrary pedigree structure and arbitrary missing marker information, Human Heredity 50 (2000) 211–223.
[19] L.E. Shapiro, T.R. Insel, Ontogeny of oxytocin receptors in rat forebrain: a quantitative study, Synapse 4 (1989) 259–266.
[20] R.S. Spielman, R.E. McGinnis, W.J. Ewens, Transmission test for linkage disequilibrium: the insulin gene region and insulin-dependent diabetes mellitus, Am. J. Human Genet. 52 (1993) 506–516.
[21] S. Wu, M. Jia, Y. Ruan, J. Liu, Y. Guo, M. Shuang, X. Gong, Y. Zhang, X. Yang, D. Zhang, Positive association of the oxytocin receptor gene (OXTR) with autism in the Chinese Han population, Biol. Psychiatry 58 (2005) 74–77.
[22] Y. Takayanagi, M. Yoshida, I.F. Bielsky, H.E. Ross, M. Kawamata, T. Onaka, T. Yanagisawa, T. Kimura, M.M. Matzuk, L.J. Young, K. Nishimori, Pervasive social deficits, but normal parturition, in oxytocin receptor-deficient mice, Proc. Natl. Acad. Sci. U.S.A. 102 (2005) 16096–16101.
[23] E. Tribollet, M. Goumaz, M. Raggenbass, J.J. Dreifuss, Appearance and transient expression of vasopressin and oxytocin receptors in the rat brain, J. Recept. Res. 11 (1991) 333–346.
[24] J. Veenstra-VanderWeele, E.H. Cook Jr., Molecular genetics of autism spectrum disorder, Mol. Psychiatry 9 (2004) 819–832.
[25] J.T. Winslow, P.L. Noble, C.K. Lyons, S.M. Sterk, T.R. Insel, Rearing effects on cerebrospinal fluid oxytocin concentration and social buffering in rhesus monkeys, Neuropsychopharmacology 28 (2003) 910–918.
[26] T. Ylisaukko-oja, M. Alarcon, R.M. Cantor, M. Auranen, R. Vanhala, E. Kempas, L. von Wendt, I. Jarvela, D.H. Geschwind, L. Peltonen, Search for autism loci by combined analysis of autism genetic resource exchange and Finnish families, Ann. Neurol. 59 (2006) 145–155.
[27] L.J. Young, L.J. Pitkow, J.N. Ferguson, Neuropeptides and social behavior: animal models relevant to autism, Mol. Psychiatry 7 (Suppl. 2) (2002) S38–S39.