

Drug-like property-driven optimization of 4-substituted 1,5-diarylanilines as potent HIV-1 non-nucleoside reverse transcriptase inhibitors against rilpivirine-resistant mutant virus

Lei Wei^a, Hui-Ling Wang^a, Li Huang^b, Chin-Ho Chen^b, Susan L. Morris-Natschke^c, Kuo-Hsiung Lee^{c,d,*}, Lan Xie^{a,c,*}

^a Beijing Institute of Pharmacology & Toxicology, 27 Tai-Ping Road, Beijing 100850, China

^b Duke University Medical Center, Box 2926, Surgical Oncology Research Facility, Durham, NC 27710, USA

^c Natural Products Research Laboratories, UNC Eshelman School of Pharmacy, University of North Carolina, Chapel Hill, NC 27599, USA

^d Chinese Medicine Research and Development Center, China Medical University and Hospital, Taichung, Taiwan

ARTICLE INFO

Article history:

Received 28 March 2017

Revised 19 April 2017

Accepted 20 April 2017

Available online 22 April 2017

Keywords:

HIV-1 NNRTIs

Diarylaniline

Drug-like property

Anti-resistance

ABSTRACT

On the basis of our prior structure-activity relationship (SAR) results, our current lead optimization of 1,5-diarylanilines (DAANs) focused on the 4-substituent (R^1) on the central phenyl ring as a modifiable position related simultaneously to improved drug resistance profiles and drug-like properties. Newly synthesized *p*-cyanovinyl-DAANs (**8a–8g**) with different R^1 side chains plus prior active *p*-cyanoethyl-DAANs (**4a–4c**) were evaluated not only for anti-HIV potency against both wild-type HIV virus and rilpivirine-resistant (E138K, E138K+M184I) viral replication, but also for multiple drug-like properties, including aqueous solubility, lipophilicity, and metabolic stability in human liver microsomes and human plasma. This study revealed that both ester and amide R^1 substituents led to low nanomolar anti-HIV potency against wild-type and rilpivirine-resistant viral strains (E138K-resistance fold changes < 3). The *N*-substituted amide- R^1 side chains were superior to ester- R^1 likely due to improved aqueous solubility, lipophilicity, and higher metabolic stability *in vitro*. Thus, three amide-DAANs **8e**, **4a**, and **4b** were identified with high potency against wild-type and rilpivirine-resistant viral strains and multiple desirable drug-like properties.

© 2017 Elsevier Ltd. All rights reserved.

HIV-1 non-nucleoside reverse transcriptase inhibitors (NNRTIs) play an important role in antiretroviral therapy (ART) for treating HIV infection and AIDS disease.¹ The early NNRTIs drugs, including delavirdine, nevirapine and efavirenz, were followed by two new-generation HIV-1 NNRTI drugs etravirine (**1**)² and rilpivirine (**2**)³ (Fig. 1), approved by the US FDA in 2008 and 2011, respectively. The new drugs **1** and **2** exhibit extremely high potency against wild-type virus and a broad spectrum of viral strains resistant to early NNRTI drugs.⁴ However, patients are now showing new drug-resistance profiles,^{5,6} with distinct viral mutations from those to the early NNRTI drugs. Therefore, to combat the new drug-resistance issues, the goal of this study is to discover and develop new

anti-HIV drugs with high potency and different chemical scaffolds as well as better drug-like properties.

In our prior studies on novel HIV-1 NNRTI agents, we discovered a new chemo-type of lead compounds diarylanilines (DAANs)⁷ (Fig. 2) and subsequently produced several series of highly potent lead compounds with low nanomolar EC_{50} values against wild-type and several early drug-resistant viral strains. By iterative structure optimizations,^{8–11} we identified several pharmacophores of DAANs as shown in Fig. 2. The *p*-cyanoaniline (A-ring) and trisubstituted phenyl (C-ring) moieties are necessary for anti-HIV potency and create a favorable “U” conformation for binding in the HIV-1 RT-NNRTI binding pocket. The amino group (NH_2) on the central phenyl ring (B-ring) provides additional H-bonds with amino acid K101 on the HIV-1 RT enzyme and, thus, is crucial for enhanced anti-HIV activity.⁷ Finally, a linear hydrophobic *p*-cyano-vinyl or *p*-cyanoethyl substituent on the C-ring, as in prior leads **3** and **4a**, helps maintain high potency against wild-type and the drug-resistant viral strains associated with early NNRTI agents. In

* Corresponding authors at: Beijing Institute of Pharmacology & Toxicology, 27 Tai-Ping Road, Beijing 100850, China and Natural Products Research Laboratories, UNC Eshelman School of Pharmacy, University of North Carolina, Chapel Hill, NC 27599, USA.

E-mail addresses: khlee@unc.edu (K.-H. Lee), lxie@email.unc.edu (L. Xie).





Fig. 1. HIV-1 NNRTI drugs approved by U.S. FDA.


Fig. 2. New-generation NNRTI DAAN leads, known SAR, and leads **3** and **4a**.

contrast, the 4-substituent (R¹) on the B-ring is modifiable; it can be changed to improve drug-like properties without losing anti-HIV potency. Recently, our structure-activity relationship (SAR) and molecular modeling studies on a series of *p*-cyanoethyl-DAAN compounds¹² revealed that the R¹ substituent can also be altered to provide activity against the drug-resistant E138K variant, a major mutation conferring resistance to **2**.^{13,14} Specifically, a carbonyl group in this substituent conjugated with the B-ring is likely a necessary structure for improved activity against the drug-resistant E138K variant.¹² Therefore, our current lead optimization focused on investigating the R¹ substituent's relationship with three factors, antiviral potency, inhibition of drug-resistance related to **2**, and multiple drug-like properties.

In our continued study, new *p*-cyanovinyl-DAAN compounds (**8a–8g**) with different R¹ substituents at the 4-position were designed and synthesized. The R¹ substituents in all seven new derivatives possessed a carbonyl group conjugated with the B-ring. The new *p*-cyanovinyl-DAAN compounds **8a–8g**, plus prior lead **3** and three related *p*-cyanoethyl-DAANs (**4a–4c**), were evaluated in parallel against wild-type and resistant viral strains E138K and

E138K+M184I resistant to new-generation NNRTI drug **2**. Their drug-like properties, including aqueous solubility, lipophilicity, and *in vitro* metabolic stability, were also assessed. With these data, we can explore the balance between high anti-HIV potency and desirable drug-like properties, with the aim of discovering and developing potential new NNRTI drug candidates.

As shown in Scheme 1, cyanovinyl-DAANs **8a–8g** were synthesized from **5**, a prior intermediate in the synthesis of DAANs.⁸ The 4-ester group on the central phenyl ring in **5** was hydrolyzed in aqueous sodium hydroxide solution to produce carboxylic acid **6**. Compound **6** was then reacted with 2-methoxyethanol or 2-cyanoethanol in the presence of 1,3-dicyclohexylcarbodiimide (DCC) and 4-(dimethylamino)pyridine (DMAP) to produce corresponding ester compounds **7a** and **7b**, respectively. In addition, compound **6** was treated with oxalyl chloride in CH₂Cl₂ followed by amidation with 2-chloroethylamine, 3-aminopropanol, or 3,3-difluoroazetidine in the presence of Et₃N to produce corresponding amide compounds **7c**, **7d**, or **7f**, respectively. Moreover, 4-chloroethylamide **7c** was treated with potassium carbonate in DMF to produce the 4-aziridinylcarbamoyl compound **7e**. Alternatively, compound **5**



Scheme 1. Reagents and conditions: i. NaOH/MeOH/THF, rt, 2 h; ii. ROH, DCC/DMAP/CH₂Cl₂, rt, 2–3 h; iii. a) oxalyl chloride/CH₂Cl₂, rt, overnight, b) Et₃N, amine, rt, 2–3 h; iv. hydrazine/EtOH, reflux; v. K₂CO₃/DMF; vi. Zn/AcOH/CH₂Cl₂, 0 °C–rt, 1 h.

was reacted with hydrazine in ethanol under reflux (80 °C) to yield 4-hydrazide compound **7g**. Finally, the nitro group in **7a–7g** was reduced by using zinc powder in the presence of acetic acid to afford corresponding new target 4-substituted DAANs **8a–8g**, respectively. The structures of the new DAANs were identified from proton NMR and mass spectrometry data.

All newly synthesized *p*-cyanovinyl-DAAN compounds **8a–8g** were first evaluated against wild-type HIV-1 (NL4-3) replication in the TZM-bl cell line. Next, selected highly potent compounds (EC_{50(wt)} < 10 nM) were tested against the NL4-3 variants with E138K and double-mutated E138K+M184I drug resistant genotypes. Lead **3** and prior active *p*-cyanoethyl-DAAN compounds **4a–4c** were also tested in parallel in the same assays with drug **2** as a reference. The data for anti-HIV potency (EC₅₀, as measured by a luciferase gene expression assay¹⁵) against wild-type and resistant viral strains as well as resistance fold changes (FCs) are summarized in Table 1. As expected, all new series-8 compounds exhibited potent anti-HIV activity against wild-type virus replication with EC₅₀ values ranging from 1.11 to 24.9 nM. Among them, *N*-hydroxypropylcarbamoyl-DAAN **8d**, with a longer R¹ side chain, was the least potent (EC₅₀ 24.9 nM) compound. Importantly, the most potent cyanovinyl- (**8a–8c**, **8e**, **3**) and cyanoethyl-DAANs (**4a–4c**) (EC_{50(NL4-3)} < 10 nM) also exhibited low resistance fold changes (FC) of 1.6 to 2.9 in relation to wild-type NL4-3 virus, while **2** had a much higher FC of 8.1. Furthermore, these same DAANs also inhibited the E138K+M184I viral strain with resistance FCs ranging from 1.4 to 6.3, better than or similar to that of **2** in the same assay. Consistent with previous SAR results, the R¹ substituent proved modifiable without loss of anti-HIV potency; compounds with both ester and amide R¹ side chains maintained high potency at a low nanomolar level against both wild-type and 2-resistant viral strains.

Three essential drug-like properties, solubility, lipophilicity, and metabolic stability, of these DAAN compounds were also assessed *in vitro*. The studies were aimed at pharmacological improvements to avoid or decrease risks in later new drug development. According to the HPLC methods described in our prior studies,⁸ aqueous solubility was measured at both pH 2.0 and pH 7.4, typical physiological


environments encountered by a drug in stomach and plasma, respectively. As shown in Table 2, these compounds were generally more soluble in acidic (pH 2.0) rather than neutral (pH 7.4) conditions, because of the presence of a free amino group. The different R¹ substituents had obvious effects on the molecular aqueous solubility at pH 2.0. In the series-8 DAANs, ester-compounds **8a** and **8b** were poorly soluble at pH 2.0 (1.64 and 0.94 µg/mL respectively), and non-substituted amide-DAAN **3** was only slightly more soluble (2.09 µg/mL). However, *N*-substituted amide-DAANs **8c–8g** displayed improved aqueous solubility ranging from moderate to high (9.53–94.2 µg/mL) compared with the remaining compounds. A similar situation was also observed with the cyanoethyl-compounds, i.e., amide-DAANs **4a** (44.7 µg/mL) and **4b** (69.4 µg/mL) were more soluble than ester-**4c** (20.1 µg/mL).

Molecular lipophilicity is another important drug-like property related to biological potency, ADME profiles, and toxicity risk. The log P values of all compounds were measured in an octanol-water system at pH 7.4 and the data are shown in Table 2. Along with increased solubility, compounds **8d–8g**, **3**, and **4a–4c** displayed better log P values, within a desirable drug-like range of 2.21 to 4.15, showing a reasonable balance between molecular lipophilicity and hydrophilicity. Conversely, compounds **8a–8c** with poorer solubility showed higher log P values (>5).


Subsequently, the metabolic stability of all DAANs, except **8d** (EC_{50(NL4-3)} 24.9 nM), was evaluated in *in vitro* assays.^{16,12} Metabolic stability is one of the most important properties related to oral bioavailability in drug discovery and often is a major liability needing improvement in a lead series. The test compounds were incubated in human liver microsomal (HLM), pooled human liver S9 fraction (S9), and human plasma (HP) assays, and then a LC/MS/MS method was used to quantitate the remaining compound at several time points in parallel with **2** and propranolol, a common reference compound with moderate metabolic stability. Human liver microsomes containing CPY and other metabolizing enzymes are the most frequently used metabolizing material in drug discovery programs. The results are given in Table 2. Ester-compounds **8a**, **8b**, and **4c** were metabolized quite quickly in the HLM assay with short t_{1/2} values of 19, 17, and 18.5 min, respec-

Table 1

Anti-HIV activities of DAANs against wild-type and drug-resistance viral strains.



Cmpd	R ¹	EC ₅₀ nM (FC ^a)			CC ₅₀ nM ^g
		NL4-3 ^b	E138K ^c	E138K+M184I ^c	
8a	-CO ₂ (CH ₂) ₂ OCH ₃	1.52 ± 0.31	4.56 ± 0.35 (2.9)	9.54 ± 3.73 (6.3)	>207
8b	-CO ₂ (CH ₂) ₂ CN	1.11 ± 0.29	2.94 ± 0.40 (2.4)	4.61 ± 1.64 (4.1)	>210
8c	-CONH(CH ₂) ₂ Cl	3.29 ± 1.13	8.50 ± 2.68 (2.6)	17.3 ± 4.54 (5.2)	>206
8d	-CONH(CH ₂) ₃ OH	24.9 ± 8.51	22.8 ± 6.43 (0.9)	ND ^d	>208
8e	-CON(CH ₂) ₂	3.33 ± 1.40	8.69 ± 3.12 (2.6)	16.7 ± 5.34 (5.0)	>223
8f	-CON(CH ₂) ₂ CF ₂	13.4 ± 4.60	ND	ND	>200
8g	-CONHNH ₂	11.6 ± 4.11	ND	ND	>228
3^e	-CONH ₂	1.82 ± 0.57	3.55 ± 0.64 (1.9)	3.55 ± 1.02 (1.9)	>23,600
4a^e	-CONHCH ₃	2.50 ± 0.66	3.87 ± 1.23 (1.6)	5.24 ± 1.62 (2.1)	>9100
4b^f	-CONHEt	6.18 ± 1.41	12.1 ± 3.09 (2.0)	8.61 ± 1.83 (1.4)	105,000
4c^e	-COOCH ₃	1.57 ± 0.50	2.95 ± 0.86 (1.9)	3.64 ± 0.75 (2.3)	>22,700
2		0.44 ± 0.11	3.55 ± 0.85 (8.1)	2.38 ± 0.71 (5.4)	>273

^a Resistance fold change (FC) is the ratio of EC₅₀(resistant viral strain)/EC₅₀(wild-type viral strain).^b HIV-1 with wild-type reverse transcriptase.^c Resistant viral strains.^d Not detected.^e Compound reported in Ref. 8.^f Compound reported in Ref. 12.^g The highest tested concentration of **8a–8g** was 100 ng/mL with at least three experiments (n = 3).**Table 2**Drug-like properties *in vitro* of DAAN compounds.

#	R ¹	aq. sol. (μg/mL) ^a		Log P ^b	t _{1/2} ^c		
		pH 2.0	pH 7.4		HLM (min)	S9 (min)	Human plasma (h)
8a	-CO ₂ (CH ₂) ₂ OCH ₃	1.64	0.77	>5	19	9.5	62
8b	-CO ₂ (CH ₂) ₂ CN	0.94	0.36	>5	17	12	32
8c	-CONH(CH ₂) ₂ Cl	9.53	1.59	>5	61	74	8
8d	-CONH(CH ₂) ₃ OH	19.9	1.01	3.22	ND	ND	ND
8e	-CON(CH ₂) ₂	94.2	1.20	2.21	158	110	12
8f	-CON(CH ₂) ₂ CF ₂	57.6	6.59	3.58	14	14	57
8g	-CONHNH ₂	89.6	2.40	2.31	36	39	8
3	-CONH ₂	2.09	0.63	3.53	90	108	65
4a	-CONHMe	44.7	8.28	2.65	112	96	72
4b	-CONHEt	69.4	1.08	3.63	139	61	91
4c	-COOMe	21.8	1.21	4.15	18.5	7.7	89
2^d		86.8	0.24	>5	108	102	92
Propranolol ^d				139	116	80	

^a General solubility guidelines for human oral absorption are <10 μg/mL, low; 10–60 μg/mL, moderate; >60 μg/mL, high.^b Measured in octanol-water at pH 7.4.^c Metabolic stability incubation assay data from at least two experiments in parallel.^d Reference compound.

tively. They were much less stable than propranolol, demonstrating that the ester groups are easily metabolized by enzymatic catalysis. The amide-DAANs, except **8f**, displayed higher metabolic

stability than ester-DAANs in the same assay. *N*-Substituted carbamoyl-DAANs **8e**, **4a**, and **4b** showed the greatest metabolic stability with t_{1/2} (HLM) values of 158, 112, and 139 min, respectively.

Their data were better than those of the non-*N*-substituted amide **3** ($t_{1/2}$ 90 min) and drug **2** ($t_{1/2}$ 108 min) and comparable with that of propranolol ($t_{1/2}$ 139 min). In addition, *N*-substituted amide-compounds **8c** and **8g** also displayed better metabolic stability ($t_{1/2}$ (HLM) 61 min and 36 min, respectively) than ester-DAANs in the HLM assay. The one exception was amide-**8f** with a *N*-substituted fluorinated four-membered ring in the R^1 side chain. It was quite quickly metabolized in the HLM assay ($t_{1/2}$ (HLM) 14 min), but the tension of the four-membered ring might be a major factor in its intrinsic structural instability. Despite some data fluctuation, similar metabolic stability patterns were found in the HLM and S9 assays. Often, $t_{1/2}$ values in the S9 assay were shorter than those in the HLM assay, because the former system contains a broader set of metabolizing enzymes than liver microsomes. Moreover, all tested compounds displayed reasonable metabolic stability in human plasma¹⁷ with $t_{1/2}$ values of at least 8 h. Thus, the compounds are sufficiently stable in human blood to stay in their original forms in circulation without being metabolized. From the above drug-like property assessments, we conclude that introducing a suitable amide substituent at the R^1 position on the B-ring can favorably increase molecular aqueous solubility at pH 2.0, improve log P values within a desirable range, and enhance molecular metabolic stability in HLM and S9 systems. Although ester- R^1 DAAN compounds showed slightly higher anti-HIV potency and lower resistance FC, on the basis of multiple property improvements, the R^1 substituents follow the rank order: *N*-substituted amides > amides > esters.

In conclusion, the R^1 substituent on the central ring of DAAN compounds is a notable moiety that can be modified to simultaneously improve both drug resistant profiles and drug-like properties. In summary, this study revealed the following findings: (1) the R^1 carbonyl conjugated with the central ring (B-ring) can be from an ester or amide for low nanomolar potency against both HIV-1 wild-type and 2-resistant (E138K, E138K + M184I) viral strains replication; (2) although compounds with ester- R^1 groups exhibited slightly higher potency, amide- R^1 side chains are considered superior because of improved aqueous solubility, lipophilicity, and metabolic stability; (3) short *N*-substituents on the amide- R^1 side chain are favored for better drug-like properties. Overall, amide-DAANs **8e**, **4a**, and **4b** displayed high potency against wild-type virus (EC_{50} (NL4-3) 3.33, 2.50, and 6.16 nM respectively), lower resistance FC (FC_{E138K} 2.6, 1.6, and 2.0, respectively) than drug **2** (FC_{E138K} 8.1), and desirable aqueous solubility (44–95 μ g/mL at pH 2.0), log P

values (2.21–3.63), and moderate metabolic stability in HLM and S9 systems. Therefore, the current study results provide useful information for further structural optimization to develop potential drug candidates with a good balance among high potency, preserved activity against drug-resistant virus, and multiple drug-like properties.

Acknowledgments

This investigation was supported by Grants 30930106 and 81120108022 from the Natural Science Foundation of China (NSFC) – China to L. Xie and NIH – United States Grants awarded to C. H. Chen (AI65310) and K. H. Lee (AI33066). This study was also supported in part by the National Megaprojects of China for Major Infectious Diseases (2013ZX10001-006) and the Taiwan Department of Health, China Medical University Hospital Cancer Research Center of Excellence (DOH100-TD-C-111-005).

A. Supplementary data

Synthesis and structure identification of new compounds, property assessment assays. Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.bmcl.2017.04.068>. These data include MOL files and InChI-Keys of the most important compounds described in this article.

References

1. Vella S, Palmisano L. *Antiviral Res.* 2000;45:1.
2. Das K, Clark Jr AD, Lewi PJ, et al. *J Med Chem.* 2004;47:2550.
3. Janssen PAJ, Lewi PJ, Arnold E, et al. *J Med Chem.* 2005;48:1901.
4. Wainberg MA. *HIV/AIDS.* 2013;5:41.
5. Ripamonti D, Bombana E, Rizzi M. *Expert Rev Anti-Infect Ther.* 2014;12:13.
6. Lambert-Niclot S, Charpentier C, Storto A, et al. *J Antimicrob Chemother.* 2013;68:1237.
7. Qin B, Jiang XK, Lu H, et al. *J Med Chem.* 2010;53:4906.
8. Sun LQ, Zhu L, Qian K, et al. *J Med Chem.* 2012;55:7219.
9. Sun LQ, Qin B, Huang L, et al. *Bioorg Med Chem Lett.* 2012;22:2376.
10. Wu ZY, Liu N, Qin B, et al. *ChemMedChem.* 2014;9:1546.
11. Liu N, Qin B, Sun LQ, et al. *Bioorg Med Chem Lett.* 2014;24:3719.
12. Liu N, Wei L, Huang L, et al. *J Med Chem.* 2016;59:3689.
13. Bourgeois A, Womack S, Newsom D, et al. *HIV Clinician.* 2012;24:12.
14. Rimsky L, Vingerhoets J, Van Eygen V, et al. *J Acquired Immune Defic Syndr.* 2012;59:39.
15. Dang Z, Lai W, Qian K, et al. *J Med Chem.* 2009;52:7887.
16. Di L, Kerns EH, Li SQ, et al. *Int J Pharm.* 2006;317:54.
17. Sofia MJ, Bao D, Chang W, et al. *J Med Chem.* 2010;53:7202.