



Identification of novel triazole inhibitors of Wnt/ β -catenin signaling based on the Niclosamide chemotype

Robert A. Mook Jr.^{a,*}, Jiangbo Wang^a, Xiu-Rong Ren^a, Hailan Piao^a, H. Kim Lyster^b, Wei Chen^{a,*}

^a Department of Medicine, Duke University Medical Center, Durham, NC 27710, United States

^b Department of Surgery, Duke University Medical Center, Durham, NC 27710, United States

ARTICLE INFO

Keywords:

Niclosamide
Wnt signaling inhibitor
Small molecule
Triazole derivatives
 β -Catenin
Cancer

ABSTRACT

Dysregulation of the Wnt signaling pathway is an underlying mechanism in multiple diseases, particularly in cancer. Until recently, identifying agents that target this pathway has been difficult and as a result, no approved drugs exist that specifically target this pathway. We reported previously that the anthelmintic drug Niclosamide inhibits the Wnt/ β -catenin signaling pathway and suppresses colorectal cancer cell growth *in vitro* and *in vivo*. In an effort to build on this finding, we sought to discover new Wnt/ β -catenin inhibitors that expanded the chemotype structural diversity. Here, we asked a specific SAR question unresolved in previous SAR studies of Niclosamide's inhibition of Wnt/ β -catenin signaling to identify a new structural class of Wnt/ β -catenin signaling inhibitors based on a triazole motif. Similar to Niclosamide, we found that the new triazole derivatives internalized Frizzled-1 GFP receptors, inhibited Wnt/ β -catenin signaling in the TOPflash assay and reduced Wnt/ β -catenin target gene levels in CRC cells harboring mutations in the Wnt pathway. Moreover, in pilot SAR studies, we found the Wnt/ β -catenin SAR trends in the anilide region were generally similar between the two chemical classes of inhibitors. Overall, these studies demonstrate the ability to use the SAR of the Niclosamide salicylanilide chemical class to expand the structural diversity of Wnt/ β -catenin inhibitors.

The Wnt signaling pathway is a fundamental growth control pathway that is essential to the regulation of many biological process and is critical to stem cell development and the proper growth and maintenance of tissues.¹ As a result, dysregulation of the pathway is an underlying factor in many diseases, particularly cancer, where mutations are found frequently in various components of the pathway.^{1–3} In colorectal cancer (CRC), more than 93% of tumors had alteration in the pathway, ~80% of which were mutations in adenomatous polyposis coli (APC) or β -catenin that result in hyperactivation of Wnt signaling.^{1,4} Given the importance of the Wnt signaling pathway, there has been an intensive effort for more than three decades to understand the mechanistic details of Wnt signaling and to develop therapies that target the pathway.^{1,5}

Our understanding of the mechanistic complexities of the signaling pathway has grown considerably during this time.^{1,3,6} In the canonical Wnt pathway, a key component is the cytosolic protein β -catenin, whose levels are controlled by a destruction complex consisting of Axin, APC, and two serine-threonine kinases, CK1 α/δ and GSK3 α/β . Signaling is activated by secreted Wnt proteins that bind the seven

transmembrane receptor Frizzled and the single transmembrane receptors LRP5/6 resulting in activation of cytosolic proteins called Dishevelled (Dvl), internalization of the Frizzled receptor, and stabilization and translocation of cytosolic β -catenin proteins into the nucleus. In the nucleus, β -catenin binds and activates the transcription factor LEF/TCF to initiate the transcription of Wnt/ β -catenin target genes. In CRC, most of the mutations observed in the pathway result in an increase in the level of β -catenin protein and activation of the pathway. Thus, agents that decrease the level of β -catenin could be useful therapeutic agents. Overall, the Wnt signaling pathway consists of many intracellular protein-protein interactions for which traditional drug discovery approaches have been difficult.^{5,7} As a result, no drugs have been approved that specifically target the pathway.⁸

Recent efforts, however, have provided opportunities to inhibit the pathway.^{1,9–18} Our contribution to this field was the discovery that the anthelmintic drug Niclosamide (Fig. 1), inhibits Wnt/ β -catenin signaling,¹⁹ a discovery that has since been confirmed by others.^{20–22} Mechanistically, we found that Niclosamide promoted Frizzled internalization, downregulated Dvl and β -catenin protein levels and

Abbreviations: APC, Adenomatous Polyposis Coli; CRC, colorectal cancer; Dvl, Dishevelled; LEF/TCF, Lymphoid enhancer factor/T cell factor; Fzd1, Frizzled1; GFP, green fluorescent protein; SAR, structure-activity relationships

* Corresponding authors.

E-mail addresses: robert.mook@duke.edu (R.A. Mook), w.chen@dm.duke.edu (W. Chen).

<https://doi.org/10.1016/j.bmcl.2018.11.022>

Received 29 June 2018; Received in revised form 2 November 2018; Accepted 10 November 2018

Available online 12 November 2018

0960-894X/ © 2018 Published by Elsevier Ltd.




Fig. 1. Chemical structures of Niclosamide, Niclosamide analogs, and the triazole class of Wnt/ β -catenin inhibitors.

inhibited colorectal cancer cell growth *in vitro* and *in vivo*.^{19,23} Initial SAR studies indicated that Niclosamide inhibition of Wnt signaling appeared unique among structurally-related anthelmintic agents, and that the potency and Wnt/ β -catenin functional response was dependent on small changes in the chemical structure of Niclosamide.²⁴

Niclosamide was approved by the FDA in 1982 for use in humans to treat tapeworm infections²⁵ and has been identified as an essential medicine by the World Health Organization.²⁶ Niclosamide is a salicylic acid derivative that belongs to the salicylanilide class of anthelmintic agents. Similar to other members of the salicylanilide class, Niclosamide has been used widely as an anthelmintic agent in livestock.²⁷ Its anthelmintic activity reported involves uncoupling of oxidative phosphorylation.^{27–30} Niclosamide has protonophore activity and contains an aryl β -hydroxy carbonyl pharmacophore that is present in a large number of natural products and drugs that have a broad range of biological activity.³¹ Thus, it is perhaps not surprising that since Niclosamide's discovery more than 50 years ago, additional biological activity has been discovered that clearly indicate Niclosamide is a multi-functional drug.³¹

Niclosamide's multi-functional activity has generated considerable interest in its use to treat cancer, rheumatoid arthritis, diabetes, and bacterial and viral infections.^{31–38} In addition to our clinical trial in colorectal cancer (NCT02687009), there are now four additional clinical trials of Niclosamide listed in [ClinicalTrials.gov](https://clinicaltrials.gov) in cancer (NCT02519582, NCT02532114, NCT02807805), and in rheumatoid arthritis (NCT03160001). Particular relevant to cancer therapy, Niclosamide inhibits the growth of cancer cells from multiple tumor types and is active against drug resistant cancers.^{23,31,39–41} In addition to its ability to inhibit the Wnt signaling pathway, Niclosamide inhibits other key oncogenic signaling pathways such as mTOR, NF- κ B, Notch and STAT-3 and has effects on metabolism, including activation of AMPK.³¹ Niclosamide is poorly absorbed upon oral administration and is cleared rapidly from systemic distribution.^{23,42,43} Although Niclosamide's pharmacokinetic properties are appropriate for its anthelmintic indication, its pharmacokinetic and solubility properties may limit its use to treat diseases where systemic exposure is required.

Given the need to identify drugs that inhibit Wnt/ β -catenin signaling and the limitations of Niclosamide's properties, we sought to expand the chemical diversity of inhibitors available to us via a classical “scaffold-hopping” strategy.⁴⁴ In our previous SAR studies of Niclosamide, we found that a trifluoromethyl group was a suitable replacement of the nitro group and resulted in an inhibitor of Wnt/ β -catenin signaling (**2**) ($IC_{50} = 0.29 \pm 0.06 \mu M$) with similar potency to Niclosamide ($IC_{50} = 0.34 \pm 0.08 \mu M$) (Fig. 1).^{9,45} However, upon insertion of one carbon atom between the anilide ring and the amide NH group of **2**, the resulting salicylic amide derivative **3** had much poorer inhibitory activity ($IC_{50} = 15.10 \pm 1.40 \mu M$).⁴⁵ To better understand if loss of activity was due to an increase in the steric demands of extending the inhibitor's structure by 1 atom and changing the position of the anilide ring or if the loss of activity resulted from the loss of conjugation of the N–H bond into the aromatic ring and a decrease in the N–H hydrogen-




Fig. 2. Synthesis of triazole derivatives.










bond donating properties, we designed a series of triazole compounds in which the salicylic scaffold of Niclosamide was simultaneously converted into a triazole scaffold (Fig. 1). This change inserts one carbon atom between the anilide ring and the amide NH group and changes the anilide ring position in a similar fashion to the less-active derivative **3** while allowing the N–H bond to maintain conjugation with the aromatic ring.

Based on SAR data indicating Niclosamide derivatives without the 2'-chloro substituent in the anilide inhibited Wnt/ β -catenin signaling⁴⁵ and the ready availability of the commercially available starting materials in our lab, we first prepared triazole **4** in three steps using published procedures (Fig. 2).⁴⁶

We then evaluated the Wnt/ β -catenin inhibitory activity of triazole **4** using assays and protocols previously reported. Upon testing this compound in the Fzd1-GFP internalization and the TOPflash Wnt3A-stimulated β -catenin gene transcription assay,^{9,24,45} we were delighted to find it induced a robust punctate pattern in the Fzd1-GFP assay indicative of internalization of the Fzd1 receptor and inhibited Wnt3A-stimulated β -catenin gene transcription with an IC_{50} of $1.41 \pm 0.34 \mu M$ (Table 1 and Fig. S1 in supplemental information). To enable better comparison of structure and inhibitory activity, we also prepared salicylamide **10**, and tested it and the imide intermediate **11** for Wnt/ β -catenin inhibitory activity. Consistent with previous results obtained with the trifluoromethyl-substituted salicylamide derivative **3**, the nitro-substituted salicylamide derivative **10** was considerably less active in both assays than the similarly substituted triazole **4**. The imide intermediate **11** however, had a different profile. Whereas its activity in the Fzd1-GFP assay was similar to DMSO control, it increased Wnt3A-stimulated signaling in the TOPFlash assay. The reasons for this result are not understood at this time. Based on the data obtained with triazole **4**, we postulate that conjugation of the N–H group and/or its hydrogen bonding donating ability was the important SAR feature for inhibitory activity.

Having identified the inhibitory activity of triazole **4**, we then conducted pilot SAR studies that varied the nitro substituent at the 4-position of the pendent aryl ring to understand if the SAR at this position, a key position in the Niclosamide series, tracked into the new series. Similar to the SAR studies conducted in the Niclosamide series, we selected substituents with a range of electron-donating and electron-withdrawing properties. Consistent with results obtained in the Niclosamide chemotype,^{24,45} a trifluoromethyl substituent was active (**5**) while substitution at the para position with an electron-donating methoxyl substituent⁴⁷ (**6**), an electron-withdrawing fluorine atom (**7**), or a hydrogen atom at this position (**8**) showed a trend toward less activity in both the Fzd1-GFP and the TOPFlash assays, although the magnitude of the difference was not as large as in the Niclosamide series. In the Niclosamide series, we noted earlier that a 2'-chloro substituent increased potency when there was a nitro group at the 4'-position.^{24,45} To test whether the addition of a corresponding 2'-chloro-substituent in the aryl ring of the triazole would produce a similar increase in potency, we prepared triazole **9**. Unfortunately, a similar increase in potency in the triazole series (**4** vs **9**) was not observed by the addition of the chloro-substituent. Overall, however, the trend in inhibitory activity of the triazole derivatives appeared consistent with the inhibitory activity of similarly substituted Niclosamide derivatives.^{24,45}

Table 1
SAR of Wnt/ β -catenin signaling.

Compound Number	Structure	Fzd1-GFP Internalization at 12.5 μ M ^a	Inhibition Wnt/ β -catenin transcription TOPFlash IC50 (μ M) \pm SE ^b
1		5	0.34 \pm 0.07
4		4	1.41 \pm 0.34
5		2	2.88 \pm 0.84
6		0	60% inhibition at 10 μ M, 14% inhibition at 5 μ M
7		1	1.91 \pm 0.60
8		0	4.02 \pm 0.98
9		5	1.84 \pm 0.60
10		0	43% inhibition at 10 μ M, 3% inhibition at 5 μ M
11		0	increased signal, 153% of control at 10 μ M

^a Fzd1-GFP assay: Internalization of Frizzled1-GFP stably expressed in U2OS cells was determined by confocal microscopy. Cells were treated with compounds for 6 h, fixed, and scored visually by the amount of punctate observed versus DMSO control. Punctate similar to control = 0, trace amount of punctate greater than control = 1, moderate = 3, strong = 5. Images in [Supplemental Information](#).

^b Inhibition of Wnt3A-stimulated Wnt/ β -catenin transcription was determined by TOPFlash assay using HEK293 cells stably expressing a TOPFlash luciferase reporter and Renilla luciferase reporter control. Cells were stimulated with Wnt3A-conditioned medium in the presence of DMSO or compounds from 0.04 to 10 μ M for 8 hrs. The TOPFlash reporter activity of Wnt3A conditioned media with DMSO treatment was set as 100%. Data were fit using GraphPad Prism (mean \pm SEM, $n \geq 3$). Percent inhibition at the two highest concentrations tested is reported for weak inhibitors that do not give a full dose response over the inhibitor concentration assayed.

To further confirm the Wnt/ β -catenin inhibitory activity of this new chemical series, compounds with significant activity in the Fzd1-GFP internalization and the Wnt3A-TOPflash assays were evaluated by immunoblot in two CRC cell lines with aberrant Wnt signaling (HCT-116, β -catenin mutant; SW480, APC mutant) for their ability to reduce the levels of Wnt/ β -catenin target gene proteins ([Fig. 3](#)). Consistent with the results of the Frizzled1-GFP and the Wnt3A-stimulated TOPflash assays, Niclosamide (**1**), and triazole derivatives **4**, **5** and **9** each reduced the levels of the Wnt/ β -catenin target genes: β -catenin, Axin2, c-Myc, Survivin and Cyclin D1 relative to controls in both cell lines, thereby demonstrating their ability to inhibit Wnt/ β -catenin signaling in CRC cells with aberrant Wnt pathway signaling.

In general, Niclosamide derivatives with IC₅₀ values in the Wnt3A-stimulated TOPflash assay in the single digit micromolar or less range evaluated to date also typically produce a robust punctate pattern in the

Fzd1-GFP internalization assay. However during the course of our studies, we have identified a few compounds that do not produce a robust response in the Fzd1-GFP internalization assay yet inhibit Wnt/ β -catenin signaling with IC₅₀s of 1–4 μ M in the TOPflash assay.^{9,24,45} Herein, compound **8** ([Table 1](#)) has this profile for which studies in HCT-116 cells confirmed the reduction in Wnt/ β -catenin target genes ([Supplemental Fig. S2](#)). The mechanistic underpinnings to account for this observation are not clear at present. Given Niclosamide has multifunctional activity, one explanation may be that these derivatives have a different mechanism or a different selectivity profile.

Overall, these studies provide insight into the SAR of the amide region that impact potency and demonstrate the ability to extend the SAR findings of the Niclosamide chemotype into a novel triazole class of Wnt/ β -catenin inhibitors. Whether the antiviral and antibiotic activity, inhibition of other signaling pathways and the effects on blood glucose




Fig. 3. Reduction of Wnt/ β -catenin target genes by Niclosamide and triazole 4, 5 and 9. Cells were treated with DMSO or compounds (5 μ M) in DMSO for 18 h, the cytosolic fraction and the whole cell lysates were probed with antibodies to the proteins indicated. β -Actin was used as a loading control. A. Western blot images B. Quantification of the protein levels of Western blots in A normalized to the β -Actin loading control and graphed as a percentage of protein level in the DMSO treated control cells.

and fatty liver also extend into this chemotype, given the structural relationship of these triazole derivatives to Niclosamide, is an open question. The compounds herein are less potent inhibitors of Wnt/ β -catenin signaling than Niclosamide. Additional studies are underway to further define the Wnt/ β -catenin inhibitory SAR to improve potency, the activity against pathways modulated by Niclosamide, and the pharmacokinetic properties of the series to support their evaluation in

CRC tumor models and models of other diseases for which Niclosamide has significant biological activity.³¹

Acknowledgements

This work was funded in part by 5 R01 CA172570 (WC), BC123280 (WC), and Clinical Oncology Research Center Development Grant

5K12-CA100639-08 (RAM). Wei Chen is a V foundation Scholar and an American Cancer Society Research Scholar. NMR instrumentation in the Duke NMR Spectroscopy Center was funded by the NIH, NSF, North Carolina Biotechnology Center and Duke University. The authors gratefully acknowledge this support and the support of Professor Eric Toone and David Gooden of the Duke Small Molecule Synthesis Facility.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.bmcl.2018.11.022>.

References

- Nusse R, Clevers H. *Cell*. 2017;169:985.
- Zhan T, Rindtorff N, Boutros M. *Oncogene*. 2017;36:1461.
- Clevers H, Nusse R. *Cell*. 2012;149:1192.
- Cancer Genome Atlas Network. *Nature*. 2012;487:330.
- Barker N, Clevers H. *Nat Rev Drug Discovery*. 2006;5:997.
- Chen W, ten Berge D, Brown J, et al. *Science*. 2003;301:1391.
- Meireles LMC, Mustata G. *Curr Top Med Chem*. 2011;11:248.
- Sebio A, Kahn M, Lenz H-J. *Expert Opin Therap Targets*. 2014;18:611.
- Mook Jr RA, Ren XR, Wang J, et al. *Bioorg Med Chem*. 1804;2017:25.
- Tabatabai R, Linhares Y, Bolos D, Mita M, Mita A. *Target Oncol*. 2017;12:623.
- Tran FH, Zheng JJ. *Protein Sci*. 2017;26:650.
- Yang K, Wang X, Zhang H, et al. *Lab Invest*. 2016;96:116.
- Lum L, Chen C. *Curr Med Chem*. 2015;22:4091.
- Li Y, Oliver PG, Lu W, et al. *Cancer Lett*. 2017;389:41.
- Li B, Orton D, Neitzel LR, Astudillo L, Shen C, Long J, Chen X, Kirkbride KC, Doundoulakis T, Guerra ML, Zaias J, Fei DL, Rodriguez-Blanco J, Thorne C, Wang Z, Jin K, Nguyen DM, Sands LR, Marchetti F, Abreu MT, Cobb MH, Capobianco AJ, Lee E, Robbins DJ. *Sci Signal*. 2017;10 eaak9916.
- You L, Zhang C, Yarravarapu N, et al. *Bioorg Med Chem Lett*. 2016;26:5891.
- Shono T, Ishikawa N, Toume K, et al. *Bioorg Med Chem Lett*. 2015;25:2735.
- Lee JH, Kim MA, Park S, Cho SH, Yun E, O YS, Kim J, Goo JI, Yun MY, Choi Y, Oh S, Song GY. *Bioorg Med Chem Lett*. 2016;26:3529.
- Chen MY, Wang JB, Lu JY, et al. *Biochemistry*. 2009;48:10267.
- Sack U, Walther W, Scudiero D, et al. *J Natl Cancer Inst*. 2011;103:1018.
- Lu W, Lin C, Roberts MJ, Waud WR, Piazza GA, Li Y. *PLoS One*. 2011;6:e29290.
- Arend RC, Londono-Joshi AI, Samant RS, et al. *Gynecol Oncol*. 2014;134:112.
- Osada T, Chen MY, Yang XY, et al. *Cancer Res*. 2011;71:4172.
- Mook Jr RA, Chen M, Lu J, Barak LS, Lyerly HK, Chen W. *Bioorg Med Chem Lett*. 2013;23:2187.
- Pearson RD, Hewlett EL. *Ann Intern Med*. 1985;102:550.
- WHO. *The selection and use of essential medicines*. Geneva: World Health Organization; 2007.
- Swan GEJS. *Afr Vet Assoc*. 1999;70:61.
- Weinbach EC, Garbus J. *Nature*. 1969;221:1016.
- Williamson RL, Metcalf RL. *Science*. 1967;158:1694.
- Frayha GJ, Smyth JD, Gobert JG, Savel J. *Gen Pharmacol*. 1997;28:273.
- Chen W, Mook Jr RA, Premont RT, Wang J. *Cell Signal*. 2018;41:89.
- Tao H, Zhang Y, Zeng X, Shulman GI, Jin S. *Nat Med*. 2014;20:1263.
- Rajamuthiah R, Fuchs BB, Conery AL, et al. *PLoS One*. 2015;10:e0124595.
- Gooyit M, Janda KD. *Sci Rep*. 2016;6:33642.
- Xu M, Lee EM, Wen Z, et al. *Nat Med*. 2016;22:1101.
- Wang YM, Lu JW, Lin CC, et al. *Antiviral Res*. 2016;135:81.
- Huang M, Zeng S, Qiu Q, et al. *Int Immunopharmacol*. 2016;31:45.
- Liang L, Huang M, Xiao Y, et al. *Inflamm Res*. 2015;64:225.
- Li Y, Li PK, Roberts MJ, Arend RC, Samant RS, Buchsbaum DJ. *Cancer Lett*. 2014;349:8.
- Liu C, Lou W, Zhu Y, et al. *Clin Cancer Res*. 2014;20:3198.
- Li R, Hu Z, Sun SY, et al. *Mol Cancer Ther*. 2013;12:2200.
- Andrews P, Thyssen J, Lorke D. *Pharmac Ther*. 1983;19:245.
- Chang Y-W, Yeh T-K, Lin K-T, et al. *Yaowu Shipin Fenxi*. 2006;14:329.
- Langdon SR, Ertl P, Brown N. *Mol Inf*. 2010;29:366.
- Mook Jr RA, Wang J, Ren XR, et al. *Bioorg Med Chem*. 2015;23:5829.
- Catarzi D, Cecchi L, Colotta V, et al. *J Med Chem*. 1995;38:2196.
- Thomas G. *Medicinal chemistry: an introduction*. West Sussex, England: John Wiley & Sons; 2007:116.