



## Functional developmental changes underlying response inhibition and error-detection processes

Wouter Braet<sup>a,b,\*</sup>, Katherine A. Johnson<sup>a</sup>, Claire T. Tobin<sup>a</sup>, Ruth Acheson<sup>a</sup>, Mark A. Bellgrove<sup>c</sup>, Ian H. Robertson<sup>a</sup>, Hugh Garavan<sup>a</sup>

<sup>a</sup> School of Psychology and Trinity College Institute of Neuroscience, Trinity College Dublin, Dublin 2, Ireland

<sup>b</sup> Laboratory of Experimental Psychology, University of Leuven, Belgium

<sup>c</sup> The University of Queensland, School of Psychology and Queensland Brain Institute, Brisbane, Australia

### ARTICLE INFO

#### Article history:

Received 30 October 2008

Received in revised form 22 July 2009

Accepted 24 July 2009

Available online 3 August 2009

#### Keywords:

Response inhibition

fMRI

Developmental

Error awareness

### ABSTRACT

This study examined the developmental trajectories associated with response inhibition and error processing as exemplar executive processes. We present fMRI data showing developmental changes to the functional networks underlying response inhibition and error-monitoring, comparing activation between adults and young adolescents performing the sustained attention to response task (SART). During successful inhibitions, we observed greater activation for the young adolescents than for the adults, in a widely distributed network including frontal, parietal and medial regions. When inhibition failed, however, adults showed increased activation compared to young adolescents in a number of regions, including bilateral parahippocampal gyrus, left and right lingual gyri, the right insula, and cerebellar regions. These differences largely remained even when the two groups were matched for performance, suggesting that performance differences are unlikely to be the driving factor behind these developmental differences. Instead, the neurodevelopmental trajectory of these important executive functions may reveal the basis for the immature executive functioning of the young adolescent.

© 2009 Elsevier Ltd. All rights reserved.

### 1. Introduction

Response inhibition, or the ability to withhold a prepotent response, is a key aspect of executive functioning, allowing us to operate in complex situations with competing stimuli that demand our attention. The dynamic control of behaviour in these situations involves a number of distinct functions, including task-monitoring, error-detection and compensatory changes to behaviour after an error has been detected. Areas involved in these functions include a number of prefrontal regions such as the anterior cingulate cortex (ACC) and dorsolateral prefrontal cortex (DLPFC) (Dehaene, Posner, & Tucker, 1994; Garavan, Ross, Murphy, Roche, & Stein, 2002). Response inhibition may be one of the first executive functions to develop, with early forms of error-monitoring, self-regulation of behaviour and response inhibition having been observed in children as young as 4 years old (Jones, Rothbart, & Posner, 2003). Whereas some cognitive processes, such as attentional orienting, develop early and show little change between childhood and

adulthood, the ongoing development of executive functions, such as response inhibition, differentiates childhood cognition from that of adulthood (Denckla, 1996). This development may occur in parallel with sequential maturation of the frontal lobes (including myelinisation, synaptic pruning and reorganisation, dendritic and axonal arborisation), which is not completed until early adulthood (Anderson, Anderson, Northam, Jacobs, & Catroppa, 2001; Gogtay et al., 2004).

Response inhibition is often measured using Go/No-Go paradigms where participants are asked to make a simple response (e.g., pressing a button) to the majority of stimuli, but to withhold this response to an infrequent (and usually unpredictable) target (No-Go) stimulus. In the current study we used the sustained attention to response test (SART), a Go/No-Go task which has previously been used to investigate 'attentional slips' in everyday life (Robertson, Manly, Andrade, Baddeley, & Yiend, 1997), as well as to investigate response inhibition in a number of clinical conditions, such as traumatic brain injury (TBI) (McAvinue, O'Keeffe, McMackin, & Robertson, 2005; but also see Whyte, Grieb-Neff, Gantz, & Polansky, 2006), attention deficit hyperactivity disorder (ADHD) (Johnson et al., 2007b), high-functioning autism (Johnson et al., 2007b), schizophrenia (Chan, Chen, Cheung, Chen, & Cheung, 2004), and sleep disorders (Fronczek, Middelkoop, van Dijk, & Lammers, 2006).

\* Corresponding author at: Laboratory of Experimental Psychology, University of Leuven, Tiensestraat 102, B 3000 Leuven, Belgium. Tel.: +32 016 32 59 57; fax: +32 016 32 60 99.

E-mail address: [Wouter.braet@gmail.com](mailto:Wouter.braet@gmail.com) (W. Braet).

Successful inhibition on the SART task elicits the classic No-Go N2/P3 complex which is a robust electrophysiological marker of inhibitory control (O'Connell et al., 2008).

In an fMRI-study of healthy adults (Fassbender et al., 2004), successful inhibitions on the SART increased activity in the right ventral frontal cortex, left DLPFC, the right inferior parietal lobe (IPL), as well as the left putamen, consistent with other studies showing a network of prefrontal and parietal regions involved in response inhibition (e.g., Garavan, Ross, & Stein, 1999; Rubia, Smith, Taylor, & Brammer, 2007; Stevens, Kiehl, Pearlson, & Calhoun, 2007). For commission errors (responding to the No-Go stimulus), which may involve processes related to (late attempts at) response inhibition and error-detection, increased activation was observed in the anterior (ACC) and posterior cingulate (PCC) cortex, bilateral inferior frontal gyri (IFG) and insulae, and (predominantly left) inferior parietal regions.

In the current study, we compared SART performance of adults and young adolescents to understand further the development of inhibitory control and its associated neural networks. To avoid performance differences between the two groups confounding the activation maps (see Murphy & Garavan, 2004), we used an event-related design which allows direct comparisons of successful and unsuccessful inhibitions. Although more recent studies (Rubia et al., 2007, 2006; also see Bunge, Dudukovic, Thomason, Vaidya, & Gabrieli, 2002) comparing response inhibition between adults and adolescents suggest increased activation of frontal and prefrontal regions in adults, earlier studies (Booth et al., 2003; Casey et al., 1997; Durston et al., 2002) showed the reverse pattern. A recent longitudinal study (Durston et al., 2006) showed activity in the right inferior frontal gyrus increased with age, whereas activity in most other regions (including precentral, superior frontal, superior temporal, and posterior cingulate gyri) decreased with age. It has been suggested (Rubia et al., 2006) that differences between these studies may be due to factors such as sample size (which were relatively small in the earlier studies), or differences in age-range (under 12 in the earlier studies, compared with adolescents aged between 10–17 years in the studies by Rubia et al.), or because some earlier studies used block rather than event-related designs (Booth et al., 2003). Differences between the specific paradigms used, however, especially when these may engage processes other than response inhibition, might also be a confounding factor. For example, the tasks used by Rubia et al. (2007, 2006) and by Bunge et al. (2002) also have a strong spatial component, in addition to engaging processes related to response selection, as participants had to choose between different responses depending on the spatial configuration of the stimuli. The SART, however, only requires participants to make a single type of response to unambiguous and pre-specified Go stimuli (all digits bar “3;” see task description below) or to withhold this response and thus may avoid contamination from other cognitive processes (see also the paradigm used by Durston et al., 2006, 2002).

The studies by Rubia et al. (2006, 2007) also differ in one other important aspect from the earlier work: whether or not the adult and adolescent groups performed tasks that were equated for difficulty in terms of stimulus properties (e.g., similar timings) or in terms of performance (e.g., by using an adaptive staircase mechanism that ensured performance at a criterion level). There are good arguments for using either method. The first, in which task properties are held constant, avoids confounding age/ability with differences in the objective difficulty-level of a task. The second method ensures that everyone performs at the same level of performance, thus avoiding confounding age/ability with processes secondary to performance such as error-related frustration. The choice of method may be consequential with one hypothesis being that equating task-difficulty may reveal greater levels of activation in young adolescents (on account of their poorer inhibitory abilities




Fig. 1. Time course of a single trial (bottom-left to top-right).

requiring more neural resources be deployed than are required by adults). On the other hand, equating performance levels may reveal greater levels of activation in adults (their superior inhibitory abilities may result in more neural resources being accessible for adults relative to those that are available for young adolescents). To disentangle the influence of ability and performance from inherent age-related differences, the present study employed a task at a set level of difficulty but also included a performance-matched analysis of a subset of adult and young adolescent participants.

In addition to activation during successful inhibitions, we also examined activation associated with unsuccessful inhibitions (commission errors). Here activation can be assumed to include processes related to error-detection and increased top-down attentional operation<sup>1</sup> (Garavan et al., 2002). Rubia et al. (2007) found increased activation of rostral ACC for adults compared to children/adolescents in a design that matched performance between the two age-groups, and this has also been observed without performance matching (Velanova, Wheeler, & Luna, 2008). To our knowledge the studies by Rubia et al. and by Velanova et al. are the only other developmental imaging studies to examine error activity during response inhibition. Accordingly, we predicted a similar pattern of activation differences between the adults and young adolescents in the current study.

## 2. Materials and methods

### 2.1. Participants

Forty right-handed participants, 20 young adolescents (all males) in the age-range 10–14 (mean age 12.4, SD 1.4) and 20 adults (15 males) in the age-range 23–35 (mean age 26.8, SD 4) were involved in this study. A subset of this sample with matched performance was subsequently selected to assess the impact of performance on activation patterns. The subsample consisted of 12 young adolescents (all male, mean age 12.7, SD 1.3) and 12 adults (including three females, mean age 25.5, SD 3.5). To ensure that this gender imbalance in the full sample did not unduly affect the results, we also report analyses using only the data from male participants.

The study was approved by the local ethics committee in accordance with the Declaration of Helsinki. All participants were reported (by themselves and/or a parent) to be free of neurological and psychiatric disorders and head trauma, and gave

<sup>1</sup> Activation changes during error trials may be related to other processes, including late attempts at inhibition, momentary lapses of attention, or increased maintenance of the go-rule. These additional processes are likely to fluctuate across trials (i.e. for any given error trial, the error might be caused by a temporary distraction, or by overreliance on automatic go-responding, or by a motor output error), and as such are conceptualized as additional noise in the MRI-signal corresponding to commission errors.




Fig. 2. Performance measures.

written consent to participate (for the young adolescents, written consent was also given by a parent).

## 2.2. Sustained attention to response task (SART)

Participants performed the random SART, in which numbers from 1 to 9 are presented in a random (i.e. non-sequential) order. In each trial (see Fig. 1) a single digit appeared on the screen for 313 ms; a mask was then presented for 125 ms, after which a response cue (a boldened cross) appeared for 63 ms, followed by a second mask for 375 ms and a fixation cross for 563 ms. The total inter-stimulus interval was 1439 ms (digit onset to digit onset). Participants were instructed to respond, using a button press, to every digit (Go-trial) except '3' (No-Go-trial). They were asked to respond when the response cue appeared on screen 125 ms after the digit was extinguished, or 438 ms from the start of the trial. The response cue was used to limit unwanted performance differences in response variability. The task was presented using E-prime (Psychology Software Tools, Pittsburgh, USA), in a single block of 450 trials (of which 50 were No-Go-trials), which included two 30 s breaks after 150 and 300 trials. The total duration of the task was approximately 12 min.

## 2.3. MRI data acquisition

All scanning was conducted on a Philips Intera Achieva 3.0 Tesla MR system. Each scanning sequence began with a reference scan to resolve sensitivity variations. A parallel Sensitivity Encoding (SENSE) approach (Pruessmann, Weiger, Scheidegger, & Boesiger, 1999) with a reduction factor of 2 was utilised for all T1-weighted image acquisitions. 180 high-resolution T1-weighted anatomic MPRAGE axial images (FOV 230 mm, thickness 0.9 mm, voxel size  $0.9 \times 0.9 \times 0.9$ ) were then acquired (total duration 325 s), to allow subsequent activation localization and spatial normalization.

Functional data were collected using a T2-weighted echo-planar imaging (EPI) sequence that acquired 32 non-contiguous (10% gap) 3.5 mm axial slices covering the entire brain (TE = 35 ms, TR = 2000 ms, FOV 224 mm,  $64 \text{ mm} \times 64 \text{ mm}$  matrix size in Fourier space). The functional scans had a total duration of 730 s.

## 2.4. fMRI-analysis

The data were analysed using AFNI (Cox, 1996; <http://afni.nimh.gov>). Images were corrected for motion (using a least-squares alignment allowing translations and rotations), and activation outside the brain was removed. Separate impulse response functions (IRFs) were estimated<sup>2</sup> for successful inhibitions and commission errors using deconvolution techniques. Gamma-variate functions were fit, voxelwise, to these IRFs using a non-linear regression programme. A percentage signal-change score (%SC) was calculated by dividing the area under the curve of these functions by the area under the baseline which, in this case, reflects tonic ongoing processes involved in Go-trial responses. Individual %SC maps were then

spatially blurred using a 3 mm rms isotropic Gaussian kernel, and transformed into MNI space using the MNI (Montréal Neurological Institute) 152-brain template.

For both commission errors and successful inhibitions, group activation maps were then determined using one-sample *t*-tests against 0 (i.e. against the null hypothesis of no change in activation compared with the baseline). Significant voxels passed a voxelwise statistical threshold ( $t(19) = 3.88, p \leq .001$ ), and were required to be part of a cluster of significant voxels with a minimum volume of 141  $\mu\text{l}$ . This minimum cluster-size was determined using Monte-Carlo simulations, with a probability of .05 (corrected) of a cluster surviving due to chance. Separate maps were generated for the adults and young adolescents, and these were subsequently combined into a single map which contained every voxel that survived thresholding in either of the two groups. These resulting maps (one for successful inhibitions, and one for commission errors) were then used as regions of interest (ROI) to extract mean activation values for each region for every participant. These data were used for independent group *t*-tests. The subsample of performance-matched participants was investigated using these same ROIs.

In addition to this, we also performed a direct voxelwise comparison between the two groups (using the same ROIs that were identified for successful inhibitions and for commission errors), to rule out a potential confound between activation amplitude and activation extent. Within these regions, significant voxels passed a voxelwise statistical threshold ( $t(38) = 3.57, p \leq .001$ ). Minimum cluster-size (to control for multiple comparisons, with a probability of .05 of a cluster surviving due to chance) was 73  $\mu\text{l}$  for successful inhibitions, and 69  $\mu\text{l}$  for commission errors. The resulting clusters identify subregions of the ROIs that differ only in activation amplitude, between the adults and the young adolescents.

## 2.5. Analysis of behavioural data

Errors of commission (responses made on the No-Go digit 3, which indicate failure of inhibitory processes) and omission (non-responses on the Go-trials, believed to reflect temporary lapses in attention) were calculated for each participant. Independent groups *t*-tests were used to investigate differences between the two groups with the alpha level set at .05. Variability of response times has been previously suggested as an index of top-down executive control (West, Murphy, Armilio, Craik, & Stuss, 2002), and ICV-scores have been previously associated with inhibitory success as well as frontal, parietal and thalamic activation during response inhibition (Bellgrove, Hester, & Garavan, 2004). We therefore also compared the mean and Intra-individual coefficient of variability (ICV) (SD Go-RT/mean Go-RT, see Stuss, Murphy, Binns, & Alexander, 2003) of the response times (RTs) on the Go-trials between the groups.

## 3. Results

### 3.1. Behavioural performance

Adults made significantly fewer errors of omission ( $t(38) = 2.25, p = .03$ ) and significantly fewer errors of commission ( $t(38) = 3.74, p = .001$ ) compared with young adolescents (see Fig. 2). There was no significant difference in RTs between the adults and young adolescents on Go-trials ( $t(38) = .81, p = .42$ ) or on commission errors ( $t(38) = 0.75, p = .46$ ). Both adults and young adolescents had faster

<sup>2</sup> Separate IRF functions were calculated for each participant, which reduces the effects of (limited) differences in the shape of the BOLD-signal that may exist between the two groups. For example, it is likely that response inhibition might be faster for adults compared to young adolescents, which might lead to a shorter interval between the stimulus and the onset of the BOLD signal. With the present method, any such differences in e.g. the onset of the BOLD signal are unlikely to affect the activation maps.

**Table 1**  
Brain regions activated during successful inhibitions; group differences in region-level activation (direction: ya, young adolescents, a, adults). The *p*-values, from left to right, refer to: activation differences (full sample, performance-matched sample). The columns on the right show clusters within these regions that differ voxelwise in activation amplitude.

Region		Centre of mass (MNI)								Amplitude differences			
		BA	Vol (μl)	X	Y	Z	p (20 vs 20)	p (12 vs 12)	Direction	Vol (μl)	X	Y	Z
Frontal lobes													
Middle frontal gyrus	R	9	11,618	35	37	36	.016	n.s.	ya > a	212	−35	30	36
Middle frontal gyrus	L	9	4,080	−34	41	34	.023	n.s.	ya > a				
Superior frontal gyrus	L	6/8	313	−21	22	59	n.s.	n.s.					
Precentral gyrus	L	6	298	−22	−17	58	n.s.	n.s.		90	−25	9	54
Superior frontal gyrus	L	6	256	−24	9	56	.001	n.s.	a > ya				
Inferior frontal gyrus	R	9/6	240	36	9	27	<.001	.005	ya > a				
Inferior frontal gyrus	L	10	209	−42	48	−2	<.001	.007	ya > a				
Superior frontal gyrus	L	8	153	−9	36	60	<.001	.005	ya > a				
Parietal lobes													
Inferior parietal lobule	R	40	18,028	51	−48	32	.01	n.s.	ya > a	576	50	−38	37
Precuneus and cuneus	R	7	2,510	9	−74	37	.007	.015	ya > a				
Inferior parietal lobule	L	40	2,023	−57	−47	26	.007	.028	ya > a				
Cuneus	L	30	1,298	−5	−62	4	.026	.025	ya > a	184	0	−63	7
Superior parietal lobe	R	7	160	33	−69	49	n.s.	n.s.					
Temporal lobes													
Middle temporal gyrus	R	20	871	49	−23	−15	.026	n.s.	ya > a	88	44	−25	−8
Medial brain regions													
Anterior cingulate	BI	6/24	31,800	6	9	44	<.001	.015	ya > a	2359	2	34	26
Insula	R	13/47	13,210	36	14	0	.011	.022	ya > a	154	27	15	−6
Insula	L	47	8,454	−33	18	−1	.002	.043	ya > a	245	4	16	30
Posterior cingulate	BI	3	3,998	3	−29	27	.003	.045	ya > a				
Brainstem	R	/	426	4	−28	−29	n.s.	n.s.					
Lentiform nucleus	L	/	263	−26	−4	13	.001	.004	ya > a				
Thalamus	R	/	257	15	−7	9	n.s.	n.s.					
Thalamus	R	/	184	8	−24	4	n.s.	n.s.					
Caudate	L	/	172	−13	10	3	.018	n.s.	ya > a				

RTs when making commission errors compared with RTs on Go-trials ( $t(19)=7.08$ ,  $p<.001$ ; and  $t(19)=5.88$ ,  $p<.001$ , respectively). Young adolescents showed higher ICVs on Go-trials compared to adults ( $t(38)=2.84$ ,  $p=.007$ ). Response variability correlated with the number of omission errors for young adolescents ( $r=.66$ ,  $p=.002$ ) but not for adults ( $r=.36$ ,  $p=.119$ ), and correlated with the number of commission errors for both young adolescents ( $r=.79$ ,  $p<.001$ ) and adults ( $r=.53$ ,  $p=.016$ ).

### 3.2. fMRI-analysis: successful inhibitions

Table 1 lists the areas that showed significant activity when participants successfully inhibited their response on No-Go-trials. Seventeen<sup>3</sup> of these 23 regions showed significant differences between the two groups, with young adolescents demonstrating higher activation in the right ( $p=.016$ ) and left ( $p=.023$ ) middle frontal gyri, right ( $p<.001$ ) and left ( $p<.001$ ) inferior frontal gyri, left posterior superior frontal gyrus ( $p<.001$ ), right ( $p=.01$ ) and left ( $p=.007$ ) inferior parietal cortex, right ( $p=.007$ ) and left ( $p=.026$ ) cuneus, right middle temporal gyrus ( $p=.026$ ), bilateral anterior ( $p<.001$ ) and posterior ( $p=.003$ ) cingulates, right ( $p=.011$ ) and left ( $p=.002$ ) insulae, left lentiform nucleus ( $p=.001$ ) and left caudate ( $p=.018$ ). Adults showed a significantly larger increase in only one region, the anterior part of the left superior frontal gyrus ( $p=.001$ ; see Fig. 3).

Nine of these regions contained clusters that showed significant voxelwise differences in activation amplitude, between the adults and the young adolescents: young adolescents demon-

strated higher activation amplitude in the left middle frontal gyrus, right inferior frontal gyrus, right inferior parietal cortex, right cuneus, right middle temporal gyrus, bilateral ACC and PCC, and left insula; adults showed higher activation amplitude in the anterior part of the left superior frontal gyrus (all  $p<.001$ ). In the remaining eight regions which showed activation differences for the region but no significant clusters of voxelwise differences, group effects may be driven by differences in activation extent (i.e. the young adolescents activating more voxels in the ROI, but with a similar amplitude as was observed for adults). It should however be noted that this is a tentative conclusion given that activation and amplitude and activation extent become confounded when one spatially blurs data to enable group averages and group comparisons.

The adults showed a significant correlation between ICV-scores and activation in the medial region of the left superior frontal gyrus ( $r=.47$ ,  $p=.035$ ), while in young adolescents ICV-scores correlated positively with activation in the left lentiform nucleus ( $r=.52$ ,  $p=.02$ ) and negatively with (the anterior region of) the left superior frontal gyrus ( $r=–.55$ ,  $p=.012$ ).


### 3.3. fMRI-analysis: commission errors

Table 2 lists the areas of significant activation when participants made commissions errors (also see Fig. 4). Of these 22 regions, nine<sup>4</sup> showed a significant difference between the adults and the young adolescents, with adults showing higher activation than young adolescents for all. This pattern was observed in the bilateral cerebellar culmen ( $p<.001$ ), right parahippocampal gyrus ( $p=.006$ ), the right

<sup>3</sup> If male participants only are compared (to avoid gender-bias), activation differences in the right insula ( $t(33)=2$ ,  $p=.053$ ), right middle frontal gyrus ( $t(33)=1.68$ ,  $p=.103$ ), left middle frontal gyrus ( $t(33)=1.88$ ,  $p=.069$ ), right middle temporal gyrus ( $t(33)=1.75$ ,  $p=.09$ ), and left caudate ( $t(33)=1.5$ ,  $p=.14$ ) are no longer significant. All remaining differences remain significant.

<sup>4</sup> If we only compare the male participants (to avoid gender-bias), the activation difference in the right parahippocampal gyrus is no longer significant ( $t(33)=2.02$ ,  $p=.052$ ). All other effects remain significant.






**Fig. 3.** Activation maps for successful inhibitions for the superior frontal gyrus (a), AC and PC (b), insula (c), and both parietal lobes (d); red: children > adults; green: adults > children; orange: no difference.

**Table 2**

Brain regions activated during commission errors; group differences in region-level activation (direction: ya = young adolescents, a = adults). The *p*-values, from left to right, refer to: activation differences (full sample, performance-matched sample). The columns on the right show clusters within these regions that differ voxelwise in activation amplitude.

Region	Centre of mass (MNI)									Amplitude differences			
	HS	BA	Vol (μl)	X	Y	Z	<i>p</i> (20 vs 20)	<i>p</i> (12 vs 12)	Direction	Vol (μl)	X	Y	Z
Frontal lobes													
Middle frontal gyrus	L	9	2,534	−33	42	33	n.s.	n.s.					
Superior frontal gyrus	R	9	1,825	29	41	37	n.s.	n.s.					
Superior frontal gyrus	L	6	186	0	24	63	n.s.	n.s.					
Temporal lobes													
Middle temporal gyrus	L	21	467	−54	−52	−1	n.s.	n.s.					
Middle temporal gyrus	R	20	253	50	−25	−18	n.s.	n.s.					
Parietal lobes													
Inferior parietal lobule	R	40	2,249	55	−48	27	n.s.	n.s.					
Inferior parietal lobule	L	40	2,214	−59	−44	28	n.s.	n.s.					
Precuneus	L	31	148	−18	−40	40	n.s.	n.s.					
Precuneus	R	7	147	12	−79	39	n.s.	n.s.					
Medial brain regions													
Anterior cingulate	BI	32/24	20,800	1	15	40	n.s.	n.s.					
Insula	R	13/47	14,207	41	12	−1	n.s.	n.s.		84	40	11	−22
Insula	L	13/47	13,974	−36	13	−2	.016	n.s.	a > ya	83	−30	−2	−14
Culmen	BI	/	4,698	−2	−32	−13	<.001	.006	a > ya	617	0	−35	−7
Posterior cingulate	BI	23	2,262	0	−25	28	n.s.	n.s.					
Caudate/thalamus	R	/	1,333	9	4	1	.016	n.s.	a > ya				
Cingulate gyrus	R	31	229	12	−34	36	n.s.	n.s.					
Parahippocampal gyrus	R	35/36	173	21	−37	−11	.006	n.s.	a > ya				
Clastrum	R	13	151	39	−16	0	<.001	.008	a > ya	70	37	−15	1
Occipital lobes													
Lingual gyrus	L	19	642	−20	−66	−11	<.001	<.001	a > ya	459	−22	−65	−6
Fusiform gyrus	R	19	613	20	−56	−13	.002	n.s.	a > ya	137	19	−49	−10
Lingual gyrus	R	19	307	22	−70	−6	.001	n.s.	a > ya				
Lingual gyrus/cuneus	L	18	217	−14	−74	−3	<.001	.011	a > ya				



**Fig. 4.** Activation maps for commission errors: vermis and lingual gyri (a), insulae (b) and uncus/amygdala (c); green indicates higher activation for adults, red indicates no difference.

lingual gyrus and two regions in the left lingual gyrus (all  $p < .001$ ), the left insula ( $p = .016$ ), the right claustrum ( $p < .001$ ), and also in the right caudate ( $p = .016$ ) and fusiform gyrus ( $p = .002$ ).

Of these regions, six contained clusters of voxels that showed a significant difference in activation amplitude: the right and left insulae, bilateral culmen, right claustrum, left lingual gyrus, as well as the right fusiform gyrus (all  $p < .001$ ). In all these regions, activation amplitude was higher for the adults, compared to the young adolescents.

Young adolescents showed significant negative correlations between response variability (ICV-scores) and %SC with a number of regions, including right ( $r = -.46$ ,  $p = .039$ ) and left ( $r = -.46$ ,  $p = .042$ ) insulae, right ( $r = -.46$ ,  $p = .039$ ) and left ( $r = -.48$ ,  $p = .031$ ) parietal cortices, right superior frontal gyrus ( $r = -.47$ ,  $p = .036$ ), and right precuneus ( $r = -.46$ ,  $p = .042$ ). A positive correlation was observed in the right claustrum ( $r = .45$ ,  $p = .048$ ). There were no significant correlations between ICV-scores and activation measures for the adults.

### 3.4. Performance-matched subsample

#### 3.4.1. Behavioural performance


The subsample was selected by excluding the adults whose number of commission errors was lower than one S.E.M. below the mean number of errors, and a corresponding number of young

adolescents who made the most commission errors. In the subsample, there were no significant differences between the young adolescents and adults on any of the performance variables, except young adolescents showed slower RTs when making errors on the No-Go-trials ( $t(22) = 2.7$ ,  $p < .001$ ) and a trend for slower RTs on Go-trials ( $t(22) = 1.8$ ,  $p = .086$ ) (all other  $t$ -values  $\leq 1.4$ ) (See Fig. 5). We observed no significant correlations between performance measures in either the young adolescents or the adults of the subsample.

#### 3.4.2. fMRI-analysis and functional correlations: successful inhibitions

In the subsample matched for performance, young adolescents had significantly greater activation compared to adults in eleven regions: right ( $p = .005$ ) and left ( $p = .007$ ) inferior frontal gyri, posterior left superior frontal gyrus ( $p = .005$ ), left inferior parietal cortex ( $p = .028$ ), right ( $p = .015$ ) and left ( $p = .025$ ) cuneus, right ( $p = .022$ ), and left ( $p = .043$ ) insulae, left lentiform nucleus ( $p = .004$ ), as well as in the ACC ( $p = .015$ ) and PCC ( $p = .045$ ). Activation in the anterior part of the left superior frontal gyrus, the left and right middle superior gyri, the right inferior parietal cortex, and the right middle temporal gyrus no longer differed reliably.

ICV-scores correlated with activation during successful inhibitions in the (anterior part of the) left superior frontal gyrus ( $r = .69$ ,  $p = .013$ ) for adults, though there were no longer reliable correlations with activation scores for the young adolescents.



**Fig. 5.** Performance measures in the performance-matched sample.

### 3.4.3. *fMRI-analysis and functional correlations: commission errors*

In the sample matched for performance, adults showed a greater increase in activation in bilateral culmen ( $p = .006$ ), two regions on the left lingual gyrus ( $p < .001$  and  $p = .011$ ), and the right claustrum ( $p = .008$ ). There were no longer significant differences in the left insula, right caudate, right parahippocampal gyrus, right fusiform gyrus, or the right lingual gyrus.

Young adolescents showed negative correlations between ICV-scores and activation changes in the left middle ( $r = -.61$ ,  $p = .036$ ) and superior ( $r = -.62$ ,  $p = .033$ ) frontal gyri, while no significant correlations with ICV-scores were seen in the adults.

## 4. Discussion

The present study confirms previous behavioural studies that found better response inhibition (fewer commission errors) and sustained attention performance (fewer errors of omission and lower response variability) in adults compared to adolescents/children (Daniel, Pelotte, & Lewis, 2000; Luna, Garver, Urban, Lazar, & Sweeney, 2004) (Clark et al., 2006; Lin, Hsiao, & Chen, 1999; though also see Karatekin, Marcus, & Couperus, 2007). Greater response variability has been previously identified in clinical groups with frontal (Stuss et al., 2003) and fronto-striatal and fronto-parietal brain pathology (e.g., ADHD, see Castellanos et al., 2005; Johnson et al., 2007a). In the current sample the greater response variability for young adolescents likely reflects that these frontal regions are not yet fully matured. This is also consistent with the young adolescents showing negative correlations between response variability and activation changes in most regions, suggesting that lower variability is associated with more 'adult-like' activations (during commission errors, as well as in the superior frontal gyrus during successful inhibitions).

During successful inhibitions, young adolescents showed increased recruitment, compared with adults, of a widely distributed network, including left (inferior, superior and middle) and right (middle and inferior) frontal gyri, left and right insulae, bilateral anterior and posterior cingulate, as well as both left and right inferior parietal cortex and left and right precune and cune. Adults showed higher activation only in the left superior frontal gyrus. The superior behavioural performance and the reduced brain activation levels of adults support the notion that the network underlying response inhibition becomes more sparsely represented as the system matures. To investigate whether this pattern of differences was likely caused by developmental changes to the underlying networks, or simply emerged because of differences in performance, we also compared two groups matched for performance. Although the same overall pattern emerged (with young adolescents showing greater activation in frontal, parietal, and medial regions), there were no longer reliable differences in a number of (predominantly frontal) regions. Thus, although we cannot rule out that performance differences affect the differences in activation maps between the adults and the young adolescents, the young adolescents still rely on a more extensive network during successful inhibitions when performance is equated. There was no indication of a shift in the direction of activation differences (i.e. increased activation for adults compared to young adolescents) when performance was matched between the groups. One limitation to this approach is that our performance matching relied on an arbitrary selection of participants, to ensure equal numbers of (both omission and commission) errors. Our focus on equating for the number of error trials (rather than any other variable, such as response times or ICV-scores) aimed to investigate the possibility that inconsistent results from prior studies (with some showing increased activation for children or adolescents compared to adults, and other studies showing the

opposite) might have been related to whether or not these studies used matched performance in terms of commission errors. While it would be very informative to investigate the relative effects of differences in error-rates, response latencies and variability, on the activation patterns that underlie response inhibition and error-monitoring, this inquiry exceeds the aims of the present study.

This pattern of results is consistent with earlier findings (Booth et al., 2003; Casey et al., 1997; Durston et al., 2002), but opposite to the findings of Rubia et al. (2007, 2006), which may be due, in part, to the latter studies using paradigms that also engage response selection and spatial judgement, in addition to response inhibition. This is suggested by a clear dichotomy in the nature of the tasks that were used in prior studies: those that found increased activation for younger, compared to older participants, used Go/No-Go tasks to probe response inhibition (Booth et al., 2003; Casey et al., 1997; Durston et al., 2006, 2002; this paper; also see Tamm, Menon, & Reiss, 2002). On the other hand, those studies that found the opposite pattern, i.e. higher activation for adults, employed tasks that required spatial discrimination (Bunge et al., 2002; Rubia et al., 2007, 2006).

In line with longitudinal findings (Durston et al., 2006) showing a progression from diffuse to more focal activation patterns, there was only one region that showed higher activation in the adults (the anterior part of the left superior frontal gyrus in our study). This supports the notion that with development of the fronto-parietal network underlying response inhibition, activation in non-critical areas is attenuated in favour of key areas, and may represent emerging cortical specialisation (see Durston et al., 2006; Kelly & Garavan, 2005; Ungerleider, Doyon, & Karni, 2002).

Activation maps relating to commission errors are likely to reflect processes such as error-monitoring and changes in top-down attentional control, as well as (late attempts at) response inhibition. Here, in contrast to the effects for successful inhibitions, we observed greater activation for adults in a number of regions, including in the right parahippocampal gyrus, the left and right lingual gyri, the left insula, and striatal and cerebellar regions including the right caudate and bilateral culmen. When comparing adults and young adolescents with equal performance, this pattern of activation was still observed in the right claustrum, left lingual gyrus, and bilateral culmen. These results, showing an opposite pattern to the relative hyperactivity of the adolescents for successful inhibitions, argue against a non-specific cognitive or vascular basis for the observed age-related changes. They suggest that the relative immaturity of the young adolescent brain translates to a diminished response for errors and requiring greater levels of activity when successfully inhibiting. Indeed, it is possible that the blunted performance monitoring of the young adolescents, as echoed in their diminished error-related response and heightened response variability, may underlie the greater effort (more diffuse activation) when successfully inhibiting. Cerebellar activation has been previously shown in adults as well as adolescents during inhibitory control in Go/No-Go tasks (e.g., Garavan, Ross, Kaufman, & Stein, 2003; Rubia et al., 2007), and this region projects to both motor and prefrontal areas through the thalamus (Kim, Ugurbil, & Strick, 1994; Middleton & Strick, 2000). Activation of the insula has been previously observed in adults during failed attempts to inhibit and, indeed, may be a key neuroanatomical substrate subserving error-detection (Magno, Foxe, Molholm, Robertson, & Garavan, 2006; Ramautar, Slagter, Kok, & Ridderinkhof, 2006). Parahippocampal activity may reflect similar processes, as this region has been implicated in anxiety and arousal (Gray & McNaughton, 2000; also see Green & Arduini, 1954) and memory (e.g., Fernández et al., 1998; also see Jansma, Ramsey, van der Wee, & Kahn, 2004). Thus, adults may be better at inhibiting the prepotent response when an unexpected target appears because when they make an error, they have a



stronger arousal-mediated response which may engage additional top-down attentional control processes to prevent or limit subsequent inhibitory failures (Hester, Barre, Murphy, Silk, & Mattingley, 2008).

Unlike prior studies (Rubia et al., 2007; Velanova et al., 2008), we did not observe differences in the ACC between the two groups for unsuccessful inhibitions, though this may be because of differences in the paradigms used, or due to lower power in our design. For example, the lack of a difference in ACC activation might be related to the use of a response cue in the SART-paradigm. Increasing time-pressure has been previously found to impair response inhibition in younger, compared to older children (see Cragg & Nation, 2008). Thus, *reducing* time-pressure might facilitate the task for the young adolescents in our study, while adults might benefit less from the cue if their performance is closer to ceiling. Given evidence that the ACC monitors response conflict (Carter et al., 1998) and/or error likelihood (Brown & Braver, 2005), the presence of the response cue may therefore reduce between-group differences in activation in this structure. Additionally, there is some evidence that differences in ACC activation during No-Go-trials, between adults and adolescents, may strongly depend on the specific age-groups that are compared. For example, Jonkman, Sniedt, and Kemner (2007, also see Jonkman, 2006) compared younger (6–7 years) and older (9–10 years) children to young adults (19–23 years) using a cued Go/No-Go task. They observed differences in ACC involvement, both in terms of an increased nogo N2 ERP response, as well as a reduced contingent negative variation (CNV) effect related to cueing, but only when comparing the youngest group to both older children and adults, while these measures did not differ between the 9–10-year olds and adults. Similarly, at least one other fMRI-study (Stevens, Kiehl, Pearson, & Calhoun, 2009) failed to find age-related differences (in the age-range 11–37) in an error-related network including the ACC, which the authors suggested might indicate that these regions are already relatively mature in adolescence.

## 5. Conclusions

In summary, previous research suggests that progressive maturation of the developing brain, and especially (pre)frontal regions which mature relatively late compared with more posterior brain regions, leads to shifts in the specific neural networks that underlie executive functions, including response inhibition. There is, however, less agreement on the direction of these differences, with different studies either showing increased or decreased activation of these regions, for children/adolescents compared with adults. We propose that a significant underlying factor for these divergent findings may be differences in the specific paradigms used, particularly with regards to whether the task taps into processes related to spatial attention, in addition to response inhibition.

The present study also provides further support for the notion that parallel to increases in executive control capacities, the underlying neural networks show a shift from more diffuse activation patterns in young adolescents, to more focal prefrontal activation in adults, with reduced reliance on other regions. Even when the two groups were matched for performance, young adolescents showed increased activation during successful inhibitions in frontal, parietal and medial regions. This suggests that the reliance of young adolescents on a more diffuse network cannot solely be attributed to differences in performance. Additionally, the present data suggest an important role for the integrated function of cerebellar, striatal and (para)hippocampal regions in modulating executive control under circumstances when inhibition has failed and greater control must be exerted.

## Acknowledgements

This study was supported by grants from the Science Foundation Ireland, the Irish Health Research Board, the Irish Higher Education Authority's Programme for Research in Third-Level Institutions, the Australian National Health and Medical Research Council Howard Florey Centenary Fellowship, and the FWO Research Foundation - Flanders. The authors would like to thank all participants, as well as the parents of the adolescents who took part in the study.

## References

- Anderson, V., Anderson, P., Northam, E., Jacobs, R., & Catroppa, C. (2001). Development of executive functions through late childhood and adolescence in an Australian sample. *Developmental Neuropsychology*, 20, 385–406.
- Bellgrove, M. A., Hester, R., & Garavan, H. (2004). The functional neuroanatomical correlates of response variability: Evidence from a response inhibition task. *Neuropsychologia*, 42(14), 1910–1916.
- Booth, J. R., Burman, D. D., Meyer, J. R., Lei, Z., Trommer, B. L., Davenport, N. D., Li, W., Parrish, T. B., Gitelman, D. R., & Mesulam, M. M. (2003). Neural development of selective attention and response inhibition. *Neuroimage*, 20, 737–751.
- Brown, J. W., & Braver, T. S. (2005). Learned predictions of error likelihood in the anterior cingulate cortex. *Science*, 307, 1118–1121.
- Bunge, S. A., Dudukovic, N. M., Thomason, M. E., Vaidya, C. J., & Gabrieli, J. D. E. (2002). Immature frontal lobe contributions to cognitive control in children: Evidence from fMRI. *Neuron*, 33, 301–311.
- Carter, C. S., Braver, T. S., Barch, D. M., Botvinick, M. M., Noll, D., & Cohen, J. D. (1998). Anterior cingulate cortex, error detection, and the online monitoring of performance. *Science*, 280, 747–749.
- Casey, B. J., Trainor, R. J., Orendi, J. L., Schubert, A. B., Nystrom, L. E., Giedd, J. N., Castellanos, F. X., Huxley, J. V., Noll, D. C., & Cohen, J. D. (1997). A developmental functional MRI study of prefrontal activation during performance of a Go-No-Go task. *Journal of Cognitive Neuroscience*, 9, 835–847.
- Castellanos, F. X., Sonuga-Barke, E. J., Scheres, A., Di Martino, A., Hyde, C., & Walters, J. R. (2005). Varieties of attention-deficit/hyperactivity disorder-related intra-individual variability. *Biological Psychiatry*, 57(11), 1416–1423.
- Chan, R. C., Chen, E. Y., Cheung, E. F., Chen, R. Y., & Cheung, H. K. (2004). A study of sensitivity of the sustained attention to response task in patients with schizophrenia. *The Clinical Neuropsychologist*, 18(1), 114–121.
- Clark, C. R., Paul, R. H., William, L. M., Arns, M., Fallahpour, K., Handmer, C., & Gordon, E. (2006). Standardized assessment of cognitive functioning during development and aging using an automated touchscreen battery. *Archives of Clinical Neuropsychology*, 21(5), 449–467.
- Cox, R. W. (1996). AFNI: Software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical Research*, 29, 162–173.
- Cragg, L., & Nation, K. (2008). Go or No-Go? Developmental improvements in the efficiency of response inhibition in mid-childhood. *Developmental Science*, 11(6), 819–827.
- Daniel, D. B., Pelotte, M., & Lewis, J. (2000). Lack of sex differences on the stroop color-word test across three age groups. *Perceptual and Motor Skills*, 90, 483–484.
- Dehaene, S., Posner, M. I., & Tucker, D. M. (1994). Localization of a neural system for error detection and compensation. *Psychological Science*, 5, 303–305.
- Denckla, M. B. (1996). A theory and model of executive function: A neuropsychological perspective. In G. Lyon, & N. Krasnegor (Eds.), *Attention, memory and executive function*. Maryland: Paul Brooks.
- Durstun, S., Davidson, M. C., Tottenham, N., Galvan, A., Spicer, J., Fossella, J. A., & Casey, B. J. (2006). A shift from diffuse to focal cortical activity with development. *Developmental Science*, 9, 1–8.
- Durstun, S., Thomas, K. M., Yang, Y. H., Uluğ, A. M., Zimmerman, R. D., & Casey, B. J. (2002). A neural basis for the development of inhibitory control. *Developmental Science*, 5, F9–F16.
- Fassbender, C., Murphy, K., Foxe, J. J., Wylie, G. R., Javitt, D. C., Robertson, I. H., & Garavan, H. (2004). A topography of executive functions and their interactions revealed by functional magnetic resonance imaging. *Cognitive Brain Research*, 20, 132–143.
- Fernández, G., Weyerts, H., Schrader-Bölsche, M., Tendolkar, I., Smid, H. G., Tempelmann, C., Hinrichs, H., Scheich, H., Elger, C. E., Mangun, G. R., & Heinze, H. J. (1998). Successful verbal encoding into episodic memory engages the posterior hippocampus: A parametrically analyzed functional magnetic resonance imaging study. *Journal of Neuroscience*, 18(5), 1841–1847.
- Fronczek, R., Middelkoop, H. A., van Dijk, J. G., & Lammers, G. J. (2006). Focusing on vigilance instead of sleepiness in the assessment of narcolepsy: High sensitivity of the sustained attention to response task (SART). *Sleep*, 29, 187–191.
- Garavan, H., Ross, T. J., Kaufman, J., & Stein, E. A. (2003). A midline dissociation between error-processing and response-conflict monitoring. *Neuroimage*, 20, 1132–1139.
- Garavan, H., Ross, T. J., Murphy, K., Roche, R. A. P., & Stein, E. A. (2002). Dissociable executive functions in the dynamic control of behavior: Inhibition, error detection, and correction. *Neuroimage*, 17, 1820–1829.
- Garavan, H., Ross, T. J., & Stein, E. A. (1999). Right hemispheric dominance of inhibitory control: An event-related functional MRI study. *Proceedings of the National Academy of Sciences of the United States of America*, 96, 8301–8306.



- Gogtay, N., Giedd, J. N., Lusk, L., Hayashi, K. M., Greenstein, D., Vaituzis, A. C., Nugent, T. F., 3rd, Herman, D. H., Clasen, L. S., Toga, A. W., Rapoport, J. L., & Thompson, P. M. (2004). Dynamic mapping of human cortical development during childhood through early adulthood. *Proceedings of the National Academy of Sciences of the United States of America*, 101, 8174–8179.
- Gray, J. A., & McNaughton, N. (2000). *The neuropsychology of anxiety: An enquiry into the functions of the septo-hippocampal system*. Oxford University Press.
- Green, J. D., & Arduini, A. A. (1954). Hippocampal electrical activity in arousal. *Journal of Neurophysiology*, 17, 533–557.
- Hester, R., Barre, N., Murphy, K., Silk, T. J., & Mattingley, J. B. (2008). Human medial frontal cortex activity predicts learning from errors. *Cerebral Cortex*, 18, 1933–1940.
- Jansma, J. M., Ramsey, N. F., van der Wee, N. J., & Kahn, R. S. (2004). Working memory capacity in schizophrenia: A parametric fMRI study. *Schizophrenia Research*, 68(2–3), 159–171.
- Johnson, K. A., Kelly, S. P., Robertson, I. H., Barry, E., Mulligan, A., Daly, M., Lambert, D., McDonnell, C., Connor, T. J., Hawi, Z., Gill, M., & Bellgrove, M. A. (2007). Absence of the 7-repeat variant of the DRD4 VNTR is associated with drifting sustained attention in children with ADHD but not in controls. *American Journal of Medical Genetics Part B: Neuropsychiatric Genetics*, doi:10.1002/ajmg.b.30718
- Johnson, K. A., Robertson, I. H., Kelly, S. P., Silk, T. J., Barry, E., Dáibhis, A., Watchorn, A., Keavey, M., Fitzgerald, M., Gallagher, L., Gill, M., & Bellgrove, M. A. (2007). Dissociation in performance of children with ADHD and high-functioning autism on a task of sustained attention. *Neuropsychologia*, 45(10), 2234–2245.
- Jones, L. B., Rothbart, M. K., & Posner, M. I. (2003). Development of executive attention in preschool children. *Developmental Science*, 6, 498–504.
- Jonkman, L. M. (2006). The development of preparation, conflict monitoring and inhibition from early childhood to young adulthood; a go/nogo ERP study. *Brain Research*, 1097, 181–193.
- Jonkman, L. M., Sniedt, F. L. F., & Kemner, C. (2007). Source localization of the nogo-N2: A developmental study. *Clinical Neurophysiology*, 118, 1069–1077.
- Karatekin, C., Marcus, D. J., & Couperus, J. W. (2007). Regulation of cognitive resources during sustained attention and working memory in 10-year-olds and adults. *Psychophysiology*, 44(1), 128–144.
- Kelly, C., & Garavan, H. (2005). Human functional neuroimaging of brain changes associated with practice. *Cerebral Cortex*, 15, 1089–1102.
- Kim, S. G., Ugurbil, K., & Strick, P. L. (1994). Activation of a cerebellar output nucleus during cognitive processing. *Science*, 265, 949–951.
- Lin, C. C. H., Hsiao, C. K., & Chen, W. J. (1999). Development of sustained attention assessed using the continuous performance test among children 6–15 years of age. *Journal of Abnormal Child Psychology*, 27(5), 403–412.
- Luna, B., Garver, K. E., Urban, T. A., Lazar, N. A., & Sweeney, J. A. (2004). Maturation of cognitive processes from late childhood to adulthood. *Child Development*, 75(5), 1357–1372.
- Magno, E., Foxe, J. J., Molholm, S., Robertson, I., & Garavan, H. (2006). The anterior cingulate and error avoidance. *The Journal of Neuroscience*, 26, 4769–4773.
- McAvinue, L., O'Keefe, F., McMackin, D., & Robertson, I. H. (2005). Impaired sustained attention and error awareness in traumatic brain injury: Implications for insight. *Neuropsychological Rehabilitation*, 15(5), 569–587.
- Middleton, F. A., & Strick, P. L. (2000). Basal ganglia and cerebellar loops: Motor and cognitive circuits. *Brain Research Reviews*, 31, 236–250.
- Murphy, K., & Garavan, H. (2004). An empirical investigation into the number of subjects required for an event-related fMRI study. *Neuroimage*, 22, 879–885.
- O'Connell, R. G., Dockree, P. M., Bellgrove, M. A., Turin, A., Ward, S., Foxe, J. J., & Robertson, I. H. (2008). Two types of action error: Electrophysiological evidence for separable inhibitory and sustained attention neural mechanisms producing error on Go/No-Go tasks. *Journal of Cognitive Neuroscience*, doi:10.1162/jocn.2009.21008
- Pruessmann, K. P., Weiger, M., Scheidegger, M. B., & Boesiger, P. (1999). SENSE: Sensitivity encoding for fast MRI. *Magnetic Resonance in Medicine*, 42(5), 952–962.
- Ramautar, J. R., Slagter, H. A., Kok, A., & Ridderinkhof, K. R. (2006). Probability effects in the stop-signal paradigm: The insula and the significance of failed inhibition. *Brain Research*, 1105(1), 143–154.
- Robertson, I. H., Manly, T., Andrade, J., Baddeley, B. T., & Yiend, J. (1997). 'Oops!': Performance correlates of everyday attentional failures in traumatic brain injured and normal subjects. *Neuropsychologia*, 35, 747–758.
- Rubia, K., Smith, A. B., Taylor, E., & Brammer, M. (2007). Linear age-correlated functional development of right inferior fronto-striato-cerebellar networks during response inhibition and anterior cingulate during error-related processes. *Human Brain Mapping*, 28, 1163–1177.
- Rubia, K., Smith, A. B., Wooley, J., Nosarti, C., Heyman, I., Taylor, E., & Brammer, M. (2006). Progressive increase in frontostriatal activation from childhood to adulthood during event-related tasks of cognitive control. *Human Brain Mapping*, 27, 973–993.
- Stevens, M. C., Kiehl, K. A., Pearson, G. D., & Calhoun, V. D. (2007). Functional neural networks underlying response inhibition in adolescents and adults. *Behavioural Brain Research*, 181, 12–22.
- Stevens, M. C., Kiehl, K. A., Pearson, G. D., & Calhoun, V. D. (2009). Brain network dynamics during error commission. *Human Brain Mapping*, 30(1), 24–37.
- Stuss, D. T., Murphy, K. J., Binns, M. A., & Alexander, M. P. (2003). Staying on the job: The frontal lobes control individual performance variability. *Brain*, 126(11), 2363–2380.
- Tamm, L., Menon, V., & Reiss, A. L. (2002). Maturation of brain function associated with response inhibition. *Journal of the American Academy of Child and Adolescent Psychiatry*, 41(10), 1231–1238.
- Ungerleider, L. G., Doyon, J., & Karni, A. (2002). Imaging brain plasticity during motor skill learning. *Neurobiology of Learning and Memory*, 78, 553–564.
- Velanova, K., Wheeler, M. E., & Luna, B. (2008). Maturation changes in anterior cingulate and frontoparietal recruitment support the development of error processing and inhibitory control. *Cerebral Cortex*, 18, 2505–2522.
- West, R., Murphy, K. J., Armilio, M. L., Craik, F. I., & Stuss, D. T. (2002). Lapses of intention and performance variability reveal age-related increases in fluctuations of executive control. *Brain and Cognition*, 49(3), 402–419.
- Whyte, J., Grieb-Neff, P., Gantz, C., & Polansky, M. (2006). Measuring sustained attention after traumatic brain injury: Differences in key findings from the sustained attention to response task (SART). *Neuropsychologia*, 44, 2007–2014.