



Contents lists available at ScienceDirect

Neurobiology of Aging

journal homepage: www.elsevier.com/locate/neuaging

Learning to remember: cognitive training-induced attenuation of age-related memory decline depends on sex and cognitive demand, and can transfer to untrained cognitive domains

Joshua S. Talboom^{a,b}, Stephen G. West^{a,b}, Elizabeth B. Engler-Chiurazzi^{a,b},
Craig K. Enders^a, Ian Crain^a, Heather A. Bimonte-Nelson^{a,b,*}

^a Department of Psychology, Arizona State University, Tempe, AZ, USA

^b Arizona Alzheimer's Consortium, Phoenix, AZ, USA

ARTICLE INFO

Article history:

Received 25 September 2013

Received in revised form 27 May 2014

Accepted 10 June 2014

Keywords:

Cognitive training

Age-related memory decline

Aging

Learning

Memory

Spatial

Enrichment

Neurobics

ABSTRACT

Aging is associated with progressive changes in learning and memory. A potential approach to attenuate age-related cognitive decline is cognitive training. In this study, adult male and female rats were given either repeated exposure to a T-maze, or no exposure to any maze, and then tested on a final battery of cognitive tasks. Two groups of each sex were tested from 6 to 18 months old on the same T-maze; Group one received a version testing spatial reference memory, and Group two received only the procedural testing components with minimal cognitive demand. Groups three and four of each sex had no maze exposure until the final battery, and were comprised of aged or young rats, respectively. The final maze battery included the practiced T-maze plus two novel tasks, one with a similar, and one with a different, memory type to the practice task. Group five of each sex was not maze tested, serving as an aged control for the effects of maze testing on neurotrophin protein levels in cognitive brain regions. Results showed that adult intermittent cognitive training enhanced performance on the practice task when aged in both sexes, that cognitive training benefits transferred to novel tasks only in females, and that cognitive demand was necessary for these effects, since rats receiving only the procedural testing components showed no improvement on the final maze battery. Further, for both sexes, rats that showed faster learning when young demonstrated better memory when aged. Age-related increases in neurotrophin concentrations in several brain regions were revealed, which were related to performance on the training task only in females. This longitudinal study supports the tenet that cognitive training can help one remember later in life, with broader enhancements and associations with neurotrophins in females.

Published by Elsevier Inc.

1. Introduction

Life expectancy is continuing to increase in tandem with the baby-boom generation growing older. Projections to the year 2050 show 19 million Americans will be living into their 80s (Shrestha, 2005; Vincent and Velkoff, 2010). As age is the number one risk factor for developing Alzheimer's disease (AD; Lindsay et al., 2002), this population faces a greater burden of both normal age-associated and pathological cognitive decline. Currently, there exist few treatments for AD, with only 2 classes of U.S. Food and Drug Administration (FDA) approved drugs which possess short-

term efficacy and do not prevent or halt the progression of AD; of note, these drugs are not prescribed for age-associated cognitive decline (Chiang and Koo, 2014; Petersen et al., 2001). Remaining cognitively active may help protect against age-related cognitive decline and development of AD. In fact, that a nonpharmacologic approach might protect against cognitive decline due to normal aging or neurodegenerative disease provides potentially new avenues for treatment.

Accumulating evidence supports the tenet that cognitive activity is related to cognitive prowess as aging ensues. For example, people 60- to 85-years-old were able to improve their ability to multitask by practicing a video game activity for one month (Anguera et al., 2013). Interestingly, these aged participants outperformed 20 year olds, enhancements persisted for at least six months, and benefits were noted in other cognitive domains (Anguera et al., 2013). Other forms of cognitive training also show benefit. Education is

* Corresponding author at: Department of Psychology, Arizona State University, PO Box 871104, Tempe, AZ 85287, USA. Tel.: +1 480 727 0766; fax: +1 480 965 8544.
E-mail address: bimonte.nelson@asu.edu (H.A. Bimonte-Nelson).

related to decreased risk of developing dementia and some symptoms of AD (Bowirrat et al., 2002; De Ronchi et al., 1998; Fritsch et al., 2002; Gatz et al., 2001; Letenneur et al., 1999; Ott et al., 1999), and more early- and mid-life cognitive activity is related to less β -amyloid deposition (Landau et al., 2012). Moreover, college professors with 21–23 years of education maintained cognitive performance as they aged, with professors in their 60s performing as well as professors in their 30s (Shimamura et al., 1995). Finally, better cognitive ability as a child, as measured on a global scale of cognitive functioning, was associated with a decreased risk of developing late-onset dementia (Whalley et al., 2000), and cognitive stimulation therapy was beneficial to people diagnosed with AD (reviewed in Ballard et al., 2011). Thus, collectively, there is evidence that multiple forms of cognitive activity attenuates age- and AD-related cognitive decline.

The research conducted in humans is largely corroborated by studies using animal models. For example, short-term working memory training in young adult mice enhanced selective attention and performance on a battery of tests (Light et al., 2010). Furthermore, long-term cognitive training in rodents benefitted later performance on tasks assessing working and reference memory, as well as selective attention (Bierley et al., 1986; Markowska and Savonenko, 2002; Matzel et al., 2011; Vicens et al., 2003); these effects do not appear to depend on visual acuity or swimming ability (Markowska and Savonenko, 2002), but do transfer to novel and/or untrained tasks and novel cognitive domains (Light et al., 2010; Matzel et al., 2011). Most of this research has evaluated males; in fact, only one rodent study has evaluated females, and there was no male comparison group (Ando and Ohashi, 1991). Hence, it is unknown whether sex differences exist in response to cognitive training, especially with regard to age-related memory changes. There is some evidence of a sex difference in the effectiveness of cognitive practice from research in young adult non-human primates; young female rhesus monkeys showed marked prior spatial training-induced enhancements on a complex spatial task while still in young adulthood, an effect not seen in males (Lacreuse et al., 2005).

The neurobiological mechanism whereby cognitive activity might attenuate age-related cognitive decline is largely unknown. One potential mechanism is the neurotrophic system, which has been found to change with age (Mufson et al., 1995). Neurotrophins, such as nerve growth factor (NGF), impact cholinergic neuron survival and maintenance (Granhölm, 2000; Levi-Montalcini, 1987; Woolf, 1991), and numerous studies have demonstrated that age-related memory deficiency is coupled with alterations in the cholinergic system (e.g., humans: Drachman and Leavitt, 1974, rats: Eckerman et al., 1980, and monkeys: Rupniak et al., 1990). Furthermore, NGF protein concentrations have been correlated with cognitive maze scores in aged rats (Bimonte-Nelson et al., 2003a), and NGF administration has been shown to improve learning and memory deficits in aged rats (Backman et al., 1996; Fischer et al., 1987; Gustilo et al., 1999; Markowska et al., 1994; Scali et al., 1994). Taken together, it is plausible that long-term cognitive training impacts NGF concentrations, which in turn could translate to an attenuation of age-related memory changes.

In the present study, we aimed to determine whether long-term cognitive training attenuates age-related cognitive decline in male and female rats, and, if so, what parameters guide these effects. Therefore, for male and female rats, this study was designed to evaluate the following: (1) whether the effects of cognitive training were due to cognitive demand and/or the procedural components of testing; and (2) whether effects were limited to the trained cognitive domain or if they transferred to an untrained cognitive domain. Two recent studies found that

cognitive demand is indeed necessary for the effects of cognitive training to be realized, and that benefits can transfer to novel tasks and novel cognitive domains (Light et al., 2010; Matzel et al., 2011); however, both studies evaluated only male mice. Furthermore, given the human literature suggesting that cognitive ability (Whalley et al., 2000) or cognitive reserve (Snowdon, 2003) impacts cognitive status when aged, we also assessed whether performance during the first cognitive training session when young related to later performance when aged. After testing on the final maze battery, NGF concentrations were assayed in several cognitive brain regions. This allowed us to evaluate whether NGF levels were altered by cognitive training and/or related to any cognitive training-induced learning and memory enhancements. We hypothesized that cognitive training would alter the trajectory of age-related cognitive decline and neurotrophic systems in a sex-specific manner. Indeed, given long-standing research suggesting that females are “buffered” against insult and show more plasticity in comparison to males (for review see Fitch and Bimonte, 2002; Juraska, 1986), we predicted that females would be most benefitted by cognitive training.

2. Methods

2.1. Subjects

Six-month-old Fisher 344 virgin rats (46 females and 50 males) were obtained from the National Institute on Aging colony at Harlan Laboratories and were pair housed in the Arizona State University rodent facility in the Department of Psychology. Rats had exposure to food and water *ad libitum* and were maintained on a 12-hour light and/or dark cycle at 23 °C. Procedures were approved by the Arizona State University Institutional Care and Use Committee (IACUC), adhered to the Guide for the Care and Use of Laboratory Rats (National Research Council, 2011), and National Institutes of Health standards. Following Markowska and Savonenko (2002), initiation of the cognitive and procedural component training schedule began at 6 months of age. Rats received 1 training session every 3 months; training continued for 12 months with rats receiving their last training session at 18 months of age. Care was taken to ensure that the T-maze, and all objects and spatial cues in the training maze room, remained unchanged for the entire 12-month training and final maze battery testing periods. After behavioral testing, all rats were euthanized, and the rapid collection of several brain regions ensued immediately. At this experimental time point, the aged groups were 23 months of age, and the young-naïve group was 8 months of age. Later, NGF protein concentrations from the collected brain regions were assayed.

To meet the goals of the experiment, we used 4 groups of rats that were 6-months-old at the beginning of the study and 21-months-old at the beginning of the final maze battery, and 1 group of young rats that were 6-months-old at the beginning of the final maze battery. This resulted in a total of 5 groups of rats, with each sex represented in each group: (1) the aged cognitive training group which assessed the effects of spatial reference memory training from 6 to 18 months of age (Aged Cognitive Training); (2) the aged procedural component group which addressed the effects of the procedural components of maze testing (Aged Swim Only); (3) the aged naïve group which had no maze exposure before the final maze battery (Aged Naïve); (4) the aged never-tested group which assessed the effects of age on NGF protein concentrations in rats that had no maze exposure (Aged Never Tested); and (5) the young 6-month-old naïve group, which we included so we could compare aged groups to a group with “optimal” learning and memory ability (Young).




Fig. 1. (A) Timeline and design of the experiment. (B) A diagram of the mazes used in this study. The spatial working and reference memory WRAM was the first novel task in a novel environment that behaviorally tested groups were exposed to during the final maze test battery. The spatial reference memory Morris maze was the second novel task in a novel environment used during the final maze test battery. The T-maze was used as the spatial reference memory training task, and was modified for use as the procedural components task, and it was tested last during the final maze test battery. Abbreviation: WRAM, water radial-arm maze.

2.1.1. Treatment groups

Fig. 1A displays each treatment group and their age, concordant with a timeline of the experimental procedures. The Aged Cognitive Training group was composed of 19 rats (8 females, 11 males). From 6 to 18 months of age, this group was tested every 3 months for 5 days, 6 trials per day, on the spatial reference memory T-maze using procedures previously published (Denenberg et al., 1990, 1991; Gibbs et al., 2004). The black Plexiglas maze (each arm was 38.1 cm × 12.7 cm) was filled with water made opaque with black nontoxic paint, and had a hidden escape platform with a wire mesh top at the end of 1 of the 2 non-start arms. Each rat had an assigned exit location (East or West) for the entire duration of training through the final maze battery. Drop off locations varied between north and south arms to minimize the possibility of rats using a motoric and/or olfactory strategy rather than a spatial strategy. After a rat located the platform, it remained there for 5 seconds; the rat was then removed and placed in its heated cage for 30 seconds until the next trial. Between trials the maze was cleaned and water disrupted to minimize the possibility of rats using an olfactory strategy. The dependent variable was the number of correct arm entries, quantified as the number of times a rat found the hidden escape platform on the first choice of each trial within a day.

The Aged Swim Only group consisted of 21 rats (10 females, 11 males). From 6 to 18 months of age, this group experienced conditions identical to the Aged Cognitive Training group, with the exception that the T-maze was modified to minimize cognitive demand. Exit arms were assigned randomly for each trial pair. Cognitive demand was minimized by keeping the nonexit, nonplatformed, arm blocked; hence, the escape arm was the only arm left open. This resulted in this group receiving similar experience as the Aged Cognitive Training group regarding being removed from the colony room and experiencing handling and maze testing. Drop off locations varied between north and south arms to maximize swimming

similarities to the Aged Cognitive Training group. Inclusion of the Aged Swim Only group addressed whether the procedural components of maze testing throughout the training period influenced age-related cognitive change.

The Aged Naïve group consisted of 20 rats (10 females, 10 males), which remained in their colony room cages from 6 to 21 months of age (i.e., the entire training period, plus the delay until the final maze test battery) and then were tested on the final maze battery. The Aged Naïve group allowed us to reference the normally aged laboratory rat that received neither cognitive training nor prior behavior testing.

The Aged Never Tested group was composed of 18 rats (9 females and 9 males) that remained in their colony room cages from 6 to 23 months of age (i.e., the entire study period) and were never behaviorally test. In fact, these rats were not handled before euthanasia aside from normal caretaking procedures (e.g., cage changes). Aged Never Tested rats were euthanized along with the other groups of rats after the final maze battery. We used this group to evaluate NGF concentrations in the cognitive brain regions of rats that were not exposed to behavioral testing.

The Young group comprised 18 (9 females and 9 males) adult rats. This group was tested only during the final maze battery, so that we could compare the aged groups with the cognitively intact, nonexperimentally manipulated, young rat. These rats were 6 months of age when tested on the final maze battery. This age was chosen for the Young group, because this is the age when the cognitive training period started for the Swim Only and Cognitive Training groups.

2.2. Vaginal and control anal swabs

After the completion of all training sessions, just before the initiation of the final maze battery, all females received vaginal smears once a day in the morning for several consecutive days following previously published methods (Acosta et al., 2009b;

Talboom et al., 2010). Vaginal cytology was classified as proestrus, estrous, metestrus, or diestrus (Goldman et al., 2007). All aged females were categorized as being in an estropause state. Nearly all of the aged females remained in anestrus; most of the aged females consistently displayed a constant diestrus smear, with overall few cells that were primarily leukocytes, consistent with anestrus and relatively stable circulating levels of ovarian hormones. Very few aged females displayed transitioning diestrus- and proestrus-like smears. The transitioning proestrus-like smear consisted of epithelial cells, intermediate cells, and noticeable mucus. All young females displayed normal 4-day cycles, indicative of normal reproductive capacity and fluctuating concentrations of circulating ovarian hormones. To control for the handling and procedure of vaginal swabs, all males received a gentle anal probe with a swab using a nearly identical procedure to the one used to perform vaginal swabs for the females.

2.3. Mazes used during the final maze battery

We began the final maze battery when the aged groups (Aged Cognitive Training, Aged Swim Only, and Aged Naïve) were 21 months of age. The final maze battery included mazes assessing both spatial working and reference memory. We chose these mazes to determine whether cognitive training on one maze (the T-maze) benefited performance only on the same maze (the T-maze) or whether effects transferred to untrained mazes (the water radial arm maze [WRAM] and Morris maze). Furthermore, because the final maze battery included mazes assessing both working and reference memory, we could also determine whether memory transfer effects to novel mazes occurred only within the trained memory domain (reference memory) or whether they transferred to an untrained memory domain (working memory) as well. The final maze battery, on which all groups except the Aged Never Tested group were evaluated, included the following mazes in the order of final battery testing: the win-shift spatial working memory and reference memory WRAM, the spatial reference memory Morris maze, and the spatial reference memory T-maze. The WRAM and Morris maze were novel tasks, assessing performance in novel environments and testing rooms. The T-maze was a familiar task in a familiar environment and testing room to the Aged Cognitive Training and Aged Swim Only groups, except cognitive demand was now required of the Aged Swim Only group. Each maze was located in a separate, distinct room that had salient, visual, extra-maze cues that remained constant throughout the testing period.

2.3.1. Water radial-arm maze

The win-shift WRAM is an 8-arm maze (Fig. 1B) which was constructed of black Plexiglas and filled with room temperature water made opaque using black nontoxic paint (Bimonte and Denenberg, 1999, 2000; Bimonte et al., 2000). Hidden escape platforms (1 cm below the water) were placed in the ends of 4 of the 8 arms. Each subject had 4 platform locations that remained fixed throughout testing. A subject was released from the start arm and had 3 minutes to locate a platform; if it did not find a platform during the 3 minutes, it was gently led using a dark rod, to the closest one. Once on a platform, the rat remained on it for 15 seconds and was then returned to its heated cage for a 30-second intertrial interval (ITI). During the ITI, the just located platform was removed from the maze. The rat was then placed again into the start arm and allowed to locate another platform. For each rat, a daily session consisted of 4 trials, with the number of platformed arms reduced by 1 on each subsequent trial. Thus, the working memory load was increasingly taxed as

trials progressed. Each rat was given 1 session a day for 12 consecutive days. Quantification and blocking procedures were based upon previous studies using the WRAM (Bimonte-Nelson et al., 2003a; Bimonte et al., 2002, 2003). Errors were quantified using orthogonal measures of working and reference memory (Jarrard et al., 1984), as done in previous WRAM studies (see Bimonte-Nelson et al., 2003a; Bimonte et al., 2000, 2002; Hyde et al., 2000). Working memory correct (WMC) errors were the number of first and repeat entries into any arm from which a platform had been removed during that day. Reference memory (RM) errors were the number of first entries into any arm that never contained a platform within a day. Working memory incorrect (WMI) errors were the number of repeat entries into an arm that never contained a platform within a day (i.e., repeat entries into a reference memory arm within a day). Number of WMC, WMI, and RM errors served as the dependent variables for statistical analysis, and data were blocked into 4-day blocks (Bimonte-Nelson et al., 2003a; Bimonte et al., 2002, 2003).

2.3.2. Morris maze

The Morris maze (see Fig. 1B for a diagram of the maze, Morris et al., 1982) evaluated spatial reference memory and consisted of a round tub (188 cm in diameter) filled with water made opaque with black nontoxic paint. The rat was placed in the maze from any of the 4 locations (North, South, East, or West) and had 60 seconds to locate a submerged hidden escape platform with a wire mesh top, which remained in a fixed location (the target Northeast quadrant, NE) throughout testing. After 15 seconds on the platform, the rat was placed into its heated cage until the next trial; the ITI was 5–8 minutes. For each rat, the testing session consisted of 4 trials/day for 5 days. A video camera recorded each rat, and a tracking system (EthoVision version 3.1, Noldus Information Technology, Wageningen, the Netherlands) analyzed each rat's path. The dependent measure was swim distance (cm), with less swim distance interpreted as better performance. To assess platform localization, a probe trial was given on an additional trial (trial 5) on the last day of testing, whereby the escape platform was removed from the maze. The dependent measures for the probe trial were percentage of total swim distance (cm) in the target NE quadrant as compared with the opposite Southwest (SW) quadrant.

2.3.3. T-maze

The procedure for the T-maze (Fig. 1B) during the final maze battery was identical to that given to the Aged Cognitive Training group during the training period, as described previously. For the rats with T-maze experience from the training sessions, the platform location remained constant (in the same arm of the maze and same place in space) during cognitive training and the final maze battery. The dependent variable was the number of correct arm entries made within a daily session.

2.4. Brain dissection and NGF analysis

Rats were anesthetized with isoflurane (VetOne, Meridian, IN, USA) and euthanized according to National Institutes of Health guidelines and American Veterinary Medical Association (AVMA) guidelines on euthanasia (2007), after which the brains were rapidly dissected. Following a stereotaxic atlas (Paxinos and Watson, 2005), right hemisphere frontal cortex, parietal cortex, temporal cortex, striatum, entorhinal cortex, and ventral hippocampus (CA1/2) were rapidly dissected, as done previously (Bimonte-Nelson et al., 2003a). Dissected tissues were immediately placed in weighed microcentrifuge tubes, frozen on dry ice,

and stored at -70°C until tissue lysates were made. NGF protein concentrations were assessed via enzyme-linked immunosorbent assay (ELISA) using commercially available kits from Promega (Madison, WI, USA). NGF assay procedures were conducted as previously described (Engler-Chiurazzi et al., 2011). Briefly, tissue lysates were processed and read in flat-bottom 96 well plates according to the kit's instructions. The range of NGF detection was between 7.8–500 pg/mL. Cross-reactivity with other trophic proteins was $<2\%$ – 3% . The concentration of NGF (pg/mg of tissue) present in the brain region of interest was the dependent variable used for statistical analysis.

2.5. Statistical methods

The present study was designed to determine whether cognitive training enhanced performance in aged animals, whether there was an age effect on each maze, and whether the cognitive training or age effects differed by sex. Therefore, we aimed to compare the Young group to the Aged Naïve group to determine age effects, and the Aged Swim Only group to the Aged Cognitive Training group to determine cognitive training effects in the aged animals, in the context of cognitive demand. As a result, comparisons were planned a priori and Type I error correction was not necessary (Keppel and Wickens, 2004). Data were analyzed using a mixed model analysis of variance (ANOVA) with either age or treatment and sex, as the between-subjects variables, and repeated measures variables as appropriate for the specific test. This statistical routine is common (e.g., Braden et al., 2011) and was necessary so that we could evaluate potentially complex higher order interactions with days and/or trials for behavior assessments. To evaluate potential relations between maze performance when young and NGF concentrations when aged, one predictor regression analysis including Pearson product-moment correlation coefficient (r) was run within the Cognitive Training group. All ANOVAs, regressions, and correlations were performed using StatView (for Windows, version 5.0.1, SAS Institute Inc, Cary, NC, USA); the data were then graphed using Prism (version 6.04, GraphPad Software, La Jolla, CA, USA).

We used linear growth modeling in the present study. Paralleling research conducted in humans (Whalley et al., 2000), we wished to evaluate whether cognitive ability when young related to cognitive ability when aged. We estimated a linear growth model from the number of correct arm entries made by the young Cognitive Training group across their first session on the cognitive training T-maze. The model was predicted as a function of the elapsed number of days from the initial cognitive training session conducted when the rats were 6 months of age (Singer and Willett, 2003). Specifically, we used growth modeling to determine if initial cognitive ability and/or performance (intercept) or rate of learning (slope) differed between males and females, and if ability or rate of learning when young was related to cognitive performance on novel tasks when aged (described in more detail in the following sections). We used Mplus (version 5.21, Muthén & Muthén, Los Angeles, CA, USA) to estimate the linear growth model.

3. Results

At the conclusion of the study, statistical analyses were performed to evaluate potential differences between the Aged Swim Only and Aged Naïve groups for behavior, and the Aged Swim Only, Aged Naïve, and Aged Never Tested groups for the NGF analysis; no significant differences would allow simplification of the analyses because these groups could be combined into one

control group for each respective analysis. There were no significant differences between the two groups for behavior for any dependent measure showing cognitive training differences, nor were there significant differences between the 3 groups for the NGF analysis. Therefore, we combined the Aged Swim Only and Aged Naïve groups into one Aged Control group for evaluation of cognitive training effects, and the Aged Swim Only, Aged Naïve, and Aged Never Tested groups into the Aged Control group for analysis of NGF concentrations.

3.1. Water radial-arm maze

Fig. 2 shows the mean \pm standard error (SE) scores for each group, collapsed across all blocks and trials. Young rats outperformed Aged Control rats for WMC ($F[1,56] = 4.40, p < 0.05$; Fig. 2A), WMI ($F[1,56] = 16.77, p < 0.0001$; Fig. 2B), and RM ($F[1,56] = 23.28, p < 0.0001$; Fig. 2C) with no age \times sex interaction. For the Aged Control and Aged Cognitive Training group analyses, there were no main effects or interactions with sex for WMC or RM. Effects were specific to WMI. Fig. 3 shows the mean \pm SE scores for each block for each group for WMI. For WMI, there was no main effect of treatment; however, there was a significant treatment \times block ($F[2,116] = 3.26, p < 0.05$) and treatment \times block \times trial interaction ($F[6,348] = 2.13, p < 0.05$). Further, WMI analysis of the last block of testing (for the asymptotic phase of performance, as done routinely to test memory after learning has occurred; e.g., Acosta et al., 2009a; Bimonte-Nelson et al., 2003b, 2004; Bimonte and Denenberg, 1999; Digby et al., 2012) revealed a significant sex \times trial interaction ($F[3,183] = 8.51, p < 0.0001$) and a marginal treatment \times sex \times trial interaction ($F[3,183] = 2.14, p = 0.09$), as shown in Fig. 4A. Because the working memory load increases as trials increase, an interaction with trial could indicate a differential ability to handle an increasing working memory load. Thus, for WMI we tested the trial at the highest working load, trial 4, as done previously using this task (Acosta et al., 2009a; Braden et al., 2010; Digby et al., 2012). As shown in Fig. 4B, for WMI Aged Cognitive Training females made fewer errors than Aged Control females when working memory load was the highest ($F[1,27] = 4.85, p < 0.05$), an effect not found in males as Aged Cognitive Training males did not differ from Aged Control males ($F[1,34] = 0.03, p = 0.86$).

3.2. Morris maze

Fig. 5A shows the mean \pm SE swim distance scores across days for each group. Young animals exhibited better performance than Aged Controls, as Young animals had lower swim distance scores than Aged Controls ($F[1,56] = 18.01, p < 0.0001$); there were no significant age \times sex interactions. For the analysis of the Aged Control and Aged Cognitive Training groups, there was no main effect of treatment, and there was a significant treatment \times day interaction ($F[4,232] = 2.99, p < 0.05$); Aged Cognitive Training rats showed better performance starting on day 3. Because these analyses revealed that the Aged Cognitive Training group demonstrated better performance starting on day 3, we evaluated overnight forgetting of the platform location during the latter block of testing (overnight forgetting for days 2–3, days 3–4, and days 4–5), by computing and testing the difference in swim distance between the first trial of the next day after the overnight interval, and the last trial of the previous day (trial 1 distance–trial 4 distance). Overnight forgetting is shown for all groups in Fig. 5B. For this assessment of overnight forgetting, there was a significant treatment \times sex interaction ($F[1,59] = 4.67, p < 0.05$); Aged Cognitive Training females outperformed Aged Control females (F




Fig. 2. Mean \pm SE number of: (A) WMC, (B) WMI, and (C) RM errors, committed on the WRAM collapsed across all blocks and trials. For each error type, Young animals outperformed the Aged Control animals; * $p < 0.05$. Abbreviations: RM, reference memory; SE, standard error; WMC, working memory correct; WMI, working memory incorrect; WRAM, water radial-arm maze.

[1,27] = 5.86, $p < 0.05$). This effect was not found in their male counterparts, as Aged Cognitive Training males did not differ from Aged Control males ($F[1,32] = 0.11$, $p = 0.74$). Female Aged Control rats had more overnight forgetting than female Young rats (female Aged Control vs. female Young: $F[1,28] = 4.08$, $p \leq 0.05$); this effect was not seen in males (male Aged Control vs. male Young: $F[1,29] = 1.90$, $p = 0.18$).

Fig. 5C shows the mean \pm SE probe trial data for each group. For the probe trial, for the Young and Aged Control comparison, all animals spent a greater percent of the total swim distance in the target NE versus opposite SW quadrant (main effect of quadrant: $F[1,55] = 182.01$, $p < 0.0001$), and there was a significant treatment \times quadrant interaction [$F(1,55) = 6.58$, $p < 0.05$]. Further analyses to understand this interaction showed that Young animals swam somewhat more in the target quadrant (percent of total swim distance in NE quadrant: Aged Control vs. Young $F[1,57] = 3.12$, $p = 0.08$), but swam less in the opposite quadrant (percent of total swim distance in SW quadrant: Aged Control vs. Young $F[1,55] = 4.99$, $p < 0.05$) when compared to Aged Control animals, and there were no sex \times quadrant interactions nor sex \times age \times quadrant interactions. For the Aged Control versus Aged Cognitive Training comparison, there was a main effect of quadrant ($F[1,57] = 147.75$, $p < 0.0001$) with both Aged Control and Aged Cognitive Training groups spending a greater percent of the total distance in the target quadrant; there were no sex \times quadrant interactions nor sex \times cognitive practice \times quadrant interactions, indicating that all aged groups were able to equally localize the platform quadrant by the end of testing.




Fig. 3. Mean \pm SE number of WMI errors committed on the WRAM across the 3 testing blocks (composed of 4 days each), collapsed across all trials, for each group. Abbreviations: SE, standard error; WMI, working memory incorrect; WRAM, water radial-arm maze.

3.3. T-maze

Fig. 6A shows the mean \pm SE number correct for each group for each day, and Fig. 6B shows the mean \pm SE number correct for each group collapsed across days. There was no main effect of age nor an age \times sex interaction, for the T-maze final battery assessment. The Aged Cognitive Training group outperformed the Aged Control group for the T-maze final battery assessment (treatment main effect: $F[1,58] = 23.73$, $p < 0.0001$), an effect which did not interact with sex.

3.4. NGF protein concentrations and correlations with behavior

Fig. 7A shows the mean \pm SE NGF protein values for each brain region for each group. Aged Control animals had higher concentrations of NGF in the frontal cortex ($F[1,71] = 6.97$, $p < 0.05$), parietal cortex ($F[1,71] = 7.52$, $p < 0.05$), temporal cortex ($F[1,71] = 4.42$, $p < 0.05$), and striatum ($F[1,70] = 5.13$, $p < 0.05$; of note, one sample was lost because of experimental error), as compared to Young animals; age did not interact with sex for any brain region. NGF concentrations did not change with cognitive training (i.e., the Aged Cognitive Training group did not differ from the Aged Control group) nor did cognitive training interact with sex for any brain region.

We found a positive correlation between how well male and female Cognitive Training rats performed on the initial exposure to the training task when young (6 months old) and their concentration of hippocampal NGF when aged at 21 months old ($b = 0.10$, $r = 0.59$, $z[19] = 2.72$, $p < 0.05$). However, analyses of males and females alone revealed that this significant correlation was carried by the females (females: $b = 0.13$, $r = 0.79$, $z[8] = 2.39$, $p < 0.05$; males: $b = 0.10$, $r = 0.37$, $z[11] = 1.09$, $p = 0.28$; Fig. 7B). In fact, when females were evaluated separately, we found another positive correlation between how well females performed on the initial exposure to the training task when young, and their concentration of frontal cortex NGF when aged ($b = 0.14$, $r = 0.71$, $z[8] = 2.00$, $p < 0.05$; Fig. 7C). This same correlation was not significant when males were evaluated alone ($b = 0.06$, $r = 0.04$, $z[10] = 0.11$, $p > 0.05$).

3.5. Growth modeling

The linear growth model evaluating the first spatial reference memory T-maze training session, when the Cognitive Training rats were young (6 months old), indicated that female rats had a steeper

Water Radial-Arm Maze: Working Memory Incorrect Errors




Fig. 4. (A) Mean \pm SE number of WMI errors committed across trials during Block 3, the last block of testing. Working memory load increases as trials progress. (B) Mean \pm SE number of WMI errors made at the trial with the greatest working memory demand, trial 4. Aged Cognitive Training females made fewer errors than Aged Control females when working memory load was the highest, whereas Aged Cognitive Training males did not show an enhanced ability to manage this highest working memory load relative to Aged Control males; * $p < 0.05$. Abbreviations: SE, standard error; WMI, working memory incorrect.

slope across days than males, suggesting that females learned the task at a faster rate compared to males ($B_3 = -0.27$, $p < 0.05$, Fig. 8). Of note, females made fewer correct arm entries on day 1 in comparison to males ($B_2 = 0.96$, $p = 0.003$; wherein, B_2 was the amount by which values of performance on day 1 differed between females and males), which would impact the slope of this line. Hence, females may have learned at a faster rate, and thereby reached the same level of performance seen in the males, by the end of the first training session.

We created a global cognitive composite measure for the novel tasks so that we could evaluate relationships between learning when young and learning when old. Specifically, the number of WRAM total errors (i.e., all 3 error measures) and

swim distance on the Morris maze, across all days and trials for both mazes were converted to z-scores. This allowed us to create a single score representing the overall performance on mazes novel to all tested groups during the final maze battery. The z-scores equally weighted the results from the two mazes even though each maze used a different dependent variable metric (i.e., errors committed and swim distance, respectively). For both sexes, rats that learned the training task faster when young demonstrated enhanced performance when aged on the composite measure of the novel tasks during the final maze battery ($B = -3.37$, $p \leq 0.05$, Fig. 8). This positive association between the learning rate of a rat when it was young and its subsequent aged memory performance was strong according to Cohen's




Fig. 5. (A) Learning: mean \pm SE swim distance (cm) across days on the Morris maze, collapsed across all trials. Young animals performed better than Aged Control and Aged Cognitive Training groups. (B) Overnight forgetting: mean \pm SE difference in swim distance (cm) from trial 1 to trial 4 (trial 1–trial 4), collapsed across days 2–5. Aged Cognitive Training females outperformed Aged Control females, an effect not found in males. (C) Probe trial: mean \pm SE percent of total swim distance (cm) in the target NE versus opposite SW quadrants during the probe trial. All groups were able to localize the platform quadrant by the end of testing; * $p < 0.05$. Abbreviations: NE, northeast; SE, standard error; SW, southwest.




Fig. 6. (A) Mean \pm SE number of correct arm choices made across all days of testing on the T-Maze, collapsed across all trials. (B) Mean \pm SE number of correct arm choices made collapsed across all days and trials. Aged Cognitive Training animals performed better than that Aged Control animals, and there were no age effects or interactions with sex. * $p < 0.05$. Abbreviation: SE, standard error.

(Cohen, 1988) conventional criteria/norms in psychological science ($r = 0.83$, $p < 0.05$).

4. Discussion

This longitudinal study supports the tenet that cognitive training can help one remember later in life. Here, we show that intermittent cognitive training throughout adulthood can attenuate age-related memory decline, but only when there is cognitive demand. The present study suggests that there are sex differences in this effect, with females showing benefits of cognitive training on novel tasks evaluated in the final maze battery, and males showing a lack of cognitive training-mediated benefits on novel tasks in the final maze battery. Hence, findings from this study indicate that females are able to transfer the training benefits to an untrained memory domain, and, at least within the parameters used here, males are not. Specifically, in the female rat, cognitive training on a spatial reference memory task every 3 months from 6 to 18 months of age enhanced later cognitive performance when aged, 21 months old, on tasks that

were new to the animal. In females, on the final maze battery, a history of cognitive training from young adulthood to old age attenuated age-related spatial working memory decline on a novel task (WRAM, as shown for the latter portion of testing for one of the two working memory measures examined on this task, WMI), protected against overnight forgetting on a novel spatial reference memory task (Morris maze), and improved performance on the practiced spatial reference memory training task (T-maze). In males, cognitive training-induced enhancements were only seen for the practiced spatial reference memory training task (T-maze). It is noteworthy that the treatment \times day interaction for the spatial reference memory Morris maze task initially indicated that cognitive training might have helped both sexes; however, noting the interactions with sex and upon further analyses, it became clear that the cognitive training benefit was due to the females showing enhanced performance for overnight retention for the latter three overnight intervals. Here, we also show that cognitive demand was necessary for cognitive training effects to be realized, since rats that received only the procedural testing components for training showed no




Fig. 7. (A) Mean \pm SE NGF protein concentration (pg/mg of tissue) for each brain region. The Young group had lower concentrations of NGF in the frontal cortex, parietal cortex, temporal cortex, and striatum. (B) Represented is the scatterplot of ventral hippocampal (CA1/2) concentrations of NGF (pg/mg tissue) in aged (23 month old) females (black filled triangles) and males (gray filled circles), and the number of correct arm choices made on the T-Maze during the first training session when young, 17 months earlier (at 6 months old). (C) Represented is the scatterplot of frontal cortex concentrations of NGF (pg/mg tissue) in aged (23 months old) females (black filled triangles) and males (gray filled circles), and the number of correct arm choices made on the T-Maze during the first training session when young, 17 months earlier (at 6 month old). Higher concentrations of NGF in the aged ventral hippocampus (CA1/2) and frontal cortex significantly correlated with better initial spatial reference memory performance when young, in females only, as represented by the black solid line; * $p < 0.05$. Abbreviations: F, females; M, males; NGF, nerve growth factor; SE, standard error.




Fig. 8. Mean \pm SE number of correct arm choices across days on the T-Maze during the first maze session. Specifically, the graph displays the learning curve of male and female Cognitive Training rats during their initial/first cognitive training session, when they were 6 months old. The linear growth model demonstrated that female rats had a steeper slope across days than males, suggesting that females learned the task at a faster rate ($B_3 = -0.27, p < 0.05$). Because females made fewer correct arm entries on day 1 in comparison to males ($B_2 = 0.96, p = 0.003$, assessing the intercept), this impacted the slope of the line. A global cognitive composite measure was calculated for the novel tasks during the final maze battery so that relationships between learning when young and learning when old, could be evaluated. The number of WRAM total errors and swim distance on the Morris maze (all days and trials for both mazes), were converted to z-scores. These z-scores equally weighted the results from the two mazes, although each maze used a different dependent variable metric (errors committed and swim distance, respectively). For both sexes, rats that learned the training task faster demonstrated enhanced performance when aged on the composite measure of the novel tasks tested during the final maze battery ($B = -3.37, p \leq 0.05$). This association between the learning rate when young and subsequent aged memory performance was strong according to Cohen's (Cohen, 1988) conventional norms in psychological science ($r = 0.83, p < 0.05$). Abbreviations: SE, standard error; WRAM, water radial-arm maze.

improvement on the final maze battery. Moreover, male and female rats that had a faster rate of learning when young demonstrated better global performance on the novel tasks when aged. We also show here that NGF concentrations increased in several brain regions as aging ensued, and NGF protein levels in the hippocampus and frontal cortex related to performance on the training task only in young females.

4.1. Sex interacts with the impact of intermittent adult cognitive training

The data presented here suggest that male and female rats differ in their response to intermittent cognitive training administered across adulthood. For the parameters used in the present study, females benefited from cognitive training and were the only sex to demonstrate enhanced performance, relative to controls, on novel tasks. That females benefit from cognitive training when aged is corroborated by studies in young non-human primates (Lacreuse et al., 2005) and aged rodents (Ando and Ohashi, 1991); however, in this rodent study, there was no male comparison group to permit sex difference evaluations. In the current study, although females did learn the training task faster than males, all rats reached and maintained near asymptotic performance by the end of their first cognitive training session, thereby limiting the possibility that the sex differences found during the final maze battery were due to differential learning of the cognitive training task. Numerous studies in males have demonstrated training-mediated enhancements of memory (e.g., Bierley et al., 1986; Light et al., 2010; Markowska and Savonenko, 2002; Matzel et al., 2011; Vicens et al., 2003); thus, it might initially appear surprising that we did not see more robust effects in males. However, most of these studies used the same or similar tasks for both the practice task and final assessment. This does correspond to our findings here, as we found that both males and females showed cognitive training-induced enhancements on the T-maze, which was used for both training and again testing on the final maze battery. Thus, it may be that both males and females

show training-induced enhancements on the same task used for training, but that females can transfer these benefits to other memory domains, showing benefits on other nonpracticed tasks. Taken together with data suggesting females exhibit greater brain plasticity (for review see Fitch and Bimonte, 2002; Juraska, 1986), it may be that the training effects found previously in males might in part be because of "practice effects" and not true training-mediated enhancements to aged performance. It remains to be determined whether sex differences would still exist with cognitive training if the cognitive training sessions were initiated more frequently or with a different level of intensity. In addition, we would be remiss if we did not acknowledge that different estropause states may have impacted the effects of cognitive training. Indeed, estropause status has been found to influence cognition in aged rats (see Markowska, 1999; Warren and Juraska, 2000). In the present study, vaginal smears demonstrated that nearly all aged females were in an anestrus state, whereas the few found to be in a non-anestrus state were distributed throughout the various aged groups. Hence, although we cannot rule out possible estropause state-mediated alterations in the effect of cognitive training, the likelihood of its impact on our cognitive training-induced sex difference findings is likely not largely accounting for the effects. However, of note, a systematic evaluation of interactions between state of estropause and efficacy of cognitive training during aging is necessary to directly test this question and could yield insight into the trajectory of sex differences in cognition as aging ensues, with and without cognitive practice.

4.2. Generalizing the impact of cognitive training: effects of adult intermittent cognitive training transfer to untrained memory domains

Cognitive training protected aged female rats against the age-related decrements when handling an increasing working memory load on a novel working memory task, the WRAM. The WRAM is sensitive to rodent age-related spatial working memory decline

(Bimonte-Nelson et al., 2003c; Bimonte et al., 2002, 2003), an effect we also demonstrate here. Spatial reference memory cognitive training also attenuated age-related memory decline on the novel spatial reference memory task, the Morris maze. This finding is corroborated by rodent and non-human primate experiments showing that cognitive training enhances later performance on tasks assessing working and reference memory, as well as selective attention (Bierley et al., 1986; Lacreuse et al., 2005; Markowska and Savonenko, 2002; Matzel et al., 2011). We also show here that spatial reference memory cognitive training enhanced performance when aged on the training task, the reference memory T-Maze. Aged Cognitive Training animals performed better on the T-Maze cognitive training task during the final maze battery when compared with Aged Control animals, and Aged Naïve animals did not differ from Aged Swim Only animals that received only the procedural components of testing, indicating that cognitive training-induced enhancements are found even when control is imposed over the procedural components of testing. Of note, an age effect was not seen on the T-Maze during the final maze battery (i.e., the Aged Control vs. Young comparison). This lack of an age effect corroborates previous research wherein no age effects were found on a land T-Maze; only the type of solving strategy differed by age (Barnes et al., 1980).

Because we found that cognitive training in females can protect against age-related decline, and that these effects were observed on mazes and environments novel to the group that had cognitive training, the effects could not simply be because of familiarity or prior experience with the specific task. The enhanced performance of the Cognitive Training group (both males and females) relative to the Control group on the T-maze during the final maze battery could be, in part, because of retention and not relearning, of the escape platform location since their last cognitive training session. Scores were essentially asymptotic with little to no variation after the first cognitive training session; this level of performance persisted to the last training session at 18 months. Collectively, these data suggest that previous cognitive training enhances performance on the training task for both males and females.

4.3. Evidence that long-term adult cognitive training-induced enhancements require cognitive demand

Our data suggest that memory protection from cognitive training is due to the cognitive demand of the task and not exposure to the procedural components of maze testing. During training, the Swim Only group had a single swim path without any choice points during the trials, thereby minimizing cognitive demand; this group “won” every time. This group, receiving only the procedural aspects of the task, including being moved from the colony to the testing room, handling required of testing, swimming, and escaping on the platform, exhibited no protection from age-related memory decline and were virtually indistinguishable from aged controls that sat in their home cages until the final maze battery (the Aged Naïve group). This finding is corroborated by recent studies where cognitive demand was necessary for enhancements of selective attention in young adult and aged male mice (Light et al., 2010; Matzel et al., 2011).

4.4. A faster rate of learning when young relates to superior global memory when old

The rate at which a male or female rat learned the cognitive training task during the first exposure at 6 months of age was related to cognitive performance when aged; indeed, a faster rate of

learning when young corresponded to better memory on the novel mazes when aged. In the context of the Hebb (1949) postulate describing the adaptation of neurons during learning and engram creation, it is tempting to speculate that learning when young strengthens brain networks so that enhanced learning and memory can occur when aged. Better learning when young might organize the brain in such a fashion that results in better learning and memory, later in life. Indeed, one might presume that such a tenet could underscore the idea that “cells that fire together, wire together” (Goodman and Shatz, 1993; Hebb, 1949), with prior learning providing greater cognitive reserve, in turn leading to better memory when challenged with a potentially cognitively-compromising situation, such as aging.

4.5. NGF protein concentrations were higher in aged rats, and concentrations in the frontal cortex and hippocampus when aged related to cognitive ability when young in females only

We evaluated NGF protein concentrations in the frontal cortex, parietal cortex, temporal cortex, striatum, entorhinal cortex, and ventral hippocampus (CA1/2), brain regions known to subserve cognitive and motor functions (Hasselmo, 2006; Woolf, 1991). Survival and maintenance of cholinergic neurons, which are critical for memory processes are dependent upon neurotrophins including NGF (Bimonte-Nelson et al., 2008, 2003a; Granholm 2000; Levi-Montalcini, 1987; Siegel and Chauhan, 2000; Woolf, 1991). We observed here an age-related increase in NGF protein concentrations in the frontal cortex, parietal cortex, temporal cortex, and striatum, although cognitive training did not alter NGF concentrations in the brain regions assayed. The current NGF findings agree with our previous data showing that both male and female aged Fischer 344 rats exhibited higher NGF protein levels in the striatum and frontal cortex when compared to young counterparts (Bimonte-Nelson et al., 2008, 2003a; Koh and Loy, 1988; Larkfors et al., 1987).

In the present study, in females only, better performance on the cognitive training task when young was associated with higher concentrations of frontal cortex and hippocampal NGF protein concentrations when aged. These findings correspond to studies showing that NGF infusions into the aged rat brain can attenuate age-related memory deficits (Markowska et al., 1996), and environmental enrichment from 2 to 14 months of age increases cortical and hippocampal NGF concomitant with enhancements in spatial memory (Ickes et al., 2000; Pham et al., 1999). It is tempting to speculate that young females who had better performance when young likely had a better cognitive reserve or had greater neuroplasticity, because those that learned faster when young had better global novel performance when aged. Hence, increased NGF protein in these animals, instead of solely responding to an age-related insult or a perturbed system (Bimonte et al., 2003; unpublished observations; Sugaya et al., 1998), might reflect a beneficial brain environment subserving plasticity and strengthening neuronal networks (see Levi-Montalcini, 1987).

Disclosure statement

None of the authors have any conflicts of interest.

Acknowledgements

This research was supported in part by funds from National Institute on Aging grants awarded to Heather A. Bimonte-Nelson (R03AG026137, R01AG028084), state of Arizona, ADHS and the Arizona Alzheimer's Disease Core Center. The authors wish to thank Dr. Clark Presson and Dr. Gene Alexander for discussion of this

project in thesis form, as well as Dr. Victor Denenberg and Dr. Gordon Winocur for critical feedback and discussion on the experimental design. They thank Donovan Terry and Britny Sundin for their excellent experimental assistance.

References

- Acosta, J.I., Mayer, L., Talboom, J.S., Tsang, C.W., Smith, C.J., Enders, C.K., Bimonte-Nelson, H.A., 2009a. Transitional versus surgical menopause in a rodent model: etiology of ovarian hormone loss impacts memory and the acetylcholine system. *Endocrinology* 150, 4248–4259.
- Acosta, J.I., Mayer, L., Talboom, J.S., Zay, C., Scheldrup, M., Castillo, J., Demers, L.M., Enders, C.K., Bimonte-Nelson, H.A., 2009b. Premarin improves memory, prevents scopolamine-induced amnesia and increases number of basal forebrain choline acetyltransferase positive cells in middle-aged surgically menopausal rats. *Horm. Behav.* 55, 454–464.
- Ando, S., Ohashi, Y., 1991. Longitudinal study on age-related changes of working and reference memory in the rat. *Neurosci. Lett.* 128, 17–20.
- Anguera, J.A., Boccanfuso, J., Rintoul, J.L., Al-Hashimi, O., Faraji, F., Janowich, J., Kong, E., Larraburo, Y., Rolle, C., Johnston, E., Gazzaley, A., 2013. Video game training enhances cognitive control in older adults. *Nature* 501, 97–101. <http://dx.doi.org/10.1038/nature12486>. <http://www.nature.com/nature/journal/v501/n7465/abs/nature12486.html#supplementary-information>.
- Association, A.V.M. 2007. AVMA Guidelines on Euthanasia (Formerly Report of the AVMA Panel on Euthanasia): Jun 2007 update. Available from http://www.avma.org/issues/animal_welfare/euthanasia.pdf.
- Backman, C., Rose, G.M., Hoffer, B.J., Henry, M.A., Bartus, R.T., Friden, P., Granholm, A.C., 1996. Systemic administration of a nerve growth factor conjugate reverses age-related cognitive dysfunction and prevents cholinergic neuron atrophy. *J. Neurosci.* 16, 5437–5442.
- Ballard, C., Gauthier, S., Corbett, A., Brayne, C., Aarsland, D., Jones, E., 2011. Alzheimer's disease. *Lancet* 377, 1019–1031.
- Barnes, C.A., Nadel, L., Honig, W.K., 1980. Spatial memory deficit in senescent rats. *Can. J. Psychol.* 34, 29–39.
- Bierley, R.A., Rixen, G.J., Troster, A.I., Beatty, W.W., 1986. Preserved spatial memory in old rats survives 10 months without training. *Behav. Neural Biol.* 45, 223–229.
- Bimonte-Nelson, H.A., Granholm, A.C., Nelson, M.E., Moore, A.B., 2008. Patterns of neurotrophin protein levels in male and female Fischer 344 rats from adulthood to senescence: how young is “young” and how old is “old”? *Exp. Aging Res.* 34, 13–26.
- Bimonte-Nelson, H.A., Hunter, C.L., Nelson, M.E., Granholm, A.-C.E., 2003a. Frontal cortex BDNF levels correlate with working memory in an animal model of Down syndrome. *Behav. Brain Res.* 139, 47–57.
- Bimonte-Nelson, H.A., Singleton, R.S., Hunter, C.L., Price, K.L., Moore, A.B., Granholm, A.C., 2003b. Ovarian hormones and cognition in the aged female rat: I. Long-term, but not short-term, ovariectomy enhances spatial performance. *Behav. Neurosci.* 117, 1395–1406.
- Bimonte-Nelson, H.A., Singleton, R.S., Nelson, M.E., Eckman, C.B., Barber, J., Scott, T.Y., Granholm, A.C., 2003c. Testosterone, but not nonaromatizable dihydrotestosterone, improves working memory and alters nerve growth factor levels in aged male rats. *Exp. Neurol.* 181, 301–312.
- Bimonte-Nelson, H.A., Singleton, R.S., Williams, B.J., Granholm, A.C., 2004. Ovarian hormones and cognition in the aged female rat: II. progesterone supplementation reverses the cognitive enhancing effects of ovariectomy. *Behav. Neurosci.* 118, 707–714.
- Bimonte, H.A., Denenberg, V.H., 1999. Estradiol facilitates performance as working memory load increases. *Psychoneuroendocrinology* 24, 161–173.
- Bimonte, H.A., Denenberg, V.H., 2000. Sex differences in vicarious trial-and-error behavior during radial arm maze learning. *Physiol. Behav.* 68, 495–499.
- Bimonte, H.A., Granholm, A.C., Seo, H., Isacson, O., 2002. Spatial memory testing decreases hippocampal amyloid precursor protein in young, but not aged, female rats. *Neurosci. Lett.* 328, 50–54.
- Bimonte, H.A., Hyde, L.A., Hoplight, B.J., Denenberg, V.H., 2000. In two species, females exhibit superior working memory and inferior reference memory on the water radial-arm maze. *Physiol. Behav.* 70, 311–317.
- Bimonte, H.A., Nelson, M.E., Granholm, A.C., 2003. Age-related deficits as working memory load increases: relationships with growth factors. *Neurobiol. Aging* 24, 37–48.
- Bowirrat, A., Friedland, R.P., Farrer, L., Baldwin, C., Korczyn, A., 2002. Genetic and environmental risk factors for Alzheimer's disease in Israeli Arabs. *J. Mol. Neurosci.* 19, 239–245.
- Braden, B.B., Talboom, J.S., Crain, I.D., Simard, A.R., Lukas, R.J., Prokai, L., Scheldrup, M.R., Bowman, B.L., Bimonte-Nelson, H.A., 2010. Medroxyprogesterone acetate impairs memory and alters the GABAergic system in aged surgically menopausal rats. *Neurobiol. Learn. Mem.* 93, 444–453.
- Braden, B.B., Garcia, A.N., Mennenga, S.E., Prokai, L., Villa, S.R., Acosta, J.I., Lefort, N., Simard, A.R., Bimonte-Nelson, H.A., 2011. Cognitive-impairing effects of medroxyprogesterone acetate in the rat: independent and interactive effects across time. *Psychopharmacology (Berl)* 218, 405–418.
- Chiang, K., Koo, E.H., 2014. Emerging therapeutics for Alzheimer's disease. *Annu. Rev. Pharmacol. Toxicol.* 54, 381–405.
- Cohen, J., 1988. *Statistical Power Analysis for the Behavioral Sciences*, second ed. L. Erlbaum Associates, Hillsdale, NJ.
- De Ronchi, D., Fratiglioni, L., Rucci, P., Paternico, A., Graziani, S., Dalmonte, E., 1998. The effect of education on dementia occurrence in an Italian population with middle to high socioeconomic status. *Neurology* 50, 1231–1238.
- Denenberg, V.H., Sherman, G.F., Schrott, L.M., Rosen, G.D., Galaburda, A.M., 1991. Spatial learning, discrimination learning, paw preference and neocortical ectopias in two autoimmune strains of mice. *Brain Res.* 562, 98–104.
- Denenberg, V.H., Talgo, N.W., Schrott, L.M., Kenner, G.H., 1990. A computer-aided procedure for measuring discrimination learning. *Physiol. Behav.* 47, 1031–1034.
- Digby, G.J., Noetzel, M.J., Bubser, M., Utley, T.J., Walker, A.G., Byun, N.E., Lebois, E.P., Xiang, Z., Sheffler, D.J., Cho, H.P., Davis, A.A., Nemirovsky, N.E., Mennenga, S.E., Camp, B.W., Bimonte-Nelson, H.A., Bode, J., Italiano, K., Morrison, R., Daniels, J.S., Niswender, C.M., Olive, M.F., Lindsley, C.W., Jones, C.K., Conn, P.J., 2012. Novel allosteric agonists of M1 muscarinic acetylcholine receptors induce brain region-specific responses that correspond with behavioral effects in animal models. *J. Neurosci.* 32, 8532–8544.
- Drachman, D.A., Leavitt, J., 1974. Human memory and the cholinergic system. A relationship to aging? *Arch. Neurol.* 30, 113–121.
- Eckerman, D.A., Gordon, W.A., Edwards, J.D., MacPhail, R.C., Gage, M.I., 1980. Effects of scopolamine, pentobarbital, and amphetamine on radial arm maze performance in the rat. *Pharmacol. Biochem. Behav.* 12, 595–602.
- Engler-Chiurazzi, E., Tsang, C., Nonnenmacher, S., Liang, W.S., Corneveaux, J.J., Prokai, L., Huentelman, M.J., Bimonte-Nelson, H.A., 2011. Tonic Premarin dose-dependently enhances memory, affects neurotrophin protein levels and alters gene expression in middle-aged rats. *Neurobiol. Aging*. <http://dx.doi.org/10.1016/j.neurobiolaging.2009.09.005>.
- Fischer, W., Victorin, K., Bjorklund, A., Williams, L.R., Varon, S., Gage, F.H., 1987. Amelioration of cholinergic neuron atrophy and spatial memory impairment in aged rats by nerve growth factor. *Nature* 329, 65–68.
- Fitch, R.H., Bimonte, H.A., 2002. Hormones, brain, and behavior: putative biological contributions to cognitive sex differences. In: McGillicuddy-De Lisi, A.V., R., D.L. (Eds.), *Biology, society, and behavior: the development of sex differences in cognition*. Ablex Publishing Westport, CT, pp. 55–91.
- Fritsch, T., McClelland, M.J., Smyth, K.A., Ogrocki, P.K., 2002. Effects of educational attainment and occupational status on cognitive and functional decline in persons with Alzheimer-type dementia. *Int. Psychogeriatr.* 14, 347–363.
- Gatz, M., Svedberg, P., Pedersen, N.L., Mortimer, J.A., Berg, S., Johansson, B., 2001. Education and the risk of Alzheimer's disease: findings from the study of dementia in Swedish twins. *J. Gerontol. B Psychol. Sci. Soc. Sci.* 56, P292–P300.
- Gibbs, R.B., Gabor, R., Cox, T., Johnson, D.A., 2004. Effects of raloxifene and estradiol on hippocampal acetylcholine release and spatial learning in the rat. *Psychoneuroendocrinology* 29, 741–748.
- Goldman, J.M., Murr, A.S., Cooper, R.L., 2007. The rodent estrous cycle: characterization of vaginal cytology and its utility in toxicological studies. *Birth Defects Res. B Dev. Reprod. Toxicol.* 80, 84–97.
- Goodman, C.S., Shatz, C.J., 1993. Developmental mechanisms that generate precise patterns of neuronal connectivity. *Cell* 72, 77–98.
- Granholm, A.C., 2000. Oestrogen and nerve growth factor—neuroprotection and repair in Alzheimer's disease. *Expert Opin. Investig. Drugs* 9, 685–694.
- Gustilo, M.C., Markowska, A.L., Breckler, S.J., Fleischman, C.A., Price, D.L., Koliatos, V.E., 1999. Evidence that nerve growth factor influences recent memory through structural changes in septohippocampal cholinergic neurons. *J. Comp. Neurol.* 405, 491–507.
- Hasselmo, M.E., 2006. The role of acetylcholine in learning and memory. *Curr. Opin. Neurobiol.* 16, 710–715.
- Hebb, D.O., 1949. *The Organization of Behavior: A Neuropsychological Theory*. Wiley, New York.
- Hyde, L.A., Sherman, G.F., Hoplight, B.J., Denenberg, V.H., 2000. Working memory deficits in BXS mice with neocortical ectopias. *Physiol. Behav.* 70, 1–5.
- Ickes, B.R., Pham, T.M., Sanders, L.A., Albeck, D.S., Mohammed, A.H., Granholm, A.C., 2000. Long-term environmental enrichment leads to regional increases in neurotrophin levels in rat brain. *Exp. Neurol.* 164, 45–52.
- Jarrard, L.E., Okaichi, H., Steward, O., Goldschmidt, R.B., 1984. On the role of hippocampal connections in the performance of place and cue tasks: comparisons with damage to hippocampus. *Behav. Neurosci.* 98, 946–954.
- Juraska, J.M., 1986. Sex differences in developmental plasticity of behavior and the brain. In: Greenough, W.T., Juraska, J.M. (Eds.), *Developmental Neuropsychobiology*. Academic Press, Inc., Orlando, FL, pp. 409–422.
- Keppel, G., Wickens, T.D., 2004. *Design and Analysis: A Researcher's Handbook*, fourth ed. Pearson Prentice Hall, Upper Saddle River, NJ.
- Koh, S., Loy, R., 1988. Age-related loss of nerve growth factor sensitivity in rat basal forebrain neurons. *Brain Res.* 440, 396–401.
- Lacrouse, A., Kim, C.B., Rosene, D.L., Kiliany, R.J., Moss, M.B., Moore, T.L., Chennareddi, L., Herndon, J.G., 2005. Sex, age, and training modulate spatial memory in the rhesus monkey (*Macaca mulatta*). *Behav. Neurosci.* 119, 118–126.
- Landau, S.M., Marks, S.M., Mormino, E.C., Rabinovici, G.D., Oh, H., O'Neil, J.P., Wilson, R.S., Jagust, W.J., 2012. Association of lifetime cognitive engagement and low beta-amyloid deposition. *Arch. Neurol.* <http://dx.doi.org/10.1001/archneurol.2011.2748>.
- Larkfors, L., Ebendal, T., Whittemore, S.R., Persson, H., Hoffer, B., Olson, L., 1987. Decreased level of nerve growth factor (NGF) and its messenger RNA in the aged rat brain. *Brain Res.* 427, 55–60.

- Letenneur, L., Gilleron, V., Commenges, D., Helmer, C., Orgogozo, J.M., Dartigues, J.F., 1999. Are sex and educational level independent predictors of dementia and Alzheimer's disease? Incidence data from the PAQUID project. *J. Neurol. Neurosurg. Psychiatry* 66, 177–183.
- Levi-Montalcini, R., 1987. The nerve growth factor 35 years later. *Science* 237, 1154–1162.
- Light, K.R., Kolata, S., Wass, C., Denman-Brice, A., Zagalsky, R., Matzel, L.D., 2010. Working memory training promotes general cognitive abilities in genetically heterogeneous mice. *Curr. Biol.* 20, 777–782.
- Lindsay, J., Laurin, D., Verreault, R., Hebert, R., Helliwell, B., Hill, G.B., McDowell, I., 2002. Risk factors for Alzheimer's disease: a prospective analysis from the Canadian Study of Health and Aging. *Am. J. Epidemiol.* 156, 445–453.
- Markowska, A.L., 1999. Sex dimorphisms in the rate of age-related decline in spatial memory: relevance to alterations in the estrous cycle. *J. Neurosci.* 19, 8122–8133.
- Markowska, A.L., Koliatsos, V.E., Breckler, S.J., Price, D.L., Olton, D.S., 1994. Human nerve growth factor improves spatial memory in aged but not in young rats. *J. Neurosci.* 14, 4815–4824.
- Markowska, A.L., Price, D., Koliatsos, V.E., 1996. Selective effects of nerve growth factor on spatial recent memory as assessed by a delayed nonmatching-to-position task in the water maze. *J. Neurosci.* 16, 3541–3548.
- Markowska, A.L., Savonenko, A.V., 2002. Protective effect of practice on cognition during aging: implications for predictive characteristics of performance and efficacy of practice. *Neurobiol. Learn. Mem.* 78, 294–320.
- Matzel, L.D., Light, K.R., Wass, C., Colas-Zelin, D., Denman-Brice, A., Waddell, A.C., Kolata, S., 2011. Longitudinal attentional engagement rescues mice from age-related cognitive declines and cognitive inflexibility. *Learn. Mem.* 18, 345–356.
- Morris, R.G., Garrud, P., Rawlins, J.N., O'Keefe, J., 1982. Place navigation impaired in rats with hippocampal lesions. *Nature* 297, 681–683.
- Mufson, E.J., Conner, J.M., Kordower, J.H., 1995. Nerve growth factor in Alzheimer's disease: defective retrograde transport to nucleus basalis. *Neuroreport* 6, 1063–1066.
- National Research Council (U.S.) Committee for the Update of the Guide for the Care and Use of Laboratory Animals, Institute for Laboratory Animal Research (U.S.) National Academies Press (U.S.), 2011. *Guide for the Care and Use of Laboratory Animals*. National Academies Press, Washington, DC xxv, 220 p.
- Ott, A., van Rossum, C.T., van Harskamp, F., van de Mheen, H., Hofman, A., Breteler, M.M., 1999. Education and the incidence of dementia in a large population-based study: the Rotterdam Study. *Neurology* 52, 663–666.
- Paxinos, G., Watson, C., 2005. *The Rat Brain in Stereotaxic Coordinates*, fifth ed. Elsevier Academic Press, Amsterdam; Boston.
- Petersen, R.C., Stevens, J.C., Ganguli, M., Tangalos, E.G., Cummings, J.L., DeKosky, S.T., 2001. Practice parameter: early detection of dementia: mild cognitive impairment (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 56, 1133–1142.
- Pham, T.M., Ickes, B., Albeck, D., Soderstrom, S., Granholm, A.C., Mohammed, A.H., 1999. Changes in brain nerve growth factor levels and nerve growth factor receptors in rats exposed to environmental enrichment for one year. *Neuroscience* 94, 279–286.
- Rupniak, N.M., Field, M.J., Samson, N.A., Steventon, M.J., Iversen, S.D., 1990. Direct comparison of cognitive facilitation by physostigmine and tetrahydroaminoacridine in two primate models. *Neurobiol. Aging* 11, 609–613.
- Scali, C., Casamenti, F., Pazzagli, M., Bartolini, L., Pepeu, G., 1994. Nerve growth factor increases extracellular acetylcholine levels in the parietal cortex and hippocampus of aged rats and restores object recognition. *Neurosci. Lett.* 170, 117–120.
- Shimamura, A., Berry, J., Mangels, J., Rustig, C., 1995. Memory and cognitive abilities in University professors. *Psychol. Sci.* 6, 271–277.
- Shrestha, L.B., 2005. *Life Expectancy in the United States*. Congressional Information Service, Library of Congress.
- Siegel, G.J., Chauhan, N.B., 2000. Neurotrophic factors in Alzheimer's and Parkinson's disease brain. *Brain Res. Brain Res. Rev.* 33, 199–227.
- Singer, J.D., Willett, J.B., 2003. *Applied longitudinal data analysis: modeling change and event occurrence*. Oxford University Press, Oxford; New York.
- Snowdon, D.A., 2003. Healthy aging and dementia: findings from the Nun Study. *Ann. Intern. Med.* 139, 450–454.
- Sugaya, K., Greene, R., Personett, D., Robbins, M., Kent, C., Bryan, D., Skiba, E., Gallagher, M., McKinney, M., 1998. Septo-hippocampal cholinergic and neurotrophin markers in age-induced cognitive decline. *Neurobiol. Aging* 19, 351–361.
- Talboom, J.S., Engler-Chiurazzi, E.B., Whiteaker, P., Simard, A.R., Lukas, R., Acosta, J.I., Prokai, L., Bimonte-Nelson, H.A., 2010. A component of Premarin® enhances multiple cognitive functions and influences nicotinic receptor expression. *Horm. Behav.* 58, 917–928.
- Vicens, P., Carrasco, M.C., Redolat, R., 2003. Effects of early training and nicotine treatment on the performance of male NMRI mice in the water maze. *Neural Plast.* 10, 303–317.
- Vincent, G.K., Velkoff, V.A., 2010. *The Next Four Decades: The Older Population in the United States: 2010 to 2050*. US Department of Commerce, Economics and Statistics Administration, US Census Bureau.
- Warren, S.G., Juraska, J.M., 2000. Sex differences and estropausal phase effects on water maze performance in aged rats. *Neurobiol. Learn. Mem.* 74, 229–240.
- Whalley, L.J., Starr, J.M., Athawes, R., Hunter, D., Pattie, A., Deary, I.J., 2000. Childhood mental ability and dementia. *Neurology* 55, 1455–1459.
- Woolf, N.J., 1991. Cholinergic systems in mammalian brain and spinal cord. *Prog. Neurobiol.* 37, 475–524.