



Developmental switch in requirement for PKA RII β in NMDA-receptor-dependent synaptic plasticity at Schaffer collateral to CA1 pyramidal cell synapses

Yupeng Yang^{3,1}, Koichi Takeuchi³, Alma Rodenas-Ruano, Yukihiro Takayasu², Michael V.L. Bennett, R. Suzanne Zukin^{*}

Dominick P. Purpura Department of Neuroscience, Kennedy Center Room 602B, Albert Einstein College of Medicine, 1300 Morris Park Avenue, Bronx, NY 10461, USA

ARTICLE INFO

Article history:

Received 21 June 2008

Received in revised form 1 August 2008

Accepted 1 August 2008

Keywords:

PKA

PKA type II regulatory subunit

PKA RII β knockout mice

NMDA receptors

Synaptic plasticity

Long-term potentiation

Long-term depression

Hippocampus

CA1 synapses

ABSTRACT

The cAMP/protein kinase A (PKA) signaling cascade is crucial for synaptic plasticity in a wide variety of species. PKA regulates Ca²⁺ permeation through NMDA receptors (NMDARs) and induction of NMDAR-dependent synaptic plasticity at the Schaffer collateral to CA1 pyramidal cell synapse. Whereas the role of PKA in induction of NMDAR-dependent LTP at CA1 synapses is established, the identity of PKA isoforms involved in this phenomenon is less clear. Here we report that protein synthesis-independent NMDAR-dependent LTP at the Schaffer collateral–CA1 synapse in the hippocampus is deficient, but NMDAR-dependent LTD is normal, in young (postnatal day 10 (P10)–P14) mice lacking PKA RII β , the PKA regulatory protein that links PKA to NMDARs at synaptic sites. In contrast, in young adult (P21–P28) mice lacking PKA RII β , LTP is normal and LTD is abolished. These findings indicate that distinct PKA isoforms may subserve distinct forms of synaptic plasticity and are consistent with a developmental switch in the signaling cascades required for LTP induction.

© 2008 Elsevier Ltd. All rights reserved.

1. Introduction

N-methyl-D-aspartate (NMDA) receptor (NMDAR)-mediated Ca²⁺ influx is essential for synaptogenesis, experience-dependent synaptic remodeling and long-lasting changes in synaptic efficacy such as NMDAR-dependent long-term potentiation and depression (Dingledine et al., 1999; Carroll and Zukin, 2002; Cull-Candy and Leszkiewicz, 2004; Malenka and Bear, 2004; Collingridge et al., 2004; Lau and Zukin, 2007). Synaptic NMDARs are localized to the postsynaptic density (PSD), where they are physically anchored and spatially restricted by scaffolding proteins such as the PSD protein of 95 kDa (PSD-95), which link upstream and downstream signaling molecules to NMDARs (Scannevin and Huganir, 2000; Lau and Zukin, 2007). Coupling of PKA and protein phosphatase-1 (PPI) to synaptic NMDARs by A-kinase adaptor proteins (AKAPs) enables bi-directional regulation of NMDAR channel activity by PKA (Westphal et al., 1999; Wong and Scott, 2004). The NMDAR-mediated rise in

postsynaptic Ca²⁺ activates a network of kinases and phosphatases that promote persistent changes in synaptic strength.

The cAMP/PKA signaling cascade is crucial for synaptic plasticity in a wide variety of species. The role of cAMP/PKA signaling in the induction (Otmakhova et al., 2000; Skeberdis et al., 2006) and late, protein synthesis-dependent phase (Huang and Kandel, 1994; Qi et al., 1996; Abel et al., 1997) of NMDAR-dependent LTP at hippocampal Schaffer collateral–CA1 (Sch–CA1) synapses has attracted considerable attention. Moreover, pharmacological blockade of PKA (Yasuda et al., 2003) reduces induction of NMDAR-dependent LTP at Sch–CA1 synapses, suggesting a “gating” role for PKA in LTP (Blitzer et al., 1998). NMDARs are molecular targets of PKA phosphorylation (Leonard and Hell, 1997; Tingley et al., 1997); however, the effects of PKA on NMDAR function are less clear. Activation of PKA or inhibition of protein phosphatase I potentiates NMDAR-mediated currents (Cerne et al., 1993; Wang et al., 1994; Raman et al., 1996; Blank et al., 1997; Snyder et al., 1998; Skeberdis et al., 2006). Moreover, targeted mutation or disruption of AKAP150, which links PKA to NMDARs and AMPARs, impairs protein synthesis-dependent (Nie et al., 2007) and independent LTP (Lu et al., 2007) at Sch–CA1 synapses. PKA promotes Ca²⁺ permeation through NMDARs, activity-dependent, NMDAR-mediated Ca²⁺ signaling in dendritic spines and induction of NMDAR-dependent LTP at Sch–CA1 synapses (Skeberdis et al., 2006). These findings link PKA-dependent synaptic plasticity to Ca²⁺ signaling in spines and

* Corresponding author. Tel.: +1 718 430 2160; fax: +1 718 430 8932.

E-mail address: zukin@acom.yu.edu (R.S. Zukin).

¹ Present address: University of Science and Technology of China, Hefei, Anhui 230026, PR China.

² Present address: Department of Otolaryngology, Gunma University Graduate School of Medicine, Maebashi, Gunma 371-8511, Japan.

³ These authors contributed equally to the paper.

provide a novel mechanism whereby PKA regulates induction of LTP and LTD.

An emerging view is that the specificity of PKA actions on multiple downstream proteins is dictated by the molecular diversity in PKA catalytic and regulatory subunits, which are differentially expressed in specific neuronal populations and spatially compartmentalized within differing microdomains of the neuron (Brandon et al., 1997). The PKA holoenzyme has four different regulatory subunits. The type II regulatory subunit of PKA (RII) localizes to the postsynaptic density (PSD) (Carr et al., 1992; Colledge and Scott, 1999; Colledge et al., 2000), where it links via A-kinase anchoring proteins (AKAPs) to NMDARs and AMPARs (Carr et al., 1992; Colledge et al., 2000; Oliveria et al., 2003) and modulates synaptic AMPAR phosphorylation and trafficking critical to synaptic plasticity and remodeling (Malinow and Malenka, 2002; Collingridge et al., 2004). Whereas the role of the PKA type I regulatory subunit in hippocampal synaptic plasticity is well-established (Brandon et al., 1995; Huang et al., 1995), the role of the type II regulatory subunit is less clear.

The present study was undertaken to define the role of the PKA RII β subunit in NMDAR-dependent synaptic plasticity at Sch–CA1 synapses of the hippocampus. Here we show that NMDAR-dependent LTD is impaired in adult (P21–28) PKA RII $\beta^{-/-}$ mice; whereas excitatory synaptic transmission (assessed by input/output relations), presynaptic function (assessed by paired pulse facilitation) and protein synthesis-independent, NMDAR-dependent LTP are normal. In contrast, in hippocampus from young (P10–P14) PKA RII $\beta^{-/-}$ mice, protein synthesis-independent LTP is impaired, whereas LTD is normal. These findings indicate that distinct PKA isoforms may subserve distinct forms of synaptic plasticity and are consistent with a developmental switch in the signaling cascades required for LTP induction.

2. Materials and methods

2.1. Animals

All animals (PKA RII β null ($^{-/-}$) mice) and wild-type (WT) littermates and non-littermates were housed and maintained in a temperature- and light-controlled environment with a 14/10-h light/dark cycle in a facility at the Institute for Animal Studies at the Albert Einstein College of Medicine, Bronx, NY. All animal experiments were carried out in accordance with the principles and guidelines of the National Institutes of Health guide for the care and use of laboratory animals. Protocols used for this study were approved by the Institutional Animal Care and Use Committee of the Albert Einstein College of Medicine. The generation of PKA RII $\beta^{-/-}$ mice was as described (Brandon et al., 1998); knockout mice were backcrossed seven or more times onto the C57BL/6 line (Taconic Farms Inc., Germantown, NY) for use in the study. Mice were a generous gift of Quentin Fischer (Yale University, New Haven, CT). To maintain a colony of knockout and wild-type mice, PKA RII $\beta^{+/-}$ heterozygous mice were bred in-house. Additional animals were obtained by crossings of PKA RII $\beta^{+/-}$ heterozygous or PKA RII $\beta^{-/-}$ homozygous mice and genotyped by PCR before use as described (Fischer et al., 2004). The PKA RII $\beta^{-/-}$ mice were viable and exhibited no blatant phenotype.

2.2. Electrophysiology

Acute hippocampal slices were prepared from postnatal day (P) 10–14 or P21–28 PKA RII $\beta^{-/-}$ mice and wild-type littermates and non-littermate C57BL/6 mice. Mice were anesthetized with isoflurane and sacrificed by decapitation. Whole brains were rapidly removed and placed in an ice-cold cutting solution containing/consisting of (in mM): 234 sucrose, 2.5 KCl, 1.25 NaH₂PO₄, 10.0 MgSO₄, 1 CaCl₂, 26 NaHCO₃, and 20 glucose, saturated with 95% O₂ and 5% CO₂. After 5 min incubation in the ice-cold sucrose solution, hippocampi were removed and glued on the stage of a DTK-1000 vibrating micro slicer (Dosaka-EM, Kyoto, Japan) with an agar block, and immersed in ice-cold cutting solution (50%) and normal external (50%) solution. Coronal slices (400 μ m thick) were cut with a micro slicer and transferred to normal external solution containing (in mM): 124 NaCl, 2.5 KCl, 2.5 CaCl₂, 1.3 MgSO₄, 26 NaHCO₃, 1 NaH₂PO₄, and 10 glucose. All solutions were saturated with 95% O₂ and 5% CO₂ (pH 7.4). Slices were incubated for at least 1 h in the external recording solution at 32 \pm 1.5 $^{\circ}$ C prior to recording. For some experiments, CA3 was removed immediately after sectioning, but there was no difference between experimental data from CA3-removed slices and those from CA3-intact slices.

For field recordings, slices were transferred to a submersion-type recording chamber mounted on the stage of an upright microscope (BX50WI, Olympus), held fixed by a grid of parallel nylon threads and perfused with external solution at a rate of 2 ml/min. Slices were maintained at 32 \pm 1.5 $^{\circ}$ C. To record field EPSPs (fEPSPs), a patch electrode (1–2 M Ω) filled with external solution were positioned in the stratum radiatum of area CA1. fEPSPs were evoked by square pulses (10–100 μ A, 200 μ s) in Schaffer collateral afferents by means of a bipolar tungsten stimulating electrode (MX21XEP, Frederick Haer). Baseline presynaptic stimulation was delivered once every 30 s using a stimulation intensity yielding 40–60% of the maximal response (for LTP and LTD experiments). The initial slope of the fEPSP was used to measure stability of synaptic responses and to quantify the magnitude of LTP and LTD.

For input/output curves, single-pulse monophasic test stimulation was applied with a Grass S88 stimulator (Grass Instruments, Quincy, MA) at 0.033 Hz, and electrode positions adjusted to maximize amplitude of the fEPSP. An input/output (I/O) relationship ranging from subthreshold to maximal response was established for 5–6 mice of each genotype. Slices in which the maximal fEPSP amplitude was <2 mV were rejected.

Paired-pulse facilitation (PPF) was assessed in 6 mice of each genotype at interstimulus intervals ranging from 0.02 to 1 s. The PPF ratio was defined as the ratio of the amplitude of the second to that of the first fEPSP amplitude elicited by pairs of stimuli. Synaptic responses were monitored with stimuli consisting of constant current pulses of 0.1 ms duration at 0.067 Hz.

LTP was induced after stable baseline recording for at least 20 min by delivery of 2 trains of stimuli (2 trains of 100 pulses at 100 Hz separated by 20 s). LTD was induced by low-frequency stimulation (LFS; 900 pulses at 1 Hz) or paired-pulse LFS (PP-LFS; 50-ms interstimulus interval). Field potentials were acquired using a Multiclamp 700A amplifier (Axon Instruments, Union City, CA, USA) and a PCI-MIO-16E-4 data acquisition device (National Instruments, Austin, TX, USA). Stimulation and acquisition were controlled by custom acquisition software run on Igor Pro 4.09A (Wavemetrics, Inc., Lake Oswego, OR, USA). To generate frequency–response curves, 900-pulse stimulation was also applied at 1, 3, 5, 10, 30 and 100 Hz. Slices that did not show a stable baseline for at least 20 min prior to stimulation were discarded.

2.3. Drugs

D-AP5 (Tocris Cookson, Ballwin, MO) and CNQX (Tocris Cookson, Ballwin, MO) were prepared as a stock solution in DMSO or H₂O, stored at –20 $^{\circ}$ C and on the day of the experiment, diluted to a final concentration of 50 μ M and 10 μ M, respectively, in external recording solution.

2.4. Data analysis

Every six responses were averaged, and fEPSP slopes were then normalized to baseline. To calculate the magnitude of long-term changes in synaptic strength (LTP, LTD), statistical comparisons were made between mean fEPSP slopes before and 45–60 min after tetanus (LTP) or 70–80 min after LFS (LTD). The data were analyzed online using IgorPro. Results are reported as mean \pm SEM. Statistical analysis was performed using both Student's paired *t*-test and two-way ANOVA at the *P* < 0.05 significance level in Origin Pro 7.0 software (Origin-Lab Corporation, Northampton, MA); these tests yielded comparable *P* values.

3. Results

3.1. Basal synaptic transmission and presynaptic short-term plasticity are normal in area CA1 of juvenile PKA RII $\beta^{-/-}$ mice

To determine whether PKA RII β is essential at Sch–CA1 synapses, we examined short- and long-term synaptic plasticity at CA1 synapses in juvenile (P10–P14) mice lacking PKA RII β . PKA RII $\beta^{-/-}$ mice are viable and fertile, but have markedly decreased white adipose tissue (Cummings et al., 1996). Moreover, PKA RII $\beta^{-/-}$ mice exhibit enhanced basal PKA activity and a compensatory increase in PKA RI α in tissues that normally express RII β (Cummings et al., 1996; Amieux et al., 1997). As at CA1 synapses in mice lacking PKA RI β (Brandon et al., 1995), CA1 synapses are functional in PKA RII $\beta^{-/-}$ mice and synaptic responses could be elicited at wild-type and mutant synapses. We first monitored excitatory synaptic transmission onto CA1 neurons in hippocampal slices from wild-type and PKA RII $\beta^{-/-}$ mice using field-potential recording. The fEPSP and the NMDA component of the fEPSP, isolated by application of CNQX (10 μ M) and low Mg²⁺ (0.05 mM), did not significantly differ at CA1 synapses of wild-type vs. mutant mice (Fig. 1A,B). The NMDA response was abolished by application of APV (50 μ M; Fig. 1A).




Fig. 1. Young PKA RII β KO mice exhibit normal synaptic transmission and short-term plasticity. (A,B) Excitatory synaptic transmission at Schaffer collateral to CA1 synapses was monitored by extracellular field recordings in area CA1 of acute hippocampal slices from WT and PKA RII β KO mice aged P10–P14. (A) Sample fEPSPs from WT and KO mice in the absence (black trace) and presence (red trace) of CNQX (10 μ M) and low Mg^{2+} (0.05 mM) to isolate NMDA-receptor-mediated responses. Application of APV (50 μ M) abolished the NMDA response (blue trace). (B) Summary data for the amplitude of the NMDA component of the fEPSP normalized to the total fEPSP for WT (left) and KO (right). The NMDA response did not differ significantly in slices of WT vs. KO mice. (C,D) The fEPSP input/output (I/O) relation was measured in area CA1 of acute hippocampal slices from WT and PKA RII β KO mice aged P10–P14. Maximum slope was plotted as a function of the presynaptic volley size for different stimulus intensities. Representative superimposed responses and summary data for WT (C) and KO (D) are shown above. The fiber volley is indicated by an arrow; initial deflections are stimulus artifact. Slopes of 3 consecutively obtained fEPSPs were averaged for each data point. The I/O function calculated from the slopes of the linear regression lines did not differ significantly between WT and PKA RII β KO mice (wild-type: 5 slices, 5 mice; knockout: 5 slices, 5 mice; 2-way ANOVA, $P < 0.05$). (E,F) Field excitatory postsynaptic potentials (fEPSPs) evoked by paired-pulse stimulation of Schaffer collaterals at interstimulus intervals ranging from 0.02 to 1 s were recorded from stratum radiatum in area CA1 of acute hippocampal slices from juvenile wild-type and PKA RII β KO mice. (E) Sample records show equal facilitation for WT (upper) and KO (lower). (F) Summary data of paired-pulse ratio as a function of interstimulus interval. The ratios did not differ significantly for WT vs. KO mice at any interstimulus interval, indicative of normal presynaptic function (wild-type: 15 slices, 15 mice; knockout: 9 slices, 9 mice; Student's t -test, $P > 0.1$). Data are reported as the mean \pm SEM. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

To more rigorously examine excitatory synaptic transmission in PKA RII $\beta^{-/-}$ mice, we undertook two experimental approaches. First, we assessed basal synaptic transmission in PKA RII $\beta^{-/-}$ mice aged P10–P14 by monitoring field excitatory postsynaptic potential (fEPSP) input/output relations in slices from knockout and wild-type mice. To reduce the variation between slices, we measured input intensity as a function of fiber volley amplitude. The input/output relation (fEPSP slope vs. fiber volley amplitude) at Sch–CA1

synapses in hippocampal slices from PKA RII β knockout mice was indistinguishable to that of wild-type mice (wild-type: $n = 5$ mice; knockout: $n = 5$ mice; $P > 0.1$; Student's t -test; Fig. 1C,D). Thus, the loss of PKA RII β (and the compensatory increase of PKA RII β ; Amieux et al., 1997) does not alter baseline synaptic transmission at Sch–CA1 synapses.

Second, we monitored field excitatory paired-pulse facilitation (PPF) at six interstimulus intervals (0.02, 0.05, 0.1, 0.2, 0.5 and 1 s) in

slices of juvenile (P10–P14) PKA $R_{II\beta}^{-/-}$ and wild-type mice. PPF is a form of short-term plasticity that is attributed to enhanced transmitter release (Wu and Saggau, 1994; Zucker and Regehr, 2002). No differences in PPF were observed between the PKA $R_{II\beta}^{-/-}$ and wild-type mice at any interstimulus interval tested (wild-type: $n = 15$ slices, 15 mice; knockout, $n = 9$ slices, 9 mice; $P > 0.05$; Fig. 1C,D) (Fig. 1E,F). Thus, the loss of PKA $R_{II\beta}$ (the compensatory increase of PKA $R_{I\beta}$; Amieux et al., 1997) does not alter short-term plasticity at Sch–CA1 synapses in juvenile mice.

3.2. LTP is impaired, whereas LTD is normal in juvenile PKA $R_{II\beta}$ KO mice

PKA is thought to play an important role in the induction (Otmakhova et al., 2000; Skeberdis et al., 2006) and late, protein synthesis-dependent phase (Huang and Kandel, 1994; Qi et al., 1996; Abel et al., 1997; Nguyen and Woo, 2003) of LTP at Sch–CA1 synapses. Most mammalian AKAPs bind PKA through the type II regulatory subunit/PKA R_{II} (Wong and Scott, 2004). In neurons, localization of PKA to dendritic structures is achieved via binding of the R_{II} subunit to AKAPs (Wong and Scott, 2004), an interaction crucial for participation of PKA in LTP (Huang et al., 2006; Nie et al., 2007; Lu et al., 2007). Consistent with this notion, mice lacking the PKA $R_{II\beta}$ subunit exhibit normal LTP (Brandon et al., 1995). To determine the role of the PKA $R_{II\beta}$ subunit in induction of NMDAR-LTP at the Sch–CA1 synapse, we examined synaptically induced/NMDAR-dependent LTP in juvenile mice lacking PKA $R_{II\beta}$, by applying high frequency stimulation (HFS, 2 trains at 100 Hz for 1 s, separated by 20 s) to acute hippocampal slices. Juvenile PKA $R_{II\beta}^{-/-}$ mice exhibited markedly impaired LTP at Sch–CA1 synapses

(wild-type: 134.77 ± 8.67 ; $n = 11$ slices, 9 mice; knockout: $106.21 \pm 4.75\%$, $n = 5$ slices, 5 mice $P < 0.01$; Fig. 2).

We next examined synaptically induced LTD in area CA1 of PKA $R_{II\beta}^{-/-}$ mice. NMDAR-dependent LTD (NMDAR-LTD) is a form of homosynaptic LTD that is mGluR-independent and is elicited in CA1 neurons by low-frequency stimulation (LFS) of Schaffer collateral axons (Mulkey and Malenka, 1992; Oliet et al., 1997; Kemp et al., 2000). The role of PKA in LTD is unclear. Pharmacological inhibition of cAMP/PKA signaling can either enhance or block the induction of NMDAR-dependent LTD at Sch–CA1 synapses (Qi et al., 1996; Kameyama et al., 1998; Santschi et al., 2006). Mice lacking PKA $R_{II\beta}$ exhibit impaired LTD induction at Sch–CA1 synapses (Brandon et al., 1995). LFS LTD at Sch–CA1 synapses is prominent at an early stage in postnatal development (P10–P14) (Oliet et al., 1997; Kemp et al., 2000). Thus, we examined whether PKA $R_{II\beta}$ subunit might also participate in LFS LTD in juvenile rats. Juvenile PKA $R_{II\beta}$ KO mice exhibited robust LTD at Sch–CA1 synapses that were indistinguishable from that of




Fig. 2. The PKA $R_{II\beta}$ subunit is required for LTP in young mice at P10–P14. (A) Sample records of fEPSPs in area CA1 for WT (left) and PKA $R_{II\beta}$ KO (right) mice aged P10–P14 before (black trace) and at 60 min after (red trace) induction of LTP (two 100 Hz tetani, 1 s in duration separated by 20 s). (B) Summary data for LTP of fEPSPs from area CA1 of young WT and PKA $R_{II\beta}$ KO mice. Open symbols represent wild-type slices; closed symbols are KO slices. KO slices were interleaved with WT. For this and subsequent panels, arrow indicates LTP induction. In WT, LTP was robust (to $137.83 \pm 7.48\%$ of baseline, measured 50–60 min after HFS; $n = 13$ slices, 10 mice). In young KO mice, potentiation decays to near baseline by 60 min after tetanus ($102.61 \pm 3.38\%$ of baseline, measured 50–60 min after HFS; $n = 7$ slices, 7 mice, $P < 0.01$ vs. WT, Student's t -test). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)




Fig. 3. The PKA $R_{II\beta}$ subunit is not required for LTD in young mice. (A) Upper, sample records and lower, summary data show LTD of fEPSPs in area CA1 in hippocampal slices from WT mice (left) and PKA $R_{II\beta}$ KO mice (right) before (black) and 60 min after (red) LFS (900 stimuli at 1 Hz). LTD was robust in WT ($63.66 \pm 3.91\%$ of baseline; $n = 8$ slices, 6 mice) and KO mice ($60.47 \pm 5.29\%$ of baseline; $n = 10$ slices, 6 mice). (B) LTD induction in young animals is NMDAR-dependent in WT ($n = 3$ slices, 3 mice) and KO ($n = 3$ slices, 3 mice), as indicated by block in the presence of APV ($50 \mu\text{M}$, application indicated by bar). As for LTP, KO slices were interleaved with WT. For this and subsequent figures an empty region in the fEPSP recordings and horizontal bar indicate time of low-frequency stimulation (LFS). (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

age-matched wild-type mice (wild-type: $63.66 \pm 3.91\%$ of baseline; $n = 8$ slices, 6 mice; knockout: $60.47 \pm 5.29\%$ of baseline; $n = 10$ slices, 6 mice; $P > 0.05$ vs. wild-type; Fig. 3A). LTD induction in young animals was NMDAR-dependent in wild-type ($n = 3$ slices, 3 mice) and knockout ($n = 3$ slices, 3 mice), as indicated by block in the presence of APV ($50 \mu\text{M}$; Fig. 3B). Moreover, in a single experiment PP-LFS induced LTD in both juvenile wild-type and knockout slices (data not illustrated).

The absence of PKA RII β might influence the magnitude of the rise in Ca^{2+} or the way that Ca^{2+} is handled after repetitive stimulation. We therefore determined whether altering the stimulus frequency, a manipulation that alters Ca^{2+} influx, could elicit LTP in the PKA RII β knockout mice. We tested different frequencies of stimulation (1, 3, 10, 30 and 100 Hz). Stimulation at 1 and 3 Hz induced robust LTD in both wild-type and KO slices (wild-type: 65.3 ± 3.8 of baseline; $n = 8$ slices, 8 mice; knockout: 64.78 ± 3.8 of baseline; 10 slices, 10 mice; $P > 0.05$ vs. 1 Hz; wild-type: 75.0 ± 6.0 of baseline; $n = 4$ slices, 4 mice; knockout: 75.04 ± 6.0 of baseline; 4 slices, 4 mice; $P > 0.05$ vs. wild-type at 3 Hz; Fig. 4A,B,F). At 10 Hz, slices of both wild-type and RII $\beta^{-/-}$ mice exhibited very mild LTD (wild-type: $80.0 \pm 7.0\%$ of baseline; $n = 5$ slices, 5 mice; knockout: 89.9 ± 5.0 of baseline; $n = 5$ slices, 5 mice; $P > 0.05$ vs. wild-type at

10 Hz; Fig. 4C,F). In wild-type mice, LTP could be triggered by repetitive stimulation at 30 Hz (Fig. 4D,F), and also at 100 Hz (Fig. 4E,F). Increasing the stimulus frequency to 100 Hz (and presumably the tetanus-induced Ca^{2+} influx) elicited an enhancement of CA1 synaptic responses as expected (data not illustrated). At both frequencies LTP was abolished in the PKA RII $\beta^{-/-}$ mice (wild-type: 131.6 ± 8.9 of baseline; $n = 5$ slices, 5 mice; knockout: 108.4 ± 6.8 of baseline; 5 slices, 5 mice; $P = 0.072$ vs. wild-type 30 Hz; wild-type: 137.8 ± 7.5 of baseline; $n = 13$ slices, 13 mice; knockout: 102.6 ± 3.4 of baseline; 7 slices, 7 mice; $P < 0.005$ vs. wild-type at 100 Hz; Fig. 4D–F). These findings argue against a role for reduced NMDAR-mediated Ca^{2+} influx in the PKA RII $\beta^{-/-}$ mice and suggest that PKA RII β plays a more complex role in NMDAR-dependent LTP.

3.3. Basal synaptic transmission is normal in area CA1 of adult (P21–P28) PKA RII $\beta^{-/-}$ mice

To examine a possible developmental regulation in the impact of RII β on synaptic plasticity, we next assessed basal synaptic transmission in slices of adult PKA RII $\beta^{-/-}$ and wild-type mice at P21–P28. No differences in basal synaptic transmission were




Fig. 4. The frequency–response curve for induction of LTP and LTD is altered in young KO mice. (A–E) Summary data for trains of stimuli given to young WT and KO mice at indicated frequencies. (Panel A is the same as the graph in panel A in Fig. 3; panel E is the same as panel B in Fig. 2.) (F) Frequency–response plot showing magnitude of LTD or LTP at 50–60 min after stimulation for wild-type (white circles) and KO mice (black circles). PKA RII β KO mice exhibited normal LTD in response to 1, 3, and 10 Hz stimulation and deficient LTP in response to 30 and 100 Hz stimulation (1 Hz: wild-type: 8 slices, 6 mice; KO: 10 slices, 6 mice; $P = 0.92$ vs. wild-type; 3 Hz: wild-type: 4 slices, 4 mice; KO: 4 slices, 4 mice; $P = 0.29$ vs. wild-type; 10 Hz: wild-type: 5 slices, 5 mice; KO: 5 slices, 5 mice; $P = 0.28$ vs. wild-type; 30 Hz: wild-type: 5 slices, 5 mice; KO: 5 slices, 5 mice; $P < 0.05$ vs. wild-type; 100 Hz: wild-type: 13 slices, 6 mice; KO: 7 slices, 6 mice; $P < 0.005$ vs. wild-type). Data reported as the mean \pm SEM normalized to baseline.

observed between the PKA $R_{II\beta}^{-/-}$ and wild-type mice (Fig. 5A,B). Second, we monitored field excitatory PPF at various interstimulus intervals (0.05, 0.1, 0.2, 0.4 s). No differences in PPF were observed between the PKA $R_{II\beta}^{-/-}$ and wild-type mice (wild-type: $n = 15$ slices, 4 mice; knockout: $n = 11$ slices, 6 mice; $P > 0.1$, Student's t -test; Fig. 5C,D). These findings indicate that in young adult rats the lack of PKA $R_{II\beta}$, a postsynaptic protein, does not affect presynaptic function.

3.4. NMDAR-dependent LTP at Sch–CA1 synapses is normal in adult PKA $R_{II\beta}^{-/-}$ mice

LTP is thought to play a role not only in learning and memory, but also in activity-dependent synapse stabilization, retraction and maturation, and neural circuit formation and refinement (Katz and Shatz, 1996; Constantine-Paton and Cline, 1998). The signaling cascades required for LTP switch with development from PKA-dependent at young ages ($<P9$) to Ca^{2+} /calmodulin-dependent protein kinase II (CaMKII)-dependent at more mature ages ($>P17$) (Yasuda et al., 2003). We next examined synaptically induced/NMDAR-dependent LTP in young adult (P21–P28) mice lacking PKA $R_{II\beta}$. Whereas the magnitude of HFS-induced LTP in juvenile PKA $R_{II\beta}^{-/-}$ mice exhibited markedly impaired LTP at Sch–CA1 synapses, LTP in young adult PKA $R_{II\beta}^{-/-}$ mice was not significantly different from that in wild-type mice (wild-type: 158.3 ± 6.5 , $n = 15$ slices, 6 mice; knockout: 144.9 ± 9.3 , $n = 9$ slices, 5 mice; $P = 0.23$;

Student's t -test, Fig. 6A). These results point to a developmental switch in the role of $R_{II\beta}$ in synaptic plasticity.

3.5. NMDAR-dependent LTD is deficient in adult PKA $R_{II\beta}^{-/-}$ mice

We next examined synaptically induced LTD in area CA1 of adult (P21–P28) PKA $R_{II\beta}^{-/-}$ mice. Whereas juvenile PKA $R_{II\beta}^{-/-}$ mice exhibited robust LTD at Sch–CA1 synapses that were indistinguishable from that of age-matched wild-type mice, LTD in adult mice was significantly impaired. In control slices, LFS (900 stimuli at 1 Hz) elicited a robust depression in the field potential (FP) slope values to 82.2 ± 3.7 of baseline, measured 35–40 min after LFS (wild-type: $n = 8$ slices, 6 mice; Fig. 6B). By contrast, the same induction protocol did not elicit detectable LTD in slices from PKA $R_{II\beta}$ knockout mice (104.0 ± 6.2 of baseline response, 35–40 min post-LFS; $n = 7$ slices, 6 mice, $P < 0.01$ vs. wild-type: Student's t -test, Fig. 6B).

As with the juvenile rats, we examined whether altering the stimulus frequency, a manipulation that alters Ca^{2+} influx, could elicit LTD in the PKA $R_{II\beta}$ knockout mice in young adult rats. We tested different frequencies of stimulation (1, 3, 5, 10 and 100 Hz). In wild-type mice, LTD could be triggered by repetitive stimulation not only at 1 Hz (Fig. 7A,F), but also at 3 Hz (Fig. 7B,F). In the PKA $R_{II\beta}^{-/-}$ mice, increasing the stimulus frequency to 3 Hz (and presumably the tetanus-induced Ca^{2+} influx) elicited an enhancement of CA1 synaptic responses as expected (data not illustrated);




Fig. 5. Adult PKA $R_{II\beta}$ KO mice exhibit normal basal neural transmission and short-term plasticity. (A,B) Input/output relations of fEPSPs were determined as in Fig. 1. Representative superimposed responses and summary data for wild-type (A) and knockout (B) mice. The fiber volley is indicated by an arrow; initial deflections are stimulus artifact. Each data point represents the average slope of 3 consecutively obtained fEPSPs. The I/O function calculated from the slopes of the linear regression lines did not differ significantly between wild-type and PKA $R_{II\beta}$ knockout mice (wild-type: 6 slices, 6 mice; knockout: 6 slices, 6 mice; 2-way ANOVA, $P > 0.05$). (C,D) PPF was monitored in the stratum radiatum of area CA1 in acute hippocampal slices from wild-type and PKA $R_{II\beta}$ knockout mice aged P21–P28 as in Fig. 1. (C) Sample records for a 0.05 s interstimulus interval show equal facilitation for wild-type (left) and knockout (right). (D) Summary data showing PPF at interstimulus intervals ranging from 0.05 to 0.4 s in wild-type and PKA $R_{II\beta}$ knockout mice. PPF in the PKA $R_{II\beta}$ knockout mice did not differ significantly from that of wild-type mice, indicative of normal presynaptic function (wild-type: 15 slices, 4 mice; knockout: 11 slices, 6 mice; Student's t -test, $P > 0.1$). Data are reported as the mean \pm SEM.




Fig. 6. LTP is normal but LTD is deficient in slices from PKA RII β KO animals at P21–P28. (A) Upper, sample records of fEPSPs for WT mice (left) and PKA RII β KO mice (right) at times indicated before (black trace) and after (red) induction of LTP (two 100 Hz tetani, 1 s in duration separated by 20 s). Lower, summary data for LTP of fEPSPs from CA1 of WT ($n = 15$ slices, 6 mice) and KO ($n = 8$ slices, 6 mice). (B) Upper, sample records of fEPSPs for WT mice (left) and PKA RII β KO mice (right) at indicated times before (black trace) and after (red trace) LTD-inducing stimulation was applied (900 stimuli at 1 Hz). Lower, summary data for LTD of fEPSPs in WT and PKA RII β KO mice ($n = 8$ slices, 6 animals). In WT, LTD was robust (to $74.3 \pm 3.7\%$ of baseline, measured 50–60 min after LFS; $n = 8$ slices, 6 mice; $P < 0.01$ vs. baseline; Student's t -test). In KO mice, LTD was absent ($99.0 \pm 6.2\%$ of baseline, measured 50–60 min post-LFS; $n = 7$ slices, 6 mice; $P < 0.01$ vs. WT; Student's t -test). As for LTP, KO slices were interleaved with WT. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

however, LTD was still abolished in the PKA RII β ^{−/−} mice (wild-type: $89.4 \pm 3.7\%$ of baseline; $n = 8$ slices, 6 mice; knockout: $98.5 \pm 4.2\%$ of baseline; 8 slices, 8 mice; $P = 0.56$ vs. 1 Hz; $P < 0.05$ vs. wild-type at 3 Hz; Fig. 7B,F). 5 Hz stimulation did not induce plasticity in either wild-type or KO slices (wild-type: $102.0 \pm 6.6\%$ of baseline; $n = 7$ slices, 6 mice; knockout: $96.6 \pm 4.8\%$ of baseline; 9 slices, 9 mice; $P > 0.05$ vs. 1 Hz; $P > 0.05$ vs. wild-type at 5 Hz;

Fig. 7C,F). At 10 Hz, slices of wild-type mice exhibited robust LTP, but slices of RII β ^{−/−} mice exhibited a significantly reduced LTP relative to that of wild-type (wild-type: $119.6 \pm 3.8\%$ of baseline; $n = 7$ slices, 6 mice; knockout: $101.2 \pm 5.4\%$ of baseline; $n = 10$ slices, 10 mice; $P < 0.01$ vs. 1 Hz; $P < 0.05$ vs. wild-type at 5 Hz; Fig. 7D,F). At 100 Hz stimulation, LTP induction was essentially normal in knockout vs. wild-type slices (Fig. 7E,F; see also Fig. 6A). These findings argue against a role for reduced NMDAR-mediated Ca²⁺ influx in the adult PKA RII β ^{−/−} mice and suggest that PKA RII β plays a more complex role in NMDAR-dependent LTD. Collectively, these findings indicate that the RII β subunit has a greater impact on long-term, postsynaptic vs. short-term, presynaptic plasticity and that the deficits in LTP and LTD observed in PKA RII β ^{−/−} mice are developmentally altered.

4. Discussion

The cAMP/PKA signaling cascade is crucial for synaptic plasticity in a wide variety of species. The role of cAMP/PKA signaling in the induction of NMDAR-dependent LTP at Sch–CA1 synapses is established (Otmakhova et al., 2000; Skeberdis et al., 2006), but the identity of the PKA isoform involved in LTP is, as yet unclear. We have shown here that in young hippocampus, PKA RII β , a regulatory subunit of PKA implicated in localization of PKA to the postsynaptic density in close proximity and direct physical association with NMDARs and AMPARs, is essential for inducing NMDAR-dependent LTP at Schaffer collateral to CA1 synapses. RII β is not required for normal NMDAR-mediated responses, excitatory synaptic transmission, presynaptic short-term plasticity or NMDAR-dependent LTD. This phenotype is unlikely to be caused by structural abnormalities, because light and electron microscopy both showed that the brain has an overall normal structure and organization in PKA RII β knockout mice (Cummings et al., 1996). In contrast, in young adult hippocampus, PKA RII β is critical to NMDAR-dependent LTD, but not LTP. These findings indicate that distinct PKA isoforms may subserve distinct forms of synaptic plasticity and are consistent with a developmental switch in the signaling cascades required for LTP induction.

Activity-dependent LTP is critical to formation, refinement and stabilization of new synapses in the young brain (Katz and Shatz, 1996; Constantine-Paton and Cline, 1998). Recent studies indicate that the signaling cascades involved in the induction phase of LTP at CA1 synapses are developmentally regulated. Whereas LTP is dependent upon CaMKII after postnatal day 20 (P20), it is dependent on PKA at P7–P8 (Yasuda et al., 2003). Our finding in the present study that PKA RII β is critical to LTP at young ages (P10–P14), but not at more mature ages (P21–P28) is consistent with this notion. Our finding extends the work of Malenka and colleagues in that it identifies for the first time a critical role for the PKA RII β subunit in LTP signaling in young, but not mature, rodent hippocampus. The PKA RII β subunit binds directly AKAP150 (Colledge et al., 2000; Oliveria et al., 2003), which in turn binds via PSD-95 and/or SAP97 and tethers the PKA holoenzyme to the NMDAR GluN2B subunit. At the time of birth, NMDARs in the hippocampus contain primarily GluN1 and GluN2B subunits. Over the course of postnatal development, there is a progressive inclusion of the GluN2A subunit (Carroll and Zukin, 2002; Cull-Candy and Leszkiewicz, 2004; Lau and Zukin, 2007). A possible scenario is that at young ages, synaptic NMDARs are GluN2B-containing and are more sensitive to regulation by PKA. At later ages, at a time when the number of spines has increased and synaptic connections have formed, NMDARs are primarily GluN2A- or GluN2A/GluN2B-containing (Lau and Zukin, 2007) and become relatively less sensitive to regulation by PKA (Skeberdis et al., 2006).

Our finding of a critical role for the PKA RII β subunit in LTP signaling in young, but not mature, rodent hippocampus is also




Fig. 7. The frequency–response curve for induction of LTP and LTD is altered in adult KO mice. (A–E) Summary data for trains of stimuli given to WT and KO mice at indicated frequencies. (Panel E is the same as the graph in panel A in Fig. 6.) (F) Frequency–response plot showing magnitude of LTD or LTP at 45–60 min after stimulation for wild-type (white circles) and KO mice (black circles). PKA RII β KO mice exhibited no LTD in response to 1, 3, 5, and 10 Hz stimulation, and normal LTP in response to 100 Hz stimulation (1 Hz: wild-type: 8 slices, 6 mice; KO: 7 slices, 6 mice; $P=0.01$ vs. wild-type; 3 Hz: wild-type: 8 slices, 6 mice; KO: 8 slices, 6 mice; $P=0.056$ vs. wild-type; 5 Hz: wild-type: 7 slices, 6 mice; KO: 9 slices, 9 mice; $P=0.38$ vs. wild-type; 10 Hz: wild-type: 7 slices, 6 mice; KO: 10 slices, 10 mice; $P=0.01$ vs. wild-type; 100 Hz: wild-type: 15 slices, 6 mice; KO: 8 slices, 5 mice; $P=0.23$ vs. wild-type). Data reported as the mean \pm SEM normalized to baseline.

consistent with findings of others that early in development, PKA phosphorylation of GluA4-long is a primary mediator of activity-dependent synaptic incorporation of AMPARs, a mechanism thought to underlie enhanced synaptic strength (Zhu et al., 2000; Esteban et al., 2003). GluA4 delivery requires PKA-dependent phosphorylation at serine residue 842 in its carboxy-terminal tail (Zhu et al., 2000; Esteban et al., 2003). Thus, a deficit in RII β , which is required for correct targeting of PKA to AMPARs could account for impaired LTP in the CA1 of RII β knockout mice. Our finding that LTP is not completely abolished even at young ages could be explained by a role for CaMKII-dependent, PKA-independent delivery of GluA2-long. In contrast, in adult hippocampus, activity drives PKA-independent insertion of GluA2-long, as well as PKA-dependent delivery of GluA1 (Man et al., 2007). These findings predict a slight impairment of LTP at CA1 synapses of adult RII β ^{−/−} mice. Thus, PKA phosphorylation of AMPARs mediates plasticity through synaptic incorporation of AMPARs early in development; in mature hippocampus, PKA is thought to play a “gating” role through phosphorylation and delivery of GluA1 (Esteban et al., 2003). Our finding that LTD is impaired in PKA RII β ^{−/−} mice in young adulthood, but not early in development is novel. Further experiments are warranted to clarify the role of PKA in LTD.

In visual cortex, regulatory subunits type I and type II of PKA differentially regulate synaptic plasticity. Whereas the PKA RI subunit is required for LTP and LTD at synapses onto layer II/III

visual cortex cells, but not ocular dominance plasticity (Hensch et al., 1998), the PKA RII β subunit is required for ocular dominance plasticity and LTD, but not LTP (Fischer et al., 2004). Pharmacological blockade of PKA abolishes all three forms of plasticity (Beaver et al., 2001; Liu et al., 2003). Targeted deletion of RII β severely reduces LTP at mossy fiber-CA3 synapses (Brandon et al., 1995) and homosynaptic LTD and depotentiation (but not LTP) at Sch-CA1 synapses of adult mice at 4–6 weeks of age (Huang et al., 1995). Our findings are consistent with and extend these studies in that we identify a role not only for RI β , but also for RII β in LTD at CA1 synapses of adult mice. These studies establish PKA subunit-specific roles in different forms of plasticity within a given region.

Our finding that LTP is impaired at developing Schaffer collateral-CA1 synapses of PKA RII β ^{−/−} mice is consistent with findings of others that LTP is impaired at developing thalamocortical synapses in the barrel cortex (Inan et al., 2006). Thus, the role of RII β in NMDAR-dependent LTP at developing synapses is not limited to CA1 synapses. Our finding that LTP is normal at CA1 synapses of PKA RII β ^{−/−} mice at 3–4 weeks of age is consistent with findings that mice deficient in AKAP150–PKA coupling exhibit normal LTP at 4 weeks, but impaired LTP at 7–12 weeks (Lu et al., 2007). Our finding that LTD is impaired at mature CA1 synapses is consistent with findings LTD is impaired at synapses of mature visual cortex of PKA RII β ^{−/−} mice (Fischer et al., 2004). Our findings extend previous findings in that we show that PKA RII β subserves distinct forms of

plasticity at CA1 synapses at different developmental stages. Interestingly, mice deficient in PKA anchoring exhibit impaired protein synthesis-dependent LTP at CA1 synapses at 8–12-weeks of age (Nie et al., 2007), suggesting a possible second “developmental switch” in LTP signaling.

The cAMP/PKA signaling cascade is crucial for synaptic plasticity in a wide variety of species. Our earlier finding that PKA activity increases Ca^{2+} permeability links Ca^{2+} signaling to NMDAR-dependent synaptic plasticity. Findings in the present study identify a role for PKA RII β in NMDAR-dependent synaptic plasticity at Sch–CA1 synapses. Our findings that 3–4 week old mice lacking PKA RII β exhibit normal LTP at Sch–CA1, but deficient LTD, indicate that distinct PKA isoforms may subserve distinct forms of synaptic plasticity. Our findings that young mice lacking PKA RII β exhibit deficient LTP and normal LTD are consistent with a developmental switch in the signaling cascades required for LTP induction. Given the widespread distribution of NMDARs and PKA throughout the CNS, the developmentally dependent, subunit-specific regulation of NMDAR-dependent synaptic plasticity by PKA potentially represents a powerful mechanism to modulate synaptic efficacy.

Acknowledgements

The authors thank Dr. Pablo E. Castillo for helpful comments on the manuscript and Ms. Adrianna Latuszek-Barrantes for help in maintaining the mice and genotyping. The PKA RII β knockout mice are a gift of Quentin Fischer (Yale University School of Medicine, New Haven, CT). This study was supported by NIH grants NS20752 (to R.S.Z.) and NS45287 (to M.V.L.B.) and Training Grant T32 DK07513. M.V.L.B. is the Sylvia and Robert S. Olnick Professor of Neuroscience and Distinguished Professor of the College.

References

- Abel, T., Nguyen, P.V., Barad, M., Deuel, T.A., Kandel, E.R., Bourtochouladze, R., 1997. Genetic demonstration of a role for PKA in the late phase of LTP and in hippocampus-based long-term memory. *Cell* 88, 615–626.
- Amieux, P.S., Cummings, D.E., Motamed, K., Brandon, E.P., Wailes, L.A., Le, K., Idzerda, R.L., McKnight, G.S., 1997. Compensatory regulation of RII α protein levels in protein kinase A mutant mice. *J. Biol. Chem.* 272, 3993–3998.
- Beaver, C.J., Ji, Q., Fischer, Q.S., Daw, N.W., 2001. Cyclic AMP-dependent protein kinase mediates ocular dominance shifts in cat visual cortex. *Nat. Neurosci.* 4, 159–163.
- Blank, T., Nijholt, I., Teichert, U., Kugler, H., Behrsing, H., Fienberg, A., Greengard, P., Spiess, J., 1997. The phosphoprotein DARPP-32 mediates cAMP-dependent potentiation of striatal N-methyl-D-aspartate responses. *Proc. Natl. Acad. Sci. U.S.A.* 94, 14859–14864.
- Blitzer, R.D., Connor, J.H., Brown, G.P., Wong, T., Shenolikar, S., Iyengar, R., Landau, E.M., 1998. Gating of CaMKII by cAMP-regulated protein phosphatase activity during LTP. *Science* 280, 1940–1942.
- Brandon, E.P., Idzerda, R.L., McKnight, G.S., 1997. PKA isoforms, neural pathways, and behaviour: making the connection. *Curr. Opin. Neurobiol.* 7, 397–403.
- Brandon, E.P., Logue, S.F., Adams, M.R., Qi, M., Sullivan, S.P., Matsumoto, A.M., Dorsa, D.M., Wehner, J.M., McKnight, G.S., Idzerda, R.L., 1998. Defective motor behavior and neural gene expression in RII β -protein kinase A mutant mice. *J. Neurosci.* 18, 3639–3649.
- Brandon, E.P., Zhuo, M., Huang, Y.Y., Qi, M., Gerhold, K.A., Burton, K.A., Kandel, E.R., McKnight, G.S., Idzerda, R.L., 1995. Hippocampal long-term depression and depotentiation are defective in mice carrying a targeted disruption of the gene encoding the RII β subunit of cAMP-dependent protein kinase. *Proc. Natl. Acad. Sci. U.S.A.* 92, 8851–8855.
- Carr, D.W., Stofko-Hahn, R.E., Fraser, I.D., Cone, R.D., Scott, J.D., 1992. Localization of the cAMP-dependent protein kinase to the postsynaptic densities by A-kinase anchoring proteins. Characterization of AKAP 79. *J. Biol. Chem.* 267, 16816–16823.
- Carroll, R.C., Zukin, R.S., 2002. NMDA-receptor trafficking and targeting: implications for synaptic transmission and plasticity. *Trends Neurosci.* 25, 571–577.
- Cerne, R., Rusin, K.I., Randic, M., 1993. Enhancement of the N-methyl-D-aspartate response in spinal dorsal horn neurons by cAMP-dependent protein kinase. *Neurosci. Lett.* 161, 124–128.
- Colledge, M., Dean, R.A., Scott, G.K., Langeberg, L.K., Huganir, R.L., Scott, J.D., 2000. Targeting of PKA to glutamate receptors through a MAGUK–AKAP complex. *Neuron* 27, 107–119.
- Colledge, M., Scott, J.D., 1999. AKAPs: from structure to function. *Trends Cell Biol.* 9, 216–221.
- Collingridge, G.L., Isaac, J.T., Wang, Y.T., 2004. Receptor trafficking and synaptic plasticity. *Nat. Rev. Neurosci.* 5, 952–962.
- Constantine-Paton, M., Cline, H.T., 1998. LTP and activity-dependent synaptogenesis: the more alike they are, the more different they become. *Curr. Opin. Neurobiol.* 8, 139–148.
- Cull-Candy, S.G., Leszkiewicz, D.N., 2004. Role of distinct NMDA receptor subtypes at central synapses. *Sci. STKE* 2004, re16.
- Cummings, D.E., Brandon, E.P., Planas, J.V., Motamed, K., Idzerda, R.L., McKnight, G.S., 1996. Genetically lean mice result from targeted disruption of the RII β subunit of protein kinase A. *Nature* 382, 622–626.
- Dingledine, R., Borges, K., Bowie, D., Traynelis, S.F., 1999. The glutamate receptor ion channels. *Pharmacol. Rev.* 51, 7–61.
- Esteban, J.A., Shi, S.H., Wilson, C., Nuriya, M., Huganir, R.L., Malinow, R., 2003. PKA phosphorylation of AMPA receptor subunits controls synaptic trafficking underlying plasticity. *Nat. Neurosci.* 6, 136–143.
- Fischer, Q.S., Beaver, C.J., Yang, Y., Rao, Y., Jakobsdottir, K.B., Storm, D.R., McKnight, G.S., Daw, N.W., 2004. Requirement for the RII β isoform of PKA, but not calcium-stimulated adenylyl cyclase, in visual cortical plasticity. *J. Neurosci.* 24, 9049–9058.
- Hensch, T.K., Gordon, J.A., Brandon, E.P., McKnight, G.S., Idzerda, R.L., Stryker, M.P., 1998. Comparison of plasticity in vivo and in vitro in the developing visual cortex of normal and protein kinase A RII β -deficient mice. *J. Neurosci.* 18, 2108–2117.
- Huang, T., McDonough, C.B., Abel, T., 2006. Compartmentalized PKA signaling events are required for synaptic tagging and capture during hippocampal late-phase long-term potentiation. *Eur. J. Cell Biol.* 85, 635–642.
- Huang, Y.Y., Kandel, E.R., 1994. Recruitment of long-lasting and protein kinase A-dependent long-term potentiation in the CA1 region of hippocampus requires repeated tetanization. *Learn. Mem.* 1, 74–82.
- Huang, Y.Y., Kandel, E.R., Varshavsky, L., Brandon, E.P., Qi, M., Idzerda, R.L., McKnight, G.S., Bourtochouladze, R., 1995. A genetic test of the effects of mutations in PKA on mossy fiber LTP and its relation to spatial and contextual learning. *Cell* 83, 1211–1222.
- Inan, M., Lu, H.C., Albright, M.J., She, W.C., Crair, M.C., 2006. Barrel map development relies on protein kinase A regulatory subunit II β -mediated cAMP signaling. *J. Neurosci.* 26, 4338–4349.
- Kameyama, K., Lee, H.K., Bear, M.F., Huganir, R.L., 1998. Involvement of a post-synaptic protein kinase A substrate in the expression of homosynaptic long-term depression. *Neuron* 21, 1163–1175.
- Katz, L.C., Shatz, C.J., 1996. Synaptic activity and the construction of cortical circuits. *Science* 274, 1133–1138.
- Kemp, N., McQueen, J., Faulkes, S., Bashir, Z.I., 2000. Different forms of LTD in the CA1 region of the hippocampus: role of age and stimulus protocol. *Eur. J. Neurosci.* 12, 360–366.
- Lau, C.G., Zukin, R.S., 2007. NMDA receptor trafficking in synaptic plasticity and neuropsychiatric disorders. *Nat. Rev. Neurosci.* 8, 413–426.
- Leonard, A.S., Hell, J.W., 1997. Cyclic AMP-dependent protein kinase and protein kinase C phosphorylate N-methyl-D-aspartate receptors at different sites. *J. Biol. Chem.* 272, 12107–12115.
- Liu, S., Rao, Y., Daw, N., 2003. Roles of protein kinase A and protein kinase G in synaptic plasticity in the visual cortex. *Cereb. Cortex* 13, 864–869.
- Lu, Y., Allen, M., Halt, A.R., Weisenhaus, M., Dallapiazza, R.F., Hall, D.D., Usachev, Y.M., McKnight, G.S., Hell, J.W., 2007. Age-dependent requirement of AKAP150-anchored PKA and GluR2-lacking AMPA receptors in LTP. *EMBO J.* 26, 4879–4890.
- Malenka, R.C., Bear, M.F., 2004. LTP and LTD: an embarrassment of riches. *Neuron* 44, 5–21.
- Malinow, R., Malenka, R.C., 2002. AMPA receptor trafficking and synaptic plasticity. *Annu. Rev. Neurosci.* 25, 103–126.
- Man, H.Y., Sekine-Aizawa, Y., Huganir, R.L., 2007. Regulation of α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor trafficking through PKA phosphorylation of the Glu receptor 1 subunit. *Proc. Natl. Acad. Sci. U.S.A.* 104, 3579–3584.
- Mulkey, R.M., Malenka, R.C., 1992. Mechanisms underlying induction of homosynaptic long-term depression in area CA1 of the hippocampus. *Neuron* 9, 967–975.
- Nguyen, P.V., Woo, N.H., 2003. Regulation of hippocampal synaptic plasticity by cyclic AMP-dependent protein kinases. *Prog. Neurobiol.* 71, 401–437.
- Nie, T., McDonough, C.B., Huang, T., Nguyen, P.V., Abel, T., 2007. Genetic disruption of protein kinase A anchoring reveals a role for compartmentalized kinase signaling in theta-burst long-term potentiation and spatial memory. *J. Neurosci.* 27, 10278–10288.
- Oliet, S.H., Malenka, R.C., Nicoll, R.A., 1997. Two distinct forms of long-term depression coexist in CA1 hippocampal pyramidal cells. *Neuron* 18, 969–982.
- Oliveria, S.F., Gomez, L.L., Dell'Acqua, M.L., 2003. Imaging kinase–AKAP79–phosphatase scaffold complexes at the plasma membrane in living cells using FRET microscopy. *J. Cell Biol.* 160, 101–112.
- Otmakhova, N.A., Otmakhov, N., Mortenson, L.H., Lisman, J.E., 2000. Inhibition of the cAMP pathway decreases early long-term potentiation at CA1 hippocampal synapses. *J. Neurosci.* 20, 4446–4451.
- Qi, M., Zhuo, M., Skalleberg, B.S., Brandon, E.P., Kandel, E.R., McKnight, G.S., Idzerda, R.L., 1996. Impaired hippocampal plasticity in mice lacking the C β 1 catalytic subunit of cAMP-dependent protein kinase. *Proc. Natl. Acad. Sci. U.S.A.* 93, 1571–1576.
- Raman, I.M., Tong, G., Jahr, C.E., 1996. Beta-adrenergic regulation of synaptic NMDA receptors by cAMP-dependent protein kinase. *Neuron* 16, 415–421.

- Santschi, L.A., Zhang, X.L., Stanton, P.K., 2006. Activation of receptors negatively coupled to adenylate cyclase is required for induction of long-term synaptic depression at Schaffer collateral–CA1 synapses. *J. Neurobiol.* 66, 205–219.
- Scannevin, R.H., Huganir, R.L., 2000. Postsynaptic organization and regulation of excitatory synapses. *Nat. Rev. Neurosci.* 1, 133–141.
- Skeberdis, V.A., Chevalleyre, V., Lau, C.G., Goldberg, J.H., Pettit, D.L., Suadcani, S.O., Lin, Y., Bennett, M.V., Yuste, R., Castillo, P.E., Zukin, R.S., 2006. Protein kinase A regulates calcium permeability of NMDA receptors. *Nat. Neurosci.* 9, 501–510.
- Snyder, G.L., Fienberg, A.A., Huganir, R.L., Greengard, P., 1998. A dopamine/D1 receptor/protein kinase A/dopamine- and cAMP-regulated phosphoprotein (Mr 32 kDa)/protein phosphatase-1 pathway regulates dephosphorylation of the NMDA receptor. *J. Neurosci.* 18, 10297–10303.
- Tingley, W.G., Ehlers, M.D., Kameyama, K., Doherty, C., Ptak, J.B., Riley, C.T., Huganir, R.L., 1997. Characterization of protein kinase A and protein kinase C phosphorylation of the *N*-methyl-D-aspartate receptor NR1 subunit using phosphorylation site-specific antibodies. *J. Biol. Chem.* 272, 5157–5166.
- Wang, L.Y., Orser, B.A., Brautigam, D.L., MacDonald, J.F., 1994. Regulation of NMDA receptors in cultured hippocampal neurons by protein phosphatases 1 and 2A. *Nature* 369, 230–232.
- Westphal, R.S., Tavalin, S.J., Lin, J.W., Alto, N.M., Fraser, I.D., Langeberg, L.K., Sheng, M., Scott, J.D., 1999. Regulation of NMDA receptors by an associated phosphatase–kinase signaling complex. *Science* 285, 93–96.
- Wong, W., Scott, J.D., 2004. AKAP signalling complexes: focal points in space and time. *Nat. Rev. Mol. Cell Biol.* 5, 959–970.
- Wu, L.G., Saggau, P., 1994. Presynaptic calcium is increased during normal synaptic transmission and paired-pulse facilitation, but not in long-term potentiation in area CA1 of hippocampus. *J. Neurosci.* 14, 645–654.
- Yasuda, H., Barth, A.L., Stellwagen, D., Malenka, R.C., 2003. A developmental switch in the signaling cascades for LTP induction. *Nat. Neurosci.* 6, 15–16.
- Zhu, J.J., Esteban, J.A., Hayashi, Y., Malinow, R., 2000. Postnatal synaptic potentiation: delivery of GluR4-containing AMPA receptors by spontaneous activity. *Nat. Neurosci.* 3, 1098–1106.
- Zucker, R.S., Regehr, W.G., 2002. Short-term synaptic plasticity. *Annu. Rev. Physiol.* 64, 355–405.