


Contents lists available at [SciVerse ScienceDirect](http://www.sciencedirect.com)

Neuropharmacology

journal homepage: www.elsevier.com/locate/neuropharm

Invited review

Reward and aversion in a heterogeneous midbrain dopamine system

Stephan Lammel¹, Byung Kook Lim¹, Robert C. Malenka*

Nancy Pritzker Laboratory, Department of Psychiatry and Behavioral Sciences, Stanford University School of Medicine, 265 Campus Drive, Stanford, CA 94305, USA

ARTICLE INFO

Article history:

Received 28 January 2013

Received in revised form

11 March 2013

Accepted 26 March 2013

Keywords:

Dopamine

Reward

Aversion

Mesocortical

Mesolimbic

Ventral tegmental area

ABSTRACT

The ventral tegmental area (VTA) is a heterogeneous brain structure that serves a central role in motivation and reward processing. Abnormalities in the function of VTA dopamine (DA) neurons and the targets they influence are implicated in several prominent neuropsychiatric disorders including addiction and depression. Recent studies suggest that the midbrain DA system is composed of anatomically and functionally heterogeneous DA subpopulations with different axonal projections. These findings may explain a number of previously confusing observations that suggested a role for DA in processing both rewarding as well as aversive events. Here we will focus on recent advances in understanding the neural circuits mediating reward and aversion in the VTA and how stress as well as drugs of abuse, in particular cocaine, alter circuit function within a heterogeneous midbrain DA system.

This article is part of a Special Issue entitled 'NIDA 40th Anniversary Issue'.

© 2013 Published by Elsevier Ltd.

1. Introduction

As investigators have delved deeper into the brain's neuronal networks underlying complex behaviors using innovative techniques such as optogenetics, it has become clear that previous notions about the functions and connectivity of individual cell types in the mammalian brain need to be revised. Nowhere is this more apparent than for midbrain dopamine (DA) cells. Over the past decade our view of the midbrain DA system has changed from

a simple organization of anatomically separate DA neurons in the substantia nigra and ventral tegmental area (VTA) to a more complex system of DA neuron subtypes with different axonal projection and inputs, distinct anatomical, molecular and electrophysiological features (Ikemoto, 2007; Lammel et al., 2008, 2011, 2012; Margolis et al., 2006, 2008) as well as additional co-transmitters such as GABA (Tritsch et al., 2012) and glutamate (El Mestikawy et al., 2011). In addition, it is well established that midbrain DA neurons are intermingled and connected with different subpopulations of GABAergic and glutamatergic neurons (Hnasko et al., 2012; Margolis et al., 2012; Yamaguchi et al., 2011). Thus, perhaps not surprisingly, DA neuron activity has been associated with a variety of brain functions including the signaling of reward, aversion, salience, uncertainty and novelty (Bromberg-Martin et al., 2010; Schultz, 2007; Ungless et al., 2010). Classic work demonstrated that midbrain DA neurons evince characteristic phasic responses to rewards and cues that predict rewards, and are inhibited by aversive events (Schultz, 1997; Ungless et al., 2004). Furthermore, it has been suggested by many investigators that drugs of abuse hijack the brain's "reward system" and that among the most important neural circuit modifications that contribute to the development of addiction are changes in the properties of excitatory synapses on midbrain VTA DA neurons (Luscher and Malenka, 2011). Here, we provide an overview of how different circuits involving distinct VTA DA neuron subpopulations may contribute to the role of DA in reward- and aversion-related behaviors. In addition, we highlight how these circuits may contribute to drug addiction.

Abbreviations: DA, dopamine; VTA, ventral tegmental area; RRF, retrorubral field; SN, substantia nigra; SNc, substantia nigra pars compacta; SNr, substantia nigra pars reticulata; ml, medial lemniscus; IPN, interpeduncular nucleus; CM, mammillary body; mVTA, medial VTA; lVTA, lateral VTA; fr, fasciculus retroflexus; PBP, parabrachial pigmented nucleus; PN, paranigral nucleus; CLi, caudal linear nucleus; IF, interfascicular nucleus; RLi, rostral linear nucleus of the raphe; MT, medial terminal nucleus of the accessory optical tract; PFC, prefrontal cortex; NAc, nucleus accumbens; BLA, basolateral amygdala; AADC, amino acid decarboxylase; VMAT2, vesicular monoamine transporter 2; DAT, dopamine transporter; D2R, dopamine D2 receptor; VGLUT2, vesicular glutamate transporter 2; KATP, ATP sensitive potassium channel; TH, tyrosine hydroxylase; GAD, glutamic acid decarboxylase; LDT, laterodorsal tegmentum; LHB, lateral habenula; RMTg, rostromedial tegmental nucleus; ChR2, channelrhodopsin 2; AMPAR, α -amino-3-hydroxy-5-methyl-4-isoxazolepropionic acid receptor; NMDAR, N-methyl-D-aspartate; MSN, medium spiny neuron.

* Corresponding author. Department of Psychiatry and Behavioral Sciences, 265 Campus Drive, Room G1021, Stanford University School of Medicine, Stanford, CA 94305, USA. Tel.: +1 650 724 2730; fax: +1 650 724 2753.

E-mail address: malenka@stanford.edu (R.C. Malenka).

¹ These authors contributed equally.

2. Heterogeneity of the mesocorticolimbic dopamine system

Midbrain DA neurons are mainly located in three nuclei, the VTA (A10), the retrorubral field (RRF, A8) and substantia nigra pars compacta (SNc, A9) as well as sparsely scattered in the substantia nigra pars reticulata (SNr). The two major subnuclei of the VTA are the parabrachial pigmented (PBP) and paranigral (PN) nuclei. Furthermore, the caudal linear nucleus (CLi), interfascicular nucleus (IF), and rostral linear nucleus of the raphe (RLi) are often also considered VTA subregions (Oades and Halliday, 1987; Swanson, 1982).

In the late 1980s and early 1990s using *in vitro* slice recordings researchers began to classify DA neurons as principal (mostly DAergic) and secondary (GABAergic) (Grace and Onn, 1989; Johnson and North, 1992) on the basis of distinct physiological and pharmacological properties as well as tyrosine hydroxylase (TH) immunohistochemistry (Grace and Onn, 1989; Johnson and North, 1992). Subsequent studies showed a third group of VTA neurons (tertiary neurons), which are hyperpolarized by serotonin and opioids but it appears that only one-third of these neurons are DAergic (Cameron et al., 1997). The neurochemical phenotype of the remaining two-third of the tertiary cells has not been clearly defined.

Based on these findings virtually all *in vitro* electrophysiological studies, many of them studying drug-induced synaptic adaptations, have considered VTA DA midbrain neurons as a single population (e.g. Argilli et al., 2008; Bellone and Luscher, 2006; Borgland et al., 2004; Chen et al., 2008; Dong et al., 2004; Engblom et al., 2008; Heikkinen et al., 2009; Liu et al., 2005; Saal et al., 2003; Stuber et al., 2008; Ungless et al., 2001). The identification of putative DA cells was based on low-frequency pacemaker activity, broad action potentials, hyperpolarization by DA via D2 receptors or the presence of the so-called Ih current, generated by hyperpolarization-activated cyclic nucleotide-regulated cation channels (HCN channels) (Kitai et al., 1999). The reliability of criteria for identification of DA neurons in slice recordings has generated some confusion (Ungless and Grace, 2012) because: (1) single-cell labeling studies have revealed that in the VTA the presence of Ih is not always consistent with a DAergic phenotype (Margolis et al., 2008; Zhang et al., 2010), (2) some VTA DA neurons do not respond to DA application (Bannon and Roth, 1983; Lammel et al., 2008) and (3) VTA DA neurons have been identified that have very small or no Ih (Ford et al., 2006; Hnasko et al., 2012; Jones and Kauer, 1999; Lammel et al., 2008, 2011; Witten et al., 2011; Zhang et al., 2010). These findings likely account for the variability in using Ih as a reliable marker for the DA phenotype (Jones and Kauer, 1999; Margolis et al., 2006; Ungless and Grace, 2012; Wanat et al., 2008; Zhang et al., 2010).

While species differences may contribute to this variability (Courtney et al., 2012) it is also likely that recordings have been performed in different subregions of the VTA (Zhang et al., 2010). Many studies that identified putative DA neurons based on their expression of a large Ih performed patch clamp recording from horizontal slices and focused on a specific subregion of the VTA defined as the region medial to the MT (medial terminal nucleus of the accessory optical tract). While in this specific VTA subregion the correlation between Ih and DA phenotype might be high, other VTA subregions (such as the PN and medial PBP of the posterior VTA) have often been ignored and may contain DA neurons with a distinct electrophysiological profile (Lammel et al., 2008). For a more complete discussion of the criteria used to identify DA neurons in the VTA and SN *in vivo* and *in vitro*, we refer readers to a recent review on the topic (Ungless and Grace, 2012).

More recent studies began to classify VTA DA neurons based on their projection targets (Beckley et al., 2013; Ford et al., 2006;

Ikemoto, 2007; Lammel et al., 2008; Margolis et al., 2006, 2008) since it is well established that midbrain DA neurons have widespread projections throughout the brain (Bjorklund and Dunnett, 2007; Ikemoto, 2007). The vast majority of VTA DA neurons appear to send projections to non-overlapping target areas and thus these ascending VTA projections can be conceptualized as independent parallel lines (Fallon, 1981; Albanese and Minciacchi, 1983; Lammel et al., 2008). Combining retrograde tracing, brain slice patch clamp recordings and single cell mRNA quantification of various DAergic marker genes revealed that the mesocorticolimbic DA system consists of several DA neuron subpopulations with different properties (Lammel et al., 2008). Importantly, some VTA DA subpopulations did not exhibit some of the classic properties that previously have been used to identify DA neurons. Injection of fluorescent retrobeads into different brain areas of adult mice and subsequent anatomical analysis of coronal brain slices demonstrated that DA neurons in medial posterior VTA (PN and medial PBP) selectively project to the NAC medial shell and core, medial prefrontal cortex (PFC), and basolateral amygdala (BLA). These “non-conventional” VTA DA neurons had no or only very little Ih and also did not exhibit typical action potential waveforms and firing patterns (Fig. 1). In contrast, DA neurons, which project to NAC lateral shell, were predominantly located in the lateral posterior and anterior VTA (lateral PBP). They exhibited a prominent Ih and firing patterns reminiscent of conventional VTA DA neurons (Hnasko et al., 2012; Lammel et al., 2008; Ungless and Grace, 2012; Zhang et al., 2010) (Fig. 2). Thus, it seems likely that the DA neurons in the lateral VTA (lateral PBP) projecting to NAC lateral shell are the same as those analyzed in studies that focused on the area medial of the MT using horizontal brain slices. The idea that “conventional” DA neurons are located in the lateral portions of the VTA is consistent with recent studies showing that VTA DA neurons have a differential latero-medial cellular expression of AADC, VMAT2, DAT, D2R and VGluT2 (Li et al., 2012). Interestingly, there is also a high degree of molecular and electrophysiological similarity between the populations of DA neurons within the lateral VTA and adjacent SN (Lammel et al., 2008; Li et al., 2012). The discovery of DA subpopulations with distinct anatomical, electrophysiological and molecular properties supports the idea that they also could be involved in mediating different behavioral responses to specific environmental stimuli (Bromberg-Martin et al., 2010).

Unlike VTA DA neurons, SNc DA neurons were thought to be a more homogenous cell population. Indeed, *in vivo* as well as *in vitro* criteria for identification of SNc DA neurons seem to be more reliable than for VTA DA neurons (Ungless and Grace, 2012). However, recent studies report that DA neurons in the SNc exhibit functional heterogeneity that may contribute to their diverse roles in behavior (Brown et al., 2009; Henny et al., 2012; Schieman et al., 2012). Specifically, SNc DA cell functional heterogeneity appears to be correlated with differences in dendrite architecture and afferent connectivity (Henny et al., 2012). Further evidence for heterogeneity in SNc DA cells comes from the observation that K-ATP channels gate bursting selectively in medial SN DA neurons projecting to the dorsomedial striatum but not in lateral SN DA neurons, which project to the dorsolateral striatum as well as VTA DA neurons (Schieman et al., 2012). DA neuronal signaling has recently become even more complex with the demonstration that SNc DA cells release GABA resulting in the inhibition of dorsal striatal medium spiny neurons (Tritsch et al., 2012). Because this GABA release is dependent on the vesicular monoamine transporter, VMAT2, other DA neuron subpopulations may also co-release GABA although this prediction needs to be tested experimentally.




Fig. 1. Dopamine neurons with unconventional electrophysiological properties are mainly located in the medial VTA of the caudal midbrain. (A) Schematic drawing showing that dopamine (DA) neurons projecting to medial prefrontal cortex (mPFC), basolateral amygdala (BLA), nucleus accumbens (NAc) core, and NAc medial shell are predominantly located in the medial VTA (IF, medial PN and medial PBP nuclei) of the caudal midbrain. These VTA subregion and projections are highlighted in green. DA neurons projecting to NAc lateral shell can be found in the lateral VTA (lateral PBP nucleus) of the caudal midbrain (highlighted in yellow). (SNr, substantia nigra pars reticulata; SNc, substantia nigra pars compacta; ml, medial lemniscus; PBP, parabrachial pigmented nucleus; PN, parafascicular nucleus; IF, interfascicular nucleus). (B) Fluorescence microscope image (TH-immunocytochemistry in red) showing the location of DA neurons in lateral (IVTA) and medial (mVTA) VTA of the caudal midbrain. (C, D, E, F) DA neurons located in the mVTA projecting to mPFC, NAc medial shell, NAc core and BLA have unconventional electrophysiological properties including: (C) lack of a prominent Ih current, (D) high maximal firing frequencies ($\sim 20\text{--}30$ Hz), (E) broad action potentials and small after hyperpolarization and (F) a high AMPAR/NMDAR ratio under basal conditions (~ 0.6).

Although this review's focus is the heterogeneity of VTA DA neurons, it is important to mention that a significant proportion of cells in what are traditionally considered DA nuclei are GABAergic or glutamatergic (Nair-Roberts et al., 2008). Furthermore, there are considerable heterogeneities within these neuronal phenotypes (Margolis et al., 2012). Approximately 20–40% of VTA cells are GABAergic (Carr and Sesack, 2000a; Margolis et al., 2012; Nair-Roberts et al., 2008), which not only form local contacts onto both DA and non-DA VTA neurons (Omelchenko and Sesack, 2009) but also project to several forebrain regions (Fields et al., 2007). Recently, it has been reported that VTA GABA neurons projecting to the NAc inhibit accumbal cholinergic interneurons in a manner that enhances stimulus-outcome learning (Brown et al., 2012). However, activation of VTA GABA neurons projecting to NAc does not alter reward consumption (van Zessen et al., 2012). Thus, VTA GABAergic neurons, like VTA DA neurons, may be a specialized and functionally heterogeneous cell population (Margolis et al., 2012).

VTA glutamate neurons selectively express VGLUT2 mRNA and can mainly be found in the medial VTA. It has been suggested that they represent approximately 2–3% of the total VTA cell population (Nair-Roberts et al., 2008). These cells rarely co-express mRNAs for tyrosine hydroxylase (TH) or glutamic acid decarboxylase (GAD) indicating that the vast majority are non-DAergic and non-GABAergic (Nair-Roberts et al., 2008; Yamaguchi et al., 2007). However, it has also been reported that a small subset of DA neurons co-release DA and glutamate (Chuhma et al., 2009; Hnasko et al., 2010; Stuber et al., 2010; Tecuapetla et al., 2010; but see Moss et al., 2011). Importantly, medial VTA glutamate neurons, which were identified in transgenic mice that express EGFP under control of the VGLUT2 promoter, exhibit a small Ih making them indistinguishable from medial VTA DA neurons, but not from lateral VTA DA neurons (Hnasko et al., 2012). VTA glutamate neurons, like VTA DA and GABAergic neurons, project to several different brain areas including NAc, lateral habenula, ventral pallidum, amygdala and medial PFC (Hnasko et al., 2012; Yamaguchi et al., 2011).




Fig. 2. Many dopamine neurons with classical electrophysiological properties are located in the lateral VTA of the rostral midbrain. (A) Schematic drawing showing that dopamine (DA) neurons projecting to nucleus accumbens (NAc) lateral shell are located in the lateral VTA of the rostral midbrain. The VTA subregion and projection is highlighted in yellow. Note, that DA neurons projecting to NAc lateral shell can also be found in the caudal midbrain (see Fig. 1). (Abbreviations are the same as in Fig. 1. fr, fasciculus retroflexus; CM, mammillary body). (B) Fluorescence microscope image (TH-immunocytochemistry in red) showing that in the rostral midbrain the lateral VTA contains many more TH-immunopositive cells than the medial VTA (compare to Fig. 1B; see Lammel et al., 2008 for a detailed anatomic analysis of DA projection neurons). (C, D, E, F) DA neurons located in the IVTA projecting to NAc lateral shell have classical electrophysiological properties including: (C) a prominent I_h current, (D) firing frequencies up to 10 Hz, (E) short action potentials with prominent after hyperpolarizations and (F) a low AMPAR/NMDAR ratio under basal conditions (~ 0.4).

3. Dopamine neuron activity during reward and aversion

For decades investigators have been studying the relationship between midbrain DA neuron activity and reward (Schultz, 1997, 2007, 2012; Wise and Rompre, 1989). The central role of DA neuron activity in reward-related processes has been well-established by electrophysiological studies. Single unit recordings in primates performing an operant task showed that putative DA neurons are phasically excited (i.e. they exhibit burst discharges of action potentials) by unexpected food rewards (Schultz, 1997). An extensive and important body of work subsequently supported the heuristically attractive hypothesis that DA neurons phasic activity encodes reward prediction errors, the discrepancy between an expected reward and the actual outcome. Thus, the omission of an expected reward had the opposite effect on putative DA neuronal activity and silent periods were associated with negative reward prediction errors (Schultz, 1997). The reward prediction error hypothesis has been confirmed in humans using fMRI (D'Ardenne et al., 2008) and more recently using optogenetic single-cell identification of VTA DA neurons in mice (Cohen et al., 2012).

While there is strong evidence in support of the hypothesis that DA neurons signal reward prediction errors, it seems unlikely that this is their only function and that all DA neurons, independent of their projection targets, subserves this single function. Indeed, it is well-established that aversive, stressful and salient events can excite midbrain DA neurons and cause DA release in target structures (Abercrombie et al., 1989; Anstrom and Woodward, 2005; Bassareo et al., 2002; Brischoux et al., 2009; Bromberg-Martin et al., 2010; Cohen et al., 2012; Guarraci and Kapp, 1999; Mantz et al., 1989; Matsumoto and Hikosaka, 2009; Ungless et al., 2010; Young, 2004). However, VTA DA neurons can also be inhibited by aversive stimuli (Mirenovic and Schultz, 1996; Ungless et al., 2004) and it can be postulated that any increase in DA neuron activity caused by an acute aversive stimulus (e.g. foot shock) might be due to the rewarding effects of terminating the stimulus (Tanimoto et al., 2004). Indeed, most DA neurons that were inhibited by foot shocks also showed a phasic excitation at the termination of the foot shock (Brischoux et al., 2009). Studies in rats using juxtacellular labeling–recording techniques supported this idea and also suggested that the VTA neurons that were excited by aversive stimuli were non-DAergic (Ungless et al., 2004). However,

subsequently the same group reported that DA neurons localized in the medioventral VTA were excited at the onset of aversive stimuli (Brischoux et al., 2009).

An unresolved question is what proportion of DA neurons are excited by aversive stimuli and whether they are a specific subpopulation that project to specific targets. While it has been argued that a relatively small proportion of DA neurons respond to aversive stimuli (Schultz, 2012), a recent study in mice found that roughly equivalent numbers of putative VTA DA neurons are activated or inhibited by tail pinch (Zweifel et al., 2011). Furthermore, the single unit activity of many optogenetically identified DA neurons in mice increased in response to an aversive stimulus (Cohen et al., 2012). It is also possible that through sampling biases VTA DA neurons that respond to aversive stimulation have been missed. Indeed, as mentioned earlier in this review, some DA subpopulations are located in a specific subregion of the caudal VTA (PN and medial PBP). Importantly, these cells have unconventional electrophysiological properties (Lammel et al., 2008) (Fig. 1). Thus, in previous single unit recording studies they might not only have been missed but also could have been misinterpreted as non-DAergic if no verification of neurochemical phenotype of the recorded cells was performed.

4. Afferent control of dopamine neuron activity

The VTA receives both excitatory and inhibitory inputs from a broad distribution of brain areas (Geisler et al., 2007; Sesack and Grace, 2010; Watabe-Uchida et al., 2012). Activation of post-synaptic NMDA receptors are particularly important for driving high-frequency bursts of action potentials (Korotkova et al., 2004) and are critical for reward-dependent learning (Zweifel et al., 2009). Clearly, an important line of research is to determine the anatomical connectivity of afferents to VTA neurons and their behavioral functions. While electrical stimulation has been used to activate known VTA inputs *in vivo* and this work has provided important information, with the advent of optogenetics and novel viral tracing strategies (Osakada et al., 2011; Tye and Deisseroth, 2012), it is now possible to determine the anatomical connectivity of specific inputs unequivocally and address their putative behavioral functions. Previous work demonstrated that electrical stimulation of the laterodorsal tegmentum (LDT) promotes burst firing of putative VTA DA neurons and increases DA release in the NAc (Forster and Blaha, 2000; Lodge and Grace, 2006). This is consistent with classical tracing work, which revealed that the LDT projects mainly to the VTA (Cornwall et al., 1990) and makes excitatory synapses on DA neurons projecting to the NAc (Omelchenko and Sesack, 2005). Interestingly, presumed inhibitory LDT inputs seems to be selective for mesoprefrontal but not mesolimbic DA neurons (Omelchenko and Sesack, 2005). In contrast, mesoprefrontal DA neurons receive excitatory input from the PFC (Carr and Sesack, 2000b).

Using *in vivo* rabies-virus mediated expression of channelrhodopsin-2 (ChR2) for optical control of LDT projection neurons combined with *in vitro* acute slice recordings from identified DA neurons with known projection targets, we found that LDT axons predominately make excitatory synaptic connections on DA neurons projecting to the NAc lateral shell (Lammel et al., 2012). Consistent with the idea that DA release in the NAc promotes reinforcement (Steinberg and Janak, 2012; Witten et al., 2011) phasic light activation of these LDT inputs within the VTA elicits conditioned place preference (CPP), which was prevented by infusion of DA receptor antagonists into the NAc (Lammel et al., 2012).

The same approach was taken to examine the connectivity and behavioral functions of the inputs to the VTA from the lateral

habenula (LHb), a brain area thought to be critical for mediating behavioral responses to aversive stimuli and when expected rewards do not occur (Hikosaka, 2010). Consistent with previous findings (Christoph et al., 1986; Hong et al., 2011; Ji and Shepard, 2007), optogenetic activation of LHb axons in acute slices generated excitatory synaptic currents in GABAergic neurons in the rostromedial tegmental nucleus (RMTg) (Lammel et al., 2012; Stamatakis and Stuber, 2012), which is also known as the tail of the VTA (Kaufling et al., 2009). Surprisingly, these same inputs made excitatory synaptic connections onto DA neurons projecting to the mPFC but connections onto other DA neuron subpopulation were not detected (Lammel et al., 2012). Furthermore, the RMTg GABAergic neurons activated by LHb inputs generated inhibitory synaptic currents in DA cells projecting to NAc lateral shell. Thus, activation of LHb inputs to the VTA would be expected to activate DA neurons projecting to mPFC and inhibit DA neurons projecting to NAc lateral shell and therefore generate a profoundly different behavioral response than LDT axon stimulation. Indeed, phasic optogenetic activation of LHb inputs to the VTA elicited conditioned place aversion (CPA), which was prevented by infusion of a DA receptor antagonist into the mPFC (Lammel et al., 2012). Activation of LHb inputs to the VTA has also been shown to elicit both passive and conditioned behavioral avoidance (Stamatakis and Stuber, 2012).

These results suggest that different afferent inputs to the VTA influence distinct DA neuron subpopulations and VTA “microcircuits” and as a consequence influence behavior in profoundly different ways. Recent anatomical tracing studies using recombinant rabies viruses have revealed that the brain areas sending projections to the VTA are much more abundant than previously envisioned (Watabe-Uchida et al., 2012). The approaches taken in this recent work (Lammel et al., 2012) will be helpful in defining the synaptic connectivity of these inputs and their behavioral functions with the long-term goal of generating a much more comprehensive understanding of the circuits in which VTA cells are embedded.

5. VTA dopamine neurons and stress-induced depression

Chronic stress is an important diathesis for depression in humans and is used to generate rodent models of depression. It has long been postulated that malfunction of the brain's reward circuitry may play an important role in mediating key symptoms of stress-elicited behaviors, including depression (Friedman et al., 2008; Nestler and Carlezon, 2006; Willner, 1991; Yadid and Friedman, 2008). Here we will briefly review the effects of stress on midbrain DA neurons. Both chronic restraint stress and repeated social defeat stress have been found to increase the spontaneous and burst firing of VTA DA neurons *in vivo* and *ex vivo* in brain slices (Anstrom and Woodward, 2005; Cao et al., 2010; Feder et al., 2009; Krishnan et al., 2007). Importantly, in the repeated social defeat stress model of depression, the increase in DA neuron activity only occurred in susceptible but not in resilient mice (Cao et al., 2010; Feder et al., 2009; Krishnan et al., 2007) and lasted for 3 weeks after the social defeat protocol (Razzoli et al., 2011). Moreover, the increased firing rates and bursting events in VTA DA neurons were largely reversed by chronic administration of the anti-depressant fluoxetine (Cao et al., 2010).

In the context of this review, a critical question is whether a specific subpopulation of VTA DA neurons exhibits this stress-induced change in firing patterns and which subpopulations contribute to the stress-induced behavioral changes. An *in vivo* microdialysis study in rats found that levels of DA in both NAc and PFC were elevated by a social defeat stress (Tidey and Miczek, 1996). Much more recently, optogenetic activation of VTA DA neuron phasic firing was shown to elicit a susceptible, depression-associated

phenotype in mice undergoing a subthreshold social defeat protocol as well as in previously resilient animals that had been subjected to repeated social defeat stress (Chaudhury et al., 2013). Surprisingly, these dramatic behavioral effects appear to be mediated by release of DA in NAc as terminal field stimulation of DA fibers in NAc, but not mPFC, induced susceptibility to social defeat stress (Chaudhury et al., 2013). In marked contrast, another group found that optogenetic activation of VTA DA neurons reversed chronic stress-induced depression-associated behaviors while inhibition of these neurons promoted the same behaviors (Tye et al., 2013). Infusion of DA receptor antagonists into the NAc prevented the consequences of VTA DA neuron activation in one key depression-associated behavioral assay suggesting that DA release specifically in the NAc is required for the anti-depressant consequences of VTA DA neuron activation.

Providing an explanation for the discrepancies in these two studies is challenging. Both studies demonstrate that activity of VTA DA neurons projecting to NAc importantly influence depression-associated behaviors but one study concludes such activity is pro-depressant while the other concludes it is anti-depressant. These findings also need to be interpreted in the context of other studies reporting that optogenetic activation of VTA DA neurons can induce CPP and facilitate the development of positive reinforcement during reward seeking (Tsai et al., 2009; Adamantidis et al., 2011). It is possible, although disconcerting, that the different stress protocols used in the two studies (i.e. repeated social defeat stress versus chronic mild stress) caused dramatically different effects on the circuits being studied. Alternatively, there may have been differences in the specific subpopulations of VTA DA neurons that were manipulated. VTA DA neurons projecting to NAc can be subdivided based on their specific projection target (i.e. core versus medial shell versus lateral shell) and these differences may be behaviorally important (Ikemoto, 2007; Lammel et al., 2008, 2011, 2012). The potential involvement of the LHB in depression (Li et al., 2011; Morris et al., 1999; Shumake and Gonzalez-Lima, 2003) and the specific synaptic connectivity of LHB inputs in the VTA (Hikosaka, 2010; Lammel et al., 2012) provide additional evidence supporting the importance of further exploring the roles of specific VTA DA neuron subpopulations and the afferent inputs that drive them in depression-associated behaviors.

6. Effects of drugs of abuse on VTA dopamine neuron excitatory synapses

The mesocorticolimbic DA system has a central role in the acquisition of behaviors that are inappropriately reinforced by drugs of abuse. Drugs of abuse such as cocaine, morphine, nicotine and amphetamine have different pharmacological effects (for recent review see Luscher and Malenka, 2011) yet they all significantly impact reward and motivation at least in part by increasing DA release in the NAc. However, there is a growing body of evidence that motivational stimuli, stress, as well as drugs of abuse evoke DA release in anatomically and functionally distinct DA projections (Abercrombie et al., 1989; Aragona et al., 2008, 2009; Bassareo et al., 2002; Ikemoto, 2007; Ito et al., 2000; Lammel et al., 2011; Lemos et al., 2012; Porter-Stransky et al., 2011; Roitman et al., 2008; Salamone and Correa, 2012; Young, 2004). Increased extracellular DA concentrations, such as that elicited by abused drugs or motivational stimuli, likely facilitate learning and memory (Luscher and Malenka, 2011; Schultz, 2012). Thus, it is not surprising that an important research topic over the last decade has been to elucidate how motivational stimuli, stress, and drugs of abuse induce adaptations of VTA DA neuron excitatory inputs, which drive spiking of VTA DA neurons (Argilli et al., 2008; Bellone and Luscher, 2006; Borgland et al., 2004; Chen et al., 2008; Dong et al., 2004; Engblom et al., 2008; Heikkinen et al., 2009; Liu et al., 2005; Luscher and

Malenka, 2011; Saal et al., 2003; Stuber et al., 2008; Ungless et al., 2001).

Briefly, these studies all support the hypothesis that all drugs of abuse as well as strong acute stress are sufficient to induce a potentiation of AMPA receptor-mediated synaptic transmission in VTA DA neurons. Natural rewards such as food can also cause LTP-like synaptic changes in VTA DA neurons but these are transient whereas the potentiation caused by self-administration of cocaine lasts at least 3 months after abstinence (Chen et al., 2008). Cocaine administration has also been reported to cause a decrease in NMDA receptor-mediated synaptic responses (Mameli et al., 2011). Importantly, these initial drug-induced synaptic adaptations in VTA DA neurons are thought to be critical for subsequent circuit modifications in downstream targets such as the NAc (Mameli et al., 2009; Wolf and Tseng, 2012).

A limitation of all of these studies is that the putative VTA DA neurons from which recordings were made were commonly identified by the presence of a large Ih, which, as reviewed above, is not present in important subpopulations of VTA DA neurons. To address this limitation, we examined the modification of excitatory synapses on identified VTA DA neuron subpopulations caused by cocaine administration as well as a strong aversive stimulus (Lammel et al., 2011). A single injection of cocaine strongly modified excitatory synapses on DA neurons projecting to the NAc medial shell but not synapses on DA neurons projecting to mPFC, as assayed by the ratio of AMPAR-mediated synaptic responses to NMDAR-mediated synaptic responses (see Kauer and Malenka, 2007 for discussion of the utility of this so-called AMPAR/NMDAR assay). An aversive stimulus (i.e. formalin injection into a single hindpaw) had the opposite effect; modifying the synapses on the medial VTA DA neurons projecting to mPFC but not those on DA neurons projecting to NAc medial shell. Both the rewarding and aversive experiences modified synapses on DA neurons projecting to NAc lateral shell. These results provide further evidence that different subpopulations of VTA DA neurons participate in related but independent circuits that contribute to the brain's response to reward, aversion and saliency.

Using optogenetics, a critical question that can now be addressed is whether distinct excitatory inputs making synapses on VTA DA neurons are modified differently by rewarding versus aversive stimuli. And if so, what are the circuit and behavioral consequences of such input-specific synaptic modifications. VTA DA neurons also receive inhibitory synaptic inputs from NAc medium spiny neurons, the RMTg as well as from local interneurons within the VTA, which themselves receive projections from the NAc. A more definitive elucidation of this inhibitory circuitry within the VTA and how these inhibitory synapses on VTA DA neuron are modified by experience will also be important for a more comprehensive understanding of the neural circuit mechanisms mediating reward- and aversion associated behaviors.

7. Conclusions

We have attempted to concisely summarize the evidence supporting the idea that VTA DA neurons are heterogeneous not only in regard to their anatomical, molecular and electrophysiological properties but also in their response to salient appetitive and aversive stimuli. Heterogeneity in DA neurons and their behavioral functions has also been observed in *Drosophila* (Claridge-Chang et al., 2009; Krashes et al., 2009; Liu et al., 2012), suggesting strong evolutionary pressure to conserve such heterogeneity as nervous systems evolved. Consistent with this view, many recent findings challenge the view that DA release in target structures has a primary or exclusive role in reward processing (Bromberg-Martin et al., 2010; Salamone and Correa, 2012) although arguments can be

made that the primary role of DA neuron phasic activity remains the generation of reward prediction error signals (Schultz, 2012). Given DA's prominent role as a neuromodulator of neuronal excitability and synaptic function, intuitively it seems likely that DA release in different target structures containing complex neural networks composed of diverse cell populations (e.g. mPFC, dorsal and ventral striatum) will have different behavioral consequences. Recent studies investigating the contribution of DA subsystems in rodent models of depression, addiction, aversive behavior and Parkinson's disease (Aragona et al., 2009; Beckley et al., 2013; Brischoux et al., 2009; Chaudhury et al., 2013; Lammel et al., 2011, 2012; Schieman et al., 2012; Tye et al., 2013) provide further evidence supporting the importance of a heterogeneous midbrain DA system in mediating the pathophysiology of prominent neuropsychiatric disorders. Over the next decade, we anticipate that further work on elucidating the organization, function and modification of midbrain DA circuits will importantly contribute to understanding how motivational systems are organized in the brain and mediate adaptive and pathological behaviors.

Acknowledgements

Work in the authors' laboratory is supported by grants from the Simons Foundation Autism Research Initiative and NIH (to R.C.M.). S.L. is supported by a fellowship from the German Academy of Sciences Leopoldina. B.K.L. is supported by a Davis Foundation Postdoctoral Fellowship in Eating Disorders Research.

References

- Abercrombie, E.D., Keefe, K.A., DiFrischia, D.S., Zigmond, M.J., 1989. Differential effect of stress on in vivo dopamine release in striatum, nucleus accumbens, and medial frontal cortex. *J. Neurochem.* 52, 1655–1658.
- Adamantidis, A.R., Tsai, H.C., Boutrel, B., Zhang, F., Stuber, G.D., Budygin, E.A., Tourino, C., Bonci, A., Deisseroth, K., de Lecea, L., 2011. Optogenetic interrogation of dopaminergic modulation of the multiple phases of reward-seeking behavior. *J. Neurosci.* 31, 10829–10835.
- Albanese, A., Minciacchi, D., 1983. Organization of the ascending projections from the ventral tegmental area: a multiple fluorescent retrograde tracer study in the rat. *J. Comp. Neurol.* 216, 406–420.
- Anstrom, K.K., Woodward, D.J., 2005. Restraint increases dopaminergic burst firing in awake rats. *Neuropsychopharmacology* 30, 1832–1840.
- Aragona, B.J., Cleaveland, N.A., Stuber, G.D., Day, J.J., Carelli, R.M., Wightman, R.M., 2008. Preferential enhancement of dopamine transmission within the nucleus accumbens shell by cocaine is attributable to a direct increase in phasic dopamine release events. *J. Neurosci.* 28, 8821–8831.
- Aragona, B.J., Day, J.J., Roitman, M.F., Cleaveland, N.A., Wightman, R.M., Carelli, R.M., 2009. Regional specificity in the real-time development of phasic dopamine transmission patterns during acquisition of a cue-cocaine association in rats. *Eur. J. Neurosci.* 30, 1889–1899.
- Argilli, E., Sibley, D.R., Malenka, R.C., England, P.M., Bonci, A., 2008. Mechanism and time course of cocaine-induced long-term potentiation in the ventral tegmental area. *J. Neurosci.* 28, 9092–9100.
- Bannon, M.J., Roth, R.H., 1983. Pharmacology of mesocortical dopamine neurons. *Pharmacol. Rev.* 35, 53–68.
- Bassareo, V., De Luca, M.A., Di Chiara, G., 2002. Differential expression of motivational stimulus properties by dopamine in nucleus accumbens shell versus core and prefrontal cortex. *J. Neurosci.* 22, 4709–4719.
- Beckley, J.T., Evins, C.E., Fedarovich, H., Gilstrap, M.J., Woodward, J.J., 2013. Medial prefrontal cortex inversely regulates toluene-induced changes in markers of synaptic plasticity of mesolimbic dopamine neurons. *J. Neurosci.* 33, 804–813.
- Bellone, C., Luscher, C., 2006. Cocaine triggered AMPA receptor redistribution is reversed in vivo by mGluR-dependent long-term depression. *Nat. Neurosci.* 9, 636–641.
- Bjorklund, A., Dunnett, S.B., 2007. Dopamine neuron systems in the brain: an update. *Trends Neurosci.* 30, 194–202.
- Borgland, S.L., Malenka, R.C., Bonci, A., 2004. Acute and chronic cocaine-induced potentiation of synaptic strength in the ventral tegmental area: electrophysiological and behavioral correlates in individual rats. *J. Neurosci.* 24, 7482–7490.
- Brischoux, F., Chakraborty, S., Brierley, D.I., Ungless, M.A., 2009. Phasic excitation of dopamine neurons in ventral VTA by noxious stimuli. *Proc. Natl. Acad. Sci. U. S. A.* 106, 4894–4899.
- Bromberg-Martin, E.S., Matsumoto, M., Hikosaka, O., 2010. Dopamine in motivational control: rewarding, aversive, and alerting. *Neuron* 68, 815–834.
- Brown, M.T., Henny, P., Bolam, J.P., Magill, P.J., 2009. Activity of neurochemically heterogeneous dopaminergic neurons in the substantia nigra during spontaneous and driven changes in brain state. *J. Neurosci.* 29, 2915–2925.
- Brown, M.T., Tan, K.R., O'Connor, E.C., Nikonenko, I., Muller, D., Luscher, C., 2012. Ventral tegmental area GABA projections pause accumbal cholinergic interneurons to enhance associative learning. *Nature* 492, 452–456.
- Cameron, D.L., Wessendorf, M.W., Williams, J.T., 1997. A subset of ventral tegmental area neurons is inhibited by dopamine, 5-hydroxytryptamine and opioids. *Neuroscience* 77, 155–166.
- Cao, J.L., Covington 3rd, H.E., Friedman, A.K., Wilkinson, M.B., Walsh, J.J., Cooper, D.C., Nestler, E.J., Han, M.H., 2010. Mesolimbic dopamine neurons in the brain reward circuit mediate susceptibility to social defeat and antidepressant action. *J. Neurosci.* 30, 16453–16458.
- Carr, D.B., Sesack, S.R., 2000a. GABA-containing neurons in the rat ventral tegmental area project to the prefrontal cortex. *Synapse* 38, 114–123.
- Carr, D.B., Sesack, S.R., 2000b. Projections from the rat prefrontal cortex to the ventral tegmental area: target specificity in the synaptic associations with mesoaccumbens and mesocortical neurons. *J. Neurosci.* 20, 3864–3873.
- Chaudhury, D., Walsh, J.J., Friedman, A.K., Juarez, B., Ku, S.M., Koo, J.W., Ferguson, D., Tsai, H.-C., Pomeranz, L., Christoffel, D.J., Nectow, A.R., Ekstrand, M., Domingos, A., Mazei-Robison, M.S., Mouzon, E., Lobo, M.K., Neve, R.L., Friedman, J.M., Russo, S.J., Deisseroth, K., Nestler, E.J., Han, M.H., 2013. Rapid regulation of depression-related behaviours by control of midbrain dopamine neurons. *Nature* 493, 532–536.
- Chen, B.T., Bowers, M.S., Martin, M., Hopf, F.W., Guillory, A.M., Carelli, R.M., Chou, J.K., Bonci, A., 2008. Cocaine but not natural reward self-administration nor passive cocaine infusion produces persistent LTP in the VTA. *Neuron* 59, 288–297.
- Christoph, G.R., Leonzio, R.J., Wilcox, K.S., 1986. Stimulation of the lateral habenula inhibits dopamine-containing neurons in the substantia nigra and ventral tegmental area of the rat. *J. Neurosci.* 6, 613–619.
- Chuhma, N., Choi, W.Y., Mingote, S., Rayport, S., 2009. Dopamine neuron glutamate cotransmission: frequency-dependent modulation in the mesoventromedial projection. *Neuroscience* 164, 1068–1083.
- Claridge-Chang, A., Roorda, R.D., Vrontou, E., Sjulson, L., Li, H., Hirsh, J., Miesenbock, G., 2009. Writing memories with light-addressable reinforcement circuitry. *Cell* 139, 405–415.
- Cohen, J.Y., Haesler, S., Vong, L., Lowell, B.B., Uchida, N., 2012. Neuron-type-specific signals for reward and punishment in the ventral tegmental area. *Nature* 482, 85–88.
- Cornwall, J., Cooper, J.D., Phillipson, O.T., 1990. Afferent and efferent connections of the laterodorsal tegmental nucleus in the rat. *Brain Res. Bull.* 25, 271–284.
- Courtney, N.A., Mamaligas, A.A., Ford, C.P., 2012. Species differences in somatodendritic dopamine transmission determine D2-autoreceptor-mediated inhibition of ventral tegmental area neuron firing. *J. Neurosci.* 32, 13520–13528.
- D'Ardenne, K., McClure, S.M., Nystrom, L.E., Cohen, J.D., 2008. BOLD responses reflecting dopaminergic signals in the human ventral tegmental area. *Science* 319, 1264–1267.
- Dong, Y., Saal, D., Thomas, M., Faust, R., Bonci, A., Robinson, T., Malenka, R.C., 2004. Cocaine-induced potentiation of synaptic strength in dopamine neurons: behavioral correlates in GluRA(−/−) mice. *Proc. Natl. Acad. Sci. U. S. A.* 101, 14282–14287.
- El Mestikawy, S., Wallen-Mackenzie, A., Fortin, G.M., Descarries, L., Trudeau, L.E., 2011. From glutamate co-release to vesicular synergy: vesicular glutamate transporters. *Nat. Rev. Neurosci.* 12, 204–216.
- Engblom, D., Bilbao, A., Sanchis-Segura, C., Dahan, L., Perreau-Lenz, S., Balland, B., Parkitna, J.R., Lujan, R., Halbout, B., Mameli, M., Parlato, R., Sprengel, R., Luscher, C., Schutz, G., Spanagel, R., 2008. Glutamate receptors on dopamine neurons control the persistence of cocaine seeking. *Neuron* 59, 497–508.
- Fallon, J.H., 1981. Collateralization of monoamine neurons: mesotelencephalic dopamine projections to caudate, septum, and frontal cortex. *J. Neurosci.* 1, 1361–1368.
- Feder, A., Nestler, E.J., Charney, D.S., 2009. Psychobiology and molecular genetics of resilience. *Nat. Rev. Neurosci.* 10, 446–457.
- Fields, H.L., Hjelmstad, G.O., Margolis, E.B., Nicola, S.M., 2007. Ventral tegmental area neurons in learned appetitive behavior and positive reinforcement. *Annu. Rev. Neurosci.* 30, 289–316.
- Ford, C.P., Mark, G.P., Williams, J.T., 2006. Properties and opioid inhibition of mesolimbic dopamine neurons vary according to target location. *J. Neurosci.* 26, 2788–2797.
- Forster, G.L., Blaha, C.D., 2000. Laterodorsal tegmental stimulation elicits dopamine efflux in the rat nucleus accumbens by activation of acetylcholine and glutamate receptors in the ventral tegmental area. *Eur. J. Neurosci.* 12, 3596–3604.
- Friedman, A., Friedman, Y., Dremencov, E., Yadid, G., 2008. VTA dopamine neuron bursting is altered in an animal model of depression and corrected by desipramine. *J. Mol. Neurosci.* 34, 201–209.
- Geisler, S., Derst, C., Veh, R.W., Zahm, D.S., 2007. Glutamatergic afferents of the ventral tegmental area in the rat. *J. Neurosci.* 27, 5730–5743.
- Grace, A.A., Onn, S.P., 1989. Morphology and electrophysiological properties of immunocytochemically identified rat dopamine neurons recorded in vitro. *J. Neurosci.* 9, 3463–3481.
- Guarraci, F.A., Kapp, B.S., 1999. An electrophysiological characterization of ventral tegmental area dopaminergic neurons during differential pavlovian fear conditioning in the awake rabbit. *Behav. Brain Res.* 99, 169–179.

- Heikkinen, A.E., Moykkynen, T.P., Korpi, E.R., 2009. Long-lasting modulation of glutamatergic transmission in VTA dopamine neurons after a single dose of benzodiazepine agonists. *Neuropsychopharmacology* 34, 290–298.
- Henny, P., Brown, M.T., Northrop, A., Faunes, M., Ungless, M.A., Magill, P.J., Bolam, J.P., 2012. Structural correlates of heterogeneous *in vivo* activity of midbrain dopaminergic neurons. *Nat. Neurosci.* 15, 613–619.
- Hikosaka, O., 2010. The habenula: from stress evasion to value-based decision-making. *Nat. Rev. Neurosci.* 11, 503–513.
- Hnasko, T.S., Chuhma, N., Zhang, H., Goh, G.Y., Sulzer, D., Palmiter, R.D., Rayport, S., Edwards, R.H., 2010. Vesicular glutamate transport promotes dopamine storage and glutamate corelease *in vivo*. *Neuron* 65, 643–656.
- Hnasko, T.S., Hjelmstad, G.O., Fields, H.L., Edwards, R.H., 2012. Ventral tegmental area glutamate neurons: electrophysiological properties and projections. *J. Neurosci.* 32, 15076–15085.
- Hong, S., Jhou, T.C., Smith, M., Saleem, K.S., Hikosaka, O., 2011. Negative reward signals from the lateral habenula to dopamine neurons are mediated by rostromedial tegmental nucleus in primates. *J. Neurosci.* 31, 11457–11471.
- Ikemoto, S., 2007. Dopamine reward circuitry: two projection systems from the ventral midbrain to the nucleus accumbens-olfactory tubercle complex. *Brain Res. Rev.* 56, 27–78.
- Ito, R., Dalley, J.W., Howes, S.R., Robbins, T.W., Everitt, B.J., 2000. Dissociation in conditioned dopamine release in the nucleus accumbens core and shell in response to cocaine cues and during cocaine-seeking behavior in rats. *J. Neurosci.* 20, 7489–7495.
- Ji, H., Shepard, P.D., 2007. Lateral habenula stimulation inhibits rat midbrain dopamine neurons through a GABA(A) receptor-mediated mechanism. *J. Neurosci.* 27, 6923–6930.
- Johnson, S.W., North, R.A., 1992. Two types of neurone in the rat ventral tegmental area and their synaptic inputs. *J. Physiol.* 450, 455–468.
- Jones, S., Kauer, J.A., 1999. Amphetamine depresses excitatory synaptic transmission via serotonin receptors in the ventral tegmental area. *J. Neurosci.* 19, 9780–9787.
- Kauer, J.A., Malenka, R.C., 2007. Synaptic plasticity and addiction. *Nat. Rev. Neurosci.* 8, 844–858.
- Kaufling, J., Veinante, P., Pawlowski, S.A., Freund-Mercier, M.J., Barrot, M., 2009. Afferents to the GABAergic tail of the ventral tegmental area in the rat. *J. Comp. Neurol.* 513, 597–621.
- Kitai, S.T., Shepard, P.D., Callaway, J.C., Scroggs, R., 1999. Afferent modulation of dopamine neuron firing patterns. *Curr. Opin. Neurobiol.* 9, 690–697.
- Korotkova, T.M., Ponomarenko, A.A., Brown, R.E., Haas, H.L., 2004. Functional diversity of ventral midbrain dopamine and GABAergic neurons. *Mol. Neurobiol.* 29, 243–259.
- Krashes, M.J., DasGupta, S., Vreede, A., White, B., Armstrong, J.D., Waddell, S., 2009. A neural circuit mechanism integrating motivational state with memory expression in *Drosophila*. *Cell* 139, 416–427.
- Krishnan, V., Han, M.H., Graham, D.L., Berton, O., Renthal, W., Russo, S.J., Laplant, Q., Graham, A., Lutter, M., Lagace, D.C., Ghose, S., Reister, R., Tannous, P., Green, T.A., Neve, R.L., Chakravarty, S., Kumar, A., Eisch, A.J., Self, D.W., Lee, E.S., Tamminga, C.A., Cooper, D.C., Gershenfeld, H.K., Nestler, E.J., 2007. Molecular adaptations underlying susceptibility and resistance to social defeat in brain reward regions. *Cell* 131, 391–404.
- Lammel, S., Hetzel, A., Hackel, O., Jones, I., Liss, B., Roeper, J., 2008. Unique properties of mesoprefrontal neurons within a dual mesocorticolimbic dopamine system. *Neuron* 57, 760–773.
- Lammel, S., Ion, D.J., Roeper, J., Malenka, R.C., 2011. Projection-specific modulation of dopamine neuron synapses by aversive and rewarding stimuli. *Neuron* 70, 855–862.
- Lammel, S., Lim, B.K., Ran, C., Huang, K.W., Betley, M.J., Tye, K.M., Deisseroth, K., Malenka, R.C., 2012. Input-specific control of reward and aversion in the ventral tegmental area. *Nature* 491, 212–217.
- Lemos, J.C., Wanat, M.J., Smith, J.S., Reyes, B.A., Hollon, N.G., Van Bockstaele, E.J., Chavkin, C., Phillips, P.E., 2012. Severe stress switches CRF action in the nucleus accumbens from appetitive to aversive. *Nature* 490, 402–406.
- Li, B., Piriz, J., Mirrione, M., Chung, C., Proulx, C.D., Schulz, D., Henn, F., Malinow, R., 2011. Synaptic potentiation onto habenula neurons in the learned helplessness model of depression. *Nature* 470, 535–539.
- Li, X., Qi, J., Yamaguchi, T., Wang, H.L., Morales, M., 2012. Heterogeneous composition of dopamine neurons of the rat A10 region: molecular evidence for diverse signaling properties. *Brain Struct. Funct.* (Epub ahead of print).
- Liu, C., Placais, P.Y., Yamagata, N., Pfeiffer, B.D., Aso, Y., Friedrich, A.B., Siwanowicz, I., Rubin, G.M., Preat, T., Tanimoto, H., 2012. A subset of dopamine neurons signals reward for odour memory in *Drosophila*. *Nature* 488, 512–516.
- Liu, Q.S., Pu, L., Poo, M.M., 2005. Repeated cocaine exposure *in vivo* facilitates LTP induction in midbrain dopamine neurons. *Nature* 437, 1027–1031.
- Lodge, D.J., Grace, A.A., 2006. The laterodorsal tegmentum is essential for burst firing of ventral tegmental area dopamine neurons. *Proc. Natl. Acad. Sci. U. S. A.* 103, 5167–5172.
- Luscher, C., Malenka, R.C., 2011. Drug-evoked synaptic plasticity in addiction: from molecular changes to circuit remodeling. *Neuron* 69, 650–663.
- Mameli, M., Bellone, C., Brown, M.T., Luscher, C., 2011. Cocaine inverts rules for synaptic plasticity of glutamate transmission in the ventral tegmental area. *Nat. Neurosci.* 14, 414–416.
- Mameli, M., Halbout, B., Creton, C., Engblom, D., Parkitna, J.R., Spanagel, R., Luscher, C., 2009. Cocaine-evoked synaptic plasticity: persistence in the VTA triggers adaptations in the NAc. *Nat. Neurosci.* 12, 1036–1041.
- Mantz, J., Thierry, A.M., Glowinski, J., 1989. Effect of noxious tail pinch on the discharge rate of mesocortical and mesolimbic dopamine neurons: selective activation of the mesocortical system. *Brain Res.* 476, 377–381.
- Margolis, E.B., Lock, H., Hjelmstad, G.O., Fields, H.L., 2006. The ventral tegmental area revisited: is there an electrophysiological marker for dopaminergic neurons? *J. Physiol.* 577, 907–924.
- Margolis, E.B., Mitchell, J.M., Ishikawa, J., Hjelmstad, G.O., Fields, H.L., 2008. Midbrain dopamine neurons: projection target determines action potential duration and dopamine D(2) receptor inhibition. *J. Neurosci.* 28, 8908–8913.
- Margolis, E.B., Toy, B., Himmels, P., Morales, M., Fields, H.L., 2012. Identification of rat ventral tegmental area GABAergic neurons. *PLoS One* 7, e42365.
- Matsumoto, M., Hikosaka, O., 2009. Two types of dopamine neuron distinctly code positive and negative motivational signals. *Nature* 459, 837–841.
- Mirenowicz, J., Schultz, W., 1996. Preferential activation of midbrain dopamine neurons by appetitive rather than aversive stimuli. *Nature* 379, 449–451.
- Morris, J.S., Smith, K.A., Cowen, P.J., Friston, K.J., Dolan, R.J., 1999. Covariation of activity in habenula and dorsal raphe nuclei following tryptophan depletion. *Neuroimage* 10, 163–172.
- Moss, J., Ungless, M.A., Bolam, J.P., 2011. Dopaminergic axons in different divisions of the adult rat striatal complex do not express vesicular glutamate transporters. *Eur. J. Neurosci.* 33, 1205–1211.
- Nair-Roberts, R.G., Chatelain-Badie, S.D., Benson, E., White-Cooper, H., Bolam, J.P., Ungless, M.A., 2008. Stereological estimates of dopaminergic, GABAergic and glutamatergic neurons in the ventral tegmental area, substantia nigra and retrorubral field in the rat. *Neuroscience* 152, 1024–1031.
- Nestler, E.J., Carlezon Jr., W.A., 2006. The mesolimbic dopamine reward circuit in depression. *Biol. Psychiatry* 59, 1151–1159.
- Oades, R.D., Halliday, G.M., 1987. Ventral tegmental (A10) system: neurobiology. 1. Anatomy and connectivity. *Brain Res.* 434, 117–165.
- Omelchenko, N., Sesack, S.R., 2005. Laterodorsal tegmental projections to identified cell populations in the rat ventral tegmental area. *J. Comp. Neurol.* 483, 217–235.
- Omelchenko, N., Sesack, S.R., 2009. Ultrastructural analysis of local collaterals of rat ventral tegmental area neurons: GABA phenotype and synapses onto dopamine and GABA cells. *Synapse* 63, 895–906.
- Osakada, F., Mori, T., Cetin, A.H., Marshel, J.H., Virgen, B., Callaway, E.M., 2011. New rabies virus variants for monitoring and manipulating activity and gene expression in defined neural circuits. *Neuron* 71, 617–631.
- Porter-Stransky, K.A., Wescott, S.A., Hershman, M., Badrinarayan, A., Vander Weele, C.M., Lovic, V., Aragona, B.J., 2011. Cocaine must enter the brain to evoke unconditioned dopamine release within the nucleus accumbens shell. *Neurosci. Lett.* 504, 13–17.
- Razzoli, M., Andreoli, M., Michielin, F., Quarta, D., Sokal, D.M., 2011. Increased phasic activity of VTA dopamine neurons in mice 3 weeks after repeated social defeat. *Behav. Brain Res.* 218, 253–257.
- Roitman, M.F., Wheeler, R.A., Wightman, R.M., Carelli, R.M., 2008. Real-time chemical responses in the nucleus accumbens differentiate rewarding and aversive stimuli. *Nat. Neurosci.* 11, 1376–1377.
- Saal, D., Dong, Y., Bonci, A., Malenka, R.C., 2003. Drugs of abuse and stress trigger a common synaptic adaptation in dopamine neurons. *Neuron* 37, 577–582.
- Salamone, J.D., Correa, M., 2012. The mysterious motivational functions of mesolimbic dopamine. *Neuron* 76, 470–485.
- Schiemann, J., Schlaudraff, F., Klose, V., Bingmer, M., Seino, S., Magill, P.J., Zaghoul, K.A., Schneider, G., Liss, B., Roeper, J., 2012. K-ATP channels in dopamine substantia nigra neurons control bursting and novelty-induced exploration. *Nat. Neurosci.* 15, 1272–1280.
- Schultz, W., 1997. Dopamine neurons and their role in reward mechanisms. *Curr. Opin. Neurobiol.* 7, 191–197.
- Schultz, W., 2007. Multiple dopamine functions at different time courses. *Annu. Rev. Neurosci.* 30, 259–288.
- Schultz, W., 2012. Updating dopamine reward signals. *Curr. Opin. Neurobiol.* 23, 229–238.
- Sesack, S.R., Grace, A.A., 2010. Cortico-Basal Ganglia reward network: microcircuitry. *Neuropsychopharmacology* 35, 27–47.
- Shumake, J., Gonzalez-Lima, F., 2003. Brain systems underlying susceptibility to helplessness and depression. *Behav. Cogn. Neurosci. Rev.* 2, 198–221.
- Stamatakis, A.M., Stuber, G.D., 2012. Activation of lateral habenula inputs to the ventral midbrain promotes behavioral avoidance. *Nat. Neurosci.* 15, 1105–1107.
- Steinberg, E.E., Janak, P.H., 2012. Establishing causality for dopamine in neural function and behavior with optogenetics. *Brain Res.* (Epub ahead of print).
- Stuber, G.D., Hnasko, T.S., Britt, J.P., Edwards, R.H., Bonci, A., 2010. Dopaminergic terminals in the nucleus accumbens but not the dorsal striatum corelease glutamate. *J. Neurosci.* 30, 8229–8233.
- Stuber, G.D., Klanker, M., de Ridder, B., Bowers, M.S., Joosten, R.N., Feenstra, M.G., Bonci, A., 2008. Reward-predictive cues enhance excitatory synaptic strength onto midbrain dopamine neurons. *Science* 321, 1690–1692.
- Swanson, L.W., 1982. The projections of the ventral tegmental area and adjacent regions: a combined fluorescent retrograde tracer and immunofluorescence study in the rat. *Brain Res. Bull.* 9, 321–353.
- Tanimoto, H., Heisenberg, M., Gerber, B., 2004. Experimental psychology: event timing turns punishment to reward. *Nature* 430, 983.
- Tecuapetla, F., Patel, J.C., Xenias, H., English, D., Tadros, I., Shah, F., Berlin, J., Deisseroth, K., Rice, M.E., Tepper, J.M., Koos, T., 2010. Glutamatergic signaling by mesolimbic dopamine neurons in the nucleus accumbens. *J. Neurosci.* 30, 7105–7110.

- Tidey, J.W., Miczek, K.A., 1996. Social defeat stress selectively alters mesocorticolimbic dopamine release: an in vivo microdialysis study. *Brain Res.* 721, 140–149.
- Tritsch, N.X., Ding, J.B., Sabatini, B.L., 2012. Dopaminergic neurons inhibit striatal output through non-canonical release of GABA. *Nature* 490, 262–266.
- Tsai, H.C., Zhang, F., Adamantidis, A., Stuber, G.D., Bonci, A., de Lecea, L., Deisseroth, K., 2009. Phasic firing in dopaminergic neurons is sufficient for behavioral conditioning. *Science* 324, 1080–1084.
- Tye, K.M., Deisseroth, K., 2012. Optogenetic investigation of neural circuits underlying brain disease in animal models. *Nat. Rev. Neurosci.* 13, 251–266.
- Tye, K.M., Mirzabekov, J.J., Warden, M.R., Ferenczi, E.A., Tsai, H.-C., Finkelstein, J., Kim, S.-Y., Adhikari, A., Thompson, K.R., Andalman, A.S., Gunaydin, L.A., Witten, I.B., Deisseroth, K., 2013. Dopamine neurons modulate neural encoding and expression of depression-related behaviour. *Nature* 493, 537–541.
- Ungless, M.A., Argilli, E., Bonci, A., 2010. Effects of stress and aversion on dopamine neurons: implications for addiction. *Neurosci. Biobehav. Rev.* 35, 151–156.
- Ungless, M.A., Grace, A.A., 2012. Are you or aren't you? Challenges associated with physiologically identifying dopamine neurons. *Trends Neurosci.* 35, 422–430.
- Ungless, M.A., Magill, P.J., Bolam, J.P., 2004. Uniform inhibition of dopamine neurons in the ventral tegmental area by aversive stimuli. *Science* 303, 2040–2042.
- Ungless, M.A., Whistler, J.L., Malenka, R.C., Bonci, A., 2001. Single cocaine exposure in vivo induces long-term potentiation in dopamine neurons. *Nature* 411, 583–587.
- van Zessen, R., Phillips, J.L., Budygin, E.A., Stuber, G.D., 2012. Activation of VTA GABA neurons disrupts reward consumption. *Neuron* 73, 1184–1194.
- Wanat, M.J., Hopf, F.W., Stuber, G.D., Phillips, P.E., Bonci, A., 2008. Corticotropin-releasing factor increases mouse ventral tegmental area dopamine neuron firing through a protein kinase C-dependent enhancement of Ih. *J. Physiol.* 586, 2157–2170.
- Watabe-Uchida, M., Zhu, L., Ogawa, S.K., Vamanrao, A., Uchida, N., 2012. Whole-brain mapping of direct inputs to midbrain dopamine neurons. *Neuron* 74, 858–873.
- Willner, P., 1991. Animal models as simulations of depression. *Trends Pharmacol. Sci.* 12, 131–136.
- Wise, R.A., Rompre, P.P., 1989. Brain dopamine and reward. *Annu. Rev. Psychol.* 40, 191–225.
- Witten, I.B., Steinberg, E.E., Lee, S.Y., Davidson, T.J., Zalocusky, K.A., Brodsky, M., Yizhar, O., Cho, S.L., Gong, S., Ramakrishnan, C., Stuber, G.D., Tye, K.M., Janak, P.H., Deisseroth, K., 2011. Recombinase-driver rat lines: tools, techniques, and optogenetic application to dopamine-mediated reinforcement. *Neuron* 72, 721–733.
- Wolf, M.E., Tseng, K.Y., 2012. Calcium-permeable AMPA receptors in the VTA and nucleus accumbens after cocaine exposure: when, how, and why? *Front. Mol. Neurosci.* 5, 72.
- Yadid, G., Friedman, A., 2008. Dynamics of the dopaminergic system as a key component to the understanding of depression. *Prog. Brain Res.* 172, 265–286.
- Yamaguchi, T., Sheen, W., Morales, M., 2007. Glutamatergic neurons are present in the rat ventral tegmental area. *Eur. J. Neurosci.* 25, 106–118.
- Yamaguchi, T., Wang, H.L., Li, X., Ng, T.H., Morales, M., 2011. Mesocorticolimbic glutamatergic pathway. *J. Neurosci.* 31, 8476–8490.
- Young, A.M., 2004. Increased extracellular dopamine in nucleus accumbens in response to unconditioned and conditioned aversive stimuli: studies using 1 min microdialysis in rats. *J. Neurosci. Methods* 138, 57–63.
- Zhang, T.A., Placzek, A.N., Dani, J.A., 2010. In vitro identification and electrophysiological characterization of dopamine neurons in the ventral tegmental area. *Neuropharmacology* 59, 431–436.
- Zweifel, L.S., Fadok, J.P., Argilli, E., Garelick, M.G., Jones, G.L., Dickerson, T.M., Allen, J.M., Mizumori, S.J., Bonci, A., Palmiter, R.D., 2011. Activation of dopamine neurons is critical for aversive conditioning and prevention of generalized anxiety. *Nat. Neurosci.* 14, 620–626.
- Zweifel, L.S., Parker, J.G., Lobb, C.J., Rainwater, A., Wall, V.Z., Fadok, J.P., Darvas, M., Kim, M.J., Mizumori, S.J., Paladini, C.A., Phillips, P.E., Palmiter, R.D., 2009. Disruption of NMDAR-dependent burst firing by dopamine neurons provides selective assessment of phasic dopamine-dependent behavior. *Proc. Natl. Acad. Sci. USA* 106, 7281–7288.