



Behavioral and biochemical sensitivity to low doses of ketamine: Influence of estrous cycle in C57BL/6 mice



Amanda M. Dossat, Katherine N. Wright, Caroline E. Strong, Mohamed Kabbaj*

Program in Neuroscience, Department of Biomedical Sciences, College of Medicine, Florida State University, 1115 W. Call Street, Tallahassee, FL 32306, United States

ARTICLE INFO

Article history:

Received 30 May 2017

Received in revised form

31 October 2017

Accepted 12 November 2017

Available online 21 November 2017

Keywords:

Estrogen

Estrous cycle

Forced swim test

Ketamine

Progesterone

Sex differences

ABSTRACT

Rationale: Low-dose ketamine is a rapid-acting antidepressant, to which female rodents are more sensitive as compared to males. However, the mechanism mediating this sex difference in ketamine sensitivity remains elusive.

Objectives: We sought to determine whether male and female mice differ in their behavioral sensitivity to low doses of ketamine, and uncover how ovarian hormones influence females' ketamine sensitivity. We also aimed to uncover some of the molecular mechanism(s) in mood-related brain regions that mediate sex differences in ketamine antidepressant effects.

Methods: Male and female mice (freely-cycling, diestrus 1 [D1], proestrus [Pro], or D1 treated with an estrogen receptor (ER) α , ER β , or progesterone receptor (PR) agonist) received ketamine (0, 1.5, or 3 mg/kg, intraperitoneally) and were tested in the forced swim test (FST) 30 min later. Ketamine's influence over synaptic plasticity markers in the prefrontal cortex (PFC) and hippocampus (HPC) of males, D1, and Pro females was quantified by Western blot 1 h post-treatment.

Results: Males, freely cycling females, D1 and Pro females exhibited antidepressant-like responses to 3 mg/kg ketamine. Pro females were the only group where ketamine exhibited an antidepressant effect at 1.5 mg/kg. D1 females treated with an agonist for ER α or ER β exhibited an antidepressant-like response to 1.5 mg/kg ketamine. Ketamine (3 mg/kg) increased synaptic plasticity-related proteins in the PFC and HPC of males, D1, and Pro females. Yet, Pro females exhibited an increase in p-Akt and p-CaMKII α in response to 1.5 and 3 mg/kg ketamine.

Conclusion: Our results indicate that females' enhanced sensitivity to ketamine during Pro is likely mediated through estradiol acting on ER α and ER β , leading to greater activation of synaptic plasticity-related kinases within the PFC and HPC.

© 2017 Elsevier Ltd. All rights reserved.

1. Introduction

The N-methyl-D-aspartate (NMDA) receptor antagonist, ketamine, has shown great promise as a rapid-acting antidepressant in humans and rodents (Abdallah et al., 2015; Kavalali and Monteggia, 2015). In clinical studies, a single infusion of low-dose of ketamine alleviates symptoms of depression within hours (Berman et al., 2000; Zarate et al., 2006). This rapid effect is superior to other antidepressants (e.g., selective serotonin-reuptake inhibitors), which can take weeks to months to provide therapeutic benefit

(Nutt, 2002; Rush et al., 2006). These exciting findings have burgeoned interest in uncovering the neural loci and molecular mechanisms of ketamine's rapid antidepressant effects. Low doses of ketamine promote the expression of downstream signaling of synaptic plasticity-related pathways within mood-related brain regions, such as the prefrontal cortex (PFC) and hippocampus (HPC) (Björkholm and Monteggia, 2016; Duman et al., 2016). While there have been great advances in our understanding of some molecular mechanisms of ketamine's rapid effects, the majority of these investigations have been conducted in males; creating a serious gap in our knowledge, since females experience a two-fold increased risk of depression as compared to males (Kessler, 2003; Van de Velde et al., 2010). In addition the disparity in risk of depression, there is evidence that pharmacodynamics and pharmacokinetics of antidepressant compounds may differ between the sexes and can

* Corresponding author. Florida State University, 3300-H, 1115 W. Call St., Tallahassee, FL 32304, United States.

E-mail address: mohamed.kabbaj@med.fsu.edu (M. Kabbaj).

be influenced by a variety of factors including levels of ovarian hormones (Bigos et al., 2009). Ovarian hormones, estrogen (E₂) and progesterone (P₄), influence many common biochemical targets to promote synaptic plasticity and increase spine density in mood-related brain regions (Frick, 2015; Woolley and McEwen, 1992; Woolley et al., 1997). Thus, it is possible for peripherally- or centrally-mediated factors to influence females' sensitivity to the antidepressant-like effects of ketamine.

Preclinical studies from our lab and others have shown female rodents exhibit behavioral sensitivity to doses of ketamine that are subthreshold for an antidepressant-like effect in males (Carrier and Kabbaj, 2013; Franceschelli et al., 2015; Sarkar and Kabbaj, 2016; Zanos et al., 2016). While most of these studies did not control for the stage of estrous, there is evidence that females' sensitivity may be mediated by the cyclic fluctuations of the ovarian hormones, E₂ and P₄ (Carrier and Kabbaj, 2013; Sarkar and Kabbaj, 2016). This is based upon evidence that surgical removal of the ovaries blocked females' sensitivity and pharmacological replacement of E₂/P₄ restored females' sensitivity to a male-subthreshold dose of ketamine (Carrier and Kabbaj, 2013). However, the molecular mechanism(s) mediating these sex differences remain elusive. In the first part of this work, we examined sensitivity to low doses of ketamine in male and female mice in which we did not control for the stage of estrous cycle (i.e., freely cycling). Then, to investigate the impact of estrous cycle on sensitivity to ketamine, we then evaluated sensitivity to ketamine in females experiencing a natural nadir (diestrus 1, D1) or a peak (proestrus, Pro) in ovarian hormone levels. To further determine which hormone receptor subtypes might be mediating ketamine sensitivity in females, D1 females were treated with an estrogen receptor (ER) α , ER β , or progesterone receptor (PR) agonist and tested 24 h later in the FST. Finally, we examined sex/estrous stage differences and ketamine-induced activation of neurotrophic signaling pathways within the prefrontal cortex (PFC) and hippocampus (HPC) of males, D1, and Pro females to explore which one(s) may mediate sex differences in ketamine sensitivity.

2. Materials and methods

2.1. Animals

One-hundred and eighty five adult male and female C57BL/6J mice (46 males and 139 females, mean weight at time of testing 23.7 and 18.8 g, respectively) from The Jackson Laboratory (Bar Harbor, Maine) aged 7 weeks at time of arrival were used in these experiments. Upon arrival, mice were pair-housed in same-sex pairs and maintained on a 12 h light:12 h dark cycle in a temperature-controlled vivarium in standard Plexiglass cages. Food and water were available *ad libitum*. Males and females were habituated to animal facilities and handling for one week prior to the start of behavioral testing. All experimental procedures were approved by the Florida State University Institutional Animal Care and Use Committee and conform to the NIH Guide for Care and Use of Laboratory Animals.

2.2. Drug treatments

Mice received a single intraperitoneal injection of sterile saline (Veh), 1.5, or 3 mg/kg of racemic ketamine hydrochloride (Ketasethesis[®], Henry Schein) and behavioral tests were conducted 30 min later. Females in D1 received a subcutaneous injection of either 4,4',4''-(4-Propyl-[1H]-pyrazole-1,3,5-triyl)trispheanol (PPT, 0.5 mg/kg, Tocris Bioscience; ER α agonist), Diarylpropionitrile (DPN, 0.5 mg/kg, Tocris Bioscience; ER β agonist) or progesterone (P₄, 0.5 mg/kg, Sigma-Aldrich), 24 h prior to behavioral testing. This

estrous stage was chosen because of the low levels of E₂ and P₄ during D1. The idea tested here is that during D1 ketamine at 1.5 mg/kg will have no antidepressant effects, however if we stimulate ERs or progesterone receptors during D1 (as to mimic what happens during Pro) ketamine may have an antidepressant-like effect. This experiment will also allow us to determine which receptors are critical in enhancing females' sensitivity to ketamine antidepressant effects.

2.3. Vaginal cytology

Female mice were gently handled and lavaged daily with 25 μ l sterile 0.9% saline as described elsewhere (McLean et al., 2012) within the first 3 h of the light cycle. For daily analysis of estrous stage, vaginal epithelial cells were deposited onto glass slides and immediately viewed with a light microscope. Degenerated, cornified epithelial cells and few stained leukocytes characterized the D1 estrous stage. The Pro stage was characterized by an abundance of rounded, nucleated epithelial cells. For crystal violet staining, slides were dried for 6 h at room temperature, stained with 0.1% crystal violet (Sigma-Aldrich) for 1 min, rinsed with DDH₂O for 1 min x2, sealed with 30 μ l Permount (Sigma-Aldrich), and coverslipped. If a female remained in single estrous stage for more than 4 days, she was excluded from the study.

2.4. Forced swim test (FST)

The FST is a sensitive measure of behavioral effects of antidepressant treatments in rodents (Porsolt et al., 1977). Mice were placed into a 4-L Pyrex glass beaker containing 3 L of water at 24 \pm 1 $^{\circ}$ C and forced to swim for 6 min. After the FST, mice were returned to their home cage and placed in a holding room until sacrifice. The FST was recorded by a video camera and videos were saved for later analysis. Beakers were cleaned with 70% EtOH and filled with fresh water between each FST. Treatments were delivered in a completely randomized design and counterbalanced such that cage mates received different treatments, and were examined in the FST in parallel. An observer who was blind to experimental assignments when the FST was conducted scored immobility time during the final 4 min of the FST. The final 4 min are commonly scored because mice are active during the onset of this test and this activity may mask the effect of the treatment (Can et al., 2012). Immobility time was defined as the duration of time in which the mouse maintained this stationary posture and only made movements necessary to keep its head above water.

2.5. Tissue collection and processing

Mice were sacrificed via rapid decapitation 24 min after the conclusion of the FST, which corresponds to 1 h post-vehicle or post-ketamine treatment. This time point was chosen due to reports that ketamine can rapidly, yet transiently influence protein expression and phosphorylation of molecules implicated in plasticity (Autry et al., 2011; Li et al., 2010). Trunk blood was collected into tubes containing chilled 0.5 M EDTA (Sigma-Aldrich) and stored on ice. Plasma was extracted via refrigerated centrifugation (2800 rpm for 20 min, at 4 $^{\circ}$ C), transferred to sterile pre-chilled microcentrifuge tubes, and stored at -80° C. Brains were rapidly removed, flash frozen in methyl butane (Sigma-Aldrich) on dry ice at -20° C and stored at -80° C. Brains underwent coronal sectioning at 100 μ M at -20° C in a cryostat, tissue punches containing the PFC (prelimbic and infralimbic cortices) and HPC (CA1, CA3, and dentate gyrus) were collected in sterile microcentrifuge tubes and stored at -80° C. Total protein was extracted from these punches using the Tri Reagent protocol (Molecular Research

Center), and concentration was determined via Bradford Assay (Bio-Rad). Whole tissue was used for western blot examination as compared to synaptoneurosome because of the limited volume of tissue that can be generated from specific regions of the mouse brain. Using whole tissue allowed for us to investigate many of the known targets of ketamine and of ovarian hormones within the same animals that underwent behavioral testing. Whole cell hippocampal extracts have also been used by other investigators to examine phosphorylation of molecular targets affected by ketamine (Autry et al., 2011; Choi et al., 2015; Gideons et al., 2014; Nosyreva et al., 2013).

2.6. Radioimmunoassay of ovarian hormones

Plasma levels of estradiol and P₄ were examined in a subset of vehicle-treated females from the FST experiment, using radioimmunoassay (RIA) kits (Cat. No. 07–238102 and 07–270102, MP Biomedicals). Samples were run in duplicate and any sample with a coefficient of variation >10% was excluded from analysis.

2.7. Western blot

Ten or 20 µg of protein per sample was resolved by a 12% acrylamide gel and subsequently transferred to a nitrocellulose membrane (Whatman, Protran BA83, GE Healthcare). Membranes were incubated with 5% milk in tris-buffered saline (TBS) for 1 h at room temperature then incubated overnight at 4 °C with a primary antibody (full list found in Table 1). After incubation with the primary antibody, membranes were rinsed once with 0.2% TBS Tween and four times with TBS on a rocking platform at room temperature, then incubated for 45 min at room temperature with a donkey anti-rabbit or goat anti-mouse secondary antibody diluted 1:10,000 (Li-COR Biosciences), rinsed once with 0.2% TBST and four times with TBS on a rocking platform, and imaged using an Odyssey infrared imaging system (Li-COR). Quantification was performed using ImageJ software (NIH). Data are shown as the ratio to vehicle-treated male values. The amount of protein for CaMKII α , GluR1, BDNF, MAPK, Akt, GSK-3 β , and mTOR was calculated as the ratio of the optical density (OD) of the target band over the OD of the GAPDH band or from total protein. Levels of p-CaMKII α , p-GluR1, p-MAPK, p-Akt, p-GSK-3 β , and p-mTOR were calculated as a ratio of the OD of the phosphorylated band over the OD of the total band. Two lanes of each gel contained standard samples, which were

each comprised of homogenized hippocampal protein from 8 male mice that received no drug treatment and were naïve to the FST. These standard samples were used to normalize protein quantities across membranes.

2.8. Statistics

Data are reported as mean + SEM. Two-way analysis of variance (ANOVA) was used to examine the effect of ketamine and sex/estrous stage. Unpaired t-tests were used to examine ovarian hormone plasma levels and to determine the effect of ketamine within each hormone receptor agonist condition. Western blot analysis of the effect of ketamine within each sex/estrous stage was examined using two-way ANOVA. When significant main effects or interaction of ketamine or of sex/estrous stage were identified in the two-way ANOVA, *post hoc* comparisons were made with Fisher's LSD. Any value that was more than two standard deviations above the group mean was considered an outlier and was removed from the analysis. *P*-values of <0.05 were considered significant. Sample size was determined based upon previous studies conducted by our group and statistical computation with power analysis that indicated a need for a sample size of 8–10 animals per group in order to obtain power values of 0.8 and higher for *p* values of 0.05.

3. Results

3.1. Effect of sex (males versus freely cycling females) on behavioral sensitivity to the antidepressant-like effect of ketamine in the FST

Thirty min following treatment with ketamine, time spent immobile in the FST was examined in male and female mice (*n* = 7–9/group) as shown in Fig. 1. There was a significant effect of ketamine treatment ($F_{(2,40)} = 5.54$, *p* = 0.0007) and a significant effect of sex ($F_{(1,40)} = 4.44$, *p* = 0.041) to influence time spent immobile. There were however no interactions ($F_{(2,40)} = 0.06$, *p* = 0.94). Fisher's *post hoc* analyses identified that 3 mg/kg ketamine significantly reduced males' and females' immobility ($t(40) = 2.22$, *p* = 0.03; $t(40) = 2.49$, *p* = 0.017, respectively).

3.2. Effect of estrous stage over behavioral sensitivity to the antidepressant-like effect of ketamine in the FST

To determine the effect of endogenous ovarian hormones to

Table 1
List of antibodies used in Western blot analysis of the prefrontal cortex and hippocampus.

Primary Antibody	Host/Type	Manufacturer	Catalog Number	Dilution
GAPDH (D16H11) XP [®]	Rabbit, mAb	Cell Signaling	5174	1:2000
Phospho-CaMKII (Thr286) (D21E4)	Rabbit, mAb	Cell Signaling	12716	1:2000
CaMKII- α (6G9)	Mouse, mAb	Cell Signaling	50049	1:2000
AMPA Receptor (GluR 1) (D4N9V)	Rabbit, mAb	Cell Signaling	13185	1:1000
Phospho-AMPA Receptor (Ser831)	Rabbit, mAb	Abcam	ab109464	1:1000
BDNF (N-20)	Rabbit, pAb	Santa Cruz	sc-546	1:500
Phospho-p44/42 MAPK (Thr202/Thr204)	Rabbit, pAb	Cell Signaling	9101	1:2000
p44/42 MAPK	Mouse, pAb	Cell Signaling	9102	1:2000
Phospho-Akt (Ser473) (D9W9U)	Mouse, mAb	Cell Signaling	12694	1:1000
Akt	Rabbit, pAb	Cell Signaling	9272	1:1000
Phospho-GSK-3 β (Ser9) (D85E12) XP [®]	Rabbit, mAb	Cell Signaling	5558	1:2000
GSK-3 β (3D10)	Mouse, mAb	Cell Signaling	9832	1:1000
Phospho-mTOR (Ser2448) (D9C2) XP [®]	Rabbit, mAb	Cell Signaling	5536	1:1000
mTOR (L27D4)	Mouse, mAb	Cell Signaling	45175	1:1000

mAb, monoclonal antibody; pAb, polyclonal antibody.




Fig. 1. Behavioral sensitivity to the antidepressant-like effect of ketamine is influenced by sex. Ketamine at 3 mg/kg reduced immobility time in males and freely-cycling females. Overall, females exhibited lower immobility time as compared to males. * $p < 0.05$ vs. vehicle, # $p < 0.05$ males vs. females. Data are means + SEM. (males: $n = 7-8$ /group, females: $n = 8$ /group).

influence behavioral sensitivity to ketamine, time spent immobile in the FST was examined 30 min following ketamine treatment in males, D1, and Pro females ($n = 7-10$ /group) as shown in Fig. 2. There was a significant effect of ketamine treatment ($F_{(2,64)} = 10.87$, $p < 0.0001$) and of sex/estrous stage to influence duration of immobility ($F_{(2,64)} = 7.36$, $p = 0.001$). There were however no interactions ($F_{(4,64)} = 1.99$, $p = 0.10$). Fisher's *post hoc* analyses revealed that males and D1 females have significantly reduced immobility time following 3 mg/kg ketamine (males: $t(64) = 2.38$, $p = 0.02$; D1: $t(64) = 3.295$, $p = 0.0016$). In contrast, Pro females display significant reductions in immobility following both 1.5 mg/kg and 3 mg/kg ketamine ($t(64) = 2.897$, $p = 0.005$; $t(64) = 2.346$, $p = 0.02$, respectively). Fisher's *post hoc* revealed that Pro females exhibited significantly less immobility as compared to D1 females in the vehicle condition ($t(64) = 2.016$, $p = 0.04$). Similarly, Pro females exhibited significantly less immobility as compared to males and D1 females following treatment with 1.5 mg/kg ketamine ($t(64) = 3.58$, $p = 0.0007$; $t(64) = 3.56$, $p = 0.0007$; respectively).




Fig. 3. Behavioral sensitivity to the antidepressant-like effect of ketamine is influenced by agonists for ER α and ER β . Ketamine at 1.5 mg/kg reduced immobility time for D1 females treated with PPT (ER α agonist) or DPN (ER β agonist), but not progesterone (P₄). * $p < 0.05$ vs. within-group vehicle; Different letters indicate statistically significant differences within the treatment group. Data are means + SEM. ($n = 7-10$ /group).

3.3. Confirmation of estrous cycle via ovarian hormone assay

As shown in Fig. 2C–D, D1 females exhibited significantly lower levels of plasma estradiol and P₄ vs. Pro females ($t(10) = 2.03$, $p = 0.03$; $t(10) = 2.87$, $p = 0.008$; respectively).

3.4. Effect of agonists for ER α , ER β , and PR over D1 females' behavioral sensitivity to the antidepressant-like effect of ketamine in the FST

To determine which specific hormone receptor subtypes may influence females' ketamine sensitivity, we examined D1 females' behavior in the FST 24 h following treatment with an agonist for ER α (PPT), ER β (DPN), or PR (P₄), and 30 min following 0 or 1.5 mg/kg ketamine ($n = 7-10$ /group). As shown in Fig. 3, two-way ANOVA revealed a significant main effect of vehicle/ketamine ($F_{(1,58)} = 15.85$, $p = 0.0002$) and a significant main effect of




Fig. 2. Behavioral sensitivity to the antidepressant-like effect of ketamine is influenced by sex and by estrous stage. (A) Ketamine at 3 mg/kg reduced immobility time in males and diestrus 1 (D1) females, 1.5 and 3 mg/kg ketamine reduced immobility time in proestrus (Pro) females (* $p < 0.05$ vs. within-group vehicle). (B) Representative vaginal lavage samples from D1 and Pro females. (C) Plasma levels of estradiol (pg/mL) and levels of progesterone (ng/mL) were significantly lower in females in D1 vs. Pro females (* $p < 0.05$, ** $p < 0.01$). Data are means + SEM. Different letters indicate statistically significant differences within the treatment group. (males: $n = 8$ /group, D1: $n = 7-9$ /group, Pro: $n = 7-10$ /group; plasma: $n = 5-7$).

hormone treatment ($F_{(3,58)} = 3.6, p = 0.019$) to influence time spent immobile. There were however no interactions ($F_{(3,58)} = 1.49, p = 0.22$). *Post hoc* analysis revealed a significant effect of ketamine to reduce immobility time only when given in combination with PPT or DPN ($t(58) = 3.317, p = 0.0016$; $t(58) = 2.85, p = 0.006$, respectively). The lack of effect of the ER α and ER β agonists within the vehicle condition indicates that these doses of hormone receptor agonist are not sufficient to influence FST behavior on their own.

3.5. Effect of sex/estrous stage and ketamine on PFC p-CaMKII α levels

One hour following ketamine treatment, we examined the level of activation of synaptic plasticity markers p-CaMKII α within the PFC ($n = 4–6$ samples/group). There was no significant effect of ketamine or of sex/estrous stage to influence phosphorylation of CaMKII α ($F_{(2,38)} = 0.27, p = 0.76$; $F_{(2,38)} = 1.04, p = 0.36$, respectively; Fig. 4A). There were no interactions ($F_{(4,38)} = 1.45, p = 0.23$).

3.6. Effect of sex/estrous stage and ketamine on PFC p-GluR1 levels

Two-way ANOVA indicated a main ketamine treatment effect ($F_{(2,45)} = 9.667, p = 0.0003$; $n = 4–7$ samples/group; Fig. 4B) and a main sex/estrous stage effect on the levels of p-GluR1 ($F_{(2,45)} = 68.49, p < 0.0001$). There were no interactions ($F_{(4,45)} = 1.33, p = 0.27$). *Post hoc* examination revealed a significant effect of 3 mg/kg ketamine to increase p-GluR1 expression in males and Pro females ($t(45) = 3.38, p = 0.0015$; $t(45) = 3.31, p = 0.0018$, respectively). Within the vehicle and 1.5 mg/kg ketamine treatment conditions, Pro females exhibited higher levels of p-GluR1 as compared to males ($t(45) = 4.89, p < 0.0001$, $t(45) = 4.19, p = 0.0001$) and D1 females ($t(45) = 6.00, p < 0.0001$, $t(45) = 5.41, p < 0.0001$). Following treatment with 3 mg/kg ketamine, D1 females exhibited significantly lower p-GluR1 levels as compared to males and Pro females ($t(45) = 3.80, p = 0.0004$; $t(45) = 8.78, p < 0.0001$, respectively); Pro females exhibited higher levels of p-GluR1 as compared to males ($t(45) = 5.13, p < 0.0001$).

3.7. Effect of sex/estrous stage and ketamine on PFC BDNF levels

Two-way ANOVA indicated a main ketamine treatment effect ($F_{(2,44)} = 7.12, p = 0.002$; $n = 4–7$ /group; Fig. 4C) and a main sex/estrous stage effect on the levels of BDNF within the PFC ($F_{(2,44)} = 23.54, p < 0.0001$). There were no interactions ($F_{(4,44)} = 2.23, p = 0.08$). *Post hoc* examination revealed a significant increase in BDNF levels following 3 mg/kg ketamine in males ($t(44) = 2.63, p = 0.01$) and Pro females ($t(44) = 3.81, p = 0.0004$). Pro females exhibited higher BDNF levels as compared to D1 females in all treatment conditions (vehicle: $t(44) = 2.62, p = 0.01$; 1.5 mg/kg: $t(44) = 3.05, p = 0.004$; 3 mg/kg: $t(44) = 6.39, p < 0.0001$). Following treatment with 1.5 mg/kg and 3 mg/kg ketamine, Pro females exhibited higher BDNF levels as compared to males ($t(44) = 2.08, p = 0.04$; $t(44) = 2.57, p = 0.01$, respectively). While males exhibited higher levels of BDNF as compared to D1 females in the 3 mg/kg ketamine treatment condition ($t(44) = 3.92, p = 0.0003$).

3.8. Effect of sex/estrous stage and ketamine on PFC p-MAPK levels

There was no significant effect of ketamine treatment to influence p-MAPK levels within the PFC ($F_{(2,47)} = 0.51, p = 0.6$; $n = 5–7$ /group; Fig. 4D), yet there was a significant main effect of sex/estrous stage ($F_{(2,47)} = 14.68, p < 0.0001$). There were no interactions ($F_{(4,47)} = 0.99, p = 0.42$). *Post hoc* analysis revealed that in

the vehicle condition, males and Pro females exhibited higher levels of p-MAPK as compared to D1 females ($t(47) = 4.47, p < 0.0001$; $t(47) = 2.51, p = 0.01$, respectively). Additionally, males exhibited higher p-MAPK levels following 3 mg/kg ketamine as compared to D1 female counterparts ($t(47) = 3.02, p = 0.004$).

3.9. Effect of sex/estrous stage and ketamine on PFC p-Akt levels

Two-way ANOVA indicated a main ketamine treatment effect ($F_{(2,53)} = 12.77, p < 0.0001$; $n = 6–8$ /group; Fig. 4E) and a main sex/estrous stage effects on the levels of p-Akt within the PFC ($F_{(2,53)} = 99.16, p < 0.0001$). There were no interactions ($F_{(4,53)} = 2.19, p = 0.08$). *Post hoc* analysis revealed a significant increase in p-Akt in males in response to 3 mg/kg ketamine ($t(53) = 2.96, p = 0.004$), and in Pro females in response to 1.5 and 3 mg/kg ketamine ($t(53) = 2.8, p = 0.007$; $t(53) = 4.69, p < 0.0001$, respectively). In all treatment conditions, D1 females exhibited lower p-Akt as compared to males and Pro females (vehicle: $t(53) = 6.83, p < 0.0001$; $t(53) = 2.89, p = 0.005$; 1.5 mg/kg: $t(53) = 7.43, p < 0.0001$; $t(53) = 4.94, p < 0.0001$; 3 mg/kg: $t(53) = 9.98, p < 0.0001$; $t(53) = 7.26, p < 0.0001$, respectively). Additionally, males exhibited higher levels of p-Akt as compared to Pro females in all treatment conditions (vehicle: $t(53) = 3.69, p = 0.0005$; 1.5 mg/kg: $t(53) = 2.64, p = 0.01$; 3 mg/kg: $t(53) = 2.16, p = 0.03$).

3.10. Effect of sex/estrous stage and ketamine on PFC p-GSK-3 β levels

There was a significant main effect of ketamine treatment to influence levels of p-GSK-3 β in the PFC ($F_{(2,46)} = 7.75, p = 0.001$; $n = 5–7$ /group; Fig. 4F), yet no effect of sex/estrous stage ($F_{(2,46)} = 1.47, p = 0.24$). There were no interactions ($F_{(4,46)} = 1.24, p = 0.31$). *Post hoc* examination revealed a significant effect of 1.5 mg/kg ketamine to increase p-GSK-3 β in males ($t(46) = 2.65, p = 0.01$), and a significant effect of 3 mg/kg ketamine to increase levels in males and Pro females ($t(46) = 2.35, p = 0.02$; $t(46) = 2.72, p = 0.009$, respectively).

3.11. Effect of sex/estrous stage and ketamine on PFC p-mTOR levels

There was a significant main effect of ketamine treatment to increase levels of PFC p-mTOR ($F_{(2,52)} = 4.01, p = 0.02$; $n = 5–8$ /group; Fig. 4G), yet no effect of sex/estrous stage ($F_{(2,52)} = 1.44, p = 0.25$), and there were no interactions ($F_{(4,52)} = 1.05, p = 0.39$). *Post hoc* examination revealed a significant increase in p-mTOR in response to 3 mg/kg ketamine in males and D1 females ($t(52) = 2.36, p = 0.02$; $t(52) = 2.34, p = 0.02$, respectively). The effects of ketamine within the PFC are summarized in Table 2 and Fig. 6A.

3.12. Effect of sex/estrous stage and ketamine on HPC p-CaMKII α levels

Following the same timeline as the one described for the PFC, we examined the phosphorylation of synaptic plasticity markers within the HPC. As shown in Fig. 5A, there were significant main effects of ketamine and of sex/estrous stage to influence levels of p-CaMKII α ($F_{(2,44)} = 9.54, p = 0.0004$; $F_{(2,44)} = 6.94, p = 0.002$, respectively; $n = 5–6$ /group) as well as a significant interaction between the two ($F_{(2,44)} = 2.59, p = 0.04$). *Post hoc* analysis revealed that 3 mg/kg ketamine increased p-CaMKII α levels in males ($t(44) = 3.22, p = 0.002$), and both 1.5 and 3 mg/kg ketamine increased levels in Pro females ($t(44) = 3.04, p = 0.0039$; $t(44) = 4.00, p = 0.0002$, respectively). *Post hoc* analysis of sex/

Prefrontal Cortex




Fig. 4. Influence of sex/estrous stage on protein expression of synaptic plasticity markers within the prefrontal cortex (PFC) following treatment with vehicle or ketamine. (A) Protein expression of p-CaMKII α /CaMKII α was not affected by ketamine nor by sex/estrous stage. (B) Males and Pro females exhibited an increased expression of p-GluR1/GluR1 and (C) increased expression of BDNF following 3 mg/kg ketamine. (D) Protein expression of p-MAPK/MAPK was not affected by ketamine, yet there was a main effect of sex/estrous stage. (E) Ketamine at 1.5 mg/kg increased expression of p-Akt/Akt in Pro females, and 3 mg/kg increased expression in males and Pro females. (F) Ketamine at 1.5 mg/kg increased expression of p-GSK-3 β /GSK-3 β in males, and in males and Pro females at 3 mg/kg. (G) Ketamine at 3 mg/kg increased p-mTOR/mTOR expression in males and D1 females. * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$, **** $p < 0.0001$ vs. within-group vehicle. Different letters indicate statistically significant differences between sex/estrous stages within the vehicle treatment group. Data are means + SEM. (n = 4–8/group).

estrous stage effects revealed significantly higher levels of p-CaMKII α in Pro females as compared to males and D1 females in response to treatment with 1.5 mg/kg ketamine ($t(44) = 6.17$, $p = 0.036$; $t(44) = 2.64$, $p = 0.01$, respectively). In contrast, D1 females exhibited lower levels of p-CaMKII α as compared to males and Pro females following 3 mg/kg ketamine ($t(44) = 3.14$, $p = 0.003$; $t(44) = 3.72$, $p = 0.0006$, respectively).

3.13. Effect of sex/estrous stage and ketamine treatment on HPC p-GluR1

As shown in Fig. 5B, there were significant main effects of ketamine and sex/estrous stage to influence levels of p-GluR1 within the HPC ($F_{(2,43)} = 9.87$, $p = 0.0003$; $F_{(2,43)} = 8.88$, $p = 0.0006$, respectively; n = 4–7/group). There were no interactions

Table 2
Summary of ketamine's effects on activation of synaptic plasticity markers in the prefrontal cortex.

Prefrontal Cortex			
	Male	Diestrus 1	Proestrus
p-CaMKII α	↔	↔	↔
p-GluR1	↑ 3 mg/kg	↔	↑ 3 mg/kg
BDNF	↑ 3 mg/kg	↔	↑ 3 mg/kg
p-MAPK	↔	↔	↔
p-Akt	↑ 3 mg/kg	↔	↑ 1.5 & 3 mg/kg
p-GSK-3 β	↑ 1.5 & 3 mg/kg	↔	↑ 3 mg/kg
p-mTOR	↑ 3 mg/kg	↑ 3 mg/kg	↔

↔ no change from vehicle-treated samples, ↑ in levels as compared to vehicle-treated samples.

($F_{(4,43)} = 1.82, p = 0.14$). *Post hoc* examination of the effect of ketamine revealed a significant increase in males and Pro females following 3 mg/kg ($t(43) = 2.82, p = 0.007; t(43) = 4.15, p = 0.0002$, respectively). *Post hoc* examination of the effect of sex/estrous stage revealed significantly higher levels of p-GluR1 in D1 females as compared to males in the vehicle condition and following 1.5 mg/kg ketamine ($t(43) = 3.18, p = 0.002; t(43) = 2.04, p = 0.04$, respectively). In contrast, following 3 mg/kg ketamine Pro females exhibited higher p-GluR1 levels as compared to males and D1 females ($t(43) = 3.2, p = 0.002; t(43) = 2.33, p = 0.02$, respectively).

3.14. Effect of sex/estrous stage and ketamine on HPC BDNF levels

There were significant main effects of ketamine and of sex/estrous stage to influence BDNF levels within the HPC ($F_{(2,44)} = 13.31, p = 0.001; F_{(2,44)} = 7.48, p < 0.0001$, respectively; $n = 5-7$ /group; Fig. 5C). There were no interactions ($F_{(4,44)} = 1.79, p = 0.14$). *Post hoc* analysis of the effect of ketamine revealed an increase in males and D1 females following 3 mg/kg ketamine ($t(44) = 2.25, p = 0.02; t(44) = 3.94, p = 0.0003$, respectively). *Post hoc* analysis of the effect of sex/estrous stage revealed significantly higher levels of BDNF in males as compared to Pro females in the vehicle condition ($t(44) = 2.12, p = 0.03$), and significantly higher levels in males and D1 vs. Pro females following treatment with 3 mg/kg ketamine ($t(44) = 3.45, p = 0.001; t(44) = 3.59, p = 0.0008$, respectively).

3.15. Effect of sex/estrous stage and ketamine on HPC p-MAPK levels

For levels of HPC p-MAPK, there were significant main effects of ketamine and of sex/estrous stage ($F_{(2,43)} = 22.51, p < 0.0001; F_{(2,43)} = 55.65, p < 0.0001$, respectively; $n = 5-6$ /group; Fig. 5D). There were no interactions ($F_{(4,43)} = 1.72, p = 0.16$). *Post hoc* analysis of the main effect of ketamine revealed that 1.5 and 3 mg/kg ketamine increased p-MAPK levels in D1 females ($t(43) = 4.19, p = 0.0001; t(43) = 5.4, p < 0.0001$, respectively) and 3 mg/kg increased levels in males and Pro females ($t(43) = 3.2, p = 0.003, t(43) = 3.04, p = 0.004$, respectively). *Post hoc* examination of the main effect of sex/estrous stage revealed significantly lower levels of p-MAPK in D1 females within all treatment conditions as compared to males (vehicle: $t(43) = 6.93, p < 0.0001; 1.5$ mg/kg: $t(43) = 4.06, p = 0.0002; 3$ mg/kg: $t(43) = 5.24, p < 0.0001$) and vs. Pro females (vehicle: $t(43) = 6.93, p < 0.0001; 1.5$ mg/kg: $t(43) = 3.69, p = 0.0006; 3$ mg/kg: $t(43) = 4.57, p < 0.0001$).

3.16. Effect of sex/estrous stage and ketamine on HPC p-Akt levels

There were significant main effects of ketamine treatment and of sex/estrous stage to influence protein expression of p-Akt in the

HPC ($F_{(2,41)} = 7.46, p = 0.0017; F_{(2,41)} = 3.49, p = 0.03$, respectively; $n = 4-7$ /group; Fig. 5E). There were no interactions ($F_{(4,41)} = 1.92, p = 0.12$). *Post hoc* analysis of the main effect of ketamine revealed a significant increase in males following 3 mg/kg ketamine ($t(44) = 2.23, p = 0.03$), and a significant increase in Pro females following 1.5 and 3 mg/kg ($t(44) = 3.08, p = 0.004; t(44) = 3.9, p = 0.0004$, respectively). *Post hoc* analysis of the main effect of sex/estrous stage revealed significantly higher levels of p-Akt in Pro females as compared to D1 females in the 1.5 mg/kg condition ($t(44) = 2.14, p = 0.04$). Additionally, following treatment with 3 mg/kg ketamine, males and Pro females exhibited higher levels of p-Akt as compared to D1 counterparts ($t(44) = 2.04, p = 0.04; t(44) = 2.47, p = 0.01$, respectively).

3.17. Effect of sex/estrous stage and ketamine on HPC p-GSK-3 β

There were significant main effects of ketamine and of sex/estrous stage to influence levels of p-GSK-3 β in the HPC ($F_{(2,43)} = 12.74, p < 0.0001; F_{(2,43)} = 17.24, p < 0.0001$, respectively; $n = 5-7$ /group; Fig. 5F). There were no interactions ($F_{(4,43)} = 1.14, p = 0.34$). *Post hoc* analysis of the main effect of ketamine revealed a significant increase in males following 1.5 mg/kg and 3 mg/kg ketamine ($t(43) = 2.77, p = 0.008; t(43) = 2.54, p = 0.01$, respectively), as well as a significant increase in D1 and Pro females following 3 mg/kg ($t(43) = 3.07, p = 0.004; t(43) = 3.12, p = 0.003$, respectively). *Post hoc* analysis of the main effect of sex/estrous stage revealed significantly lower levels of p-GSK-3 β in D1 females as compared to males and Pro females in the vehicle condition ($t(43) = 3.05, p = 0.004; t(43) = 2.67, p = 0.01$, respectively) and following 3 mg/kg ketamine ($t(43) = 2.51, p = 0.01; t(43) = 2.98, p = 0.005$, respectively). In response to 1.5 mg/kg ketamine, males exhibited higher levels of p-GSK-3 β as compared to D1 and Pro females ($t(43) = 4.28, p = 0.0001; t(43) = 2.43, p = 0.02$, respectively).

3.18. Effect of sex/estrous stage and ketamine on HPC p-mTOR

There were significant main effects of ketamine and of sex/estrous stage to influence expression of p-mTOR in the HPC ($F_{(2,41)} = 15.97, p < 0.0001; F_{(2,41)} = 13.4, p < 0.0001$, respectively; $n = 4-6$ /group) as well as a significant interaction between the two ($F_{(2,41)} = 4.73, p = 0.003$; Fig. 5G). *Post hoc* analysis of the ketamine effect revealed a significant increase in males and in Pro females following 3 mg/kg ($t(41) = 5.66, p < 0.0001; t(41) = 2.04, p = 0.04$, respectively). *Post hoc* analysis of the main effect of sex/estrous stage to influence p-mTOR revealed significantly higher levels in males as compared to Pro females treated with vehicle ($t(41) = 2.41, p = 0.02$). Additionally, there were group differences in the 3 mg/kg ketamine condition, such that males exhibited higher levels as compared to all female counterparts (D1: $t(41) = 4.365, p < 0.0001$; Pro: $t(41) = 6.34, p < 0.0001$). The effects of ketamine within the HPC are summarized in Table 3 and Fig. 6B.

4. Discussion

In this study, we demonstrate that when examined only by sex, male and female mice exhibit similar behavioral sensitivity to ketamine in the FST. In contrast, when stage of estrous is controlled for, D1 females and males exhibit similar ketamine sensitivity. Yet Pro females exhibit an antidepressant-like response to a dose of ketamine that is subthreshold for effect for the other groups (i.e., 1.5 mg/kg). Furthermore, we show that pharmacological activation of ER α or ER β is sufficient to render D1 females sensitive to the Pro-effective dose of ketamine, which suggests that E $_2$ action at these receptor subtypes may mediate

Hippocampus




Fig. 5. Influence of sex/estrous stage on protein expression of synaptic plasticity markers within the hippocampus (HPC) following treatment with vehicle or ketamine. (A) Protein expression of p-CaMKII α /CaMKII α was increased by 1.5 mg/kg ketamine in Pro females, and in males and Pro females by 3 mg/kg. (B) Ketamine at 3 mg/kg increased expression of p-GluR1/GluR1 in males and Pro females, (C) and increased expression of BDNF in males and D1 females. (D) Protein expression of p-MAPK/MAPK was significantly lower in D1 females in all treatment conditions. Ketamine at 1.5 mg/kg increased expression of p-MAPK in D1 females, and in all groups at 3 mg/kg. (E) Ketamine at 1.5 mg/kg increased expression of p-Akt/Akt in Pro females, and 3 mg/kg increased expression in males and Pro females. (F) Ketamine at 1.5 mg/kg increased expression of p-GSK-3 β /GSK-3 β in males, and in all groups at 3 mg/kg. (G) Ketamine at 3 mg/kg increased p-mTOR/mTOR expression in males and D1 females. * $p < 0.05$, ** $p < 0.005$, *** $p < 0.0005$, **** $p < 0.0001$ vs. within-group vehicle. Different letters indicate statistically significant differences between sex/estrous stages within the vehicle treatment group. Data are means + SEM. (n = 4–7/group).

females' behavioral sensitivity to ketamine. Investigation into the molecular mechanism(s) that may mediate different ketamine sensitivity revealed that at behaviorally effective doses of ketamine (1.5 and 3 mg/kg), Pro females exhibited activation of CaMKII α within the HPC, and of Akt in the PFC and HPC in response to a male- and D1-subthreshold dose of ketamine.

4.1. Behavioral effects of ketamine and ovarian hormones in the FST

In the present study, we observed similar behavioral sensitivity to ketamine in males and freely cycling female mice in the FST when examined 30 min post-treatment. This was unexpected, due to previous reports in rats (Carrier and Kabbaj, 2013; Sarkar and

Table 3
Summary of ketamine's effects on activation of synaptic plasticity markers in the hippocampus.

Hippocampus			
	Male	Diestrus 1	Proestrus
p-CaMKII α	↑ 3 mg/kg	↔	↑ 1.5 & 3 mg/kg
p-GluR1	↑ 3 mg/kg	↔	↑ 3 mg/kg
BDNF	↑ 3 mg/kg	↑ 3 mg/kg	↔
p-MAPK	↑ 3 mg/kg	↑ 1.5 & 3 mg/kg	↑ 3 mg/kg
p-Akt	↑ 3 mg/kg	↔	↑ 1.5 & 3 mg/kg
p-GSK3- β	↑ 1.5 & 3 mg/kg	↑ 3 mg/kg	↑ 3 mg/kg
p-mTOR	↑ 3 mg/kg	↔	↑ 3 mg/kg

↔ no change from vehicle, ↑ in levels as compared to vehicle-treated samples.

Kabbaj, 2016) and in mice (Franceschelli et al., 2015; Zanos et al., 2016) which demonstrated that females respond to a male-subthreshold dose of ketamine. While unexpected, this difference is not unwarranted since when not controlling for the stage of estrous, the females within the test group may be comprised of more Pro vs. D1 females depending upon the group make up on the test day (Goldman et al., 2007). Furthermore, without assessing estrous cyclicity in females, it is impossible to determine if the females are experiencing natural fluctuations in ovarian hormones, or experiencing a period of anestrus due to housing in an all-female environment (Cooper et al., 1993). To control for this potential confound, the goal of the present study was to examine females' ketamine sensitivity during stages of the estrous cycle characterized by low and high ovarian hormone levels (D1 and Pro, respectively), because it is well known that ovarian hormones (E₂ and P₄) influence depression-like behavior (Frye and Walf, 2002; Mahmoud et al., 2016), can impact behavioral sensitivity to antidepressants (Carrier and Kabbaj, 2013; Estrada-Camarena et al., 2008; Kokras et al., 2015), as well as influence neurochemical

responses to antidepressants (Allen et al., 2012; Kendall et al., 1981; Wilson et al., 1989). In line with the known antidepressant-like effects of E₂ and P₄ (Frye, 2011; Walf and Frye, 2009, 2010), we found that vehicle-treated D1 females exhibit higher indices of behavioral despair in the FST as compared to Pro counterparts. Males and D1 females exhibited similar sensitivity to ketamine in the FST, where both groups showed an antidepressant-like response 30 min following treatment with 3 mg/kg ketamine. These results are in line with a report by Autry et al. (2011) that demonstrated male mice exhibit sensitivity to 3 mg/kg ketamine at 30 min, 3 h, 24 h, and 1 week post-treatment (Autry et al., 2011), but see (Franceschelli et al., 2015; Zanos et al., 2016). In the present study, Pro females exhibited an antidepressant-like response to both 1.5 and 3 mg/kg ketamine, indicating enhanced behavioral sensitivity. To our knowledge, this is the first report of the effect of estrous stage to influence sensitivity to the antidepressant-like effects of ketamine. These findings are consistent with a recent report, which demonstrated females' responsivity to ketamine's reinforcing effects is dependent upon estrous stage (Wright et al., 2017). Our findings are also in agreement with findings that ovariectomized (OVX) females treated with E₂/P₄ exhibit behaviorally sensitivity to 2.5 mg/kg ketamine, which is subthreshold for effect in OVX females without E₂/P₄ and in males (Carrier and Kabbaj, 2013; Saland et al., 2016; Wright et al., 2017). Adding support to our findings in Pro females, when D1 females were treated with an agonist for ER α or ER β , but not for PR, they exhibited an antidepressant-like response to 1.5 mg/kg ketamine. Taken together, these results suggest that endogenous ovarian hormones, specifically E₂, can act at either ER α or ER β subtypes to enhance females' sensitivity to ketamine.

4.2. Neurotrophic effects of ketamine and ovarian hormones within the PFC and HPC

The present study aimed to elucidate potential molecular mechanisms that mediate sex/estrous stage differences in behavioral sensitivity to antidepressant doses of ketamine. The rapid antidepressant-like effects of ketamine are paralleled, and supported by, the expression and activation of neuroplasticity-related signaling pathways within the PFC and HPC (Björkholm and Monteggia, 2016; Duman et al., 2016). This includes increased expression and release of BDNF and activation of mTOR within mood-related brain regions. Ketamine, E₂, and P₄ share many biochemical targets, through which they support synaptic plasticity and increase spine density in mood-related brain regions (Frick, 2015; Woolley and McEwen, 1992; Woolley et al., 1997). Based upon this evidence we hypothesized that if females are experiencing a natural peak in levels of E₂/P₄, then enhanced activation of these shared plasticity-related signaling pathways may be one mechanism through which Pro females are more sensitive to ketamine as compared to males and D1 females.

Ketamine rapidly increases glutamate release and neuronal depolarization (Höflich et al., 2017; Li et al., 2017; Moghaddam et al., 1997). Preclinical evidence suggests that ketamine's induction of neuronal plasticity-related functions likely underlies its rapid antidepressant-like effects (Lepack et al., 2015; Li et al., 2010, 2011). Postsynaptic depolarization is accompanied by an influx of calcium, which promotes long-term potentiation and synaptic plasticity through its phosphorylation of CaMKII α (Herring and Nicoll, 2016; Lisman et al., 2012). Previous reports have shown that ketamine can rapidly increase levels of p-CaMKII α in the HPC of male rats (Choi et al., 2015), which we confirmed in the present investigation in male mice. As has been shown for ketamine, there is evidence that E₂ also promotes the phosphorylation of CaMKII α (Hasegawa et al., 2015). In line with this role of E₂, we found that




Fig. 6. Schematic representation of the hypothesized shared influence of ketamine and ovarian hormones within mood related brain regions.

(A) A simplified schematic of the intracellular signaling pathways that are influenced by ketamine and ovarian hormones within the PFC and (B) the HPC. Yellow circles indicate pathways activated by behaviorally-effective doses of ketamine in Pro females. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

Pro females exhibited HPC CaMKII α activation in a manner that was consistent with their behavioral ketamine sensitivity (i.e., increased by 1.5 and 3 mg/kg). Of the many functions carried out by p-CaMKII α to support synaptic plasticity, one important task is the phosphorylation of the GluR1 subunit of AMPA receptors (Barria et al., 1997). This supports GluR1 insertion into the synapse and can enhance channel conductivity (Derkach et al., 2007). Previous reports posited that ketamine's up-regulation of AMPA receptor expression may underlie enhanced glutamatergic sensitivity and synaptic plasticity (Kavalali and Monteggia, 2012; Li et al., 2010). Consistent with this effect of ketamine, we observed a significant increase in levels of p-GluR1 in the PFC and HPC of males and Pro females in 1 h following treatment with 3 mg/kg ketamine.

Antidepressant doses of ketamine rapidly increase BDNF levels in the PFC and HPC (Autry et al., 2011; Garcia et al., 2008; Lepack et al., 2015; Zhou et al., 2014b). In line with these previous reports in males, we observed this effect of ketamine in the male PFC and HPC. We extend previous reports and demonstrate an increase in BDNF within the PFC of Pro females, and in the HPC of D1 females. Consistent with the role of E₂ and P₄ to promote BDNF expression/release (Kaur et al., 2007; Scharfman et al., 2003; Segal and Murphy, 2001), Pro females exhibited higher levels of BDNF within the PFC as compared to D1 females in all treatment conditions. However, we observed no ovarian hormone-related difference in BDNF levels within the HPC. This may be due to our collection of the entire CA1/CA3 region, since the estradiol-related expression of BDNF exhibits subregion specificity (e.g., CA3) (Spencer-Segal et al., 2012).

There are many intracellular signaling pathways that are downstream of BDNF and are activated in response to antidepressant doses of ketamine. Levels of phosphorylated MAPK have been shown to rapidly rise in response to antidepressant doses of ketamine in the PFC of male rodents (Li et al., 2010). Unexpectedly, we observed no effect of ketamine over activation of this kinase within the PFC. This lack of effect may be due to methodological differences between the studies such as rodent species (rat vs. mouse) or dose of ketamine (3 vs. 10 mg/kg). Additionally, protein extraction technique (synaptoneurosomal vs. total protein) may have impacted our ability to detect ketamine-induced phosphorylation of MAPK, as we used total protein and observed no effect of ketamine, consistent with another report (Réus et al., 2014). It is important to note that other studies have reported ketamine-induced elevations in p-MAPK within the HPC using total protein (Choi et al., 2015; Park et al., 2014), which we confirmed in the present study. E₂/P₄ exert neurotrophic functions through their ability to promote activation of MAPK within the HPC (Boonyaratankornkit, 2011; Leonhardt et al., 2003; Orr et al., 2012; Tuscher et al., 2016). In line with these data, we demonstrate that D1 females exhibited lower levels of HPC p-MAPK as compared to Pro females. Interestingly, while 1.5 mg/kg ketamine was sub-threshold for a behavioral effect in D1 females, this dose did significantly increase p-MAPK, which may be due to the low baseline levels of p-MAPK.

The phosphoinositide 3-kinase/protein kinase B (PI3K-Akt) pathway is another downstream intracellular signaling target of BDNF. Consistent with previous reports (Caffino et al., 2016; Li et al., 2010), we demonstrate that an antidepressant dose of ketamine increases levels of p-Akt within the PFC and HPC of male rodents. E₂/P₄ promote activation of the PI3K-Akt pathway and this is one mechanism through which they promote cell survival and neuronal plasticity (Briz and Baudry, 2014; Fortress et al., 2013; Singh, 2001). This is consistent with our finding that Pro females exhibit an increase in PFC and HPC p-Akt in a manner that directly mirrored their behavioral sensitivity to ketamine (i.e., 1.5 and 3 mg/kg). Following phosphorylation, p-Akt promotes synaptic plasticity via

inhibition of the pro-apoptotic signaling molecule, GSK-3 β (Beaulieu, 2012; Beurel et al., 2015). In line with the relationship between Akt and GSK-3 β , ketamine promotes the phosphorylation of GSK-3 β in the PFC and HPC of male rodents (Beurel et al., 2011; Zhou et al., 2014a), and this phosphorylation is required for ketamine's rapid antidepressant-like effects (Beurel et al., 2011). Consistent with these reports we demonstrate that ketamine induced an increase in p-GSK-3 β in the male and Pro PFC, as well as the HPC of all groups. It is important to note that in addition to the PI3K-Akt pathway, there are many ketamine-activated kinases that promote phosphorylation of GSK-3 β (e.g., p70 S6 kinase) (Beurel et al., 2015). Thus, our finding that there was no effect of ketamine to affect phosphorylation of Akt in the HPC of D1 females, yet an effect of ketamine to influence levels of p-GSK-3 β is somewhat surprising, but not unfounded.

mTOR is another neurotrophic signaling molecule that has garnered much attention in the field of antidepressant research, and is also downstream of BDNF and its associated signaling pathways (e.g., Akt and GSK-3 β) (Beurel et al., 2011; Liu et al., 2013; Navé et al., 1999; Zoncu et al., 2011). Phosphorylation of mTOR and its ability to promote the translation of synaptic plasticity-related proteins in the PFC are required for the rapid antidepressant-like effects of ketamine in males (Li et al., 2010). Indeed, we found that a behaviorally effective dose of ketamine induced activation of mTOR in the PFC of males and D1 females, and in the HPC of males and Pro females.

The present study aimed to determine doses of ketamine that exert antidepressant-like effects in mice and to elucidate the contribution of endogenous ovarian hormones to mediate sex differences in ketamine sensitivity. We demonstrated that females experiencing a natural peak in levels of the ovarian hormones E₂ and P₄ exhibited heightened behavioral and biochemical sensitivity to ketamine. Taking this a step further, we demonstrate that pharmacological activation of ER α or ER β was sufficient to render D1 females sensitive to a dose of ketamine that was only effective in Pro females. This finding indicates that action of E₂ at either of these ER subtypes may mediate Pro females heightened behavioral sensitivity to ketamine. Investigation into the potential molecular mechanism(s) underlying Pro females' ketamine sensitivity revealed some interesting clues. First, we confirmed that in male mice behaviorally effective dose of ketamine increased the expression and/or activation of many neurotrophic pathways within the PFC and HPC. Then, we identified ketamine-sensitive biochemical pathways in these same mood-related brain regions in D1 and Pro females. Overall, when compared to D1 counterparts, Pro females were more sensitive to the biochemical effects of ketamine in both the PFC and HPC, which is in line with evidence that both of these mood-related brain regions are robustly influenced by ovarian hormones and display estrous cycle-related changes in synaptogenesis and neuronal excitability (Scharfman and MacLusky, 2006). Furthermore, we found that Pro females exhibited activation of Akt and CaMKII α within mood-related brain regions in response to 1.5 mg/kg ketamine; indicating that ovarian hormones may prime these signaling pathways to render females sensitive to male-subthreshold doses of ketamine. While these results provide some insight into the molecular mechanisms of ketamine within low and high ovarian hormone conditions, future studies need to examine the effects of ketamine in these hormone states in animals not exposed to an FST, as the acute stress associated with this test has the potential to influence expression of synaptic plasticity markers.

The present study goes some way to determine the molecular mechanisms underlying sex differences in behavioral sensitivity to ketamine, yet it is important to note that other mechanisms may also contribute to the observed effects. Sex differences in

pharmacokinetics and pharmacodynamics are well established (Bigos et al., 2009; Soldin and Mattison, 2009; Waxman and Holloway, 2009), and recent work by Zanos et al. (2016), demonstrated that female mice have three-fold higher levels of the ketamine metabolite, (2S,6; 2R6R)hydroxynorketamine, up to 1 h following treatment with ketamine. Taken together, these studies suggest that females may take longer to clear metabolites of ketamine, which may explain sex differences in ketamine antidepressant-like effects. Furthermore, ovarian hormones can influence excitatory receptor expression and/or receptor sensitivity (Gazzaley et al., 1996; Woolley et al., 1997), which may influence sensitivity to the antidepressant effects of ketamine. Thus, further investigation into the impact of sex and hormonal status over sensitivity to the rapid-acting antidepressant ketamine is warranted.

Conflicts of interest

The authors declare no conflict of interest.

Funding

This work was supported by the National Institute of Mental Health grant to M.K. (R01 MH087583 and R01 MH099085).

Acknowledgements

We thank Dania Tawfiq for her outstanding assistance throughout the duration of the experiment.

Appendix A. Supplementary data

Supplementary data related to this article can be found at <https://doi.org/10.1016/j.neuropharm.2017.11.022>.

References

- Abdallah, C.G., Sanacora, G., Duman, R.S., Krystal, J.H., 2015. Ketamine and rapid-acting antidepressants: a window into a new neurobiology for mood disorder therapeutics. *Annu. Rev. Med.* 66, 509–523.
- Allen, P.J., D'Anci, K.E., Kanarek, R.B., Renshaw, P.F., 2012. Sex-specific antidepressant effects of dietary creatine with and without sub-acute fluoxetine in rats. *Pharmacol. Biochem. Behav.* 101, 588–601.
- Autry, A.E., Adachi, M., Nosyreva, E., Na, E.S., Los, M.F., Cheng, P.F., Kavalali, E.T., Monteggia, L.M., 2011. NMDA receptor blockade at rest triggers rapid behavioural antidepressant responses. *Nature* 475, 91–95.
- Barria, A., Muller, D., Derkach, V., Griffith, L.C., Soderling, T.R., 1997. Regulatory phosphorylation of AMPA-type glutamate receptors by CaM-KII during long-term potentiation. *Science* 276, 2042–2045.
- Beaulieu, J.M., 2012. A role for Akt and glycogen synthase kinase-3 as integrators of dopamine and serotonin neurotransmission in mental health. *J. Psychiatry Neurosci.* 37, 7–16.
- Berman, R.M., Cappiello, A., Anand, A., Oren, D.A., Heninger, G.R., Charney, D.S., Krystal, J.H., 2000. Antidepressant effects of ketamine in depressed patients. *Biol. Psychiatry* 47, 351–354.
- Beurel, E., Grieco, S.F., Jope, R.S., 2015. Glycogen synthase kinase-3 (GSK3): regulation, actions, and diseases. *Pharmacol. Ther.* 148, 114–131.
- Beurel, E., Song, L., Jope, R.S., 2011. Inhibition of glycogen synthase kinase-3 is necessary for the rapid antidepressant effect of ketamine in mice. *Mol. Psychiatry* 16, 1068–1070.
- Bigos, K.L., Pollock, B.G., Stankevich, B.A., Bies, R.R., 2009. Sex differences in the pharmacokinetics and pharmacodynamics of antidepressants: an updated review. *Gend. Med.* 6, 522–543.
- Björkholm, C., Monteggia, L.M., 2016. BDNF - a key transducer of antidepressant effects. *Neuropharmacology* 102, 72–79.
- Boonyaratankornkit, V., 2011. Scaffolding proteins mediating membrane-initiated extra-nuclear actions of estrogen receptor. *Steroids* 76, 877–884.
- Briz, V., Baudry, M., 2014. Estrogen regulates protein synthesis and actin polymerization in hippocampal neurons through different molecular mechanisms. *Front. Endocrinol. (Lausanne)* 5, 22.
- Caffino, L., Di Chio, M., Giannotti, G., Venniro, M., Mutti, A., Padovani, L., Cheung, D., Fumagalli, G.F., Yew, D.T., Fumagalli, F., Chiamulera, C., 2016. The modulation of BDNF expression and signalling dissects the antidepressant from the reinforcing properties of ketamine: effects of single infusion vs. chronic self-administration in rats. *Pharmacol. Res.* 104, 22–30.
- Can, A., Dao, D.T., Arad, M., Terrillion, C.E., Piantadosi, S.C., Gould, T.D., 2012. The mouse forced swim test. *J. Vis. Exp.* 59, 36338.
- Carrier, N., Kabbaj, M., 2013. Sex differences in the antidepressant-like effects of ketamine. *Neuropharmacology* 70, 27–34.
- Choi, M., Lee, S.H., Wang, S.E., Ko, S.Y., Song, M., Choi, J.S., Kim, Y.S., Duman, R.S., Son, H., 2015. Ketamine produces antidepressant-like effects through phosphorylation-dependent nuclear export of histone deacetylase 5 (HDAC5) in rats. *Proc. Natl. Acad. Sci. U. S. A.* 112, 15755–15760.
- Cooper, R., Goldman, J., Vandenberg, J., 1993. Monitoring of Estrous Cyclicity in the Laboratory Rodent by Vaginal Lavage. Academic Press, New York.
- Derkach, V.A., Oh, M.C., Guire, E.S., Soderling, T.R., 2007. Regulatory mechanisms of AMPA receptors in synaptic plasticity. *Nat. Rev. Neurosci.* 8, 101–113.
- Duman, R.S., Aghajanian, G.K., Sanacora, G., Krystal, J.H., 2016. Synaptic plasticity and depression: new insights from stress and rapid-acting antidepressants. *Nat. Med.* 22, 238–249.
- Estrada-Camarena, E., Rivera, N.M., Berlanga, C., Fernández-Guasti, A., 2008. Reduction in the latency of action of antidepressants by 17 beta-estradiol in the forced swimming test. *Psychopharmacol. Berl.* 201, 351–360.
- Fortress, A.M., Fan, L., Orr, P.T., Zhao, Z., Frick, K.M., 2013. Estradiol-induced object recognition memory consolidation is dependent on activation of mTOR signaling in the dorsal hippocampus. *Learn Mem.* 20, 147–155.
- Franceschelli, A., Sens, J., Herchick, S., Thelen, C., Pitychoutis, P.M., 2015. Sex differences in the rapid and the sustained antidepressant-like effects of ketamine in stress-naïve and “depressed” mice exposed to chronic mild stress. *Neuroscience* 290, 49–60.
- Frick, K.M., 2015. Molecular mechanisms underlying the memory-enhancing effects of estradiol. *Horm. Behav.* 74, 4–18.
- Frye, C.A., 2011. Progesterone attenuates depressive behavior of younger and older adult C57/BL6, wildtype, and progesterone receptor knockout mice. *Pharmacol. Biochem. Behav.* 99, 525–531.
- Frye, C.A., Walf, A.A., 2002. Changes in progesterone metabolites in the hippocampus can modulate open field and forced swim test behavior of proestrous rats. *Horm. Behav.* 41, 306–315.
- Garcia, L.S., Comim, C.M., Valvassori, S.S., Réus, G.Z., Barbosa, L.M., Andreazza, A.C., Stertz, L., Fries, G.R., Gavioli, E.C., Kapczynski, F., Quevedo, J., 2008. Acute administration of ketamine induces antidepressant-like effects in the forced swimming test and increases BDNF levels in the rat hippocampus. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 32, 140–144.
- Gazzaley, A.H., Weiland, N.G., McEwen, B.S., Morrison, J.H., 1996. Differential regulation of NMDAR1 mRNA and protein by estradiol in the rat hippocampus. *J. Neurosci.* 16, 6830–6838.
- Gideons, E.S., Kavalali, E.T., Monteggia, L.M., 2014. Mechanisms underlying differential effectiveness of memantine and ketamine in rapid antidepressant responses. *Proc. Natl. Acad. Sci. U. S. A.* 111, 8649–8654.
- Goldman, J.M., Murr, A.S., Cooper, R.L., 2007. The rodent estrous cycle: characterization of vaginal cytology and its utility in toxicological studies. *Birth Defects Res. B Dev. Reprod. Toxicol.* 80, 84–97.
- Hasegawa, Y., Hojo, Y., Kojima, H., Ikeda, M., Hotta, K., Sato, R., Oishi, Y., Yoshiya, M., Chung, B.C., Yamazaki, T., Kawato, S., 2015. Estradiol rapidly modulates synaptic plasticity of hippocampal neurons: involvement of kinase networks. *Brain Res.* 1621, 147–161.
- Herrington, B.E., Nicoll, R.A., 2016. Long-term potentiation: from CaMKII to AMPA receptor trafficking. *Annu. Rev. Physiol.* 78, 351–365.
- Höflich, A., Hahn, A., Küblböck, M., Kranz, G.S., Vanicek, T., Ganger, S., Spies, M., Windischberger, C., Kasper, S., Winkler, D., Lanzemberger, R., 2017. Ketamine-dependent neuronal activation in healthy volunteers. *Brain Struct. Funct.* 222, 1533–1542.
- Kaur, P., Jodhka, P.K., Underwood, W.A., Bowles, C.A., de Fiebre, N.C., de Fiebre, C.M., Singh, M., 2007. Progesterone increases brain-derived neurotrophic factor expression and protects against glutamate toxicity in a mitogen-activated protein kinase- and phosphoinositide-3 kinase-dependent manner in cerebral cortical explants. *J. Neurosci. Res.* 85, 2441–2449.
- Kavalali, E.T., Monteggia, L.M., 2012. Synaptic mechanisms underlying rapid antidepressant action of ketamine. *Am. J. Psychiatry* 169, 1150–1156.
- Kavalali, E.T., Monteggia, L.M., 2015. How does ketamine elicit a rapid antidepressant response? *Curr. Opin. Pharmacol.* 20C, 35–39.
- Kendall, D.A., Stancel, G.M., Enna, S.J., 1981. Imipramine: effect of ovarian steroids on modifications in serotonin receptor binding. *Science* 211, 1183–1185.
- Kessler, R.C., 2003. Epidemiology of women and depression. *J. Affect. Disord.* 74, 5–13.
- Kokras, N., Antoniou, K., Mikail, H.G., Kafetzopoulos, V., Papadopoulou-Daifoti, Z., Dalla, C., 2015. Forced swim test: what about females? *Neuropharmacology* 99, 408–421.
- Leonhardt, S.A., Boonyaratankornkit, V., Edwards, D.P., 2003. Progesterone receptor transcription and non-transcription signaling mechanisms. *Steroids* 68, 761–770.
- Lepack, A.E., Fuchikami, M., Dwyer, J.M., Banasr, M., Duman, R.S., 2015. BDNF release is required for the behavioral actions of ketamine. *Int. J. Neuropharmacol.* 18.
- Li, M., Demenescu, L.R., Colic, L., Metzger, C.D., Heinze, H.J., Steiner, J., Speck, O., Fejtova, A., Salvadore, G., Walter, M., 2017. Temporal dynamics of antidepressant ketamine effects on glutamine cycling follow regional fingerprints of AMPA and NMDA receptor densities. *Neuropsychopharmacology* 42, 1201–1209.

- Li, N., Lee, B., Liu, R.J., Banasr, M., Dwyer, J.M., Iwata, M., Li, X.Y., Aghajanian, G., Duman, R.S., 2010. mTOR-dependent synapse formation underlies the rapid antidepressant effects of NMDA antagonists. *Science* 329, 959–964.
- Li, N., Liu, R.J., Dwyer, J.M., Banasr, M., Lee, B., Son, H., Li, X.Y., Aghajanian, G., Duman, R.S., 2011. Glutamate N-methyl-D-aspartate receptor antagonists rapidly reverse behavioral and synaptic deficits caused by chronic stress exposure. *Biol. Psychiatry* 69, 754–761.
- Lisman, J., Yasuda, R., Raghavachari, S., 2012. Mechanisms of CaMKII action in long-term potentiation. *Nat. Rev. Neurosci.* 13, 169–182.
- Liu, R.J., Fuchikami, M., Dwyer, J.M., Lepack, A.E., Duman, R.S., Aghajanian, G.K., 2013. GSK-3 inhibition potentiates the synaptogenic and antidepressant-like effects of subthreshold doses of ketamine. *Neuropsychopharmacology* 38, 2268–2277.
- Mahmoud, R., Wainwright, S.R., Chaiton, J.A., Lieblich, S.E., Galea, L.A., 2016. Ovarian hormones, but not fluoxetine, impart resilience within a chronic unpredictable stress model in middle-aged female rats. *Neuropharmacology* 107, 278–293.
- McLean, A.C., Valenzuela, N., Fai, S., Bennett, S.A., 2012. Performing vaginal lavage, crystal violet staining, and vaginal cytological evaluation for mouse estrous cycle staging identification. *J. Vis. Exp.* 15 e43889.
- Moghaddam, B., Adams, B., Verma, A., Daly, D., 1997. Activation of glutamatergic neurotransmission by ketamine: a novel step in the pathway from NMDA receptor blockade to dopaminergic and cognitive disruptions associated with the prefrontal cortex. *J. Neurosci.* 17, 2921–2927.
- Navé, B.T., Ouwens, M., Withers, D.J., Alessi, D.R., Shepherd, P.R., 1999. Mammalian target of rapamycin is a direct target for protein kinase B: identification of a convergence point for opposing effects of insulin and amino-acid deficiency on protein translation. *Biochem. J.* 344 (Pt 2), 427–431.
- Nosyreva, E., Szabla, K., Autry, A.E., Ryazanov, A.G., Monteggia, L.M., Kavalali, E.T., 2013. Acute suppression of spontaneous neurotransmission drives synaptic potentiation. *J. Neurosci.* 33, 6990–7002.
- Nutt, D.J., 2002. The neuropharmacology of serotonin and noradrenaline in depression. *Int. Clin. Psychopharmacol.* 17 (Suppl. 1), S1–S12.
- Orr, P.T., Rubin, A.J., Fan, L., Kent, B.A., Frick, K.M., 2012. The progesterone-induced enhancement of object recognition memory consolidation involves activation of the extracellular signal-regulated kinase (ERK) and mammalian target of rapamycin (mTOR) pathways in the dorsal hippocampus. *Horm. Behav.* 61, 487–495.
- Park, S.W., Lee, J.G., Seo, M.K., Lee, C.H., Cho, H.Y., Lee, B.J., Seol, W., Kim, Y.H., 2014. Differential effects of antidepressant drugs on mTOR signalling in rat hippocampal neurons. *Int. J. Neuropsychopharmacol.* 17, 1831–1846.
- Porsolt, R.D., Bertin, A., Jalfre, M., 1977. Behavioral despair in mice: a primary screening test for antidepressants. *Arch. Int. Pharmacodyn. Ther.* 229, 327–336.
- Rush, A.J., Trivedi, M.H., Wisniewski, S.R., Nierenberg, A.A., Stewart, J.W., Warden, D., Niederehe, G., Thase, M.E., Lavori, P.W., Lebowitz, B.D., McGrath, P.J., Rosenbaum, J.F., Sackeim, H.A., Kupfer, D.J., Luther, J., Fava, M., 2006. Acute and longer-term outcomes in depressed outpatients requiring one or several treatment steps: a STAR*D report. *Am. J. Psychiatry* 163, 1905–1917.
- Réus, G.Z., Vieira, F.G., Abelaira, H.M., Michels, M., Tomaz, D.B., dos Santos, M.A., Carlessi, A.S., Neotti, M.V., Matias, B.I., Luz, J.R., Dal-Pizzol, F., Quevedo, J., 2014. MAPK signaling correlates with the antidepressant effects of ketamine. *J. Psychiatr. Res.* 55, 15–21.
- Saland, S.K., Schoepfer, K.J., Kabbaj, M., 2016. Hedonic sensitivity to low-dose ketamine is modulated by gonadal hormones in a sex-dependent manner. *Sci. Rep.* 6, 21322.
- Sarkar, A., Kabbaj, M., 2016. Sex differences in effects of ketamine on behavior, spine density, and synaptic proteins in socially isolated rats. *Biol. Psychiatry*.
- Scharfman, H.E., MacLusky, N.J., 2006. Estrogen and brain-derived neurotrophic factor (BDNF) in hippocampus: complexity of steroid hormone-growth factor interactions in the adult CNS. *Front. Neuroendocrinol.* 27, 415–435.
- Scharfman, H.E., Mercurio, T.C., Goodman, J.H., Wilson, M.A., MacLusky, N.J., 2003. Hippocampal excitability increases during the estrous cycle in the rat: a potential role for brain-derived neurotrophic factor. *J. Neurosci.* 23, 11641–11652.
- Segal, M., Murphy, D., 2001. Estradiol induces formation of dendritic spines in hippocampal neurons: functional correlates. *Horm. Behav.* 40, 156–159.
- Singh, M., 2001. Ovarian hormones elicit phosphorylation of Akt and extracellular-signal regulated kinase in explants of the cerebral cortex. *Endocrine* 14, 407–415.
- Soldin, O.P., Mattison, D.R., 2009. Sex differences in pharmacokinetics and pharmacodynamics. *Clin. Pharmacokinet.* 48, 143–157.
- Spencer-Segal, J.L., Tsuda, M.C., Mattei, L., Waters, E.M., Romeo, R.D., Milner, T.A., McEwen, B.S., Ogawa, S., 2012. Estradiol acts via estrogen receptors alpha and beta on pathways important for synaptic plasticity in the mouse hippocampal formation. *Neuroscience* 202, 131–146.
- Tuscher, J.J., Luine, V., Frankfurt, M., Frick, K.M., 2016. Estradiol-mediated spine changes in the dorsal Hippocampus and medial prefrontal cortex of ovariectomized female mice depend on ERK and mTOR activation in the dorsal Hippocampus. *J. Neurosci.* 36, 1483–1489.
- Van de Velde, S., Bracke, P., Levecque, K., 2010. Gender differences in depression in 23 European countries. Cross-national variation in the gender gap in depression. *Soc. Sci. Med.* 71, 305–313.
- Walf, A.A., Frye, C.A., 2009. Effects of two estradiol regimens on anxiety and depressive behaviors and trophic effects in peripheral tissues in a rodent model. *Gen. Med.* 6, 300–311.
- Walf, A.A., Frye, C.A., 2010. Estradiol reduces anxiety- and depression-like behavior of aged female mice. *Physiol. Behav.* 99, 169–174.
- Waxman, D.J., Holloway, M.G., 2009. Sex differences in the expression of hepatic drug metabolizing enzymes. *Mol. Pharmacol.* 76, 215–228.
- Wilson, M.A., Dwyer, K.D., Roy, E.J., 1989. Direct effects of ovarian hormones on antidepressant binding sites. *Brain Res. Bull.* 22, 181–185.
- Woolley, C.S., McEwen, B.S., 1992. Estradiol mediates fluctuation in hippocampal synapse density during the estrous cycle in the adult rat. *J. Neurosci.* 12, 2549–2554.
- Woolley, C.S., Weiland, N.G., McEwen, B.S., Schwartzkroin, P.A., 1997. Estradiol increases the sensitivity of hippocampal CA1 pyramidal cells to NMDA receptor-mediated synaptic input: correlation with dendritic spine density. *J. Neurosci.* 17, 1848–1859.
- Wright, K.N., Strong, C.E., Addonizio, M.N., Brownstein, N.C., Kabbaj, M., 2017. Reinforcing properties of an intermittent, low dose of ketamine in rats: effects of sex and cycle. *Psychopharmacol. Berl.* 234, 393–401.
- Zanos, P., Moaddel, R., Morris, P.J., Georgiou, P., Fischell, J., Elmer, G.I., Alkondon, M., Yuan, P., Pribut, H.J., Singh, N.S., Dossou, K.S., Fang, Y., Huang, X.P., Mayo, C.L., Wainer, I.W., Albuquerque, E.X., Thompson, S.M., Thomas, C.J., Zarate, C.A., Gould, T.D., 2016. NMDAR inhibition-independent antidepressant actions of ketamine metabolites. *Nature* 533, 481–486.
- Zarate, C.A., Singh, J.B., Carlson, P.J., Brutsche, N.E., Ameli, R., Luckenbaugh, D.A., Charney, D.S., Manji, H.K., 2006. A randomized trial of an N-methyl-D-aspartate antagonist in treatment-resistant major depression. *Arch. Gen. Psychiatry* 63, 856–864.
- Zhou, W., Dong, L., Wang, N., Shi, J.Y., Yang, J.J., Zuo, Z.Y., Zhou, Z.Q., 2014a. Akt mediates GSK-3 β phosphorylation in the rat prefrontal cortex during the process of ketamine exerting rapid antidepressant actions. *Neuroimmunomodulation* 21, 183–188.
- Zhou, W., Wang, N., Yang, C., Li, X.M., Zhou, Z.Q., Yang, J.J., 2014b. Ketamine-induced antidepressant effects are associated with AMPA receptors-mediated upregulation of mTOR and BDNF in rat hippocampus and prefrontal cortex. *Eur. Psychiatry* 29, 419–423.
- Zoncu, R., Efeyan, A., Sabatini, D.M., 2011. mTOR: from growth signal integration to cancer, diabetes and ageing. *Nat. Rev. Mol. Cell Biol.* 12, 21–35.