



Selective serotonin 5-HT_{2C} receptor activation suppresses the reinforcing efficacy of cocaine and sucrose but differentially affects the incentive-salience value of cocaine- vs. sucrose-associated cues

Kathryn A. Cunningham^{a,*}, Robert G. Fox^a, Noelle C. Anastasio^a, Marcy J. Bubar^a, Sonja J. Stutz^a, F. Gerard Moeller^b, Scott R. Gilbertson^c, Sharon Rosenzweig-Lipson^{d,1}

^a Center for Addiction Research, Department of Pharmacology and Toxicology, University of Texas Medical Branch, Galveston, TX, USA

^b Department of Psychiatry and Behavioral Sciences, University of Texas Health Sciences Center, Houston, TX, USA

^c Department of Chemistry, University of Houston, Houston, TX, USA

^d Neuroscience, Pfizer Global Research and Development, Groton, CT, USA

ARTICLE INFO

Article history:

Received 30 November 2010

Received in revised form

26 April 2011

Accepted 29 April 2011

Keywords:

5-HT_{2C} receptor

Addiction

Cocaine

Reward

Self-administration

Serotonin

Sucrose

WAY 163909

ABSTRACT

Serotonin (5-HT) controls affective and motivational aspects of palatable food and drug reward and the 5-HT_{2C} receptor (5-HT_{2C}R) has emerged as a key regulator in this regard. We have evaluated the efficacy of a selective 5-HT_{2C}R agonist, WAY 163909, in cocaine and sucrose self-administration and reinstatement assays employing parallel experimental designs in free-fed rats. WAY 163909 dose-dependently reduced the reinforcing efficacy of cocaine (ID₅₀ = 1.19 mg/kg) and sucrose (ID₅₀ = 0.7 mg/kg) as well as reinstatement (ID₅₀ = 0.5 mg/kg) elicited by exposure to cocaine-associated contextual cues, but not sucrose-associated contextual cues. The ID₅₀ of WAY 163909 predicted to decrease the reinforcing efficacy of cocaine or sucrose as well as reinstatement upon exposure to cocaine-associated cues was ~5–12-fold lower than that predicted to suppress horizontal ambulation (ID₅₀ = 5.89 mg/kg) and ~2–5-fold lower than that predicted to suppress vertical activity (ID₅₀ = 2.3 mg/kg). Thus, selective stimulation of the 5-HT_{2C}R decreases the reinforcing efficacy of cocaine and sucrose in freely-fed rats, but differentially alters the incentive-salience value of cocaine- vs. sucrose-associated cues at doses that do not impair locomotor activity. Future research is needed to tease apart the precise contribution of 5-HT_{2C}R neurocircuitry in reward and motivation and the learning and memory processes that carry the encoding for associations between environmental cues and consumption of rewarding stimuli. A more complete preclinical evaluation of these questions will ultimately allow educated proof-of-concept trials to test the efficacy of selective 5-HT_{2C}R agonists as adjunctive therapy in chronic health maladies including obesity, eating disorders and drug addiction.

This article is part of a Special Issue entitled 'Serotonin: The New Wave'.

© 2011 Published by Elsevier Ltd.

1. Introduction

Survival is ensured by neural circuits engaged by basic needs (e.g., food, water) and states (e.g., hunger, thirst) that motivate the organism to sustain important biological functions. Brain motivational systems are sensitive to endogenous and exogenous stimuli

that become linked to natural and drug rewards, and neuroplasticity in these circuits engender affective states that powerfully drive such behaviors as bingeing on palatable food (e.g., fat, sucrose) or abused drugs (e.g., cocaine). A complex appetitive process links the motivation to consume (e.g., hunger) with the palatability and incentive salience of the rewarding stimulus (drug or food) and its associated conditioned cue properties as well as with the satiety signals that terminate further intake (Halford and Harrold, 2008). Studies over the last 20 years have identified cellular and behavioral mechanisms of these appetitive processes to include dopamine, glutamate, and their intracellular signaling webs in the limbic–cortico-striatal–hypothalamic circuit (Kelley, 2004). Serotonin (5-HT) is additionally important in the control over the affective and motivational aspects of palatable food and

Abbreviations: 5-HT, serotonin; 5-HT_{2A}R, 5-HT_{2A} receptor; CHO, Chinese Hamster Ovary; FR, fixed ratio; i.p., intraperitoneal; Veh, vehicle; WAY, WAY 163909.

* Corresponding author. University of Texas Medical Branch, Galveston, TX 77550-0615, USA. Tel.: +1 409 772 9629; fax: +1 409 772 7050.

E-mail address: kcunning@utmb.edu (K.A. Cunningham).

¹ Current address: In Vivo Solutions, East Brunswick, NJ, USA.

drug reward, which has been described to occur at the level of satiety (Hewitt et al., 2002; Lyness et al., 1980; Blundell et al., 1980) as well as palatability or reinforcing efficacy (Wogar et al., 1991). In particular, there is a growing literature based upon genetic and pharmacological manipulations that the serotonin 5-HT_{2C} receptor (5-HT_{2C}R) signaling regulates such neurobehavioral processes which may underlie important chronic health maladies including obesity, eating disorders and drug addiction (Halford et al., 2010; Bubar and Cunningham, 2008; Steiger, 2004).

The selective activation of signaling through the 5-HT_{2C}R has been associated with reduced feeding and decreased body weight in animals (Fletcher et al., 2009; Clifton et al., 2000) and most recently in humans (Smith et al., 2009, 2010). The constitutive knockout of the 5-HT_{2C}R in mice results in hyperphagia (Tecott et al., 1995) the phenotype also includes adult-onset obesity and depressed metabolic rates (Tecott et al., 1995). Selective 5-HT_{2C}R antagonists have been shown to increase baseline food intake (Bonhaus et al., 1997; Thomsen et al., 2008) (but see, Hewitt et al., 2002) in the same dose range shown to block the effects of a 5-HT_{2C}R selective agonist to decrease feeding (Thomsen et al., 2008; Grottick et al., 2000). Analogous to these findings with palatable food reward, rates for responding for cocaine in a self-administration task were suppressed by pretreatment with a preferential 5-HT_{2C}R agonist and enhanced by a selective 5-HT_{2C}R antagonist (Fletcher et al., 2008; Grottick et al., 2000), suggesting an important role for the 5-HT_{2C}R in control of the overt rewarding efficacy of cocaine. A preferential 5-HT_{2C}R agonist also suppressed intake of nicotine (Grottick et al., 2001) and alcohol (Tomkins et al., 2002). The mechanisms are complex but a role for the 5-HT_{2C}R to enhance satiety and/or suppress incentive-motivational aspects of appetitive food or drug reinforcers are both supported (Fletcher et al., 2010; Burbassi and Cerro, 2008; Rocha et al., 2002; Vickers et al., 1999; Hewitt et al., 2002).

There is much less known about the role of serotonin in general, and the 5-HT_{2C}R in specific, to regulate drug- or food-seeking behavior upon exposure to environmental stimuli previously associated with drug or palatable food, respectively. Loss of 5-HT neurons after intraventricular infusion of a 5-HT neurotoxin decreased cocaine-seeking but enhanced sucrose-seeking during extinction in freely-fed rats previously trained to self-administer cocaine or sucrose, respectively (Tran-Nguyen et al., 2001). The selective 5-HT reuptake inhibitor fluoxetine was shown to suppress palatable food intake most potently in freely-fed female rats pre-exposed to food cues (Cifani et al., 2009). The preferential 5-HT_{2C}R agonists MK 212 (Neisewander and Acosta, 2007) and Ro 60-0175 (Fletcher et al., 2008) have been shown to suppress cue-evoked reinstatement in cocaine self-administration. However, the interpretation of findings across studies and the development of an overarching appreciation of how 5-HT_{2C}R neurocircuitry controls drug or palatable food intake or “relapse” in the face of reward-related cues are hampered by two issues. The first challenge is that selective agonists with singular high affinity and efficacy for the 5-HT_{2C}R have only recently become commercially available. The second challenge is that few studies have established the sensitivity of palatable reward vs. reward-related cues to a specific manipulation of 5-HT_{2C}R function employing comparable assay methodologies in a given species [for a recent review of these methodologies, see (Nair et al., 2009)].

The 5-HT_{2C} receptor (5-HT_{2C}R) shares high homology with the two other members (5-HT_{2A}R, 5-HT_{2B}R) of the 5-HT₂R family of G-protein-coupled receptors. Until recently, only “preferential” 5-HT_{2C}R ligands which frequently display affinity (agonists, antagonist) and/or efficacy (agonists) at the 5-HT_{2A}R and 5-HT_{2B}R have been available and some experimental outcomes with non-selective 5-HT_{2C}R ligands have led to ambiguous conclusions concerning

the biological roles for this receptor. Furthermore, as 5-HT_{2A}R or 5-HT_{2B}R agonists would be expected to evoke hallucinations (Nichols, 2004) or cardiac valvulopathy (Fitzgerald et al., 2000; Roth, 2007), respectively, therapeutically-useful 5-HT_{2C}R agonists must not have demonstrable efficacy at 5-HT_{2A}R or 5-HT_{2B}R *in vivo*. Such selectivity has been difficult to achieve, providing challenges to the careful preparation of preclinical analyses in support of ultimate proof-of-concept studies of selective 5-HT_{2C}R agonists in humans for treatment of obesity, eating disorders or addiction.

In a compound series recently developed at Wyeth Research, vabicaserin (SCA-136) was identified as a selective 5-HT_{2C}R full agonist ($K_i = 3$ nM; efficacy 100% relative to 5-HT), a 5-HT_{2B}R antagonist ($IC_{50} = 29$ nM) and a very weak 5-HT_{2A}R antagonist ($IC_{50} = 1650$ nM) (Rosenzweig-Lipson et al., 2007a; Tong et al., 2010a; Tong et al., 2010b). Vabicaserin has been in clinical trials to evaluate antipsychotic potential (www.clinicaltrials.gov). WAY 163909 is chemically-similar to vabicaserin with high affinity ($K_i = 10.5$ nM) and full efficacy (90% relative to 5-HT) at the 5-HT_{2C}R. WAY 163909 exhibits a low affinity ($K_i = 212$ nM) at the 5-HT_{2A}R, and no efficacy and is a weak partial agonist at the 5-HT_{2B}R at concentrations 23-fold greater (Dunlop et al., 2005). In the present study, we have evaluated the efficacy of WAY 163909 to suppress the reinforcing effects of the abused psychostimulant cocaine and the palatable food sucrose as well as cue-evoked reinstatement in parallel experimental designs in freely-fed rats. To date, the only other preferential 5-HT_{2C}R agonists to be profiled in a somewhat similar fashion are MK 212 and Ro 60-0175; Table 1 provides a comparison of affinity and efficacy at the three 5-HT₂R for MK 212, Ro 60-0175 and WAY 163909.

The present study investigated the hypothesis that WAY 163909 would suppress cocaine or sucrose self-administration or cue-evoked reinstatement in freely-fed male rats under conditions that minimized potential confounds imposed by food (or water) restriction [see (Grottick et al., 2000; Fletcher et al., 2002; Nic Dhonnchadha et al., 2009), for discussion] and operant pretraining for an appetitive reinforcer (Grottick et al., 2000; Fletcher et al., 2002; Nic Dhonnchadha et al., 2009). The doses of cocaine for reliable self-administration and reinstatement are well-established

Table 1

Affinity and efficacy of MK 212, Ro 60-0175 and WAY 163909 for 5-HT_{2A}R, 5-HT_{2B}R and 5-HT_{2C}R

Agonist	5-HT ₂ Receptor Family ^a					
	h5-HT _{2A} R		h5-HT _{2B} R		h5-HT _{2C} R	
	Affinity K_i (nM)	Efficacy E_{max}	Affinity K_i (nM)	Efficacy E_{max}	Affinity K_i (nM)	Efficacy E_{max}
MK 212	1023	<50%	617	75%	98	100%
Ro 60-0175	36.3	69%	5.4	79%	6.0	84%
WAY 163909	212	0%	2101	40%	10.5	90%

^a Published studies employed radioligand binding assays to establish the affinity (K_i) and efficacy (E_{max}) of MK 212 (Knight et al., 2004; Cussac et al., 2002; Porter et al., 1999) and Ro 60-0175 for the 5-HT₂R subtypes in h5-HT_{2A}R- or h5-HT_{2C}R-transfected Human Embryonic Kidney (HEK)-293 (Knight et al., 2004) and h5-HT_{2A}R-, h5-HT_{2B}R- or h5-HT_{2C}R-transfected Chinese Hamster Ovary (CHO)-K1 clonal cell lines (Porter et al., 1999; Cussac et al., 2002) while the affinity of WAY 163909 for each h5-HT₂R subtype was established in CHO-K1 cells (Dunlop et al., 2005). The efficacy of MK 212 (Porter et al., 1999), Ro 60-0175 (Porter et al., 1999) and WAY 163909 (Dunlop et al., 2005) was established in agonist-stimulated mobilization of intracellular calcium in CHO-K1 cells with a fluorometric imaging plate reader; the E_{max} is expressed as a percentage of the maximal response to 5-HT. MK 212 did not exhibit efficacy ($IC_{50} > 1$ μM) for the 5-HT_{2A}R (Knight et al., 2004). Ro 60-0175 did not exhibit efficacy ($IC_{50} > 1$ μM) for a multitude of proteins with the highest affinity for β_2 adrenoceptor (β_2 -noradrenergic receptor; $IC_{50} = 251$; Martin et al., 1998). WAY 163909 exhibited modest affinity for the dopamine D₄R ($K_i = 245$ nM), 5-HT₇R ($K_i = 343$ nM) and α_1 adrenoceptor ($K_i = 665$ nM; Dunlop et al., 2005).

in the literature (Nic Dhonnchadha et al., 2009; Carroll and Lac, 1997; Pentkowski et al., 2010; Grottick et al., 2000; Burbassi and Cervo, 2008; Fletcher et al., 2002) and we chose to compare cocaine to a sucrose reinforcer under the same assay conditions because preliminary observations indicated that freely-fed rats would establish levels of active lever presses for sucrose pellets under an ultimate fixed ratio (FR) 5 schedule that were comparable to those seen for a cocaine self-administration. Employing these analogous assay methodologies in the rat allowed us to establish the differential sensitivity of a natural (sucrose) and drug (cocaine) reward vs. reward-related cues to pretreatment with a highly-selective and efficacious 5-HT_{2C}R agonist and to compare with the body of literature available for the preferential 5-HT_{2C}R agonists MK 212 (Neisewander and Acosta, 2007) and Ro 60-0175 (Higgins et al., 2001; Grottick et al., 2001; Fletcher et al., 2008, 2010). We report that cocaine-taking and especially cocaine-seeking are highly sensitive to suppression by WAY 163909. Interestingly, sucrose-taking is also potently suppressed by WAY 163909, but sucrose-seeking is relatively insensitive to the selective 5-HT_{2C}R agonist in freely-fed rats.

2. Methods

2.1. Animals

A total of 91 male Sprague–Dawley rats (Harlan, Inc., Indianapolis, IN, USA) weighing 225–325 g at the start of the experiments were used. Rats were allowed to acclimate for 5–7 days in a colony room at a constant temperature (21–23 °C) and humidity (45–50%) on a 12 h light–dark cycle (lights on 0700–1900 h). Rats were housed either two (self-administration studies) or four rats per cage (locomotor activity studies) with food and water *ad libitum* throughout all phases of the studies. All experiments were carried out in accordance with the National Institutes of Health *Guide for the Care and Use of Laboratory Animals* (National Research Council, 1996) and with the approval of the Institutional Animal Care and Use Committee at University of Texas Medical Branch. All efforts were made to minimize animal suffering, to reduce the number of animals used, and to utilize alternatives to *in vivo* techniques, when available.

2.2. Drugs

WAY 163909 [(7bR, 10aR)-1,2,3,4,8,9,10,10a-octahydro-7bH-cyclopenta-[b][1,4]diazepino [6,7,1hi]indole; Wyeth Research, Princeton, NJ, USA) was dissolved in 0.9% NaCl. SB 242084 [6-chloro-5-methyl-1-[[2-(2-methylpyrid-3-yloxy) pyrid-5-yl]carbomoyl]indolinedihydrochloride; Sigma Chemical Co., St. Louis, MO, USA) was dissolved in 0.9% NaCl containing 10 mM citric acid (Sigma Chemical Co.) and 8% 2-hydroxypropyl-β-cyclodextrin (Trappsol® Hydroxypropyl Beta Cyclodextrin, Pharmaceutical Grade, Cyclodextrin Technologies Development Inc., High Springs, FL, USA) with the final pH of the solution adjusted to 5.6. (–)-Cocaine (National Institutes on Drug Abuse, Research Triangle Park, NC, USA) was dissolved in 0.9% NaCl.

2.3. Locomotor activity studies

2.3.1. Apparatus

Locomotor activity was monitored and quantified under low light conditions using a modified open field activity system (San Diego Instruments, San Diego, CA, USA). Clear Plexiglass chambers (40 × 40 × 40 cm) were surrounded by a 4 × 4 photobeam matrix positioned 4 cm from chamber floor. Consecutive photobeam breaks within the central 16 × 16 cm of the activity monitor were recorded as central ambulation. Peripheral ambulation was counted as consecutive beam breaks in the surrounding perimeter. Central and peripheral ambulations were summed to provide a measure of total horizontal ambulation. Vertical activity was also recorded using a row of 16 photobeams, positioned 16 cm from the activity monitor floor; breaks in these beams indicated vertical (rearing) activity.

2.3.2. Procedures

Rats in locomotor activity studies (*n* = 16) were acclimated to the colony room 5–7 days prior to the start of handling protocols. Following seven days of handling, rats were habituated to the activity monitors for 60 min. Twenty four hours later, rats received the first dose of vehicle (saline, 1 ml/kg, i.p.) or WAY 163909 (0.3, 1, 3 or 10 mg/kg, i.p.) immediately prior to placement in activity monitors; activity was assessed for 60 min. Using a repeated-measures design, rats were tested every three days such that all rats received vehicle and all three doses of WAY 163909 in a counterbalanced manner.

2.4. Self-administration studies

2.4.1. Surgery

Implantations of intravenous catheters with back mounts were performed under anesthesia with a cocktail containing 8.6 mg/kg of xylazine, 1.5 mg/kg of acepromazine, and 43 mg/kg of ketamine in bacteriostatic saline. Small incisions were made in the right posterior neck and just below the left shoulder blade to expose the jugular vein and to insert the catheter base, respectively. A subcutaneous burrow was made between the two incisions, and the catheter apparatus was pulled through this burrow. The catheter was composed of Silastic tubing (Dow Corning, Midland MI, USA) connected to a bent 22-gauge metal cannula encased within a plastic screw connector (Plastics One, Roanoke, VA, USA) at one end and affixed with a small ball of silicone 3.5 cm from the other end. The distal end of the catheter was inserted into the jugular vein until flush with the silicone ball, terminating outside the right atrium. The catheter was secured to the vein with sutures (SP116 braided silk non-absorbable 4.0; Surgical Specialties Corporation, Reading, PA, USA) on both sides of the silicone ball. The incision was then sutured (Monomid nylon non-absorbable 4.0; CP Medical, Portland, OR, USA). The cannula base was encased with dental acrylic and a small mesh circle (3 cm × 3 cm; Polypro mesh 500 micron, Small Parts, Inc., Seattle, WA, USA) affixed to the bottom, which was pulled thru the small incision in the back. The remaining opening was then sutured (Monomid nylon non-absorbable 4.0; CP Medical). Daily flushes with a solution of 0.1 mL of bacteriostatic saline containing heparin sodium (10 U/mL; American Pharmaceutical Partners, East Schaumburg, IL, USA), streptokinase (0.67 mg/mL; Sigma Chemical) and ticarcillin disodium (66.67 mg/mL; Research Products International, Mt. Prospect, IL, USA) were performed after each cocaine self-administration session to maintain catheter patency. Proper catheter function was verified periodically throughout the experiment by intravenous administration of 10 mg/kg of methohexital sodium (Monarch Pharmaceuticals Inc., Bristol, TN, USA), a dose sufficient to briefly anesthetize the animal only when administered intravenously. All rats were allowed at least five days of recovery after surgery before initiation of self-administration training. Rats trained in sucrose self-administration were naïve at the start of the experiment.

2.4.2. Apparatus

Standard operant conditioning chambers (Med-Associates, Inc., St. Albans, VT, USA) housed in ventilated sound-attenuating cubicles with fans (Med-Associates, Inc.) were utilized for both the cocaine and sucrose self-administration studies. Each chamber was equipped with two retractable response levers, a stimulus light above each response lever, a house light opposite the levers, and a magazine-type pellet dispenser. Cocaine infusions were delivered by a syringe attached to an infusion pump (Med-Associates, Inc.) located outside the cubicle. The infusion pumps were connected to liquid swivels (Instech, Plymouth Meeting, PA, USA) that were fastened to the catheters via polyethylene 20 tubing encased inside a metal spring leash (Plastics One). Sucrose pellets (45 mg; Bio-Serv, Frenchtown, NJ, USA) were delivered into a pellet receptacle located between the two levers.

2.4.3. Experimental design

2.4.3.1. Effects of WAY 163909 and SB 242084 during cocaine self-administration. Cocaine self-administration training consisted of daily 60-min sessions during which rats (*n* = 8) were trained to press the active lever to obtain a cocaine infusion (0.25 mg/kg/0.1 mL infusion). Animals were not food restricted or trained on an operant task prior to commencement of self-administration and no cocaine priming infusions were given during the experiment. Schedule completions on the active lever resulted in simultaneous illumination of the house and stimulus lights, followed 1 s later by activation of the infusion pump. Rats received a 0.1 mL cocaine infusion delivered over a 6-s period, after which the pump and stimulus light (conditioned stimuli paired with delivery of cocaine) were inactivated simultaneously. The house light remained illuminated for a 20-s timeout period, during which lever presses had no scheduled consequences. Responses on the inactive lever were recorded but had no scheduled consequences. Rats were initially trained on an FR 1 schedule of cocaine reinforcement until they met a criterion of seven infusions/hr for at least three sessions with <10% variation in the number of infusions received for three consecutive sessions. After reaching this acquisition criterion and demonstrating stable cocaine self-administration on an FR1 schedule of reinforcement, rats progressed to an FR 3 schedule of reinforcement, and then on to an FR 5 schedule of reinforcement once the same criterion levels were met. Once stable cocaine self-administration on an FR 5 schedule was achieved (i.e., <10% variation in the number of infusions received for three consecutive sessions), the efficacy and selectivity of WAY 163909 to alter response for cocaine self-administration was assessed. Rats were administered vehicle (saline; 1 mL/kg; 15 min) or WAY 163909 (0.5, 1 or 2 mg/kg, i.p.; 15 min) prior to self-administration sessions, according to a within-subjects design. To investigate the selectivity of WAY 163909 for the 5-HT_{2C} receptor, the same rats were administered the 5-HT_{2C} receptor antagonist SB 242084 (0.5 mg/kg, i.p.; 30 min) plus vehicle (saline; 1 mL/kg; 15 min) or WAY 163909 (2 mg/kg, i.p.; 15 min) prior to self-administration sessions, according to a within-subjects design. The order of injections was counterbalanced and a minimum of two intervening sessions of cocaine self-administration (0.25 mg/kg/0.1 mL infusion) occurred between each drug challenge to assure stability of baseline responding.

2.4.3.2. Effects of WAY 163909 on cue-induced reinstatement in rats trained to self-administer cocaine. A separate cohort of rats ($n = 34$) was utilized to assess the effects of WAY 163909 on cue-induced reinstatement following cocaine self-administration. Cocaine self-administration training consisted of daily 180-min sessions during which rats were trained to press the active lever to obtain a cocaine infusion (0.75 mg/kg/0.1 mL infusion) on an FR 1 schedule of reinforcement before progressing to an FR 5 schedule. Animals were not food restricted or trained on an operant task prior to commencement of self-administration and no cocaine priming infusions were given during the experiment. Rats that met the criterion for stable self-administration ($<10\%$ variation in the number of infusions received for three consecutive sessions) were subjected to daily 60-min extinction sessions, during which active and inactive lever presses were recorded but had no scheduled consequences (i.e., did not activate the infusion pump or result in presentation of the stimulus light). Once rats achieved the extinction criterion of response rates <15 total responses/hr for three consecutive sessions, the effects of WAY 163909 on cue-induced reinstatement of cocaine-seeking behavior were examined. The criterion of <15 responses/hr as indicative of extinction level responding is commonly employed in self-administration studies and corresponds to approximately 15% of responding during active cocaine self-administration (Leri et al., 2002; Nic Dhonnchadha et al., 2009). To initiate reinstatement sessions, one non-contingent presentation of the cocaine-paired cues (pump and stimulus light) was given. During the 60-min test session, responses on the active lever were reinforced by presentation of the cocaine-paired cues on an FR 1 schedule. To assess the effects of WAY 163909 on cue-induced reinstatement of cocaine-seeking behavior, rats were injected with saline (1 mL/kg) or WAY 163909 (0.5, 1, or 1.5 mg/kg, i.p.) 15 min before the start of the reinstatement session according to a between-subjects, counterbalanced design ($n = 8$ –9/group).

2.4.3.3. Effects of WAY 163909 and SB 242084 on sucrose self-administration. Sucrose self-administration training consisted of daily 60-min sessions during which rats ($n = 8$) were trained to press the active lever to obtain a 45 mg sucrose pellet. Experimental parameters were identical to those described in Section 2.4.3.1, above, except for substitution of a single 45 mg sucrose pellet as the reinforcer. Animals were not food restricted or trained on an operant task prior to commencement of self-administration and no sucrose priming events were given during the experiment. Upon reaching stability on an FR 5 schedule, the efficacy and selectivity of WAY 163909 to alter response for sucrose self-administration was assessed, as described above (Section 2.4.3.1).

2.4.3.4. Effects of WAY 163909 on cue-induced reinstatement in rats trained to self-administer sucrose pellets. A separate cohort of rats ($n = 25$) was utilized to assess the effects of WAY 163909 on cue-induced reinstatement following sucrose self-administration. Sucrose self-administration training consisted of daily 180-min sessions during which rats were trained to press the active lever to obtain a 45 mg sucrose pellet on an FR 1 schedule of reinforcement before progressing to an FR 5 schedule. Experimental parameters were identical to those described in Section 2.4.3.2, above, except for substitution of a single 45 mg sucrose pellet as the reinforcer. Animals were not food restricted or trained on an operant task prior to commencement of self-administration and no sucrose priming events were given during the experiment. To assess the effects of WAY 163909 on cue-induced reinstatement of sucrose-seeking behavior, rats were injected with saline (1 mL/kg) or WAY 163909 (0.5 or 1.5 mg/kg, i.p.) 15 min before the start of the reinstatement session according to a between-subjects, counterbalanced design ($n = 8$ –9/group).

2.5. Statistical analysis

Locomotor activity data were analyzed and are presented as mean total horizontal ambulation or total vertical activity (\pm SEM) over the 60-min session (Fig. 1A Inset, 1B Inset) or within the 15, 30, 45 and 60 min time bins (Fig. 1A, B). The main effect of treatment (WAY 163909 dose) on total horizontal ambulation or total vertical activity was analyzed using a repeated-measures, one-way analysis of variance (ANOVA) using the general linear model (GLM) procedure (SAS for Windows, Version 8.2, SAS Institute, Inc., Cary, NC, USA). A two-way ANOVA for repeated measures for the factors of treatment (WAY 163909 dose) and time (15, 30, 45 or 60 min) was also conducted. Subsequent *a priori* comparisons between means of total horizontal ambulation or total vertical activity were made using the Dunnett's procedure, with vehicle as the comparator. The four parameter logistic nonlinear regression model equation (Sigma Plot, Version 11.0, Systat Software, Inc., Chicago, IL, USA) was used to estimate the dose of WAY 163909 predicted to decrease horizontal ambulation or vertical activity by 50% (ID_{50}) (Table 2) (Ratkowsky and Reedy, 1986; Tallarida and Murray, 1987). All statistical analyses were conducted with an experiment wise error rate of $\alpha = 0.05$.

The data from the self-administration, extinction and reinstatement phases were analyzed separately and presented as mean \pm SEM. A one-way ANOVA (SAS for Windows, V8.2) for repeated measures was used to analyze the dependent measures of the total number of 1) active lever responses/session, 2) inactive lever presses/session, and 3) number of cocaine infusions or sucrose pellets/session during the self-administration phase and the total number of responses/session on the 4) previously-active, or 5) -inactive lever during the extinction phase. A one-way




Fig. 1. WAY 163909 evokes a dose-dependent suppression of horizontal ambulation and vertical activity. [A] The time course of horizontal ambulation is divided into 15 min time bins across the 60 min session; [A, Inset] the mean total horizontal ambulation (counts/60 min) (\pm SEM) is represented following administration of vehicle (saline; Veh) or WAY 163909 (WAY; 0.3, 1, 3 or 10 mg/kg, i.p.). [B] The time course of vertical activity is divided into 15 min time bins across the 60 min session; [B, Inset] the mean total vertical activity (counts/60 min) (\pm SEM) is represented following administration of vehicle (saline; Veh) or WAY 163909 (WAY; 0.3, 1, 3, or 10 mg/kg, i.p.). * $p < 0.05$ vs. Veh.

ANOVA for repeated measures (cocaine- or sucrose-taking) or independent groups (cocaine- or sucrose-seeking) was used to analyze the effects of pretreatment with WAY 163909 on the total number of 1) responses/session on the previously-active, or 2) -inactive lever, and 3) cocaine infusions/session or sucrose pellets/session, and 4) the latency to respond on the previously-active lever during the self-administration and the cue-induced reinstatement phase. *A priori* comparisons for WAY 163909 pretreatment were carried out with Student–Newman–Keuls test with the alpha level set at $p < 0.05$. A repeated-measures one-way ANOVA was employed to compare responding on the previously-active lever over the last three sessions of extinction with responding observed following vehicle pretreatment

during reinstatement sessions. The four parameter logistic nonlinear regression model equation (Sigma Plot, Version 11, Systat Software, Inc.) was used to estimate the dose of WAY 163909 predicted to decrease lever presses during cocaine or sucrose self-administration and reinstatement sessions by 50% (ID_{50}), for each (Ratkowsky and Reedy, 1986; Tallarida and Murray, 1987).

3. Results

3.1. WAY 163909 evokes a dose-dependent suppression of locomotor activity

A main effect of treatment ($F_{(4,66)} = 10.33$, $p < 0.001$), time ($F_{(3,66)} = 253.42$, $p < 0.001$) and a treatment \times time interaction ($F_{(12,198)} = 5.23$, $p < 0.001$) was observed for horizontal ambulation divided into four 15 min time bins (Fig. 1A). WAY 163909 at 0.3 mg/kg significantly reduced horizontal ambulation vs. vehicle during the third interval (45 min), while 3 mg/kg of WAY 163909 significantly reduced horizontal ambulation vs. vehicle at the first (15 min), second (30 min) and third intervals (45 min) ($p < 0.05$; Fig. 1A). WAY 163909 at 10 mg/kg significantly reduced horizontal ambulation vs. vehicle during each of the four 15 min intervals examined ($p < 0.05$; Fig. 1A). A main effect of WAY 163909 treatment was observed for horizontal ambulation ($F_{(4,66)} = 10.65$, $p < 0.001$; Fig. 1A, Inset) summed across the 60-min test session; *a priori* comparisons using the Dunnett's procedure revealed that 3 and 10 mg/kg of WAY 163909 significantly reduced horizontal ambulation vs. vehicle ($p < 0.05$; Fig. 1A, Inset). The dose of WAY 163909 predicted to decrease horizontal ambulation by 50% (ID_{50}) was 5.89 mg/kg (see Table 2).

A main effect of treatment ($F_{(4,66)} = 6.16$, $p < 0.001$), time ($F_{(3,66)} = 146.02$, $p < 0.001$) and a significant treatment \times time interaction ($F_{(12,198)} = 2.73$, $p < 0.01$) was observed for vertical activity divided into four 15 min intervals (Fig. 1B). WAY 163909 at 10 mg/kg significantly reduced vertical activity vs. vehicle during the first three 15 min intervals ($p < 0.05$; Fig. 1B). A main effect of WAY 163909 treatment was observed for vertical activity ($F_{(4,66)} = 5.66$, $p < 0.001$; Fig. 1B, Inset) summed across the 60-min test session; *a priori* comparisons using the Dunnett's procedure revealed that 10 mg/kg of WAY 163909 significantly reduced vertical activity vs. vehicle ($p < 0.05$; Fig. 1B, Inset). The dose of WAY 163909 predicted to decrease vertical activity by 50% (ID_{50}) was 2.3 mg/kg (see Table 2).

3.2. WAY 163909 dose-dependently suppresses cocaine self-administration

Rats ($n = 8$) readily acquired cocaine self-administration (0.25 mg/kg/0.1 ml infusion) to stability (i.e., seven infusions/hr on an FR 5 schedule for at least three sessions) and displayed $<10\%$ variation in the number of infusions received (i.e., cocaine intake) (Fig. 2A). Across the last three sessions of stable self-administration, there was no main effect of session on the number of active [$F_{(2,21)} = 0.00$, ns] or inactive lever responses [$F_{(2,17)} = 1.75$, ns], or the number of infusions received [$F_{(2,21)} = 0.00$, ns] (Fig. 2A). Average daily cocaine intake over the last three sessions of training was 4.4 ± 0.7 mg/kg.

An one-way ANOVA revealed a main effect of pretreatment on the number of active lever presses [$F_{(5,42)} = 2.92$; $p < 0.05$; Fig. 2B, left panel] and number of cocaine infusions received across the entire 60-min session [$F_{(5,42)} = 2.79$; $p < 0.05$; Fig. 2C, left panel]. No main effect was observed for the number of inactive lever presses [$F_{(5,42)} = 0.12$; ns; Fig. 2B]. *A priori* comparisons revealed that pretreatment with 2 mg/kg of WAY 163909 significantly reduced the number of active lever presses ($p < 0.05$; Fig. 2B, left panel) as well as the number of cocaine infusions ($p < 0.05$; Fig. 2C, left panel) without altering the number of inactive lever presses




Fig. 2. WAY 163909 dose-dependently suppresses cocaine self-administration. [A] Mean responses (\pm SEM) on the active (black circles) or inactive lever (white circles), and total number of cocaine infusions (\pm SEM; gray circles) are presented for the acquisition phase of cocaine self-administration training under FR1 – FR5 schedules of reinforcement. Active lever responses resulted in the delivery of an intravenous cocaine infusion (0.25 mg/kg/0.1 mL infusion) and simultaneous presentation of the cocaine-paired cues. [B] Mean responses (\pm SEM) on the active (black bars) or inactive lever (white bars) or [C] mean total cocaine infusions (\pm SEM; gray bars) following pretreatment with vehicle (saline; Veh) or WAY 163909 (0.5, 1 or 2 mg/kg, i.p.) (left panel) or following pretreatment with the 5-HT_{2C}R antagonist SB 242084 (0.5 mg/kg, i.p.) plus vehicle (saline; Veh) or WAY 163909 (2 mg/kg, i.p.) (right panel) during the cocaine self-administration maintenance phase. Active lever responses resulted in the delivery of an intravenous cocaine infusion (0.25 mg/kg/0.1 mL infusion) according to an FR 5 schedule and simultaneous presentation of cocaine-paired cues. * $p < 0.05$ vs. Veh; ^ $p < 0.05$ vs. WAY 163909 (2 mg/kg).

(ns; Fig. 2B). The dose of WAY 163909 predicted to decrease lever presses for cocaine by 50% (ID_{50}) is 1.19 mg/kg (see Table 2). Pretreatment with the selective 5-HT_{2C}R antagonist SB 242084 (0.5 mg/kg) completely blocked the effects of WAY 163909 (2 mg/kg) on the number active lever presses ($p < 0.05$; Fig. 2B, right panel) and number of cocaine infusions received ($p < 0.05$; Fig. 2C, right panel).

3.3. WAY 163909 dose-dependently suppresses cocaine cue-evoked reinstatement

Rats ($n = 34$) readily acquired cocaine self-administration (0.75 mg/kg/0.1 ml infusion) in daily 180 min sessions to stability (i.e., seven infusions/hr on an FR 5 schedule for at least three sessions) and displayed $<10\%$ variation in the number of infusions received (i.e., cocaine intake) (Fig. 3A). Across the last three sessions of stable self-administration on an FR 5 schedule, there was no main effect of session on the number of active [$F_{(2,99)} = 2.75$, ns] or inactive lever responding [$F_{(2,99)} = 0.63$, ns] or the number of

infusions received [$F_{(2,99)} = 2.42$, ns]. The rats were then separated into four groups for reinstatement sessions. Cocaine intake did not differ significantly across groups of rats assigned to receive pretreatment with vehicle (34.6 ± 1.8 infusions/session) or WAY 163909 at doses of 0.5 mg/kg (33.5 ± 1.8 infusions/session), 1 mg/kg (32 ± 1.2 infusions/session), or 1.5 mg/kg (31.9 ± 2.8 infusions/session) [$F_{(3,30)} = 0.62$; ns].

A decrease in presses on the previously-active lever was observed across extinction sessions with all rats achieving the extinction criterion (<15 active lever responses/session for three consecutive sessions) by session 12 of extinction training (data not shown). A repeated-measures ANOVA revealed a main effect of extinction session on active [$F_{(11,265)} = 78.8$; $p < 0.05$] and inactive lever presses [$F_{(11,265)} = 4.88$; $p < 0.05$] for all rats ($n = 34$) with both active and inactive lever presses decreasing across the extinction sessions (data not shown). The “extinction baseline” was calculated as the mean total lever presses of all rats on the active (7.8 ± 0.6) or inactive lever (4.9 ± 0.9) during the final 60 min extinction session (Fig. 3B).

Fig. 3B illustrates the effects of vehicle or WAY 163909 (0.5–1.5 mg/kg) pretreatment on reinstatement of responding supported by cues previously associated with cocaine availability. Cocaine-trained rats readily reinstated active lever pressing across the 60-min reinstatement session (Fig. 3B). A main effect of pretreatment on active lever responding was observed [$F_{(4,63)} = 43.02$; $p < 0.001$; Fig. 3B]. Significant decreases in active lever presses relative to vehicle were observed after pretreatment with WAY 163909 at doses of 0.5, 1 and 1.5 mg/kg ($p < 0.05$; Fig. 3B). There was no main effect of pretreatment on inactive lever responding [$F_{(4,61)} = 1.66$; ns; Fig. 3B]. The latency to the first lever press following pretreatment with WAY 163909 was unaltered [$F_{(3,30)} = 0.95$; ns] (data not shown). The dose of WAY 163909 predicted to decrease lever presses for cocaine-paired cues by 50% (ID_{50}) is 0.5 mg/kg (see Table 2).

3.4. WAY 163909 dose-dependently suppresses sucrose self-administration

Rats ($n = 8$) readily acquired sucrose self-administration (45 mg sucrose pellet/response) to stability (i.e., 10 pellets/hr on an FR 5 schedule for at least three sessions) and displayed $<10\%$ variation in the number of pellets received (i.e., pellet intake) (Fig. 4A). Across the last three sessions of stable self-administration on an FR5 schedule, there was no main effect of session on the number of active [$F_{(2,21)} = 0.01$, ns] or inactive operant lever responses [$F_{(2,21)} = 1.8$, ns] or the number of pellets received [$F_{(2,21)} = 0.03$, ns]. Average daily sucrose intake over the last three sessions of training was 23.0 ± 3.4 pellets.

An one-way ANOVA revealed a main effect of pretreatment on the number of active lever presses [$F_{(5,42)} = 9.88$; $p < 0.05$; Fig. 4B, left panel] and number of sucrose pellets received across the 60-min session [$F_{(5,42)} = 10.73$; $p < 0.05$; Fig. 4C, left panel]. No main effect was observed for the number of inactive lever presses [$F_{(5,42)} = 1.66$; ns; Fig. 4B]. *A priori* comparisons revealed that pretreatment with 1 or 2 mg/kg of WAY 163909 significantly reduced the number of active lever presses ($p < 0.05$; Fig. 4B, left panel) as well as the number of sucrose pellets received ($p < 0.05$; Fig. 4C, left panel), without altering the number of inactive lever presses (ns; Fig. 4B). The dose of WAY 163909 predicted to decrease lever presses for sucrose by 50% (ID_{50}) is 0.7 mg/kg (see Table 2). Pretreatment with the selective 5-HT_{2C}R antagonist SB 242084 (0.5 mg/kg) completely blocked the effects of WAY 163909 (2 mg/kg) on the number active lever presses ($p < 0.05$; Fig. 4B, right panel) and number of sucrose pellets received ($p < 0.05$; Fig. 4C, right panel).




Fig. 3. WAY 163909 dose-dependently suppresses cocaine cue-evoked reinstatement. [A] Mean responses (\pm SEM) on the active (black circles) or inactive lever (white circles), and the total number of cocaine infusions (\pm SEM; gray circles) are presented for the acquisition phase of cocaine self-administration training under FR1–FR5 schedules of reinforcement. Active lever responses resulted in the delivery of an intravenous cocaine infusion (0.75 mg/kg/0.1 mL infusion) and simultaneous presentation of the cocaine-paired cues. [B] Mean (\pm SEM) total number of responses on the active (black bars) or inactive lever (white bars) following extinction (“Extinction Baseline”) and upon pretreatment with vehicle (saline; Veh) or WAY 163909 (0.5, 1, or 1.5 mg/kg, i.p.) on the test day for reinstatement of extinguished cocaine-seeking behavior. Reinstatement was initiated by a single non-contingent presentation of cocaine-paired cues (pump and stimulus light). Each active lever press resulted in the presentation of the conditioned stimuli in the absence of cocaine delivery. * $p < 0.05$ vs. Extinction; ^ $p < 0.05$ vs. Veh.




Fig. 4. WAY 163909 dose-dependently suppresses sucrose self-administration. [A] Mean responses (±SEM) on the active (black circles) or inactive lever (white circles), and total number of sucrose pellets (±SEM; gray circles) are presented for the acquisition phase of cocaine self-administration training under FR1–FR5 schedules of reinforcement. Active lever responses resulted in the delivery of a single sucrose pellet (45 mg) and simultaneous presentation of the sucrose-paired cues. [B] Mean responses (±SEM) on the active (black bars) or inactive lever (white bars) or [C] mean total sucrose pellets (±SEM; gray bars) following pretreatment with vehicle (saline; Veh) or WAY 163909 (0.5, 1 or 2 mg/kg, i.p.) (left panel) or following pretreatment with the 5-HT_{2C}R antagonist SB 242084 (0.5 mg/kg, i.p.) plus vehicle (saline; Veh) or WAY 163909 (2 mg/kg, i.p.) (right panel) during the sucrose self-administration maintenance phase. Active lever responses resulted in the delivery of a single sucrose pellet (45 mg) according to an FR5 schedule of reinforcement and simultaneous presentation of the sucrose-paired cues. * $p < 0.05$ vs. Veh; ^ $p < 0.05$ vs. WAY 163909 (2 mg/kg).

3.5. WAY 163909 does not alter sucrose cue-evoked reinstatement

Rats ($n = 25$) readily acquired sucrose self-administration (45 mg pellet) in daily 180 min sessions to stability (i.e., minimum 20 pellets/session on an FR 5 schedule for at least three sessions) and displayed $<10\%$ variation in the number of pellets received (i.e., sucrose intake) (Fig. 5A). Across the last three sessions of stable self-administration on an FR 5 schedule, there was no main effect of session on the number of active [$F_{(2,72)} = 0.54$, ns] or inactive lever responding [$F_{(2,72)} = 0.42$, ns] or the number of pellets received [$F_{(2,72)} = 0.32$, ns]. The rats were then separated into three groups for reinstatement sessions. Sucrose intake did not differ significantly across groups of rats assigned to receive pretreatment with vehicle (26.4 ± 5.2 pellets/session) or WAY 163909 at doses of 0.5 mg/kg (24.2 ± 3.9 pellets/session), or 1.5 mg/kg (22.6 ± 3.9 pellets/session) [$F_{(2, 22)} = 0.34$; ns].

A decrease in presses on the previously-active lever was observed across extinction sessions with all rats achieving the extinction criterion (<15 active lever responses/session for three consecutive sessions) by session 34 of extinction training (data not shown). A repeated-measures ANOVA revealed a main effect of




Fig. 5. WAY 163909 does not alter sucrose cue-evoked reinstatement. [A] Mean responses (±SEM) on the active (black circles) or inactive lever (white circles), and the total number of sucrose pellets (±SEM; gray circles) are presented for the acquisition phase of sucrose self-administration under FR1–FR5 schedules of reinforcement. Active lever responses resulted in the delivery of a single sucrose pellet (45 mg) and simultaneous presentation of the sucrose-paired cues. [B] Mean (±SEM) total number of responses on the active (filled bars) or inactive levers (open bars) following extinction (“Extinction Baseline”) and upon pretreatment with vehicle (saline; Veh) or WAY 163909 (0.5 or 1.5 mg/kg, i.p.) on the test day for reinstatement of extinguished sucrose-seeking behavior. Reinstatement was initiated by a single non-contingent presentation of sucrose-paired cues (pump and stimulus light). Each active lever press resulted in the presentation of the conditioned stimuli in the absence of sucrose pellet delivery. * $p < 0.05$ vs. Extinction.

extinction session on active [$F_{(33,311)} = 19.19$; $p < 0.05$] and inactive lever presses [$F_{(33,311)} = 3.99$; $p < 0.05$] across the extinction sessions for all rats (data not shown). The “extinction baseline” was calculated as the mean total lever presses of all rats on the active (7.9 ± 0.9) or inactive lever (5 ± 1.4) during the final 60 min extinction session (Fig. 5B).

Fig. 5B illustrates the effects of WAY 163909 (0.5 and 1.5 mg/kg) pretreatment on reinstatement of responding supported by cues previously associated with sucrose availability. Sucrose-trained rats readily reinstated active lever pressing across the 60-min reinstatement session (Fig. 5B). A main effect of pretreatment on active lever responding was observed across the 60-min reinstatement session [$F_{(3,45)} = 11.83$; $p < 0.001$; Fig. 5B]. *A priori* comparisons failed to reveal any significant differences in active lever presses relative to vehicle after pretreatment with any doses of WAY 163909 tested (ns; Fig. 5B). The one-way ANOVA revealed no main effect of pretreatment on inactive lever responding [$F_{(3,46)} = 2.61$; ns; Fig. 5B]. The latency to the first lever press following pretreatment with WAY 163909 was unaltered [$F_{(2,22)} = 0.82$; ns] (data not shown).

4. Discussion

Our results demonstrate that the highly-selective and efficacious 5-HT_{2C}R agonist WAY 163909 dose-dependently reduced the reinforcing efficacy of cocaine and sucrose as assessed in parallel self-administration tasks conducted in freely-fed rats. We also show for the first time that WAY 163909 dose-dependently reduced reinstatement elicited by exposure to cocaine-associated contextual cues, but failed to alter reinstatement of cue-evoked sucrose-seeking in freely-fed rats. Thus, selective stimulation of the 5-HT_{2C}R signaling decreases the reinforcing efficacy of cocaine and sucrose, but differentially alters the incentive-salience value of cocaine- vs. sucrose-associated cues.

The dose of WAY 163909 predicted to decrease the reinforcing efficacy of cocaine ($ID_{50} = 1.19$ mg/kg) by 50% was ~5-fold and ~2-fold lower than that predicted to suppress horizontal ambulation ($ID_{50} = 5.89$ mg/kg) and vertical activity ($ID_{50} = 2.3$ mg/kg), respectively (Table 2). Although 5-HT_{2C}R agonists are reported to suppress both horizontal and vertical activity measures (Halford et al., 1997; Halberstadt et al., 2009), most pharmacological studies of 5-HT_{2C}R agonists (e.g., Grottick et al., 2000; Fletcher et al., 2002) typically report horizontal activity as a comparative, independent measure in pharmacological assessments. Horizontal ambulation and vertical activity often correlate, however, these measures have been behaviorally (Walsh and Cummins, 1976; Lever et al., 2006) and pharmacologically dissociated (e.g., Wu et al., 2005) and appear to be mediated by different brain regions (Jackson et al., 1975; Lever et al., 2006). In general, vertical activity is more sensitive to pharmacological manipulation than is horizontal activity (Prut and Belzung, 2003) and, consistent with the current study, Brookshire and Jones (2009) demonstrated that the IC_{50} for the preferential 5-HT_{2C}R

agonist MK 212 to suppress vertical activity ($IC_{50} = \sim 0.93$ mg/kg) was lower than that predicted to suppress horizontal activity ($IC_{50} = \sim 2.29$ mg/kg). In a review of the literature, we could not identify previous publications in which doses of a selective 5-HT_{2C}R agonist were assessed in parallel for efficacy to suppress both horizontal and vertical activity as well as operant responding for an abused drug and palatable reinforcer in freely-fed rats. We uniquely present these observations that, coupled with the failure of WAY 163909 to significantly alter inactive lever presses at any dose, rule out the interpretation that the inhibitory effects of WAY 163909 on drug-seeking were due to a generalized suppression of behavior. Thus, WAY 163909 presents an efficacy window between its effects on cocaine-taking and effects on locomotor activity which may reflect its greater selectivity as a 5-HT_{2C}R agonist relative to 5-HT_{2A}R and 5-HT_{2B}R (see below). Furthermore, these data illustrate that the effects of MK 212 and Ro 60-0175 (Table 1) which have been profiled to suppress cocaine-taking are consistent with their actions preferentially as 5-HT_{2C}R agonists.

A primary therapeutic indication for selective 5-HT_{2C}R agonists is obesity and 5-HT_{2C}R agonists have been shown to suppress food intake and reduce weight gain in animals (Clifton et al., 2000; Dunlop et al., 2005; Fletcher et al., 2010; Hayashi et al., 2005; Hewitt et al., 2002) and most recently in humans (Smith et al., 2009, 2010). We observed a potent WAY 163909-induced suppression of sucrose intake in freely-fed rats. The ID_{50} for WAY 163909 (0.7 mg/kg) in the present study is much lower than the effective dose reported to suppress feeding on powdered chow in fasted Sprague–Dawley rats ($ID_{50} = 2.93$ mg/kg) and rats with diet-induced obesity ($ID_{50} = 5.19$ mg/kg), but closer to the effective doses in obese Zucker rats ($ID_{50} = 1.4$ mg/kg) (Dunlop et al., 2005). Similarly, Ro 60-0175 suppressed sucrose intake at doses much lower than those observed to suppress feeding in fasted rats (Hayashi et al., 2005). Skinnerian tasks such as those employed here rely on the reinforcer (i.e., sucrose) to maintain operant responding and do not simply assay food consumption. In the present study, the motivation to lever press for sucrose is based upon the efficacy and palatability of this reinforcer and the motivational state of rats that have free access to food. Previous studies suggest that preferential 5-HT_{2C}R agonists reduce food intake by enhancing satiety (Hewitt et al., 2002; Somerville et al., 2007), an effect which may account for our observations, particularly in freely-fed rats in an operant task. Stimulation of the 5-HT_{2C}R could also reduce the reinforcing efficacy of sucrose and thus the incentive-salience value of sucrose. Most likely, stimulation of the 5-HT_{2C}R engages neural processes which control both satiety and reinforcing efficacy (Fletcher et al., 2010). Future studies are required to understand how the 5-HT_{2C}R signaling integrates neural process involved in motivation, satiety, palatability and reinforcing efficacy of food.

Associations between environmental (contextual) cues and consumption of rewarding stimuli (food or drug) become encoded memories and it is theorized that such cues can play an important role in relapse to drug use (Lu et al., 2005), and possibly also in non-homeostatic eating, i.e., food intake not driven by energy deficits (Berthoud, 2004). In the present study, WAY 163909 potentially suppressed reinstatement of cocaine-seeking with an 11-fold lower ID_{50} dose (0.5 mg/kg) relative to the ID_{50} dose (5.89 mg/kg) for suppression of horizontal ambulation and 5-fold lower ID_{50} dose relative to the ID_{50} (2.3 mg/kg) for suppression of vertical activity. Interestingly, neither WAY 163909 (present study) nor Ro 60-1075 (Burbassi and Cervo, 2008) inhibited sucrose-seeking during reinstatement. However, a 5-HT_{2C}R agonist was shown to dose-dependently suppress cue-evoked food seeking in food-restricted mice (Somerville et al., 2007). The pharmacological suppression of drug-seeking during reinstatement is most often interpreted as

Table 2

Efficacy of WAY 163909 on locomotor activity, cocaine and sucrose self-administration, and reinstatement of cocaine and sucrose seeking.

Suppression of Horizontal Ambulation (ID_{50})	Suppression of Vertical Activity (ID_{50})	Suppression of Cocaine (ID_{50})		Suppression of Sucrose (ID_{50})	
		Taking	Seeking	Taking	Seeking
5.89 mg/kg	2.3 mg/kg	1.19 mg/kg	0.5 mg/kg	0.7 mg/kg	NE

The dose predicted to decrease basal horizontal ambulation, vertical activity and both cocaine and sucrose-taking and -seeking by 50% (ID_{50}) was estimated using the four parameter logistic nonlinear regression model equation (see Methods). NE, no effect.

reflecting decrements in the incentive-motivational value of the cue-elicited cocaine-seeking behavior (Fletcher et al., 2008; Nic Dhonnchadha et al., 2009). Assessment and interpretation of the incentive-motivational value of non-drug-associated cues may be dependent on the palatability of the reinforcer (i.e., sucrose vs. food) as well as motivational state of rats (i.e., freely-fed vs. food-restricted) and additional studies are needed to elucidate this complex relationship. This would suggest that while selective serotonin 5-HT_{2C}R activation suppresses the reinforcing efficacy of cocaine and sucrose, this signaling pathway is differentially engaged in the control of the incentive-motivational value of cocaine- and sucrose-associated cues.

The attenuation of cocaine- and sucrose-taking and cocaine-seeking behavior by WAY 163909 is attributable to a selective stimulation of the 5-HT_{2C}R based upon several lines of evidence. WAY 163909 is fully efficacious as a 5-HT_{2C}R agonist, does not stimulate the 5-HT_{2A}R, and is only a weak 5-HT_{2B}R partial agonist at much greater concentrations than those that activate the 5-HT_{2C}R (Dunlop et al., 2005) (Table 1). A low dose of the selective 5-HT_{2C}R antagonist SB 242084 completely reversed the inhibitory effects of WAY 163909 on cocaine- and sucrose-taking providing empirical evidence of behaviorally-relevant actions at the 5-HT_{2C}R. Rosenzweig-Lipson and colleagues have demonstrated the suppressive potency of WAY 163909 and the ability of SB 242084 to block WAY 163909-generated behavioral effects in several assay systems, including those modeling the psychiatric disorders of anxiety, depression and schizophrenia (Rosenzweig-Lipson et al., 2007b; Navarra et al., 2008; Grauer et al., 2009; Dunlop et al., 2006). In addition, WAY 163909 blocked cocaine-evoked hyperlocomotion as well as the associated enhancement of dopamine efflux in the nucleus accumbens (Hughes et al., 2006). Interestingly, the high potency of WAY 163909 (0.5–2 mg/kg) to suppress cocaine-taking and -seeking, as well as sucrose-taking, is similar to that seen to suppress impulsivity assessed in a five-choice serial reaction time task (Navarra et al., 2008) as well as aggressive behavior in rodents (Rosenzweig-Lipson et al., 2007b). Cocaine use is associated with increased impulsivity in animals (Anastasio et al., 2011; Stoffel and Cunningham, 2008) and cocaine-dependent subjects often present with impulsivity (Moeller et al., 2001, 2002, 2004; Kjome et al., 2010) and there is recent evidence to suggest that impulsivity contributes to the development and maintenance of obesity and binge eating disorder (Mobbs et al., 2010). Thus, selective, high efficacy 5-HT_{2C}R agonists may prove therapeutically useful in disorders characterized by impulsive traits.

The present observations support the assertion that selective activation of the 5-HT_{2C}R signaling may prove therapeutically useful to reduce craving for cocaine and extend abstinence (Bubar and Cunningham, 2008). However, the progress of selective 5-HT_{2C}R agonists from preclinical evidence of efficacy to verification of safety and efficacy as psychotherapeutic medications in humans has been slow. Clinical trials evaluating the first selective 5-HT_{2C}R agonist, lorcaserin (APD 356, Arena Pharmaceuticals), for weight reduction were successfully completed (Smith et al., 2009, 2010). Lorcaserin, which exhibits ~15-fold and ~100-fold greater potency to stimulate the 5-HT_{2C}R over the homologous 5-HT_{2A}R and 5-HT_{2B}R, respectively, effectively suppressed food intake, reduced weight gain, and was well tolerated by obese subjects (Smith et al., 2009, 2010). Further, when evaluated in polydrug users, the doses shown to be therapeutic in obesity (Smith et al., 2009, 2010) were well-tolerated, exhibited low abuse liability, and were not associated with notable neurocognitive or perceptual effects (Schram et al., 2009). The selective 5-HT_{2C}R agonist vabicaserin has been evaluated in clinical trials to establish its efficacy and safety in subjects with an acute exacerbation of schizophrenia. The

development of two selective 5-HT_{2C}R agonists which advanced to clinical trials opens the door to implementation of clinical research to evaluate these compounds as innovative pro-abstinence, anti-relapse therapeutics for stimulant addiction, eating disorders and obesity.

In conclusion, the 5-HT_{2C}R is a functionally important regulator of the neural substrates that control the rewarding efficacy of a drug (cocaine) and natural (sucrose) reinforcer as well as the incentive-salience value of cocaine-associated cues. WAY 163909 is a potent, selective and efficacious 5-HT_{2C}R agonist which presents a favorable efficacy index between doses useful in inhibiting self-administration/reinstatement and doses which suppress locomotor activity. This compound will prove useful to further tease apart the precise involvement of 5-HT_{2C}R neurocircuitry in reward and motivation as well as the learning and memory processes that carry the encoding for associations between environmental (contextual) cues and consumption of rewarding stimuli (food or drug). A more complete preclinical evaluation of these questions will ultimately allow educated proof-of-concept trials to test the concept that selective 5-HT_{2C}R agonists may be useful as adjunctive therapy in such chronic disorders marked by impulsive drug and/or food intake.

Acknowledgments

This research was supported by the National Institute on Drug Abuse grants P20 DA024157 (KAC, SRG, FGM), R01 DA006511 (KAC), K02 DA000403 (FGM), K05 DA020087 (KAC), and the Jeanne B. Kempner Postdoctoral Scholar Award (NCA).

References

- Anastasio, N.C., Stoffel, E.C., Fox, R.G., Bubar, M.J., Rice, K.C., Moeller, F.G., Cunningham, K.A., 2011. The serotonin 5-HT_{2A} receptor: association with inherent and cocaine-evoked behavioral disinhibition in rats. *Behav. Pharmacol.* 22, 248–261.
- Berthoud, H.R., 2004. Neural control of appetite: cross-talk between homeostatic and non-homeostatic systems. *Appetite* 43, 315–317.
- Blundell, J.E., Tombros, E., Rogers, P.J., Latham, C.J., 1980. Behavioural analysis of feeding: implications for the pharmacological manipulation of food intake in animals and man. *Prog. Neuropsychopharmacol.* 4, 319–326.
- Bonhaus, D.W., Weinhardt, K.K., Taylor, M., DeSouza, A., Mcneely, P.M., Szczepanski, K., Fontana, D.J., Trinh, J., Rocha, C.L., Dawson, M.W., Flippin, L.A., Eglen, R.M., 1997. RS-102221: a novel high affinity and selective, 5-HT_{2C} receptor antagonist. *Neuropharmacology* 36, 621–629.
- Brookshire, B.R., Jones, S.R., 2009. Direct and indirect 5-HT receptor agonists produce gender-specific effects on locomotor and vertical activities in C57 BL/6J mice. *Pharmacol. Biochem. Behav.* 94, 194–203.
- Bubar, M.J., Cunningham, K.A., 2008. Prospects for serotonin 5-HT_{2R} pharmacotherapy in psychostimulant abuse. *Prog. Brain Res.* 172, 319–346.
- Burbassi, S., Cervo, L., 2008. Stimulation of serotonin(2C) receptors influences cocaine-seeking behavior in response to drug-associated stimuli in rats. *Psychopharmacology (Berl)* 196, 15–27.
- Carroll, M.E., Lac, S.T., 1997. Acquisition of IV amphetamine and cocaine self-administration in rats as a function of dose. *Psychopharmacology* 129, 206–214.
- Cifani, C., Zanoncelli, A., Tessari, M., Righetti, C., Di, F.C., Ciccocioppo, R., Massi, M., Melotto, S., 2009. Pre-exposure to environmental cues predictive of food availability elicits hypothalamic-pituitary-adrenal axis activation and increases operant responding for food in female rats. *Addict. Biol.* 14, 397–407.
- Clifton, P.G., Lee, M.D., Dourish, C.T., 2000. Similarities in the action of Ro 60-0175, a 5-HT_{2C} receptor agonist and D-fenfluramine on feeding patterns in the rat. *Psychopharmacology (Berl)* 152, 256–267.
- Cussac, D., Newman-Tancredi, A., Quentric, Y., Carpentier, N., Poissonnet, G., Parmentier, J.G., Goldstein, S., Millan, M.J., 2002. Characterization of phospholipase C activity at h5-HT_{2C} compared with h5-HT_{2B} receptors: influence of novel ligands upon membrane-bound levels of [3H]phosphatidylinositols. *Naunyn Schmiedeberg's Arch. Pharmacol.* 365, 242–252.
- Dunlop, J., Sabb, A.L., Mazandarani, H., Zhang, J., Kalgaonker, S., Shukhina, E., Sukoff, S., Vogel, R.L., Stack, G., Schechter, L., Harrison, B.L., Rosenzweig-Lipson, S., 2005. WAY-163909 ((7bR,10aR)-1,2,3,4,8,9,10,10a-octahydro-7bH-cyclopenta-[b][1,4]diazepino[6,7,1hi]indole): a novel 5-HT_{2C} receptor selective agonist with anorectic activity. *J. Pharmacol. Exp. Ther.* 313, 862–869.
- Dunlop, J., Marquis, K.L., Lim, H.K., Leung, L., Kao, J., Cheesman, C., Rosenzweig-Lipson, S., 2006. Pharmacological profile of the 5-HT(2C) receptor agonist WAY-163909; therapeutic potential in multiple indications. *CNS Drug Rev.* 12, 167–177.

- Fitzgerald, L.W., Burn, T.C., Brown, B.S., Patterson, J.P., Corjay, M.H., Valentine, P.A., Sun, J.H., Link, J.R., Abbaszade, I., Hollis, J.M., Largent, B.L., Hartig, P.R., Hollis, G.F., Meunier, P.C., Robichaud, A.J., Robertson, D.W., 2000. Possible role of valvular serotonin 5-HT(2B) receptors in the cardiopathy associated with fenfluramine. *Mol. Pharmacol.* 57, 75–81.
- Fletcher, P.J., Grottick, A.J., Higgins, G.A., 2002. Differential effects of the 5-HT_{2A} receptor antagonist M100,907 and the 5-HT_{2C} receptor antagonist SB242,084 on cocaine-induced locomotor activity, cocaine self-administration and cocaine-induced reinstatement of responding. *Neuropsychopharmacol.* 27, 576–586.
- Fletcher, P.J., Rizos, Z., Sinyard, J., Tampakeras, M., Higgins, G.A., 2008. The 5-HT(2C) receptor agonist RO 60-0175 reduces cocaine self-administration and reinstatement induced by the stressor yohimbine and contextual cues. *Neuropsychopharmacol.* 33, 1402–1412.
- Fletcher, P.J., Tampakeras, M., Sinyard, J., Slassi, A., Isaac, M., Higgins, G.A., 2009. Characterizing the effects of 5-HT(2C) receptor ligands on motor activity and feeding behaviour in 5-HT(2C) receptor knockout mice. *Neuropharmacology* 57, 259–267.
- Fletcher, P.J., Sinyard, J., Higgins, G.A., 2010. Genetic and pharmacological evidence that 5-HT_{2C} receptor activation, but not inhibition, affects motivation to feed under a progressive ratio schedule of reinforcement. *Pharmacol. Biochem. Behav.* 97, 170–178.
- Grauer, S.M., Graf, R., Navarra, R., Sung, A., Logue, S.F., Stack, G., Huselton, C., Liu, Z., Comery, T.A., Marquis, K.L., Rosenzweig-Lipson, S., 2009. WAY-163909, a 5-HT_{2C} agonist, enhances the preclinical potency of current antipsychotics. *Psychopharmacology (Berl)* 204, 37–48.
- Grottick, A.J., Fletcher, P.J., Higgins, G.A., 2000. Studies to investigate the role of 5-HT(2C) receptors on cocaine- and food-maintained behavior. *J. Pharmacol. Exp. Ther.* 295, 1183–1191.
- Grottick, A.J., Corrigan, W.A., Higgins, G.A., 2001. Activation of 5-HT_{2C} receptors reduces the locomotor and rewarding effects of nicotine. *Psychopharmacology (Berl)* 157, 292–298.
- Halberstadt, A.L., Iris van, dH., Ruderman, M.A., Risbrough, V.B., Gingrich, J.A., Geyer, M.A., Powell, S.B., 2009. 5-HT(2A) and 5-HT(2C) receptors exert opposing effects on locomotor activity in mice. *Neuropsychopharmacol.* 34, 1958–1967.
- Halford, J.C., Harrold, J.A., 2008. Neuropharmacology of human appetite expression. *Dev. Disabil. Res. Rev.* 14, 158–164.
- Halford, J.C., Lawton, C.L., Blundell, J.E., 1997. The 5-HT₂ receptor agonist MK-212 reduces food intake and increases resting but prevents the behavioural satiety sequence. *Pharmacol. Biochem. Behav.* 56, 41–46.
- Halford, J.C., Boyland, E.J., Blundell, J.E., Kirkham, T.C., Harrold, J.A., 2010. Pharmacological management of appetite expression in obesity. *Nat. Rev. Endocrinol.* 6, 255–269.
- Hayashi, A., Suzuki, M., Sasamata, M., Miyata, K., 2005. Agonist diversity in 5-HT(2C) receptor-mediated weight control in rats. *Psychopharmacology (Berl)* 178, 241–249.
- Hewitt, K.N., Lee, M.D., Dourish, C.T., Clifton, P.G., 2002. Serotonin 2C receptor agonists and the behavioural satiety sequence in mice. *Pharmacol. Biochem. Behav.* 71, 691–700.
- Higgins, G.A., Ouagazzal, A.M., Grottick, A.J., 2001. Influence of the 5-HT(2C) receptor antagonist SB242,084 on behaviour produced by the 5-HT(2) agonist Ro60-0175 and the indirect 5-HT agonist dexfenfluramine. *Br. J. Pharmacol.* 133, 459–466.
- Hughes, Z.A., Mabrouk, O.S., Kandebo, M., Pulicicchio, C., Malberg, J.E., Rosenzweig-Lipson, S., 2006. The 5-HT_{2C} Receptor Agonist Prevents Cocaine-induced Increases in Locomotor Activity and Accumens Dopamine. Program No. 691.9, 2006 Neuroscience Meeting Planner. Society for Neuroscience, Atlanta, GA.
- Jackson, D.M., Anden, N.E., Dahlstrom, A., 1975. A functional effect of dopamine in the nucleus accumbens and in some other dopamine-rich parts of the rat brain. *Psychopharmacologia* 45, 139–149.
- Kelley, A.E., 2004. Memory and addiction: shared neural circuitry and molecular mechanisms. *Neuron* 44, 161–179.
- Kjome, K.L., Lane, S.D., Schmitz, J.M., Green, C., Ma, L., Prasla, I., Swann, A.C., Moeller, F.G., 2010. Relationship between impulsivity and decision making in cocaine dependence. *Psychiatry Res.* 178, 299–304.
- Knight, A.R., Misra, A., Quirk, K., Benwell, K.R., Revell, D.F., Kennett, G.A., Bickerdike, M.J., 2004. Pharmacological characterisation of the agonist radioligand binding site of 5-HT_{2A}, 5-HT_{2B} and 5-HT_{2C} receptors. *Naunyn-Schmiedeberg's Arch. Pharmacol.* 370, 114–123.
- Leri, F., Flores, J., Rodaros, D., Stewart, J., 2002. Blockade of stress-induced but not cocaine-induced reinstatement by infusion of noradrenergic antagonists into the bed nucleus of the stria terminalis or the central nucleus of the amygdala. *J. Neurosci.* 22, 5713–5718.
- Lever, C., Burton, S., O'Keefe, J., 2006. Rearing on hind legs, environmental novelty, and the hippocampal formation. *Rev. Neurosci.* 17, 111–133.
- Lu, L., Hope, B.T., Dempsey, J., Liu, S.Y., Bossert, J.M., Shaham, Y., 2005. Central amygdala ERK signaling pathway is critical to incubation of cocaine craving. *Nat. Neurosci.* 8, 212–219.
- Lyness, W.H., Friedle, N.M., Moore, K.E., 1980. Increased self-administration of D-amphetamine after destruction of 5-hydroxytryptaminergic neurons. *Pharmacol. Biochem. Behav.* 12, 937–941.
- Martin, J.R., Bos, M., Jenck, F., Moreau, J.L., Mutel, V., Sleight, A.J., Wichmann, J., Andrews, J.S., Berendsen, H.H.G., Broekkamp, C.L.E., Ruigt, G.S.F., Kohler, C., Van Delft, A.M.L., 1998. 5-HT_{2C} receptor agonists: pharmacological characteristics and therapeutic potential. *J. Pharmacol. Exp. Ther.* 286, 913–924.
- Mobbs, O., Crepin, C., Thiery, C., Golay, A., Van der, L.M., 2010. Obesity and the four facets of impulsivity. *Patient Educ. Couns.* 79, 372–377.
- Moeller, F.G., Dougherty, D.M., Barratt, E.S., Schmitz, J.M., Swann, A.C., Grabowski, J., 2001. The impact of impulsivity on cocaine use and retention in treatment. *J. Subst. Abuse Treat.* 21, 193–198.
- Moeller, F.G., Dougherty, D.M., Barratt, E.S., Oderinde, V., Mathias, C.W., Harper, R.A., Swann, A.C., 2002. Increased impulsivity in cocaine dependent subjects independent of antisocial personality disorder and aggression. *Drug Alcohol Depend.* 68, 105–111.
- Moeller, F.G., Barratt, E.S., Fischer, C.J., Dougherty, D.M., Reilly, E.L., Mathias, C.W., Swann, A.C., 2004. P300 event-related potential amplitude and impulsivity in cocaine-dependent subjects. *Neuropsychobiology* 50, 167–173.
- Nair, S.G., ms-Deutsch, T., Epstein, D.H., Shaham, Y., 2009. The neuropharmacology of relapse to food seeking: methodology, main findings, and comparison with relapse to drug seeking. *Prog. Neurobiol.* 89, 18–45.
- Navarra, R., Comery, T.A., Graf, R., Rosenzweig-Lipson, S., Day, M., 2008. The 5-HT(2C) receptor agonist WAY-163909 decreases impulsivity in the 5-choice serial reaction time test. *Behav. Brain Res.* 188, 412–415.
- Neisewander, J.L., Acosta, J.L., 2007. Stimulation of 5-HT_{2C} receptors attenuates cue and cocaine-primed reinstatement of cocaine-seeking behavior in rats. *Behav. Pharmacol.* 18, 791–800.
- Nic Dhonnchadha, B.A., Fox, R.G., Stutz, S.J., Rice, K.C., Cunningham, K.A., 2009. Blockade of the serotonin 5-HT_{2A} receptor suppresses cue-evoked reinstatement of cocaine-seeking behavior in a rat self-administration model. *Behav. Neurosci.* 123, 382–396.
- Nichols, D.E., 2004. Hallucinogens. *Pharmacol. Ther.* 101, 131–181.
- Pentkowski, N.S., Duke, F.D., Weber, S.M., Pockros, L.A., Teer, A.P., Hamilton, E.C., Thiel, K.J., Neisewander, J.L., 2010. Stimulation of medial prefrontal cortex serotonin 2C 5-HT_{2C} receptors attenuates cocaine-seeking behavior. *Neuropsychopharmacol.* 35, 2037–2048.
- Porter, R.H., Benwell, K.R., Lamb, H., Malcolm, C.S., Allen, N.H., Revell, D.F., Adams, D.R., Sheardown, M.J., 1999. Functional characterization of agonists at recombinant human 5-HT_{2A}, 5-HT_{2B} and 5-HT_{2C} receptors in CHO-K1 cells. *Br. J. Pharmacol.* 128, 13–20.
- Prut, L., Belzung, C., 2003. The open field as a paradigm to measure the effects of drugs on anxiety-like behaviors: a review. *Eur. J. Pharmacol.* 463, 3–33.
- Ratkowsky, D.A., Reedy, T.J., 1986. Choosing near-linear parameters in the four-parameter logistic model for radioligand and related assays. *Biometrics* 42, 575–582.
- Rocha, B.A., Goulding, E.H., O'Dell, L.E., Mead, A.N., Coufal, N.G., Parsons, L.H., Tecott, L.H., 2002. Enhanced locomotor, reinforcing, and neurochemical effects of cocaine in serotonin 5-hydroxytryptamine 2C receptor mutant mice. *J. Neurosci.* 22, 10039–10045.
- Rosenzweig-Lipson, S., Dunlop, J., Marquis, K.L., 2007a. 5-HT_{2C} receptor agonists as an innovative approach for psychiatric disorders. *Drug News Perspect.* 20, 565–571.
- Rosenzweig-Lipson, S., Sabb, A., Stack, G., Mitchell, P., Lucki, I., Malberg, J.E., Grauer, S., Brennan, J., Cryan, J.F., Sukoff Rizzo, S.J., Dunlop, J., Barrett, J.E., Marquis, K.L., 2007b. Antidepressant-like effects of the novel, selective, 5-HT_{2C} receptor agonist WAY-163909 in rodents. *Psychopharmacology (Berl)* 192, 159–170.
- Roth, B.L., 2007. Drugs and valvular heart disease. *N. Engl. J. Med.* 356, 6–9.
- Schram, M.J., Schoedel, K.A., Barlett, C., Shazer, R.L., Sellers, E.M., 2009. Abuse potential evaluation of lorcaserin, a selective and potent agonist at the serotonin 2C (5-HT_{2C}) receptor, in healthy male and female recreational polydrug users. *Amer. Coll. Neuropsychopharmacol. 48th Annu. Meet.*
- Smith, S.R., Prosser, W.A., Donahue, D.J., Morgan, M.E., Anderson, C.M., Shanahan, W.R., Study Group, 2009. Lorcaserin (APD356), a selective 5-HT(2C) agonist, reduces body weight in obese men and women. *Obes. (Silver Spring)* 17, 494–503.
- Smith, S.R., Weissman, N.J., Anderson, C.M., Sanchez, M., Chuang, E., Stubbe, S., Bays, H., Shanahan, W.R., 2010. Multicenter, placebo-controlled trial of lorcaserin for weight management. *N. Engl. J. Med.* 363, 245–256.
- Somerville, E.M., Horwood, J.M., Lee, M.D., Kennett, G.A., Clifton, P.G., 2007. 5-HT(2C) receptor activation inhibits appetitive and consummatory components of feeding and increases brain c-fos immunoreactivity in mice. *Eur. J. Neurosci.* 25, 3115–3124.
- Steiger, H., 2004. Eating disorders and the serotonin connection: state, trait and developmental effects. *J. Psychiatry Neurosci.* 29, 20–29.
- Stoffel, E.C., Cunningham, K.A., 2008. The relationship between the locomotor response to a novel environment and behavioral disinhibition in rats. *Drug Alcohol Depend.* 92, 69–78.
- Tallarida, R.J., Murray, R.B., 1987. *Manual of Pharmacologic Calculations with Computer Programs*. Springer-Verlag, New York.
- Tecott, L.H., Sun, L.M., Akana, S.F., Strack, A.M., Lowenstein, D.H., Dallman, M.F., Julius, D., 1995. Eating disorder and epilepsy in mice lacking 5-HT_{2C} serotonin receptors [see comments]. *Nature* 374, 542–546.
- Thomsen, W.J., Grottick, A.J., Menzaghi, F., Reyes-Saldana, H., Espitia, S., Yuskin, D., Whelan, K., Martin, M., Morgan, M., Chen, W., Al-Shamma, H., Smith, B., Chalmers, D., Behan, D., 2008. Lorcaserin, a novel selective human 5-hydroxytryptamine_{2C} agonist: in vitro and in vivo pharmacological characterization. *J. Pharmacol. Exp. Ther.* 325, 577–587.
- Tomkins, D.M., Joharchi, N., Tampakeras, M., Martin, J.R., Wichmann, J., Higgins, G.A., 2002. An investigation of the role of 5-HT(2C) receptors in modifying ethanol self-administration behaviour. *Pharmacol. Biochem. Behav.* 71, 735–744.

- Tong, Z., Chandrasekaran, A., Demaio, W., Espina, R., Lu, W., Jordan, R., Scatina, J., 2010a. Metabolism of vabicaserin in mice, rats, dogs, monkeys, and humans. *Drug Metab. Dispos.* 38, 2266–2277.
- Tong, Z., Chandrasekaran, A., Demaio, W., Jordan, R., Li, H., Moore, R., Poola, N., Burghart, P., Hultin, T., Scatina, J., 2010b. Species differences in the formation of vabicaserin carbamoyl glucuronide. *Drug Metab. Dispos.* 38, 581–590.
- Tran-Nguyen, L.T., Bellew, J.G., Grote, K.A., Neisewander, J.L., 2001. Serotonin depletion attenuates cocaine seeking but enhances sucrose seeking and the effects of cocaine priming on reinstatement of cocaine seeking in rats. *Psychopharmacology (Berl)* 157, 340–348.
- Vickers, S.P., Clifton, P.G., Dourish, C.T., Tecott, L.H., 1999. Reduced satiating effect of *d*-fenfluramine in serotonin 5-HT_{2C} receptor mutant mice. *Psychopharmacology* 143, 309–314.
- Walsh, R.N., Cummins, R.A., 1976. The open-field test: a critical review. *Psychol. Bull.* 83, 482–504.
- Wogar, M.A., Bradshaw, C.M., Szabadi, E., 1991. Evidence for an involvement of 5-hydroxytryptaminergic neurones in the maintenance of operant behaviour by positive reinforcement. *Psychopharmacology* 105, 119–124.
- Wu, J., Zou, H., Strong, J.A., Yu, J., Zhou, X., Xie, Q., Zhao, G., Jin, M., Yu, L., 2005. Bimodal effects of MK-801 on locomotion and stereotypy in C57BL/6 mice. *Psychopharmacology* 177, 256–263.