

# $\alpha$ 5 Subunit-containing GABA<sub>A</sub> receptors mediate a slowly decaying inhibitory synaptic current in CA1 pyramidal neurons following Schaffer collateral activation

Mariana Vargas-Caballero<sup>a,\*</sup>, Loren J. Martin<sup>b</sup>, Michael W. Salter<sup>c,d,e</sup>,  
Beverly A. Orser<sup>b,c,f,g</sup>, Ole Paulsen<sup>a</sup>

<sup>a</sup> Department of Physiology, Anatomy and Genetics, University of Oxford, Sherrington Building, Parks Road, Oxford OX1 3PT, United Kingdom

<sup>b</sup> Institute of Medical Science, University of Toronto, Medical Sciences Building University of Toronto, Toronto, M5S 1A8, Canada

<sup>c</sup> Department of Physiology, University of Toronto, Medical Sciences Building University of Toronto, Toronto, M5S 1A8, Canada

<sup>d</sup> Program in Neurosciences & Mental Health, Hospital for Sick Children, 555 University Avenue, Toronto, Ontario M5G 1X8, Canada

<sup>e</sup> University of Toronto Centre for the Study of Pain, University of Toronto, 124 Edward Street Room 374, Toronto, Ontario M5G 1G6, Canada

<sup>f</sup> Department of Anesthesia, University of Toronto, Canada, 150 College Street Toronto, Ontario M5S 3E2, Canada

<sup>g</sup> Department of Anesthesia, Sunnybrook Health Science Centre, 2075 Bayview Avenue, Toronto, Ontario M4N 3M5, Canada

## ARTICLE INFO

### Article history:

Received 6 October 2009

Received in revised form

13 November 2009

Accepted 16 November 2009

### Keywords:

$\alpha$ 5  
GABA<sub>A</sub> receptor  
Hippocampus  
Inhibition  
Mouse  
Rat  
Schaffer collateral

## ABSTRACT

GABA<sub>A</sub> receptors that contain the  $\alpha$ 5 subunit ( $\alpha$ 5GABA<sub>A</sub>Rs) are highly expressed in the hippocampus, and have been implicated in learning and memory processes. They generate a tonic form of inhibition that regulates neuronal excitability. Recently it was shown that  $\alpha$ 5GABA<sub>A</sub>Rs also contribute to slow phasic inhibition of CA1 pyramidal neurons following local stimulation in the *stratum lacunosum moleculare*. However, it is unknown whether  $\alpha$ 5GABA<sub>A</sub>Rs can also be recruited indirectly by stimulation of Schaffer collaterals. Here, we studied GABAergic currents evoked by stimulation in the *stratum radiatum* of CA1 in the presence and absence of CNQX to block AMPA receptor-mediated excitation. We tested their sensitivity to gabazine and two drugs acting at the benzodiazepine site of  $\alpha$ 1/ $\alpha$ 2/ $\alpha$ 3 or  $\alpha$ 5GABA<sub>A</sub>Rs (400 nM zolpidem and 20 nM L-655,708, respectively). IPSCs evoked by stimulation in the *stratum radiatum* in the presence of CNQX were potentiated by zolpidem, blocked by 1  $\mu$ M gabazine and were relatively insensitive to L-655,708 consistent with the lack of  $\alpha$ 5GABA<sub>A</sub>Rs. In contrast, IPSCs evoked by stimulation of Schaffer collaterals had a significant gabazine-insensitive component. This component was attenuated by L-655,708 and enhanced by burst stimulation. Furthermore, the L-655,708-sensitive current was absent in recordings from mice lacking  $\alpha$ 5GABA<sub>A</sub>Rs (*gabra5*<sup>−/−</sup> mice). These results show that  $\alpha$ 5GABA<sub>A</sub>R-mediated phasic inhibition is activated by the Schaffer collateral pathway and provide evidence for activity pattern-dependent participation of  $\alpha$ 5GABA<sub>A</sub>Rs in inhibition.

© 2009 Elsevier Ltd. All rights reserved.

## 1. Introduction

Synaptic inhibition in the hippocampus plays a crucial role in balancing and synchronising the activity of excitatory cells.  $\gamma$ -aminobutyric acid (GABA) released by inhibitory interneurons activates GABA<sub>A</sub> receptors (GABA<sub>A</sub>Rs), and in most mature neurons, GABA causes a reduction of the postsynaptic cell excitability via hyperpolarising and/or shunting inhibition (for review, see Mann and Paulsen, 2007). GABA<sub>A</sub>Rs are Cl<sup>−</sup> permeable, pentameric ionic channels that are formed from the combination of distinct subunits

( $\alpha$ 1–6,  $\beta$ 1–3,  $\gamma$ 1–3,  $\delta$ ,  $\epsilon$ ,  $\theta$ ,  $\pi$ ,  $\rho$ 1–3), and the majority of native combinations identified to date have a common 2:2:1  $\alpha/\beta/\gamma$  stoichiometry (reviewed in Wafford, 2005).

The targeting of pyramidal cells by inhibitory interneurons follows a highly organised pattern, and the vast majority of GABAergic interneurons target either the perisomatic or specific dendritic domains of pyramidal cells (Klausberger and Somogyi, 2008). However, the role of specific GABA<sub>A</sub>R subtypes expressed in distinct CA1 pyramidal cell compartments is still poorly understood. There is some evidence of a high correlation between presynaptic interneuron type and their specific GABA<sub>A</sub>Rs subunit targets (Nusser et al., 1996; Thomson et al., 2000). For example, in the neocortex, recordings from synaptically-connected pairs between GABAergic interneurons and pyramidal cells have demonstrated that dendritic targeting inhibitory neurons preferentially activate  $\alpha$ 5GABA<sub>A</sub>Rs,

\* Corresponding author. Tel.: +44 (0) 1865 282 160, +44 (0) 7975 623 420 (mobile); fax: +44 1865 272 469.

E-mail address: [mariana.vargas@dpg.ox.ac.uk](mailto:mariana.vargas@dpg.ox.ac.uk) (M. Vargas-Caballero).

whereas those targeting the soma activate  $\alpha 1$ GABA<sub>A</sub>Rs (Ali and Thomson, 2008).

By identifying the GABA<sub>A</sub>R subtypes in different inhibitory pathways, it may be possible to pharmacologically target specific GABAergic networks in the hippocampus. Recent studies have started to dissect the contribution of GABA<sub>A</sub>Rs subtypes to different behaviours. Pharmacological tools in this area include the use of benzodiazepine derivatives or genetic modifications targeted at the benzodiazepine site, which is located at the interface of  $\gamma 2$  and  $\alpha$  subunits (reviewed in Wafford, 2005). By altering the kinetics of single GABA<sub>A</sub>R channels, benzodiazepines enhance the effect of GABA and a behavioural readout can be obtained to interpret the function of targeted  $\alpha$  subunits. For example, mice with a point mutation in the  $\alpha 1$ GABA<sub>A</sub>R subunit (His101Arg), which rendered it insensitive to diazepam, did not display the sedative and amnesic effects of benzodiazepines (McKernan et al., 2000; Rudolph et al., 1999). Conversely, inverse agonists acting at the benzodiazepine site (Tenen and Hirsch, 1980) inhibit the effect of GABA, and have effects opposite to those of the classical benzodiazepines. Using this approach, systemic application of the  $\alpha 3$  inverse-agonist  $\alpha 3$ IA promoted anxiety-related behaviours in rodents (Atack et al., 2005).

$\alpha 5$ GABA<sub>A</sub>Rs are of particular interest as they are highly expressed in the adult hippocampus both at synaptic and extrasynaptic sites (Sperk et al., 1997; Sur et al., 1999), in stark contrast to low expression levels in other brain areas. Consistent with their hippocampal localisation, behavioural studies using  $\alpha 5$  subunit-specific inverse agonists and  $\alpha 5$  subunit knock out mice strongly implicated  $\alpha 5$ GABA<sub>A</sub>Rs in the modulation of learning and memory (Collinson et al., 2002; Atack et al., 2006; Ballard et al., 2009). Therefore,  $\alpha 5$ GABA<sub>A</sub>Rs are currently considered as relevant targets for memory blocking drugs (Martin et al., 2009) and cognitive enhancing drugs with clinical applications such as in Alzheimer's disease patients, whose  $\alpha 5$ GABA<sub>A</sub>Rs are well preserved (Howell et al., 2000). However, the precise mechanisms underlying the regulation of hippocampal function by  $\alpha 5$ GABA<sub>A</sub>Rs are not known.

It is well established that extrasynaptic  $\alpha 5$ GABA<sub>A</sub>Rs can mediate a large component of tonic inhibition in the hippocampus (Caraiscos et al., 2004; Scimemi et al., 2005; Glykys and Mody, 2006; Prenosil et al., 2006). In contrast, the role of  $\alpha 5$ GABA<sub>A</sub>Rs in phasic inhibition remains poorly understood. Studies comparing spontaneous and locally-evoked inhibition between mice lacking  $\alpha 5$ GABA<sub>A</sub>Rs (*gabra5*<sup>−/−</sup>) and wild type (WT) mice suggested a negligible contribution of  $\alpha 5$ GABA<sub>A</sub>Rs to phasic inhibition (Collinson et al., 2002; Glykys and Mody, 2006). Other studies have suggested that  $\alpha 5$ GABA<sub>A</sub>Rs mediate a slowly decaying component of synaptic inhibition (GABA<sub>A,slow</sub>) (Prenosil et al., 2006; Zarnowska et al., 2009). Evoked GABA<sub>A,slow</sub> potentials have only been observed following local extracellular stimulation at or near the *stratum lacunosum moleculare* (SLM) of the hippocampus (Pearce, 1993; Ouardouz and Lacaille, 1997; Zarnowska et al., 2009). Thus, GABA<sub>A,slow</sub> has been proposed to modulate the activity of distal dendrites in hippocampal CA1 pyramidal neurons, and to mediate a component of synaptic inhibition activated by the direct input from the entorhinal cortex to the hippocampus at the SLM (Banks et al., 2000).

In order to understand the underlying mechanisms of  $\alpha 5$ GABA<sub>A</sub>Rs targeting cognitive enhancing drugs, it becomes important to establish whether in addition to their SLM activation,  $\alpha 5$ GABA<sub>A</sub>Rs are recruited by CA3 output via Schaffer collateral activity. Local stimulation at the *stratum radiatum* (SR) in CA1 has been reported to produce fast decaying IPSCs via perisomatic targeting inhibitory cells (Ouwardouz and Lacaille, 1997) mediated by  $\alpha 1/\alpha 2/\alpha 3$ GABA<sub>A</sub>Rs (Thomson et al., 2000). However, under conditions of local stimulation, excitatory synaptic transmission is usually blocked with glutamate receptor antagonists. Feed-forward inhibition requires activation of afferent fibres to interneurons

which in turn release GABA onto pyramidal cells (Alger and Nicoll, 1982). The Schaffer collaterals are likely to stimulate directly or indirectly a wide variety of interneurons that would not be reached by local stimulation. In the present study, we compared locally-evoked and Schaffer collateral-stimulated inhibitory currents. To determine whether  $\alpha 5$ GABA<sub>A</sub>Rs contribute to the evoked IPSCs we used the inverse-agonist L-655,708 in rats and *gabra5*<sup>−/−</sup> mice. The results show that stimulation of Schaffer collaterals can activate a slowly decaying component of GABAergic inhibition, mediated by  $\alpha 5$  subunit-containing GABA<sub>A</sub> receptors, particularly following bursts of high-frequency stimulation of Schaffer collateral afferent input.

## 2. Methods

Animals were housed in groups with access to food and water *ad libitum*. The holding facilities maintained a temperature of approximately 22 °C, humidity of 60–70%, and a 12-h light/dark cycle. All animal care and experimental procedures were in accordance with the UK Home Office regulations under the Animals (Scientific Procedures) Act of 1986, and the Animal Care Committee of the University of Toronto.

### 2.1. Tissue preparation

Parasagittal slices containing the hippocampus were obtained from male Sprague Dawley rats (supplied by Harlan, Bicester, UK), or from *gabra5*<sup>−/−</sup> mice (Collinson et al., 2002) and wild type (WT) littermates ranging from postnatal day 14 to 28. Rodents were anaesthetized with 5% isoflurane until breathing slowed down to approximately one breath per second, and stimulation of the limb withdrawal reflex no longer elicited a response. After decapitation, the brain was quickly removed into ice-cold artificial cerebrospinal fluid (aCSF), containing (in mM): NaCl, 126; KCl, 2.5; NaHCO<sub>3</sub>, 26; CaCl<sub>2</sub>, 2; MgCl<sub>2</sub>, 2; NaH<sub>2</sub>PO<sub>4</sub>, 1.25; glucose, 10, saturated with 95% O<sub>2</sub>/5% CO<sub>2</sub>, with a final pH of 7.2–7.4. Slices were prepared at 350  $\mu$ m thickness using a Leica VT1000S microtome. Slices containing the hippocampal formation were trimmed from other brain regions and were maintained and recorded at room temperature (22–27 °C).

### 2.2. Electrophysiological recordings

After transferring a single slice to a submerged-style recording chamber, a monopolar stainless steel stimulation electrode (A-M Systems, Sequim, WA, USA) was placed into the SR of CA1 50–100  $\mu$ m away from the *stratum pyramidale* (SP) for synaptic stimulation. Stimulation in the SR was carried out under two different conditions: firstly, to record GABA<sub>A,local</sub> AMPA receptor-mediated excitation was blocked with CNQX while recording from a pyramidal cell. The stimulation electrode was placed approximately 100  $\mu$ m lateral to the recorded cell to ensure stimulation of local interneurons. Secondly, to record inhibition elicited by the Schaffer collaterals (GABA<sub>A,SC</sub>) CNQX was not included. The stimulation electrode was placed approximately 300  $\mu$ m lateral to the recorded cell to reduce local stimulation of GABAergic neurons in addition to afferent stimulation.

Experiments were performed in voltage clamp mode. The intracellular solution contained (in mM): Gluconic acid 70; CsCl 10; NaCl 5; BAPTA free acid 10; Hepes 10; QX-314 10; GTP 0.3; Mg-ATP 4; pH was titrated to 7.25  $\pm$  0.05 with CsOH. The estimated final Cs concentration for the intracellular solution was  $\sim$  120 mM. The final osmolality was 280  $\pm$  5 mOsmol l<sup>−1</sup>. BAPTA was used to prevent Ca<sup>2+</sup> dependent changes while measuring synaptic activity at depolarised membrane potentials. QX-314 blocks GABA<sub>B</sub> receptor-mediated currents in addition to Na<sup>+</sup> channels (Nathan et al., 1990). All voltage values were corrected for the liquid junction potential measured as 13 mV.

Whole-cell patch clamp recordings were obtained with 2–4 M $\Omega$  borosilicate pipettes from putative CA1 pyramidal cells identified by their location in the SP and by their shape.

The calculated E<sub>Cl</sub> at room temperature was −56 mV, and AMPA receptor-mediated currents reversed near 0 mV. For this reason GABAergic currents were recorded at 0 mV, both for local and Schaffer collateral stimulation, so as to isolate them from AMPA receptor-mediated currents in the latter case. For recordings at voltages other than −70 mV, a voltage step from −70 mV to the test potential started 5 s before synaptic stimulation. Whole-cell recordings were made using an Axon Multiclamp 700B amplifier (Molecular Devices, Union City, CA, USA). Recordings were low-pass filtered at 2 kHz and digitised at 20 kHz with a National Instruments A/D board (Austin, TX, USA) using Ginp 1.0 software (courtesy of Hugh P. C. Robinson) for acquisition from within Matlab (Mathworks Ltd, Natick, MA, USA). Postsynaptic currents were evoked using a stimulus isolator unit (ISO-flex, A.M.P.I. Jerusalem, Israel) which delivered pulses of 100  $\mu$ s duration in current mode; stimulation intensities ranged between 20  $\mu$ A and 70  $\mu$ A and the computer-controlled stimulation interval was 60 s. Drugs were applied after a stable baseline of 6–10 min

(<10% drift allowed). Series resistance was not compensated during recordings. Series resistance was measured before each stimulation with a 5 mV, 50 ms step pulse. Recordings were terminated if series resistance ( $16 \pm 6 \text{ M}\Omega$ ) changed by more than 20%.

### 2.3. Data analysis

Data analysis was done using Matlab. Statistical testing was done using Matlab and SPSS software. Charge transfer was calculated by integrating the current responses from 5 to 750 ms following synaptic stimulation after leak subtraction (or 5–35 ms and 50–750 ms to separate early and late components in Fig. 1Ci). For comparison across experiments, synaptic peak current values or total charge transfer were normalised relative to the mean of baseline values obtained 5 min before drug application. For statistical comparison of drug effects, the average value of the last 4 min of recording was used. Data are presented as mean  $\pm$  standard error of the mean (SEM) and are displayed in two-minute bin intervals. *N* values refer to the number of slices recorded. Example traces are the average of 3–5 traces (Gaussian-filtered at a corner frequency of 2 kHz). Statistical significance was assessed using Student's two-sample two-tailed *t*-test, one-way ANOVA or repeated measures (RM) ANOVA, with Bonferroni post-hoc corrections for multiple comparisons where appropriate.  $P < 0.05$  was considered statistically significant.

### 2.4. Drugs


20  $\mu\text{M}$  DL-2-amino-5-phosphonopentanoic acid was used to block *N*-methyl-D-aspartic acid (NMDA) receptors in all experiments. CNQX (6-cyano-7-nitro-quinoline-2,3-dione disodium) was used to block AMPA receptor-mediated synaptic transmission for local stimulation of GABAergic interneurons. Gabazine (SR 95531 hydrobromide; 6-imino-3-(4-methoxyphenyl)-1(6*H*)-pyridazinebutanoic acid hydrobromide) is a competitive antagonist at GABA<sub>A</sub>Rs. L-655,708 (11,12,13,13a-

tetrahydro-7-methoxy-9-oxo-9*H*-imidazo[1,5-*a*]pyrrolo[2,1-*c*][1,4]benzodiazepine-1-carboxylic acid, ethyl ester) is a partial inverse agonist at the benzodiazepine binding site of  $\alpha 5\text{GABA}_A\text{R}$ ; L-655,708 was initially dissolved to 10 mM in 1 N HCl, then diluted to 10  $\mu\text{M}$  by adding  $\text{H}_2\text{O}$ , and stored in frozen aliquots until used. Zolpidem (*N,N*,6-trimethyl-2-(4-methylphenyl)imidazo[1,2-*a*]pyridine-3-acetamide) is a benzodiazepine acting primarily at  $\alpha 1\text{GABA}_A\text{R}$ , and with some affinity for  $\alpha 2/\alpha 3\text{GABA}_A\text{R}$  at 400 nM. Zolpidem was dissolved at 100 mM in ethanol, frozen in aliquots, and diluted in aCSF just before use; equivalent amounts of ethanol were added in corresponding control experiments. All drugs described above were purchased from Tocris (Bristol, UK). Other chemicals were purchased from Sigma Aldrich (St. Louis, MO, USA).

## 3. Results

To measure synaptic GABA<sub>A</sub>R-mediated currents in CA1 pyramidal neurons in hippocampal slices, single cells were voltage clamped and brief extracellular stimuli delivered in the SR. First, to activate local GABAergic currents (referred to as GABA<sub>A,local</sub>), AMPA receptor-mediated excitation was blocked using 10  $\mu\text{M}$  CNQX. GABA<sub>A,local</sub> reversed close to  $E_{\text{Cl}}$  at  $-54 \pm 3 \text{ mV}$  ( $n = 5$ ; data not shown; for example traces see Fig. 1Ai) and was seen as an outward current at 0 mV. At this holding potential, IPSCs had a time constant of decay ( $\tau_{\text{decay}}$ ) of  $163 \pm 16 \text{ ms}$  ( $n = 12$ ).

Next, IPSCs were elicited by stimulating the Schaffer collaterals. For these recordings, CNQX was omitted from the extracellular solution. Biphasic responses comprising both glutamatergic and



**Fig. 1.** Locally-evoked and Schaffer collateral-stimulated GABA<sub>A</sub>R-mediated currents are differentially sensitive to gabazine. (Ai) Current traces showing voltage dependence of GABA<sub>A</sub>R currents evoked by synaptic stimulation in two different recording conditions. *Top traces*: inhibitory currents recorded at 0 mV; GABA<sub>A,local</sub> in the presence of 10  $\mu\text{M}$  CNQX and GABA<sub>A,SC</sub>. *Bottom traces*: Synaptic currents recorded at different holding potentials between  $-80$  and  $+40 \text{ mV}$  in steps of  $20 \text{ mV}$ . (Aii) Expanded traces for Schaffer collateral-stimulated currents (holding potentials of  $-70$ ,  $-50$ ,  $-30$ ,  $-10$ ,  $+10$  and  $+40 \text{ mV}$ ) and current voltage relationship for estimated AMPAR (inverted triangles) and GABA<sub>A</sub>R (squares) mediated components ( $n = 5$ ). Currents were measured at times indicated by vertical lines in example traces. (Bi) Concentration-response plot for GABA<sub>A,local</sub> and GABA<sub>A,SC</sub>. Response is the charge transfer normalised to pre-gabazine control values. GABA<sub>A,SC</sub>(gz) indicates the  $1 \mu\text{M}$  gabazine-resistant component of GABA<sub>A,SC</sub>. *Inset*: Example traces of Schaffer collateral-stimulated currents in  $1 \mu\text{M}$  gabazine with holding potentials as in Ai. (Bii) Superimposed current traces, with and without  $1 \mu\text{M}$  gabazine. Normalised traces for GABA<sub>A,local</sub>, GABA<sub>A,SC</sub>, and GABA<sub>A,SC</sub>(gz). (Ci) Concentration-response plot for the early and late components of GABA<sub>A,SC</sub> elicited with burst stimulation (3 stimuli at  $100 \text{ Hz}$ ). GABA<sub>A,SC</sub>(5–35 ms) and GABA<sub>A,SC</sub>(50–750 ms) were the measured charge transfers in the indicated time windows after stimulation. (Cii) Example traces of GABA<sub>A,SC</sub>(burst) recorded at  $0 \text{ mV}$  with and without gabazine. Normalised traces on the right. Scale bars in Bi-ii and Cii:  $250 \text{ ms}$ ,  $400 \text{ pA}$ .



GABAergic currents were observed (Fig. 1Ai and Aii). At  $-70$  mV, the slowly decaying inhibitory current was fully blocked by  $10 \mu\text{M}$  gabazine leaving only a fast decaying AMPA current blocked by  $10 \mu\text{M}$  CNQX. The excitatory component reversed near  $0$  mV as expected, and the inhibitory component reversed at  $-45 \pm 1$  mV ( $n = 5$ ; Fig. 1Aii). The isolated GABA<sub>A</sub>R-mediated component was recorded at a holding potential of  $0$  mV (referred to as GABA<sub>A,SC</sub>, Fig. 1Ai). The decay time was significantly longer than that for GABA<sub>A,local</sub> (GABA<sub>A,SC</sub>,  $\tau_{\text{decay}} = 277 \pm 29$  ms,  $n = 10$ ;  $t$ -test,  $P < 0.001$ ). The ratio of current observed at  $500$  ms over  $15$  ms after stimulation ( $I_{500\text{ms}}/I_{15\text{ms}}$ ) showed that GABA<sub>A,local</sub> current had decayed to approximately 5% of its peak value after  $500$  ms, while a substantial fraction of GABA<sub>A,SC</sub> could still be observed (GABA<sub>A,local</sub>,  $I_{500\text{ms}}/I_{15\text{ms}} = 0.05 \pm 0.01$ ,  $n = 11$ ; GABA<sub>A,SC</sub>,  $I_{500\text{ms}}/I_{15\text{ms}} = 0.32 \pm 0.06$ ,  $n = 6$ ;  $t$ -test,  $P < 0.001$ ).

Both GABA<sub>A,local</sub> and GABA<sub>A,SC</sub> decayed significantly faster at  $-70$  mV than at  $0$  mV ( $\tau_{\text{decay}} = 70 \pm 5$  ms,  $n = 11$ , and  $180 \pm 10$  ms,  $n = 6$ , respectively,  $t$ -test for both,  $P < 0.01$ ). The slower decay of GABA<sub>A,local</sub> and GABA<sub>A,SC</sub> at  $0$  mV is consistent with previous reports showing that GABA<sub>A</sub>R currents decay more slowly at depolarised potentials as receptors unbind agonist at a slower rate (Mellor and Randall, 1998; Burgard et al., 1999). The currents observed at  $-70$  mV were slower than those reported in some previous studies. Two factors might have contributed to the slower kinetics observed in our experiments: the use of BAPTA and the use of Cs-gluconate. Slower IPSC decay has been observed both during recording with BAPTA (Banks and Pearce, 2000) and with Cs-gluconate (Stepanyuk et al., 2002).

$\alpha 5$ GABA<sub>A</sub>Rs are widely expressed in the SR subfield of the hippocampal CA1 area (Sperk et al., 1997; Sur et al., 1999), however their synaptic contribution has not been observed while stimulating locally in the SR. We therefore asked whether  $\alpha 5$ GABA<sub>A</sub>Rs can contribute to GABA<sub>A,SC</sub>. As a first pharmacological approach we tested the gabazine sensitivity of GABA<sub>A</sub>R currents, as it has been previously reported that a tonic inhibitory conductance mediated by  $\alpha 5$ GABA<sub>A</sub>Rs is resistant to  $1 \mu\text{M}$  gabazine (Bai et al., 2001; Caraiscos et al., 2004).

To compare the gabazine sensitivity of locally-evoked and Schaffer collateral-stimulated currents, IPSCs were recorded in the presence of increasing concentrations of gabazine (Fig. 1Bi). GABA<sub>A,local</sub> currents were abolished by  $1 \mu\text{M}$  gabazine (initial amplitude,  $863 \pm 144$  pA,  $n = 11$ ) while a significant fraction of GABA<sub>A,SC</sub> remained under these conditions ( $19 \pm 3\%$  of GABA<sub>A,SC</sub> of control charge transfer remaining in  $1 \mu\text{M}$  gabazine;  $n = 6$ , initial amplitude,  $1261 \pm 57$  pA). This component is referred to as GABA<sub>A,SC(gz)</sub>. Example traces are shown in Fig. 1Bii.


The time from stimulation to peak (Table 1) was significantly different for GABA<sub>A,local</sub>, GABA<sub>A,SC</sub>, and GABA<sub>A,SC(gz)</sub> (one-way ANOVA;  $F_{2,21} = 128$ ;  $P < 0.001$ ; post-hoc  $P < 0.01$  for all comparisons). Both GABA<sub>A,SC</sub> and GABA<sub>A,SC(gz)</sub> showed slower kinetics compared to GABA<sub>A,local</sub> (Table 1, 10–90% rise time; one-way ANOVA;  $F_{2,21} = 51$ ;  $P < 0.001$ ; post-hoc  $P < 0.05$  for all comparisons). The longer time-to-peak seen for GABA<sub>A,SC(gz)</sub> was additionally due to longer latency (Table 1, time from stimulus to 10% amplitude; one-way ANOVA;  $F_{2,21} = 75$ ;  $P < 0.001$ ; post-hoc comparisons, GABA<sub>A,SC(gz)</sub> versus GABA<sub>A,local</sub> and GABA<sub>A,SC(gz)</sub> versus GABA<sub>A,SC</sub>, both  $P < 0.001$ ; GABA<sub>A,local</sub> versus GABA<sub>A,SC</sub>,  $P = 0.22$ ).

**Table 1**  
Time-to-peak and rise time for GABA<sub>A</sub>R currents (in ms).

	Time-to-peak (stimulus to peak)	Time from stimulus to 10% amplitude	Time from 10 to 90% amplitude
GABA <sub>A,local</sub>	$17 \pm 1.8$	$4.1 \pm 0.1$	$7.5 \pm 0.3$
GABA <sub>A,SC</sub>	$32 \pm 3.1$	$7.8 \pm 0.3$	$14.2 \pm 0.8$
GABA <sub>A,SC(gz)</sub>	$75 \pm 3.5$	$27.4 \pm 1.1$	$30 \pm 0.9$

As bursting input from the Schaffer collateral could be necessary for the firing of dendritic targeting interneurons (Maccaferri and Dingledine, 2002), we next asked whether bursting activity in this pathway could enhance GABA<sub>A,SC(gz)</sub>. Indeed this component became prominent with burst stimulation (3–4 stimuli at  $100$  Hz) as  $49 \pm 7\%$  of inhibitory charge transfer remained in  $1 \mu\text{M}$  gabazine (Fig. 1Ci). Remarkably, the early component (5–35 ms after stimulation) of burst responses was almost as sensitive to  $1 \mu\text{M}$  gabazine as GABA<sub>A,local</sub> (fraction remaining in  $1 \mu\text{M}$  gabazine: GABA<sub>A,local</sub>,  $0.4 \pm 1\%$ ,  $n = 6$ ; GABA<sub>A,SC</sub> (burst, 5–35 ms),  $1 \pm 5\%$ ,  $n = 6$ ;  $t$ -test,  $P = 0.73$ ; example traces in Fig. 1Cii). A significant effect of gabazine inhibition was observed for all recording conditions (GABA<sub>A,local</sub>, GABA<sub>A,SC</sub> and GABA<sub>A,SC(gz)</sub>; RM ANOVA. Charge transfer as the between-subjects factor, and dose as within-subject factor;  $F_{2,19} = 25$ ;  $P < 0.001$ ; post-hoc comparisons showed that at  $1 \mu\text{M}$  gabazine, each group was different from the others,  $P < 0.001$ ).

To test whether  $\alpha 5$ GABA<sub>A</sub>Rs contribute to the inhibitory currents described above we next investigated the effect of L-655,708, which is an inverse-agonist selective for  $\alpha 5$ GABA<sub>A</sub>R subunits in rats, mice and humans at concentrations below  $20$  nM (Atack et al., 2006). After a stable baseline recording,  $20$  nM L-655,708 was added to the extracellular solution. The effects of L-655,708 on charge transfer and peak amplitude were estimated for GABA<sub>A,local</sub>, GABA<sub>A,SC</sub> and GABA<sub>A,SC(gz)</sub>, with either single or burst stimulation for GABA<sub>A,SC(gz)</sub> (Fig. 2A). In all conditions, the average charge transfer after addition of L-655,708 was significantly different from baseline, but stronger effects were observed for GABA<sub>A,SC(gz)</sub> for both single and burst stimulation (percentage of reduction: GABA<sub>A,local</sub>,  $13 \pm 5\%$ ,  $n = 7$ ; GABA<sub>A,SC</sub>,  $12 \pm 5\%$ ,  $n = 9$ ; GABA<sub>A,SC(gz)</sub> (single),  $30 \pm 2\%$ ,  $n = 6$ ; GABA<sub>A,SC(gz)</sub> (burst),  $33 \pm 3\%$ ,




**Fig. 2.** The  $\alpha 5$  inverse-agonist L-655,708 inhibits GABA<sub>A,SC(gz)</sub>. (A) Superimposed example traces of GABA<sub>A</sub>R currents before (grey) and after (black) application of L-655,708. (B) Normalised charge transfer (i) and peak current (ii) measured during baseline and after bath application of L-655,708 at time = 0. \*\*\* $P < 0.001$ .


$n = 7$ ; Fig. 2B). A significant effect on peak was also observed for  $GABA_{A,SC}(gz)$  (single) ( $t$ -test,  $P < 0.01$ ), but did not reach significance for  $GABA_{A,SC}(gz)$  (burst) ( $t$ -test,  $P = 0.07$ ).

To corroborate our findings on the effects of L-655,708 on  $GABA_A$ R-mediated currents, and to test the specificity of L-655,708 for  $\alpha 5GABA_A$ Rs, we next used  $gabra5^{-/-}$  mice and WT littermate control mice (Collinson et al., 2002). In this set of experiments,  $GABA_{A,SC}$  currents from WT mice showed a significant reduction from baseline following application of 20 nM L-655,708 ( $GABA_{A,SC}$ ,  $33 \pm 4\%$  reduction,  $n = 5$ ;  $GABA_{A,SC}(gz)$ ,  $36 \pm 4\%$  reduction,  $n = 7$ ;  $t$ -test,  $P < 0.001$  for both), whereas L-655,708 did not significantly alter the charge transfer for  $GABA_{A,SC}$  and  $GABA_{A,SC}(gz)$  currents in  $gabra5^{-/-}$  slices (Fig. 3A, B;  $GABA_{A,SC}$ ,  $n = 7$ ;  $t$ -test,  $P = 0.11$ ;  $GABA_{A,SC}(gz)$ ,  $n = 7$ ;  $t$ -test,  $P = 0.20$ ).

We additionally tested whether L-655,708 could be used as a selective inhibitor of  $\alpha 5GABA_A$ Rs at concentrations higher than 20 nM during our recording conditions. We found that L-655,708 at 1  $\mu$ M concentration, and even at concentrations as low as 50 nM, inhibited the  $GABA_A$ R current in WT mice to a similar extent to that seen in  $gabra5^{-/-}$  mice (no significant difference WT versus  $gabra5^{-/-}$ ,  $n = 7$  and  $n = 5$ , respectively: 50 nM,  $P = 0.33$ , and 1 mM,  $P = 0.16$ ; Fig. 3C), suggesting that L-655,708 is not selective for  $\alpha 5GABA_A$ Rs at concentrations above 20 nM, and thus limiting the selective inhibition of the  $\alpha 5GABA_A$ R-mediated current to no more than  $\sim 30\%$ . Further analysis to compare  $GABA_{A,SC}$  currents recorded from WT and  $gabra5^{-/-}$  mice revealed that the fraction of  $GABA_{A,SC}$  charge transfer that persisted after the application of 1  $\mu$ M gabazine was larger in WT mice than in  $gabra5^{-/-}$  mice ( $34.4 \pm 4\%$ ,  $n = 11$  versus  $20.6 \pm 2\%$ ,  $n = 10$ , respectively,  $t$ -test,  $P < 0.01$ ; Fig. 4Ai, Aii, Bi). Also, the peak current was significantly different between  $gabra5^{-/-}$  and WT mice ( $t$ -test,  $P < 0.05$ ; Fig. 4Ai, Aii, Bii).



**Fig. 3.** L-655,708 (20 nM) is specific for the  $GABA_A$  receptor  $\alpha 5$  subunit. (A) Superimposed example traces of  $GABA_{A,SC}$  (i) and  $GABA_{A,SC}(gz)$  (ii) before and after application of 20 nM L-655,708 in wild type and  $gabra5^{-/-}$  mice. (B) Normalised values for  $GABA_A$ R charge transfer measured during baseline and following application of L-655,708 at time 0 for  $GABA_{A,SC}$  (i) and  $GABA_{A,SC}(gz)$  (ii). (C) Concentration-response plot of  $GABA_{A,SC}$  for 20, 50 and 1000 nM L-655,708 in wild type and  $gabra5^{-/-}$  mice. \*\*\* $P < 0.001$ .




**Fig. 4.**  $GABA_A$  receptor-mediated currents in  $gabra5^{-/-}$  mice and WT mice are differentially sensitive to 1  $\mu$ M gabazine. (A) Superimposed example traces of  $GABA_{A,SC}$  currents before (grey) and after (black) application of 1  $\mu$ M gabazine. Recordings from WT (i), and  $gabra5^{-/-}$  mice (ii). (B) Normalised values for charge transfer (i) and peak current (ii) measured during baseline and following application of 1  $\mu$ M gabazine at time 0. \* $P < 0.05$ ; \*\* $P < 0.01$ .

Finally, to test whether the  $GABA_A$ R currents described above are mediated in part by  $\alpha 1/\alpha 2/\alpha 3GABA_A$ Rs receptors we tested the effects of 400 nM zolpidem (Fig. 5). As expected, zolpidem increased the charge transfer of  $GABA_{A,local}$  currents ( $48 \pm 6\%$ ,  $n = 7$ ). Also, both  $GABA_{A,SC}$  and  $GABA_{A,SC}(gz)$  charge transfer was enhanced ( $GABA_{A,SC}$  by  $33.3 \pm 4\%$ ,  $n = 5$ , and  $GABA_{A,SC}(gz)$  by  $16 \pm 5\%$ ,  $n = 5$ ; Fig. 5A, B). Zolpidem had a significantly different effect on currents under all recording conditions ( $GABA_{A,local}$ ,  $GABA_{A,SC}$  and  $GABA_{A,SC}(gz)$ ; one-way ANOVA,  $F_{2,32} = 21.6$ ,  $P < 0.001$ ; post-hoc comparisons with significant differences as shown in Fig. 5Bi).

A slight increase in peak amplitude from baseline values was observed in all three recording conditions with no significant difference among them (Fig. 5Bii). The large increase in charge transfer and small change in peak amplitude in response to zolpidem have been previously observed for miniature postsynaptic potentials (Perrais and Ropert, 1999). In total, our results suggest that  $\alpha 5GABA_A$ Rs contribute significantly to  $GABA_{A,SC}$ .

#### 4. Discussion

Previous work investigating the subunit composition of synaptic  $GABA_A$  receptors has measured spontaneous or stimulus-evoked GABAergic inhibition in the hippocampus while blocking glutamatergic excitation. In the present study, we describe the distinct pharmacology of locally-evoked versus Schaffer collateral-stimulated  $GABA_A$ R currents.  $GABA_{A,local}$  currents were eliminated by 1  $\mu$ M gabazine and were markedly enhanced by 400 nM zolpidem, whereas  $GABA_{A,SC}$  currents were less sensitive to 1  $\mu$ M gabazine



**Fig. 5.** The enhancing effect of zolpidem is larger on  $GABA_{A,local}$  and  $GABA_{A,SC}$  than on  $GABA_{A,SC(gz)}$ . (A) Superimposed example traces of  $GABA_A$  currents before and after application of zolpidem (400 nM). From left to right:  $GABA_{A,local}$ ,  $GABA_{A,SC}$ , and  $GABA_{A,SC(gz)}$  with corresponding symbols in B. (B) Normalised values of charge transfer (i) and peak current (ii) measured during baseline and following application of zolpidem at time 0. Significant difference of charge transfers but not peak currents between baseline and the last 4 min of recording for the three recording conditions. \* $P < 0.05$ ; \*\*\* $P < 0.001$ .

and zolpidem. The gabazine-insensitive component of  $GABA_{A,SC}$  showed the greatest reduction by L-655,708 and was relatively insensitive to zolpidem, suggesting a significant proportion of  $\alpha 5GABA_A$ R. Furthermore, the gabazine-insensitive current was markedly enhanced following burst stimulation of Schaffer collaterals. Finally, we confirmed the specificity of L-655,708 on  $GABA_{A,SC}$  and the gabazine-insensitive currents by recording from WT and *gabra5*<sup>-/-</sup> mice.

IPSCs recorded in CA1 pyramidal neurons evoked by local stimulation at the SR were likely generated by perisomatic targeting interneurons, including basket cells (Ouardouz and Lacaille, 1997). This  $GABA_{A,local}$  current had a  $\tau_{decay}$  of 163 ms at 0 mV. In experiments designed to study Schaffer collateral-stimulated currents, the IPSCs decayed with a strikingly slow time constant (277 ms).  $GABA_{A,local}$  and  $GABA_{A,SC}$  also showed significantly different time-to-peak, and 10–90% rise time. Previous studies have already shown evidence for such fast and slow components of synaptic inhibition (Pearce, 1993; Banks et al., 1998). Whereas there was no significant difference in latency between  $GABA_{A,local}$  and  $GABA_{A,SC}$ , suggesting that stimulation at the SR could also trigger direct activation of interneurons,  $GABA_{A,SC(gz)}$  also showed a longer latency, suggesting that under these conditions, a delay in activation of presynaptic interneurons contributed to the longer time-to-peak.

Several possible explanations could account for the late peak and surprisingly slow decay of  $GABA_{A,SC}$  compared to  $GABA_{A,local}$ . Firstly, using fast perfusion on excised patches from cells expressing recombinant receptors, it has been shown that subunit composition can affect decay time (Tia et al., 1996; Burgard et al., 1999). Secondly, the time course of GABA concentration at the

release sites would affect the response kinetics, for example the rise time responses for extrasynaptic receptors could be slowed down as has been suggested for glutamatergic synapses (Scimemi et al., 2004). Thirdly, long exposure to GABA, either in the synaptic cleft or in extrasynaptic space, could produce the reactivation of synaptic receptors as previously shown by modifying GABA uptake kinetics (Roepstorff and Lambert, 1994). Recent studies using somatic recordings in anatomically-identified connected cell pairs suggest that the time course of inhibitory responses can indeed be determined both by subunit composition and by distinct transmitter release transients (Szabadics et al., 2007; Ali and Thomson, 2008), but we cannot exclude the possibility that dendritic filtering contributes to the slow kinetics as recorded at the soma. A further possibility with potential physiological significance is that the slow rise and decay kinetics we observed for  $GABA_{A,SC}$  is due to the pattern of activation of  $GABA_A$  interneurons. For example, late-firing dendritic targeting interneurons have been shown to generate slow  $GABA_A$  events in CA1 pyramidal neurons (Maccaferri and Dingledine, 2002). Furthermore, bursts of GABA release produced by several action potentials or slow asynchronous release (Hefft and Jonas, 2005) could contribute to slow kinetics.

$GABA_{A,SC}$  and  $GABA_{A,local}$  showed different sensitivities to gabazine. We consistently found that approximately 20% of  $GABA_{A,SC}$  charge transfer evoked by single stimulation was insensitive to 1  $\mu$ M gabazine, and this fraction increased to ~50% when using burst stimulation. We interpreted this finding as an indication of a component mediated by  $\alpha 5GABA_A$ Rs as it has been previously shown that the tonic current mediated by these is not sensitive to 1  $\mu$ M gabazine (Bai et al., 2001; Caraiscos et al., 2004). We obtained further evidence that  $\alpha 5GABA_A$ Rs are activated by Schaffer collateral stimulation by studying the effects of L-655,708 on the isolated  $GABA_A$  components described above.  $GABA_{A,SC(gz)}$  produced either by single or burst stimulation was reduced by 30% and there was no significant difference in the amount of reduction using either type of stimulation ( $P = 0.27$ ) as would be expected if in both cases the synaptically evoked current is mediated by a similar proportion of  $\alpha 5GABA_A$ Rs. L-655,708 also produced a significant but smaller effect on  $GABA_{A,SC}$  and  $GABA_{A,local}$ . In previous studies, spontaneous  $GABA_{A,local}$  currents were found not to contain an  $\alpha 5GABA_A$ R-mediated component (Caraiscos et al., 2004; Glykys and Mody, 2006; Zarnowska et al., 2009), however, it is likely that the sorting of populations of spontaneous IPSCs by their decay time for analysis could limit the detection of a slow component in  $GABA_{A,local}$  currents. Furthermore, it is likely that the extracellular stimulation used to produce  $GABA_{A,local}$  under our experimental conditions would recruit interneurons other than perisomatic targeting interneurons. It is also possible that gabazine, as a competitive antagonist, could be displaced from  $GABA_A$  receptors by released GABA under our recording conditions for  $GABA_{A,SC(gz)}$ . Although we do not have evidence to discard this possibility, the effects of 20 nM L-655,708 on  $GABA_{A,SC}$  in both rat and mouse recordings without the use of gabazine strongly support a specific  $\alpha 5GABA_A$ R-mediated component.

We corroborated the specificity of 20 nM L-655,708 on  $GABA_{A,SC}$  and  $GABA_{A,SC(gz)}$  currents using *gabra5*<sup>-/-</sup> mice and WT littermate controls. There was a one third reduction in the  $GABA_{A,SC}$  and  $GABA_{A,SC(gz)}$  currents from initial control values following application of L-655,708 in WT mice, and no significant reduction in *gabra5*<sup>-/-</sup> mice. *In vitro* analysis of L-655,708 activity on recombinant human  $GABA_A$ Rs showed a maximum inhibition of 20% of the  $\alpha 5GABA_A$ R current (Atack et al., 2006). Consistent with this we found in both rat and mouse recordings that 20 nM L-655,708 inhibited approximately 30% of  $GABA_{A,SC(gz)}$ .

As a final test to probe the composition of  $GABA_A$ Rs mediating the early and late components of inhibition, we used 400 nM



zolpidem which is a highly potent and selective benzodiazepine, with  $\alpha 5$ GABA<sub>A</sub>R sparing properties ( $\alpha 5$ GABA<sub>A</sub>R,  $K_i > 15 \mu\text{M}$ , reviewed in Sieghart, 1995). GABA<sub>A,local</sub> was markedly enhanced after application of zolpidem with a three fold increase in charge transfer compared to GABA<sub>A,SC(gz)</sub>. In comparison, GABA<sub>A,SC</sub> showed only a two fold increase compared to GABA<sub>A,SC(gz)</sub>. We interpret this result to suggest that local stimulation in SR near SP in the absence of fast synaptic excitation mostly activates perisomatic targeting interneurons, which selectively activate  $\alpha 1$ ,  $\alpha 2$  and  $\alpha 3$ GABA<sub>A</sub>Rs (Thomson et al., 2000), and that activation of Schaffer collaterals not only stimulates perisomatic targeting cells but also dendritic targeting interneurons, some of which specifically target  $\alpha 5$ GABA<sub>A</sub>Rs. Therefore, the combined results using 400 nM zolpidem and 20 nM L-655,708 suggest that GABA<sub>A,SC</sub> contains a significant population of  $\alpha 5$ GABA<sub>A</sub>Rs. Thus, the present observations highlight the strong activation of  $\alpha 5$ GABA<sub>A</sub>Rs following Schaffer collateral stimulation in the hippocampus and the pharmacological analysis of GABA<sub>A,SC(gz)</sub> suggests a particularly robust activation during bursting activity.

Our results are consistent with previous suggestions that slow inhibition is mediated by a specific subpopulation of interneurons and molecularly distinct receptors (Banks et al., 1998; Pearce, 1993; Zarnowska et al., 2009). The widespread localisation of  $\alpha 5$ GABA<sub>A</sub>Rs at dendritic sites (Sperk et al., 1997) suggests that they are involved in gating dendritic excitability, for example they could be involved in regulating the generation of dendritic spikes such as those observed *in vivo* during sharp waves (Kamondi et al., 1998). Furthermore, the time course similarity of GABA<sub>A,SC</sub> and GABA<sub>A,SC(gz)</sub> to the decay kinetics of NMDA receptors (Vicini et al., 1998) makes them well suited for inhibition of the induction of long-term potentiation by providing a shunting inhibitory effect on the NMDAR current (Staley and Mody, 1992).

The higher acquisition rate in associative memory tasks observed in mice after systemic application of the inverse-agonist L-655,708 or in *gabra5*<sup>-/-</sup> mice could be the expression of plasticity produced by enhanced dendritic excitation. In order to understand the cellular and network mechanisms that lead to enhanced learning after reduction of  $\alpha 5$ GABA<sub>A</sub>Rs function it will be necessary to study the requirement for activation/silencing of inputs targeting  $\alpha 5$ GABA<sub>A</sub>Rs on hippocampal pyramidal neurons by afferent inputs to the hippocampus and their timing in relation to the timing of somatic and dendritic spikes. Here we provide evidence of activity pattern-dependent feed-forward activation of  $\alpha 5$ GABA<sub>A</sub>Rs in CA1. Testing the role of  $\alpha 5$ GABA<sub>A</sub>Rs in controlling the generation of dendritic spikes or in the fine tuning of long-term potentiation awaits the broader availability of more potent and selective drugs acting on  $\alpha 5$ GABA<sub>A</sub>Rs, such as RO4938581 (Ballard et al., 2009) or the specific activation of  $\alpha 5$ GABA<sub>A</sub>R-targeting interneurons *in vivo*.

## Acknowledgements

We thank Dr. Elizabeth Tunbridge for helpful suggestions with analysis. M.V.-C. was funded by a Wellcome Trust International Research Fellowship. L.J.M. was supported by a Canadian Institutes of Health Research (CIHR), Canada Graduate Scholarship. B.A.O. is supported by CIHR operating grants and a Canada Research Chair. M.W.S. (CIHR MT-12682); M.W.S. holds a Canada Research Chair; M.W.S. is an International Research Scholar of the Howard Hughes Medical Institute. The original breeding pairs of *gabra5*<sup>-/-</sup> mice were generously supplied by Merck, Sharp and Dohme, Harlow, UK.

## References

Alger, B.E., Nicoll, R.A., 1982. Feed-forward dendritic inhibition in rat hippocampal pyramidal cells studied *in vitro*. *J. Physiol.* 328, 105–123.

- Ali, A.B., Thomson, A.M., 2008. Synaptic  $\alpha 5$  subunit-containing GABA<sub>A</sub> receptors mediate IPSPs elicited by dendrite-preferring cells in rat neocortex. *Cereb. Cortex* 18, 1260–1271.
- Attack, J.R., Bayley, P.J., Seabrook, G.R., Wafford, K.A., McKernan, R.M., Dawson, G.R., 2006. L-655,708 enhances cognition in rats but is not proconvulsant at a dose selective for  $\alpha 5$ -containing GABA<sub>A</sub> receptors. *Neuropharmacology* 51, 1023–1029.
- Attack, J.R., Hutson, P.H., Collinson, N., Marshall, G., Bentley, G., Moyes, C., Cook, S.M., Collins, L., Wafford, K., McKernan, R.M., Dawson, G.R., 2005. Anxiogenic properties of an inverse agonist selective for  $\alpha 3$  subunit-containing GABA<sub>A</sub> receptors. *Br. J. Pharmacol.* 144, 357–366.
- Ballard, T.M., Knoflach, F., Prinssen, E., Borroni, E., Vivian, J.A., Basile, J., Gasser, R., Moreau, J.L., Wettstein, J.G., Buettelmann, B., Knust, H., Thomas, A.W., Trube, G., Hernandez, M.C., 2009. RO4938581, a novel cognitive enhancer acting at GABA<sub>A</sub>  $\alpha 5$  subunit-containing receptors. *Psychopharmacology (Berl.)* 202, 207–223.
- Bai, D., Zhu, G., Pennefather, P., Jackson, M.F., MacDonald, J.F., Orser, B.A., 2001. Distinct functional and pharmacological properties of tonic and quantal inhibitory postsynaptic currents mediated by  $\gamma$ -aminobutyric acid<sub>A</sub> receptors in hippocampal neurons. *Mol. Pharmacol.* 59, 814–824.
- Banks, M.I., Li, T.B., Pearce, R.A., 1998. The synaptic basis of GABA<sub>A,slow</sub>. *J. Neurosci.* 18, 1305–1317.
- Banks, M.I., Pearce, R.A., 2000. Kinetic differences between synaptic and extra-synaptic GABA<sub>A</sub> receptors in CA1 pyramidal cells. *J. Neurosci.* 20, 937–948.
- Banks, M.I., White, J.A., Pearce, R.A., 2000. Interactions between distinct GABA<sub>A</sub> circuits in hippocampus. *Neuron* 25, 449–457.
- Burgard, E.C., Haas, K.F., Macdonald, R.L., 1999. Channel properties determine the transient activation kinetics of recombinant GABA<sub>A</sub> receptors. *Brain Res. Mol. Brain Res.* 73, 28–36.
- Caraiscos, V.B., Elliott, E.M., You-Ten, K.E., Cheng, V.Y., Belelli, D., Newell, J.G., Jackson, M.F., Lambert, J.J., Rosahl, T.W., Wafford, K.A., MacDonald, J.F., Orser, B.A., 2004. Tonic inhibition in mouse hippocampal CA1 pyramidal neurons is mediated by  $\alpha 5$  subunit-containing  $\gamma$ -aminobutyric acid type A receptors. *Proc. Natl. Acad. Sci. U.S.A.* 101, 3662–3667.
- Collinson, N., Kuenzi, F.M., Jarolimek, W., Maubach, K.A., Cotterill, R., Sur, C., Smith, A., Otu, F.M., Howell, O., Attack, J.R., McKernan, R.M., Seabrook, G.R., Dawson, G.R., Whiting, P.J., Rosahl, T.W., 2002. Enhanced learning and memory and altered GABAergic synaptic transmission in mice lacking the  $\alpha 5$  subunit of the GABA<sub>A</sub> receptor. *J. Neurosci.* 22, 5572–5580.
- Glykys, J., Mody, I., 2006. Hippocampal network hyperactivity after selective reduction of tonic inhibition in GABA<sub>A</sub> receptor  $\alpha 5$  subunit-deficient mice. *J. Neurophysiol.* 95, 2796–2807.
- Hefft, S., Jonas, P., 2005. Asynchronous GABA release generates long-lasting inhibition at a hippocampal interneuron-principal neuron synapse. *Nat. Neurosci.* 8, 1319–1328.
- Howell, O., Attack, J.R., Dewar, D., McKernan, R.M., Sur, C., 2000. Density and pharmacology of  $\alpha 5$  subunit-containing GABA<sub>A</sub> receptors are preserved in hippocampus of Alzheimer's disease patients. *Neuroscience* 98, 669–675.
- Kamondi, A., ACSady, L., Buzsaki, G., 1998. Dendritic spikes are enhanced by cooperative network activity in the intact hippocampus. *J. Neurosci.* 18, 3919–3928.
- Klausberger, T., Somogyi, P., 2008. Neuronal diversity and temporal dynamics: the unity of hippocampal circuit operations. *Science* 321, 53–57.
- Maccaferri, G., Dingledine, R., 2002. Control of feedforward dendritic inhibition by NMDA receptor-dependent spike timing in hippocampal interneurons. *J. Neurosci.* 22, 5462–5472.
- Mann, E.O., Paulsen, O., 2007. Role of GABAergic inhibition in hippocampal network oscillations. *Trends Neurosci.* 30, 343–349.
- Martin, L.J., Oh, G.H.T., Orser, B.A., 2009. Etomidate targets  $\alpha 5$   $\gamma$ -aminobutyric acid subtype A receptors to regulate synaptic plasticity and memory blockade. *Anesthesiology* 111, 1025–1035.
- McKernan, R.M., Rosahl, T.W., Reynolds, D.S., Sur, C., Wafford, K.A., Attack, J.R., Farrar, S., Myers, J., Cook, G., Ferris, P., Garrett, L., Bristow, L., Marshall, G., Macaulay, A., Brown, N., Howell, O., Moore, K.W., Carling, R.W., Street, L.J., Castro, J.L., Ragan, C.L., Dawson, G.R., Whiting, P.J., 2000. Sedative but not anxiolytic properties of benzodiazepines are mediated by the GABA<sub>A</sub> receptor  $\alpha 1$  subtype. *Nat. Neurosci.* 3, 587–592.
- Mellor, J.R., Randall, A.D., 1998. Voltage-dependent deactivation and desensitization of GABA responses in cultured murine cerebellar granule cells. *J. Physiol.* 506, 377–390.
- Nathan, T., Jensen, M.S., Lambert, J.D., 1990. The slow inhibitory postsynaptic potential in rat hippocampal CA1 neurones is blocked by intracellular injection of QX-314. *Neurosci. Lett.* 110, 309–313.
- Nusser, Z., Sieghart, W., Benke, D., Fritschy, J.M., Somogyi, P., 1996. Differential synaptic localization of two major  $\gamma$ -aminobutyric acid type A receptor subunits on hippocampal pyramidal cells. *Proc. Natl. Acad. Sci. U.S.A.* 93, 11939–11944.
- Ouardouz, M., Lacaille, J.C., 1997. Properties of unitary IPSCs in hippocampal pyramidal cells originating from different types of interneurons in young rats. *J. Neurophysiol.* 77, 1939–1949.
- Pearce, R.A., 1993. Physiological evidence for two distinct GABA<sub>A</sub> responses in rat hippocampus. *Neuron* 10, 189–200.
- Perrais, D., Ropert, N., 1999. Effect of zolpidem on miniature IPSCs and occupancy of postsynaptic GABA<sub>A</sub> receptors in central synapses. *J. Neurosci.* 19, 578–588.
- Prenosil, G.A., Schneider Gasser, E.M., Rudolph, U., Keist, R., Fritschy, J.M., Vogt, K.E., 2006. Specific subtypes of GABA<sub>A</sub> receptors mediate phasic and tonic forms of inhibition in hippocampal pyramidal neurons. *J. Neurophysiol.* 96, 846–857.

- Roepstorff, A., Lambert, J.D., 1994. Factors contributing to the decay of the stimulus-evoked IPSC in rat hippocampal CA1 neurons. *J. Neurophysiol.* 72, 2911–2926.
- Rudolph, U., Crestani, F., Benke, D., Brunig, I., Benson, J.A., Fritschy, J.M., Martin, J.R., Bluethmann, H., Mohler, H., 1999. Benzodiazepine actions mediated by specific  $\gamma$ -aminobutyric acid<sub>A</sub> receptor subtypes. *Nature* 401, 796–800.
- Scimemi, A., Fine, A., Kullmann, D.M., Rusakov, D.A., 2004. NR2B-containing receptors mediate cross talk among hippocampal synapses. *J. Neurosci.* 24, 4767–4777.
- Scimemi, A., Semyanov, A., Sperk, G., Kullmann, D.M., Walker, M.C., 2005. Multiple and plastic receptors mediate tonic GABA<sub>A</sub> receptor currents in the hippocampus. *J. Neurosci.* 25, 10016–10024.
- Sieghart, W., 1995. Structure and pharmacology of  $\gamma$ -aminobutyric acid<sub>A</sub> receptor subtypes. *Pharmacol. Rev.* 47, 181–234.
- Sperk, G., Schwarzer, C., Tsunashima, K., Fuchs, K., Sieghart, W., 1997. GABA<sub>A</sub> receptor subunits in the rat hippocampus I: immunocytochemical distribution of 13 subunits. *Neuroscience* 80, 987–1000.
- Staley, K.J., Mody, I., 1992. Shunting of excitatory input to dentate gyrus granule cells by a depolarizing GABA<sub>A</sub> receptor-mediated postsynaptic conductance. *J. Neurophysiol.* 68, 197–212.
- Stepanyuk, A., Chvanov, M., Ivanov, A., Boychuk, Y., Pivnevà, T., Belan, P., 2002. Prolonged decay of evoked inhibitory postsynaptic currents in hippocampal neurons is not shaped by asynchronous release. *Neurophysiology* 34, 239–242.
- Sur, C., Fresu, L., Howell, O., McKernan, R.M., Atack, J.R., 1999. Autoradiographic localization of  $\alpha 5$  subunit-containing GABA<sub>A</sub> receptors in rat brain. *Brain Res.* 822, 265–270.
- Szabadics, J., Tamas, G., Soltesz, I., 2007. Different transmitter transients underlie presynaptic cell type specificity of GABA<sub>A,slow</sub> and GABA<sub>A,local</sub>. *Proc. Natl. Acad. Sci. U.S.A.* 104, 14831–14836.
- Tenen, S.S., Hirsch, J.D., 1980.  $\beta$ -Carboline-3-carboxylic acid ethyl ester antagonizes diazepam activity. *Nature* 288, 609–610.
- Thomson, A.M., Bannister, A.P., Hughes, D.I., Pawelzik, H., 2000. Differential sensitivity to Zolpidem of IPSPs activated by morphologically identified CA1 interneurons in slices of rat hippocampus. *Eur. J. Neurosci.* 12, 425–436.
- Tia, S., Wang, J.F., Kotchabhakdi, N., Vicini, S., 1996. Distinct deactivation and desensitization kinetics of recombinant GABA<sub>A</sub> receptors. *Neuropharmacology* 35, 1375–1382.
- Vicini, S., Wang, J.F., Li, J.H., Zhu, W.J., Wang, Y.H., Luo, J.H., Wolfe, B.B., Grayson, D.R., 1998. Functional and pharmacological differences between recombinant N-methyl-D-aspartate receptors. *J. Neurophysiol.* 79, 555–566.
- Wafford, K.A., 2005. GABA<sub>A</sub> receptor subtypes: any clues to the mechanism of benzodiazepine dependence? *Curr. Opin. Pharmacol.* 5, 47–52.
- Zarnowska, E.D., Keist, R., Rudolph, U., Pearce, R.A., 2009. GABA<sub>A</sub> receptor  $\alpha 5$  subunits contribute to GABA<sub>A,slow</sub> synaptic inhibition in mouse hippocampus. *J. Neurophysiol.* 101, 1179–1191.