



Auditory verbal hallucinations and cognitive functioning in healthy individuals

Kirstin Daalman ^{a,*}, Martine van Zandvoort ^{a,b}, Florian Bootsman ^a, Marco Boks ^{a,c},
René Kahn ^a, Iris Sommer ^a

^a Department of Psychiatry, Neuroscience Division, University Medical Center Utrecht & Rudolf Magnus Institute for Neuroscience, Heidelberglaan 100, 3584 CX Utrecht, The Netherlands

^b Helmholtz Research Institute, Utrecht University, P.O. Box 80125, 3508 TC Utrecht, The Netherlands

^c Julius Centre for Health Sciences and Primary Care, University Medical Center Utrecht, Universiteitsweg 100, 3584 CG Utrecht, The Netherlands

ARTICLE INFO

Article history:

Received 5 April 2011

Received in revised form 6 July 2011

Accepted 10 July 2011

Available online 12 August 2011

Keywords:

Auditory verbal hallucinations

Non-psychotic individuals

Cognitive tasks

Voices

ABSTRACT

Auditory verbal hallucinations (AVH) are a characteristic symptom in schizophrenia, and also occur in the general, non-clinical population. In schizophrenia patients, several specific cognitive deficits, such as in speech processing, working memory, source memory, attention, inhibition, episodic memory and self-monitoring have been associated with auditory verbal hallucinations. Such associations are interesting, as they may identify specific cognitive traits that constitute a predisposition for AVH. However, it is difficult to disentangle a specific relation with AVH in patients with schizophrenia, as so many other factors can affect the performance on cognitive tests. Examining the cognitive profile of healthy individuals experiencing AVH may reveal a more direct association between AVH and aberrant cognitive functioning in a specific domain.

For the current study, performance in executive functioning, memory (both short- and long-term), processing speed, spatial ability, lexical access, abstract reasoning, language and intelligence performance was compared between 101 healthy individuals with AVH and 101 healthy controls, matched for gender, age, handedness and education.

Although performance of both groups was within the normal range, not clinically impaired, significant differences between the groups were found in the verbal domain as well as in executive functioning. Performance on all other cognitive domains was similar in both groups. The predisposition to experience AVH is associated with lower performance in executive functioning and aberrant language performance. This association might be related to difficulties in the inhibition of irrelevant verbal information.

© 2011 Elsevier B.V. All rights reserved.

1. Introduction

Auditory verbal hallucinations (AVH) are a characteristic symptom of schizophrenia but have also been described in the general population (Tien, 1991; Verdoux and Van Os, 2002). Thus far, the pathophysiology of AVH is still largely unknown. Many theories concerning the origin of AVH have been postulated of which several have implicated specific cognitive dysfunctions as the core abnormality to cause AVH. For example, Frith and Done (1988) hypothesized a failure in self-monitoring as the basic deficit in AVH whereas Verkammen et al. (2008) stated that increased top-down processing plays an important role in the vulnerability to experience AVH. In support of such cognitive deficits or traits, hypothesized to underlie AVH, a number of studies found prominent impairments in several

cognitive functions such as speech processing (Hoffman et al., 1999), working memory (Hoffman et al., 1999), episodic memory (Berenbaum et al., 2008), source memory (Brébion et al., 2007), attention (Berman et al., 1997), inhibition (Waters et al., 2003) and self-monitoring (Seal et al., 2004; Waters et al., 2010). However, patients with schizophrenia suffer from various other symptoms besides AVH, among which avolition, lack of motivation and a general decline in cognitive functioning. Therefore, decreased performance on specific tests is not necessarily a reflection of their tendency to hallucinate. A more specific reflection of AVH may be provided by cognitive differences that occur in non-psychotic individuals with AVH, who are free of negative symptoms and have only sub-clinical levels of positive symptoms (Sommer et al., 2010a,b). The fact that these healthy individuals with AVH function at a normal level, were able to finish their education, are medication naïve and have no history of admission to hospital is an additional advantage. Although differences in AVH have been found between healthy individuals and psychotic patients, regarding for instance frequency and emotional content, several similarities remain: no differences were found between location of AVH, loudness, number of voices and personification (Daalman et al., 2011). Based on these results one cannot conclude that both types of AVH are different. Furthermore, Diederens et al.

* Corresponding author at: Neuroscience Division, University Medical Center Utrecht & Rudolf Magnus Institute for Neuroscience, B01.206, Heidelberglaan 100, 3584 CX, Utrecht, The Netherlands. Tel.: +31 8 8755 6370; fax: +31 8 8755 5509.

E-mail addresses: K.Daalman@umcutrecht.nl (K. Daalman), M.J.E.vanZandvoort@umcutrecht.nl (M. van Zandvoort), F.Bootsman@umcutrecht.nl (F. Bootsman), M.P.M.Boks@umcutrecht.nl (M. Boks), R.Kahn@umcutrecht.nl (R. Kahn), I.Sommer@umcutrecht.nl (I. Sommer).

(2011) found no significant differences in brain activation during the experience of AVH between healthy individuals with AVH and patients.

In order to measure cognitive functioning in non-psychotic individuals with AVH, a group of 101 persons with AVH who were screened for axis I or II pathology was compared to a matched control group with a battery of neuropsychological tests. These tests focus primarily on cognitive domains that were previously found to be affected in patients with a psychotic disorder experiencing AVH. The most important cognitive domains in that perspective were included: memory, language, executive functioning, processing speed, spatial ability, verbal and non-verbal reasoning.

The aim of the present study was to establish a cognitive profile of healthy individuals with AVH. Compared to healthy individuals without AVH, this group might show deviant cognitive performance. These cognitive differences will then provide clues for a potential cognitive mechanism that could underlie AVH since these individuals are otherwise healthy.

2. Method

2.1. Participants

A total of 101 healthy individuals with AVH were compared to 101 healthy individuals without AVH. Hallucinating individuals that were free of a DSM-IV diagnosis, as assessed by an independent psychiatrist using the Comprehensive Assessment of Symptoms and History (CASH) interview (Andreasen et al., 1992) and the Structured Clinical Interview for Personality Disorder (SCID-II, First et al., 1995), were included. Depressive disorder in complete remission was not an exclusionary criterion. Urine samples were used to screen for cannabis, amphetamine, cocaine, methadone or heroine abuse, which was an exclusion criterion. Additional exclusion criteria for both groups were alcohol abuse and IQ below 80.

For the healthy individuals with AVH, the minimum frequency to experience AVH was once every three months and the minimum duration since onset of AVH was one year.

Both healthy controls and healthy individuals with AVH were recruited with the help of a Dutch website called 'explore your mind' (www.verkenuwgeest.nl). For more details about selection and assessment procedure see previous studies by our group (Sommer et al., 2010a, b; Daalman et al., 2011). The control group was matched for gender, age, handedness and education and did not differ significantly on these variables, as shown in Table 1. All participants had four Dutch grandparents. The study was approved by the Humans Ethics Committee of the University Medical Center Utrecht. After complete description of the study to the participants, written informed consent was obtained.

2.1.1. Phenomenology of AVH in healthy individuals

To establish the phenomenological characteristics of AVH, the PSYRATS Auditory Hallucination Rating Scale (AHRs, Haddock et al., 1999) was administered. This questionnaire describes 11 characteristics of AVH. Each item of this scale is evaluated on a 5-point Likert Scale

ranging from 0 to 4. For the use of this questionnaire in healthy individuals, the range of the frequency scale is extended to 0–6 (also covering options 'at least once every month' and 'at least once every three months' since AVH are experienced less often than once a week (the original minimum score of this item). This questionnaire was administered by trained psychologists.

Due to high correlations between several of these items, two new variables were computed (see also Daalman et al., 2011). The variable 'emotional valence of content' was operationalized as the sum of three items from the AHRs: 'amount of negative content of voices', 'degree of negative content' and 'amount of distress'; i.e. an ordinal variable expressing overall burden of voices with negative content. The variable 'total distress' was operationalized as the sum of two items from the AHRs: 'intensity of distress', and 'disruption to life caused by voices'. As a result, the following items were used in this study: frequency, duration, location, loudness, beliefs re-origin of voices, controllability, emotional valence of content and total distress.

2.2. Measures

2.2.1. Neuropsychological assessments

The neuropsychological tests used in this study cover the domains in which impaired functioning in psychotic patients with AVH is found, and are thus candidates for examining the relationship between AVH in healthy individuals and cognitive functioning (Table 2).

Tests were administered in a fixed order and all examiners were extensively trained and supervised in the use of the tests. To rule out language deficits (aphasia, language expression and comprehension difficulties), the Boston naming task (Kaplan et al., 1983) and Token test (De Renzi and Vignolo, 1962) was administered. Participants who showed impaired language functioning, as measured by these tasks, were excluded.

2.3. Statistical analyses

Between-group comparison on the above described cognitive measures was achieved through multivariate analysis of variance (MANOVA), applying a General Linear Model. The independent variable was group (experiencing AVH or not i.e. controls) and the dependent variables were the raw scores on the fourteen cognitive tasks as described above. The Step-Up Hochberg correction was used to adjust P-values because of multiple testing (Westfall and Young, 1993; Benjamini and Hochberg, 1995). All data were analyzed with the Statistical package for the Social Sciences (SPSS, 2006).

3. Results

Table 3 describes how often AVH were experienced in the healthy individuals. Table 4 provides more information about the phenomenology of AVH in this group.

As expected, individuals of both groups performed the tests within the normal range, when compared to the norm reference scores of each test. There was a statistically significant difference between the groups on the combined dependent variables: $F(14,187) = 3.65$, $P < 0.001$; Pillai's Trace 0.22. When the results for the dependent variables were considered separately (Table 5), the individuals experiencing AVH were more sensitive to distraction as reflected in lower performance on the Stroop interference measure, had a lower verbal working memory capacity (reflected in lower Digit-span backward performance), underperformed, compared to controls, on a task for vocabulary (Vocabulary test, WAIS III subtask) and for judging verbal similarities (Similarities test, WAIS III subtask). In addition the individuals with AVH performed slightly lower on the NART, an estimate of verbal intelligence. No differences were found on tasks tapping verbal and nonverbal memory, attention span, nor on

Table 1
Demographic characteristics of the participants: healthy individuals with AVH and healthy controls.

Group	Individuals with AVH	Controls	Difference (significance)
n	101	101	
Male (%)	33.7	29.7	$\chi^2 = .366$ ($P = 0.545$)
Right handed (%)	78.2	84.2	$\chi^2 = 1.167$ ($P = 0.280$)
Mean age (s.d.)	43.78 (12.50)	43.30 (14.23)	$t = .257$ ($P = 0.797$)
Mean years of education (s.d.)	13.39 (2.18)	13.76 (2.40)	$t = -1.17$ ($P = 0.245$)

Table 2
Description of neuropsychological tests and measured domains.

Task	Measured domain
Stroop Color-Word Task (Stroop, 1935) Card 1: Subject is asked to read names of colors Card 2: Subject is asked to name color of ink Card 3: Subject is asked to name color of ink while written word states different color	Executive functioning: response inhibition and selective attention (time card 3–time card 2) Processing speed (time card 1, time card 2)
Backward digit span-task, WAIS III subtask (Wechsler, 1997) Numbers are presented and subject has to repeat them backwards. Sequence of numbers increases after two trials	Verbal working memory Executive functioning and verbal working memory, requiring executive manipulation of verbal presented stimuli
Forward digit span, WAIS III subtask (Wechsler, 1997) Numbers are presented and subject has to repeat them in same order. Sequence of numbers increases after two trials	Attention span
California Verbal Learning Test (CVLT, Delis et al., 1987) (Dutch version: VLGT, Mulder et al., 1996) Recall of list of words after 25 min. Words can be grouped into categories	Long-term verbal memory
Complex figure of Rey–Osterrieth (Rey–O, Knight and Kaplan, 2003) Subject has to copy a complex figure. After 25 min, subject is asked to draw this figure again from memory	Spatial ability Non-verbal long-term memory (recall after 25 min)
Vocabulary test, WAIS III subtask (Wechsler, 1997) Subject is asked to give the definition of words	Lexical access
Similarities test, WAIS III subtask (Wechsler, 1997) Subject is asked to state the similarity between two concepts	(Verbal) abstract reasoning
Dutch version of the controlled oral word association test, COWAT (Lezak et al., 2004) Subject is asked to name as many words within 1 min, starting with a specific letter	Executive functioning Phonemic fluency (letters N and A, each 1 min), verbal retrieval and recall, self-monitoring, self-initiation
Semantic fluency (Lezak et al., 2004) Subject is asked to name as many items from a category within 2 min	Semantic memory Categorical fluency (animals and occupations, each 2 min). Verbal retrieval and recall, self monitoring, Semantic associations in the lexicon
National Reading Test for Adults (NART, Crawford et al., 1989; Blair and Spreen, 1989).: Dutch adaptation NLV, Schmand et al., 1992) Subject is asked to read aloud a list of words with irregular pronunciation	Verbal IQ
Raven's Advanced progressive Matrices (Raven et al., 1998) Subject has to choose which pattern is missing from the overall matrix on the page. With increasing difficulty after each item	Non-verbal IQ and abstract reasoning

verbal fluencies. The level of nonverbal reasoning and non-verbal IQ was also similar in both groups.

4. Discussion

We compared cognitive functioning in various domains between 101 non-psychotic individuals with auditory verbal hallucinations (AVH) and 101 healthy controls matched for age, gender, handedness and education. As expected, task performance of both groups was within the non-pathological range, both groups scored within the top 50% to 25% on all the tests.

The cognitive profile of healthy individuals experiencing AVH was largely similar to that of healthy controls without AVH. However, they underperformed on several domains compared to the controls. Healthy individuals with AVH showed poorer performance on verbal distractibility, inhibition (Stroop interferences), verbal working memory and on tasks tapping lexical access and reasoning (WAIS backward digit-span, vocabulary and similarities, respectively). Thus, it appears that a specific combination of decreased executive functioning, consisting of verbal inhibition, distractibility and verbal working memory in particular, and

a reduced level of verbal intellectual performance is associated with the tendency to hallucinate in the auditory verbal domain.

Although the groups were matched for total years of education and also had similar levels of nonverbal abstract reasoning (nonverbal intelligence estimate), individuals with AVH had significantly lower scores on a verbal estimate for level of intellectual functioning (NART). Lower scores in the AVH group on the vocabulary test and the similarities test may be related to this lower verbal intelligence, since the NART scores correlate highly with WAIS-R Verbal-IQ as well as with WAIS-R Vocabulary scores (Carswell et al., 1997; Uttl, 2002). Interestingly, no difference between both groups was found on phonological and semantic verbal fluency tasks. However, these tasks also assess the ability to generate concepts and to associate which is, apparently, similar in both groups.

Table 4
Characteristics of AVH in the healthy individuals.

Characteristic of auditory verbal hallucinations	Mean (s.d.)	Description of closest anchor
Age at onset	14.5 (14.3)	Mean age of onset voices is 14.5 years old
Frequency (0–6)	3.31 (1.2)	At least once a week
Duration (0–4)	1.47 (0.65)	A few seconds
Location (0–4)	2.41 (1.18)	Voices outside the head. Inside the head may also be present
Loudness (0–4)	1.91 (0.60)	About same loudness as own voice
Beliefs of re-origin (0–4)	3.04 (1.09)	Holds $\geq 50\%$ conviction (but $< 100\%$) that voices originate from external causes
Controllability (0–4)	2.00 (1.6)	Some control over their voices, approximately half of the time
Emotional valence (0–12)	1.25 (2.53)	Hardly any negative content
Total distress (0–8)	0.51 (1.38)	No distress due to voices

Table 3
Frequency of AVH.

Frequency of AVH	Percentage	Frequency (N = 101)
At least once every 3 months	3.0	6
At least once a month	11.4	23
At least once a week	17.3	35
At least daily	11.4	23
At least once per hour	4.5	9
Continuous	2.5	5

Table 5

Comparison of cognitive measures in healthy individuals experiencing AVH and controls.

Measures	AVH (n = 101) Mean (s.d.)	Controls (n = 101) Mean (s.d.)	F(1,200), P-value
Executive function and working memory			
Stroop interference	35.42 (19.66)	28.98 (12.95)	7.55 (0.007 ^a)
Digit-span backward	6.35 (2.06)	7.13 (2.01)	7.46 (0.007 ^a)
Attention			
Digit-span forward	8.75 (1.67)	9.17 (1.88)	2.77 (0.098)
Memory			
CVLT delayed recall	12.45 (2.82)	12.31 (2.56)	0.13 (0.715)
Rey-O delayed recall	19.72 (6.73)	19.41 (6.32)	0.12 (0.731)
Processing speed			
Stroop card 1	46.41 (7.75)	44.96 (7.79)	1.75 (0.188)
Stroop card 2	57.96 (10.66)	56.54 (10.60)	0.90 (0.345)
Spatial ability			
Rey-O copy	32.71 (3.46)	32.74 (2.89)	0.00 (0.947)
Lexical access and abstract reasoning			
Vocabulary test	47.43 (10.83)	53.75 (6.52)	25.29 (<0.0005 ^a)
Similarities test	26.28 (4.33)	28.01 (3.66)	9.44 (0.002 ^a)
Verbal fluency			
Letter fluency total	28.87 (9.02)	28.77 (8.58)	0.01 (0.936)
Semantic fluency total	71.72 (16.43)	68.99 (14.91)	1.53 (0.217)
Intelligence correlates			
Raven's matrices	9.14 (2.14)	9.27 (2.18)	0.18 (0.673)
National adult reading test	84.12 (10.07)	89.37 (7.36)	17.87 (<0.0005 ^a)

^a Significant after Step-Up Hochberg correction for multiple testing.

Speculating, this reduced inhibition and increased distractibility in the verbal domain could render individuals less apt to inhibit irrelevant verbal information. Decreased inhibition could prevent them from focusing on the appropriate information, resulting in reduced performance on executive tasks within the verbal domain. In support of this hypothesis, healthy individuals with AVH were found to have increased levels of positive formal thought disorder as compared to controls without AVH, including peculiar word and sentence usage and peculiar logic (Sommer et al., 2010a,b). Possibly, irrelevant verbal associations which are not adequately inhibited may not be recognized as self-generated and, as such, be attributed to an external source, resulting in an auditory verbal hallucination.

To our knowledge, this is the only study assessing cognitive performance in healthy individuals with AVH. However, other groups have assessed cognitive functioning in related groups, such as healthy individuals with an increased tendency to experience hallucinations, healthy siblings of patients with schizophrenia, individuals at increased genetic risk to develop psychosis and individuals with schizotypal traits. Individuals with high scores on the Launay Slade Hallucination Scale, and thus with a high predisposition to hallucinate were found to have poorer intentional inhibition of memories (Paulik et al., 2007). This is in line with our finding that the presence of AVH is associated with aberrant inhibition. As these individuals were not extensively screened to investigate their hallucinations, it remains unclear whether their experience can be truly classified as, for example, an AVH and whether they have experienced this once or more frequently. Sibling studies show deficits in most cognitive domains, including executive functions, attention, (working) memory, spatial ability, language and performance speed (Snitz et al., 2006; Kuha et al., 2007). For individuals at high genetic risk for psychosis this profile is somewhat similar albeit less extensive: poorer performance on executive function, on global intellectual function, on learning and memory (Byrne et al., 2003). These observed cognitive deficits in siblings and high-risk individuals can be related to the genetic predisposition for schizophrenia in general, rather than to AVH specifically, since there were no differences in spatial ability, processing speed and (working) memory in our sample. Healthy individuals with AVH have significantly higher scores on the Schizotypal

Personality Questionnaire (SPQ; Raine, 1991) (Sommer et al., 2010a,b)), situating them on a continuum somewhere between healthy controls on one end and individuals with schizotypal personality disorder at the other. While they did experience perceptual abnormalities and some degree of suspicion, their social abilities were generally good and they functioned both socially and professionally within the normal range. Studies on cognitive functioning in individuals with schizotypal traits, as measured by the SPQ, may therefore show partial overlap with our results. With respect to verbal subtasks of the WAIS-R, poorer cognitive performance was found in healthy individuals with high scores on the SPQ (Noguchi et al., 2007). Executive working memory was also found to be lower (Matheson and Landon, 2008) in the subjects with schizotypal traits, but not executive functioning (Noguchi et al., 2007). In the latter study a relationship between schizotypal traits and an inductive reasoning component of IQ was found, whereas in our study, this domain appears unaffected. Decreased 'cognitive inhibition' is also often found in individuals with high schizotypy (Beech et al., 1989; Peters et al., 1994; Moritz and Mass, 1997). The fact that our healthy individuals with AVH show cognitive deviations that partly fit these findings of individuals with high schizotypal traits comes as no surprise since these two groups show considerable overlap in features of perceptual abnormalities and the tendency to suspicion.

Our results should be interpreted with caution as they provide no information with respect to causality. It could be hypothesized that aberrant verbal and executive functioning underlies the predisposition to hallucinate, but alternatively the experience of AVH may lower performance on these specific tasks. As most participants did not experience AVH during cognitive testing, the first explanation appears stronger, although the second cannot be ruled out. Another limitation is that the individuals that participated in this study are a highly selected group as only participants who were willing to visit us, and in the case of the healthy individuals with AVH, who were willing to talk openly about their AVH, could be included. This might also be the reason for the overrepresentation of women in the sample, although auditory hallucinations are more prevalent in women than in men (Rector and Seeman, 1992).

In conclusion, while their cognitive performance is within the normal range and thus not clinically impaired, healthy individuals experiencing AVH do show reduced levels of executive functioning and verbal (intellectual) performance. This deviation compared to individuals without AVH suggests that experiencing AVH is directly associated with difficulties in the inhibition of irrelevant verbal information.

Role of funding source

This work was supported by the Nederlandse Wetenschappelijke Organisatie (Dutch Scientific Research Organisation) (grant number 916.56.172); the Nederlandse Wetenschappelijke Organisatie had no further role in study design; in the collection analysis and interpretation of data; in the writing of the report; and in the decision to submit the paper for publication.

Contributors

Authors FB and MvZ designed the study. Author KD contributed to the design of the study, collected the subjects, organized the acquisition of data, analyzed the data and wrote the manuscript. All authors contributed to interpretation of the results and have contributed to and approved the final manuscript.

Conflict of interest

The authors report no conflicts of interest.

Acknowledgments

The authors wish to thank Willemijn van Gastel, Anne Lotte Meijering, Eveline Hendriks and Monique te Beek for their time and effort in collecting the data.

References

- Andreasen, N.C., Flaum, M., Arndt, S., 1992. The Comprehensive Assessment of Symptoms and History (CASH). An instrument for assessing diagnosis and psychopathology. Arch. Gen. Psychiatry. 49 (8), 615–623.
- Beech, A., Baylis, G.C., Smithson, P., Claridge, G., 1989. Individual differences in schizotypy as reflected in measures of cognitive inhibition. Br J Clin Psychol. 28, 117–129.

- Benjamini, Y., Hochberg, Y., 1995. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J. R. Stat. Soc. B* 57 (1), 289–300.
- Berenbaum, H., Kerns, J.G., Vernon, L.L., Gomez, J.J., 2008. Cognitive correlates of schizophrenia signs and symptoms: III. Hallucinations and delusions. *Psychiatry Res.* 159, 163–166.
- Berman, I., Viegner, B., Merson, A., Allan, E., Pappas, D., Green, A., 1997. Differential relationships between positive and negative symptoms and neuropsychological deficits in schizophrenia. *Schizophr. Res.* 25, 1–10.
- Blair, R.J., Spreen, O., 1989. Predicting premorbid IQ: a revision of the national adult reading test. *The Clinical Neuropsychologist* 3 (2), 129–136.
- Brébion, G., David, A.S., Jones, H.M., Ohlsen, R., Pilowsky, L.S., 2007. Temporal context discrimination in patients with schizophrenia: associations with auditory hallucinations and negative symptoms. *Neuropsychologia* 45, 817–823.
- Byrne, M., Clafferty, B.A., Cosway, R., Grant, E., Hodges, A., Whalley, H.C., Lawrie, S.M., Cunningham Owens, D.G., Johnstone, E.C., 2003. Neuropsychology, genetic liability, and psychotic symptoms in those at high risk of schizophrenia. *J. Abnorm. Psychol.* 112, 38–48.
- Carswell, L.M., Graves, R.E., Snow, W.G., Tierney, M.C., 1997. Postdicting Verbal IQ of elderly individuals. *J. Clin. Exp. Neuropsychol.* 19 (6), 914–921.
- Crawford, J.R., Parker, D., Stewart, L., Besson, J., De Lacey, G., 1989. Prediction of WAIS IQ with the National Adult Reading Test: cross-validation and extension. *Br. J. Clin. Psychol.* 28, 267–273.
- Daalman, K., Boks, M.P.M., Diederer, K.M.J., de Weijer, A.D., Blom, J.D., Kahn, R.S., Sommer, I.E.C., 2011. The same or different? A phenomenological comparison of auditory verbal hallucinations in healthy and psychotic individuals. *J. Clin. Psychiatry* 72 (3), 320–325.
- De Renzi, E., Vignolo, L.A., 1962. The token test: a sensitive test to detect disturbances in aphasics. *Brain* 85, 665–678.
- Delis, D., Kramer, J., Kaplan, E., Ober, B., 1987. California Verbal Learning Test. The Psychological Corporation, San Antonio, TX.
- Diederer, K.M.J., Daalman, K., de Weijer, A.D., Neggers, S.F., van Gastel, W., Blom, J.D., Kahn, R.S., Sommer, I.E.C., 2011. Auditory hallucinations elicit similar brain activation in psychotic and nonpsychotic individuals. *Schizophr. Bull.* (Apr 28 [Epub ahead of print]).
- First, M.B., Spitzer, R.L., Gibbon, M., Williams, J.B.W., 1995. The structured clinical interview for DSM-III-R personality-disorders (Scid-II).1 Description. *J. Pers. Disord.* 9, 83–91.
- Frith, C.D., Done, D.J., 1988. Toward a neuropsychology of schizophrenia. *Br. J. Psychiatry* 153, 437–443.
- Haddock, G., McCarron, J., Tarrier, N., Faragher, E.B., 1999. Scales to measure dimensions of hallucinations and delusions: the psychotic symptom rating scales (PSYRATS). *Psychol. Med.* 29 (4), 879–889.
- Hoffman, R.E., Rapaport, J., Mazure, C.M., Quinlan, D.M., 1999. Selective speech perception alterations in schizophrenic patients reporting hallucinated “voices”. *Am. J. Psychiatry* 156, 393–399.
- Kaplan, E.F., Goodglass, H., Weintraub, S., 1983. The Boston Naming Test 2nd edition. Lea & Febiger, Philadelphia.
- Knight, J., Kaplan, E., 2003. The Handbook of Rey–Osterrieth Complex Figure Usage: Clinical and Research Applications. Psychological Assessment Resources, Inc., Lutz, FL.
- Kuha, A., Tuulio-Hendriksson, A., Eerola, M., Perälä, J., Suvisaari, J., Partonen, T., Lönngqvist, J., 2007. Impaired executive performance in healthy siblings of schizophrenia patients in a population-based study. *Schizophr. Res.* 92, 142–150.
- Lezak, M.D., Loring, D.W., Howieson, D.B., 2004. Neuropsychological Assessment 4th Ed. Oxford University Press, New York.
- Matheson, S., Landon, R., 2008. Schizotypal traits impact upon executive working memory and aspects of IQ. *Psychiatr Res.* 159, 207–214.
- Moritz, S., Mass, R., 1997. Reduced cognitive inhibition in schizotypy. *Br J Clin Psychol.* 36, 365–376.
- Mulder, J., Dekker, R., Dekker, P., 1996. Handleiding Verbale Leer en Geheugen Test. Swets & Zeitlinger, Lisse.
- Noguchi, H., Hori, H., Kunugi, H., 2007. Schizotypal traits and cognitive functioning in healthy adults. *Psychiatr Res.* 161, 162–169.
- Paulik, G., Badcock, J.C., Maybery, M.T., 2007. Poor intentional inhibition in individuals predisposed to hallucinations. *Cogn. Neuropsychiatry* 12 (5), 457–470.
- Peters, E.R., Pickering, A.D., Hemsley, D.R., 1994. ‘Cognitive inhibition’ and positive symptomatology in schizotypy. *Br J Clin Psychol.* 33, 33–48.
- Raine, A., 1991. The SPQ: a scale for the assessment of schizotypal personality based on DSM-III-R criteria. *Schizophr Bull.* 17, 555–564.
- Raven, J., Raven, J.C., Court, J.N., 1998. Manual for Raven’s Progressive Matrices and Vocabulary Scales Section 4: the Advanced Progressive Matrices. Harcourt Assessment, San Antonio, TX.
- Rector, N.A., Seeman, M.V., 1992. Auditory hallucinations in women and men. *Schizophr Res.* 7, 233–236.
- Schmand, B., Lindeboom, J., Harskamp, F., 1992. NLV, Nederlandse Leestest voor Volwassenen. Handleiding. Swets & Zeitlinger, Lisse.
- Seal, M.L., Aleman, A., McGuire, P.K., 2004. Compelling imagery, unanticipated speech and deceptive memory: neurocognitive models of auditory verbal hallucinations in schizophrenia. *Cogn. Neuropsychiatry* 9 (1/2), 43–72.
- Snitz, B.E., MacDonald, A.W., Carter, C.S., 2006. Cognitive deficits in unaffected first-degree relatives of schizophrenia patients: a meta-analytic review of putative endophenotypes. *Schizophr. Bull.* 32 (1), 179–194.
- Sommer, I.E., Daalman, K., Rietkerk, T., Diederer, K.M., Bakker, S., Wijkstra, J., Boks, M.P.M., 2010a. Healthy individuals with auditory verbal hallucinations; who are they? Psychiatric assessments of a selected sample of 103 subjects. *Schizophr. Bull.* 36 (3), 633–641.
- Sommer, I.E., Derwort, A.M.C., Daalman, K., de Weijer, A.D., Liddle, P.F., Boks, M.P.M., 2010b. Formal thought disorder in non-clinical individuals with auditory verbal hallucinations. *Schizophr. Res.* 118, 140–145.
- SPSS, 2006. SPSS 15.0 for Windows Version 15.0 Chicago. SPSS Inc, IL.
- Stroop, J.R., 1935. Studies of interference in serial-verbal reaction. *J. Exp. Psychol.* 18, 643–663.
- Tien, A.Y., 1991. Distributions of hallucinations in the population. *Soc. Psychiatry Psychiatr. Epidemiol.* 26, 287–292.
- Uttl, B., 2002. North American Adult Reading Test: age norms, reliability, and validity. *J. Clin. Exp. Neuropsychol.* 24 (8), 1123–1137.
- Verdoux, H., van Os, J., 2002. Psychotic symptoms in non-clinical populations and the continuum of psychosis. *Schizophr. Res.* 54, 59–65.
- Verkammen, E., de Haan, E.H.F., Aleman, A., 2008. Hearing a voice in the noise: auditory hallucinations and speech perception. *Psychol. Med.* 38, 1177–1184.
- Waters, F.A.V., Badcock, J.C., Maybery, M.T., Michie, P.T., 2003. Inhibition in schizophrenia: associations with auditory hallucinations. *Schizophr. Res.* 62, 275–280.
- Waters, F.A.V., Woodward, T., Allen, P., Aleman, A., Sommer, I.E.C., 2010. Self-recognition deficits in schizophrenia patients with auditory hallucinations: a meta-analysis of the literature. *Schizophr Bull.* 2010 Dec 8. [Epub ahead of print].
- Wechsler, D., 1997. Wais-III Administration and Scoring Manual. The Psychological Corporation, San Antonio, TX.
- Westfall, P.H., Young, S.S., 1993. Resampling-based Multiple Testing: Examples and Methods for P-value Adjustment. J. Wiley, New York.