



Microplastic contamination in the San Francisco Bay, California, USA



Rebecca Sutton ^{a,*}, Sherri A. Mason ^b, Shavonne K. Stanek ^{a,1}, Ellen Willis-Norton ^{a,2},
Ian F. Wren ^c,Carolynn Box ^d

^a San Francisco Estuary Institute, 4911 Central Avenue, Richmond, CA 94804, USA

^b State University of New York at Fredonia, 280 Central Avenue, Science Complex 340, Fredonia, NY 14063, USA

^c San Francisco Baykeeper, 1736 Franklin Street, Suite 800, Oakland, CA 94612, USA

^d The 5 Gyres Institute, 3131 Olympic Blvd, Suite 302, Santa Monica, CA 90404, USA

ARTICLE INFO

Article history:

Received 8 December 2015

Received in revised form 13 May 2016

Accepted 28 May 2016

Available online 8 June 2016

Keywords:

Plastic pollution

Marine debris

San Francisco Bay

Microplastic

Microbeads

Wastewater

ABSTRACT

Despite widespread detection of microplastic pollution in marine environments, data describing microplastic abundance in urban estuaries and microplastic discharge via treated municipal wastewater are limited. This study presents information on abundance, distribution, and composition of microplastic at nine sites in San Francisco Bay, California, USA. Also presented are characterizations of microplastic in final effluent from eight wastewater treatment plants, employing varying treatment technologies, that discharge to the Bay. With an average microplastic abundance of 700,000 particles/km², Bay surface water appears to have higher microplastic levels than other urban waterbodies sampled in North America. Moreover, treated wastewater from facilities that discharge into the Bay contains considerable microplastic contamination. Facilities employing tertiary filtration did not show lower levels of contamination than those using secondary treatment. As textile-derived fibers were more abundant in wastewater, higher levels of fragments in surface water suggest additional pathways of microplastic pollution, such as stormwater runoff.

© 2016 Elsevier Ltd. All rights reserved.

1. Introduction

While plastic pollution of the marine environment has been reported for decades, only recently have estuaries and freshwater systems been a focal point of similar studies (Dubaish and Liebezeit, 2013; Eriksen et al., 2013; Castañeda et al., 2014; Free et al., 2014; Yonkos et al., 2014; Davis and Murphy, 2015). A key component of this pollution, microplastic describes fragments of plastic that are smaller than 5 mm (Thompson et al., 2009; Masura et al., 2015). Sources of microplastic to the environment include microbeads used in personal care products, pre-production pellets used as precursors to manufacture plastic products, fibers derived from clothes and fabrics made with synthetic materials (e.g., polyester and acrylic) or fishing line, fragments from the photodegradation of larger plastic items, and plastic foam particles from polystyrene products or cigarette filters (Fendall and Sewell, 2009; Browne et al., 2011; Eriksen et al., 2013; Free et al., 2014; van Franeker and Law, 2015). Microplastic can enter the aquatic environment through wind advection, stormwater runoff, or illegal dumping of plastic materials (Eriksen et al., 2013). Additionally, both microbeads

from personal care products and fibers from synthetic clothing can be washed down the drain and enter wastewater treatment plants, where their small size, buoyancy, and lack of reactivity limits removal, resulting in release via treated wastewater (Browne et al., 2011; NYS OAG, 2015).

Microplastic particles pose risks to wildlife because the particles may be mistaken for food and ingested (Wright et al., 2013). The particles are also small enough that they can be ingested by planktonic organisms and other filter feeders (Browne et al., 2008; Cole et al., 2013). The hydrophobicity and high surface area to volume ratio of microplastic particles leads to sorption of persistent organic pollutants such as polycyclic aromatic hydrocarbons (Teuten et al., 2007). Organisms that ingest microplastic particles may thus receive higher doses of sorbed contaminants, potentially causing additional harm (Wright et al., 2013). Ingestion of microplastic can block the digestive tract, reduce growth rates, block enzyme production, lower steroid hormone levels, affect reproduction, and may lead to greater exposure to plastic additives with toxic properties (Wright et al., 2013).

Despite widespread detection of microplastic pollution in the marine environment, data describing microplastic abundance in urban estuaries and microplastic discharge via treated municipal wastewater are limited. This initial, screening study characterized microplastic in treated wastewater effluent from eight facilities employing a range of treatment technologies and discharging to San Francisco Bay, hereafter referred to as the Bay. Treated wastewater is considered an important

* Corresponding author.

E-mail address: RebeccaS@sfei.org (R. Sutton).

¹ Present address: Oberlin College, Oberlin, OH 44074 USA.

² Present address: University of California at Santa Cruz, 100 Shaffer Rd., Santa Cruz, CA 95060 USA.

pathway for microplastic to enter receiving waters, but only a few studies of this matrix are available (Carr et al., 2016; Mason et al., in review). In addition, this study provides data on microplastic in surface waters of the Bay, the largest estuary on the west coast of North America, which is surrounded by a dense urban population and drains roughly 40% of the waters of California.

2. Materials and methods

2.1. Wastewater

Treated wastewater is discharged to San Francisco Bay from more than 30 different discharge locations. Eight facilities, providing approximately 60% of measured wastewater flows directly to the Bay, permitted researcher access to final effluent sinks or other available ports, allowing us to collect samples. Samples of microplastic discharged from wastewater treatment plants were collected by passing flows of final, treated effluent through 8-in. diameter stacked Tyler sieves with 0.355 mm and 0.125 mm stainless steel mesh, typically for 2 hours during each facility's peak flow. The 0.125 mm mesh has been found to be particularly useful for retention of microbeads discharged to the sewer via use of personal care products (Napper et al., 2015; Carr et al., 2016). A single set of two samples, differentiated by sieve mesh size, was collected in the fall of 2014 at each of the eight facilities. Facilities participated voluntarily, and were selected based on multiple factors, including higher discharge levels, geographic diversity, and range of treatment technologies (secondary vs. tertiary filtration; Table 1). Rate of flow at the point of collection was measured before and after each sample was obtained (to ensure consistency), allowing calculation of number of particles per volume of treated wastewater. Each facility provided the 24-h discharge flow rate for the day of sample collection, allowing estimation of the number of particles discharged to the Bay per day.

In order to remove labile organic material, samples were processed via a wet peroxide oxidation (WPO) based upon a National Oceanic and Atmospheric Administration method (Masura et al., 2015), which

has been tested to ensure that the most common plastic materials survive. Briefly, samples were reacted with a 30% hydrogen peroxide solution in the presence of an iron (II) catalyst in order to oxidize natural organic material, leaving the synthetic plastic material behind. Wastewater samples were processed as individual samples according to the collected size classification (i.e., 0.125–0.355 mm or >0.355 mm).

After processing, samples were once again filtered through a stacked sieve set (0.355 mm and 0.125 mm) and rinsed using deionized (DI) water into petri dishes. Given their density relative to that of DI water and most natural materials, floating particles within this medium are assumed to be plastic, a common technique within this field of research (Hidalgo-Ruz et al., 2012; Rocha-Santos and Duarte, 2014). Using a dissection microscope, plastic particles were removed, enumerated, and categorized into five classifications: fragment, pellet (spherical particle), fiber/line, film or foam (Free et al., 2014; McCormick et al., 2014). While instrumental analysis methods such as infrared or Raman spectroscopy are necessary for polymeric identification (i.e., polyethylene versus polypropylene), numerous studies have employed only visual identification for microplastic classification (e.g., Bond et al., 2014; Lavers et al., 2014; Devriese et al., 2015; Rochman et al., 2015; Romeo et al., 2015; Fossia et al., 2016; Hammer et al., 2016; Miranda and Carvalho-Souza, 2016; Nicolau et al., 2016; Peters and Bratton, 2016). Given the source (i.e., wastewater), fibers obtained in this processing would presumably be anthropogenic and derived from textiles, though a portion of fibers observed in wastewater may not be plastic, instead derived from other anthropogenic sources (Remy et al., 2015; Nirmela Arsem, personal communication).

2.2. Surface water

Single surface water microplastic samples were collected from each of nine sites in San Francisco Bay over the course of 2 days in January 2015 (Fig. 1). Central and southern portions of the Bay contain higher levels of litter, including macroplastic debris, than northern stretches, and were the focus of this study (Rubissow-Okamoto, 2014). During sample collection, conditions were calm: the sea state on the Beaufort

Table 1
Microplastic particles present in treated wastewater, and estimates of discharge per liter and per day.

| Wastewater treatment plant | Flow ^a (MLD) | Highest level of treatment | Size category (mm) | No. plastic particles by type | | | | | | No. plastic particles | |
|--|----------------------------|-------------------------------|-----------------------|-------------------------------|--------|-------|------|------|-------|---------------------------|----------------------|
| | | | | Fragment | Pellet | Fiber | Film | Foam | Total | Per liter ^b | Per day ^a |
| San José-Santa Clara | 310 | Tertiary filtration | 0.125–0.354 | 0 | 0 | 26 | 0 | 0 | 26 | 0.047 | 15,000,000 |
| | | | ≥0.355 | 0 | 0 | 33 | 0 | 0 | 33 | | |
| | | | total | 0 | 0 | 59 | 0 | 0 | 59 | | |
| East Bay Municipal Utilities District (EBMUD) | 170 | Secondary | 0.125–0.354 | 1 | 0 | 11 | 1 | 0 | 13 | 0.071 | 12,000,000 |
| | | | ≥0.355 | 7 | 0 | 5 | 3 | 0 | 15 | | |
| | | | total | 8 | 0 | 16 | 4 | 0 | 28 | | |
| Palo Alto | 76 | Tertiary filtration | 0.125–0.354 | 3 | 0 | 24 | 0 | 0 | 27 | 0.13 | 9,600,000 |
| | | | ≥0.355 | 8 | 0 | 23 | 2 | 0 | 33 | | |
| | | | total | 11 | 0 | 47 | 2 | 0 | 60 | | |
| Central Contra Costa | 110 | Secondary | 0.125–0.354 | 21 | 0 | 28 | 0 | 0 | 49 | 0.072 | 8,100,000 |
| | | | ≥0.355 | 5 | 0 | 10 | 0 | 0 | 15 | | |
| | | | total | 26 | 0 | 38 | 0 | 0 | 64 | | |
| Fairfield-Suisun | 45 | Tertiary Filtration | 0.125–0.354 | 2 | 0 | 43 | 0 | 0 | 45 | 0.092 | 4,100,000 |
| | | | ≥0.355 | 2 | 0 | 50 | 2 | 0 | 54 | | |
| | | | total | 4 | 0 | 93 | 2 | 0 | 99 | | |
| East Bay Dischargers Association (EBDA) | 190 | Secondary | 0.125–0.354 | 1 | 0 | 11 | 0 | 0 | 12 | 0.022 | 4,100,000 |
| | | | ≥0.355 | 1 | 0 | 9 | 0 | 0 | 10 | | |
| | | | total | 2 | 0 | 20 | 0 | 0 | 22 | | |
| San Mateo | 31 | Tertiary filtration | 0.125–0.354 | 20 | 0 | 24 | 0 | 3 | 47 | 0.064 | 2,000,000 |
| | | | ≥0.355 | 7 | 0 | 21 | 3 | 0 | 31 | | |
| | | | total | 27 | 0 | 45 | 3 | 3 | 78 | | |
| San Francisco Airport Sanitary (SFO) | 2.3 | Secondary | 0.125–0.354 | 5 | 0 | 49 | 0 | 0 | 54 | 0.19 | 460,000 |
| | | | ≥0.355 | 4 | 0 | 42 | 0 | 1 | 47 | | |
| | | | total | 9 | 0 | 91 | 0 | 1 | 101 | | |
| Total count | | | total | 87 | 0 | 409 | 11 | 4 | 511 | | |
| Percentage by type | | | total | 17% | 0% | 80% | 2% | 1% | 100% | | |

^a Measured discharge on day of sample collection, used to calculate plant discharge per day.

^b Calculated using average flow rate at point of sample collection, see Supplementary Content.




Fig. 1. Estimated abundance of microplastic particles in surface water at nine sites in San Francisco Bay. Site numbers correspond to those listed in Table 2; circles are located at trawl midpoints.

Scale remained between 0 and 2 for all sites. Samples were collected using a Manta Trawl with a rectangular opening 16 cm high by 61 cm wide, and a 3 m long 333 μ m mesh net with a 30 \times 10 cm² collecting bag, a sample collection technique now in common use (e.g., Eriksen et al., 2013; Free et al., 2014; Masura et al., 2015). Sampling took place over the course of 30 min at each site, with tow speeds from 2 to 4 knots and the vessel maintaining a consistent heading. Coordinates were recorded at the start and finish of each trawl to establish tow length; this length, multiplied by the width of the trawl, provided the surface area sampled, allowing calculation of standardized values per square kilometer.

Most samples were collected during flood tides, with two samples, at sites 5 and 9, collected during slack tides. For the central Bay, where tidal exchanges are significant, flood tides may result in reduction of microplastic abundance due to the addition of ocean water with lower levels of urban contaminants; therefore, microplastic abundances measured in the central Bay should be considered lower-bound estimates. Abundances measured in southern stretches of the Bay, where oceanic exchange is significantly reduced, are not expected to be significantly affected by tidal flushing. All samples were placed in sample jars and preserved with 70% isopropyl alcohol.

Nine small fish were collected as accidental by-catch at one site (6). Although fish collection was not planned, the abundance of microplastic in these fish was determined after thorough rinsing to remove external contaminants; results are provided in Supplementary Content.

Surface water samples then went through a pre-processing stage where each sample was visually inspected to remove any obvious larger (i.e., > 1 mm) plastic debris, as well as any larger vegetative debris (such as leaves, twigs, seeds, feathers, and insects), which would be difficult to

chemically process (as described below). All materials were thoroughly rinsed prior to removal and vegetative debris was discarded. Samples were then subjected to the WPO treatment to remove seston and other labile organic matter, sieved into three different size classifications using 8-in. diameter stacked Tyler sieves of 0.355 mm, 1.00 mm and 4.75 mm stainless steel mesh, and rinsed thoroughly with deionized (DI) water. After processing, all samples and all size classifications were analyzed using a dissection microscope and plastic particles within were removed, enumerated, and categorized as described above.

A single field blank sample, created by rinsing sieves employed during both Bay and wastewater sample collection, and four lab blank samples in which DI water was stored within sample containers for periods of 1–14 days, were processed in the same manner as all other samples. While the field blank sample revealed the presence of four fibers measuring less than 1 mm, the lab blank samples were not found to have any microplastic contamination. The fibers within the field blank are thus most likely derived from sample cross-contamination (e.g., particles that adhered to the mesh bag or the sieves themselves), as lab blank samples would indicate that contamination from the containers, lab or processing was negligible. Nevertheless, field blank contents were subtracted from all sample counts.

3. Results

3.1. Wastewater

The eight San Francisco Bay wastewater treatment plants discharged an average of 0.086 microplastic particles per liter (or 0.33 particles per gallon) and 7 million microplastic particles per day (Table 1). These levels were higher than those found in effluent from nine municipal wastewater treatment facilities in the Midwest and Northeast U.S. analyzed using the same method (0.050 particles per liter and 2 million particles per day; Mason et al., in review), as well as eight southern California facilities analyzed with a different analytical method (all facilities discharged <0.001 particles per liter; Carr et al., 2016). In total, the eight Bay facilities, representing approximately 60% of treated wastewater flows to San Francisco Bay, discharged 56 million microplastic particles per day. If levels of microplastic in effluent from these eight facilities are assumed to be generally representative of the region, an estimated 90 million microplastic particles per day may be discharged into San Francisco Bay. This estimate does not include contributions of microplastic discharged by treatment plants into the Sacramento-San Joaquin River Delta, which ultimately drains to the Bay.

Fibers were the dominant form of particulate pollution in effluent, followed by fragments (Table 1). The San José-Santa Clara Regional Wastewater Facility, which employs tertiary granular filtration, discharged only fibers during sample collection. However, overall there were no consistent differences between the proportions of fibers and fragments discharged by facilities employing secondary treatment only (EBMUD, Central Contra Costa, EBDA, SFO) versus those that included tertiary filtration as well (San José-Santa Clara, Palo Alto, Fairfield-Suisun, San Mateo). Likewise, plants employing tertiary granular filtration did not display consistently lower concentrations of overall microplastic per liter than those employing only secondary treatment technologies.

Comparable numbers of smaller (0.125–0.355 mm) and larger (>0.355 mm) microplastic particles were discharged by most facilities (Table 1). The Central Contra Costa Sanitary District facility was an exception, discharging three times more smaller particles than larger ones.

3.2. Surface water

All nine surface water samples contained microplastic pollution (Table 2). Average abundance for Bay samples was 700,000 \pm 600,000 particles/km² (range: 15,000–2,000,000 particles/km²), higher than average measurements from the Great Lakes (Eriksen et al., 2013),

Table 2

Count, tow length, and abundance of microplastic particles in surface waters at nine sites within San Francisco Bay.

| Site | Total count | Tow length, meters | Abundance, particles/km ² |
|-----------------------------------|-------------|--------------------|--------------------------------------|
| <i>Central San Francisco Bay</i> | | | |
| 1 | 26 | 2855 | 15,000 |
| 2 | 107 | 1708 | 100,000 |
| 3 | 994 | 1776 | 920,000 |
| 4 | 188 | 1655 | 190,000 |
| <i>Southern San Francisco Bay</i> | | | |
| 5 | 438 | 2302 | 310,000 |
| 6 | 1192 | 2017 | 970,000 |
| 7* | 3641 | 2959 | 2,000,000 |
| 8* | 1247 | 2799 | 730,000 |
| 9* | 1665 | 2643 | 1,000,000 |

* Samples included considerable bulky vegetation.

the Chesapeake Bay (Yonkos et al., 2014), and the Salish Sea (Davis and Murphy, 2015). All samples contained fragments, fibers, and to a lesser extent pellets in the smallest (0.355–0.999 mm) size class (see Supplementary Content). One site (1) lacked foamed microplastic particles in the smallest size class, while another (site 2) lacked films in the smallest size class. Larger size classes were most often dominated by fragments and fibers (Table 3). Preliminary data from nine small, prey fish obtained as accidental by-catch suggested high levels of microplastic contamination, particularly by particles in the 0.355–0.999 mm size class (see Supplementary Content).

Sites in the southern portion of San Francisco Bay had higher average microplastic abundance, 1,000,000 particles/km², than sites in the central Bay, which averaged 310,000 particles/km². Flood tides occurring during sampling in the central Bay could contribute to reduced abundances measured at these sites. Southern sites typically contained higher levels of small fragments (Table 3); three of these samples also included large amounts of bulky vegetation (Table 2).

4. Discussion

Overall levels of microplastic pollution measured in this initial, screening study were greater than comparable measurements available for other urbanized areas of the U.S., including treated wastewater from municipal wastewater treatment facilities located in other parts of the U.S. (Carr et al., 2016; Mason et al., in review) and surface waters of the Great Lakes, Chesapeake Bay, and the Salish Sea (Eriksen et al., 2013; Yonkos et al., 2014; Davis and Murphy, 2015). Bay Area wastewater facilities investigated here typically serve large populations that

have implemented significant water conservation measures due to severe drought, in contrast to the facilities from other regions for which consistently measured data are available (Mason et al., in review); these differences provide a potential explanation for the increased concentrations of microplastic particles in Bay Area wastewater, as the same overall amounts of urban contaminants would be more concentrated if released with lesser amounts of water. The apparent elevation of San Francisco Bay surface water microplastic pollution can be at least partially explained by a dense urban population surrounding a small body of water with limited interchange with the Pacific Ocean.

Southern Bay levels of microplastic were generally higher than those of the central San Francisco Bay. A similar regional pattern has been observed for a number of contaminants derived largely from treated wastewater (e.g., Klosterhaus et al., 2013). Surface waters in the southern Bay receive a large volume of treated wastewater and urban stormwater, have the highest hydraulic residence time relative to other portions of the Bay, and experience the least amount of dilution.

However, the observation that small fragments drove the higher microplastic levels measured at southern Bay sites, while wastewater discharges were primarily composed of fibers, suggests that treated wastewater was not the only source of microplastic pollution for southern stretches of the Bay. Instead, microplastic from other pollution pathways, such as stormwater, and in situ processes including the fragmentation of larger plastic debris, in combination with the long residence times, may have contributed to higher levels of microplastic contamination.

An additional qualitative observation was that some of the largest counts of plastic were associated with Bay samples containing substantial quantities of vegetation (Table 2). The potential for vegetation to entrain microplastic merits exploration in future studies. Of particular concern, the presence of surface vegetation can result in more concentrated feeding activity nearby. If microplastic levels are higher in these zones, wildlife may be more likely to ingest these particles.

Comparison of wastewater treatment plants employing typical secondary treatment versus those equipped with additional tertiary filtration did not indicate a significant effect on microplastic particle discharge or distribution of particle types. Additional monitoring is needed to explore this finding, which is consistent with observations from a study of New York State treatment facilities focusing exclusively on plastic pellets (NYS OAG, 2015). Assessment of microplastic content in associated influent and biosolids would better elucidate the impacts of different treatment types on microplastic particles in wastewater. A study exploring microplastic levels at different points in the wastewater treatment train found considerable removal via skimming during initial stages of treatment (Carr et al., 2016). Nevertheless, the limited data currently available suggest that granular tertiary filtration may not be an effective means of controlling microplastic pollution.

Fibers were the dominant particle type found in treated wastewater from Bay Area facilities, a finding common in similar microplastic analyses of wastewater in other locations (e.g., Browne et al., 2011; Mason et al., in review). An alternative study, which utilized a different method for effluent processing and analysis, did not detect anthropogenic fibers, only microbially derived detritus (Carr et al., 2016). A recent examination of artificial fibers found in the digestive tracts of aquatic invertebrates suggests some portion of these fibers may be derived from cellulose rather than plastic (Remy et al., 2015). Some cellulose-derived fibers can survive the wet peroxide oxidation process that all samples were subjected to in this study (Nirmela Arsem, personal communication); as fibers were not subjected to Raman nor FTIR spectroscopy to confirm their identity, it is possible that the presence of anthropogenic, cellulose-derived fibers resulted in overestimates of the overall levels of plastic fibers in these samples. However, given previous studies indicating high levels of plastic fibers in synthetic clothing wash water as well as in wastewater samples subjected to greater levels of spectroscopic examination (Browne et al., 2011), synthetic plastic is likely to be a dominant source of fibers in the wastewater examined in this study.

Table 3

Average abundance and type of particles in three size classes in central and southern San Francisco Bay surface water samples.

| Central San Francisco Bay | 0.355–0.999 mm | 1.000–4.749 mm | ≥4.75 mm | % of Total |
|-----------------------------|----------------|----------------|----------|------------|
| Fragment | 68,000 | 35,000 | 2100 | 34% |
| Pellet | 3100 | 970 | 0 | 1% |
| Fiber | 80,000 | 67,000 | 1200 | 48% |
| Film | 8200 | 22,000 | 3500 | 1% |
| Foam | 2300 | 12,000 | 460 | 5% |
| Total Count/km ² | 160,000 | 140,000 | 7200 | |
| % of Total | 53% | 45% | 2% | |
| Southern San Francisco Bay | 0.355–0.999 mm | 1.000–4.749 mm | ≥4.75 mm | % of Total |
| Fragment | 450,000 | 150,000 | 5400 | 60% |
| Pellet | 17,000 | 2500 | 0 | 2% |
| Fiber/ine | 140,000 | 86,000 | 2800 | 22% |
| Film | 25,000 | 37,000 | 6700 | 7% |
| Foam | 35,000 | 52,000 | 2300 | 9% |
| Total Count/km ² | 670,000 | 330,000 | 17,000 | |
| % of Total | 66% | 32% | 2% | |

Of note, fibers are more reliably identified through visual inspection as plastic than fragments (Lenz et al., 2015).

Multi-colored, spherical plastic pellets in the size class <1 mm, likely derived from rinse-off personal care products, were detected at low levels in all Bay sites, but in none of the treated wastewater samples. While the absence of these clearly identifiable pellets in treated wastewater was somewhat unexpected, it does not indicate a lack of personal care product-derived contamination. Most of the microplastic particles in personal care products are rough and irregular in shape and therefore classified as fragments (Napper et al., 2015; Carr et al., 2016), with less than 10% consisting of spherical pellets (unpublished data, S.A. Mason). It is likely that a significant portion of the microplastic fragments detected in treated wastewater samples in this study were derived from personal care products containing microbeads.

5. Conclusion

The results of this initial, screening study indicate that microplastic contamination, a global concern, may be higher in San Francisco Bay than in other urban areas in North America for which data are available, including the Great Lakes (Eriksen et al., 2013), the Chesapeake Bay (Yonkos et al., 2014), and the Salish Sea (Davis and Murphy, 2015). Effluent samples from Bay Area treatment facilities also showed higher levels of contamination than seen in other facilities in the U.S. (Carr et al., 2016; Mason et al., in review).

Results from this study indicate the need for method development and standardization to assess microplastic found in common pollution pathways. The increased proportion of fragments over fibers in Bay water samples relative to treated wastewater suggested that other pathways, such as stormwater, may be important contributors to microplastic pollution in the Bay. As yet, there are no established methods for measuring microplastic in stormwater discharges; until such methods are developed, a hypothesis about the relative contributions of stormwater versus wastewater to overall microplastic pollution in receiving waters cannot be tested. The method employed in this study to characterize effluent was originally developed for ambient receiving waters and could be further refined, particularly given the wide range of anthropogenic fibers that may be found at higher concentrations in wastewater. In addition, 24-hour effluent samples could provide a more comprehensive picture of microplastic pollution in treated wastewater, in particular because peak personal care product use follows distinct diurnal patterns.

Ultimately, concerns about microplastic pollution are driven by potential impacts to wildlife or humans. Recent detection of microplastic, including fibers, within sport fish from coastal California as well as Indonesia (Rochman et al., 2015) indicates exposure is occurring; at this time, studies linking microplastic exposure to adverse impacts in controlled laboratory settings have not resulted in development of specific aquatic or tissue-based toxicity thresholds. By-catch prey fish collected at a single site in this study also contained microplastic, especially fibers (Supplementary Content). Of note, most microplastic monitoring in wildlife focuses on microorganisms, invertebrates, or sport fish (e.g., Wright et al., 2013; Rochman et al., 2015), with relatively few data available for smaller, prey fish such as those examined here. Additional study of small, prey fish species is recommended to develop an understanding of microplastic exposure and potential impacts throughout the food web.

Acknowledgments

Funding for this study was provided by the Regional Monitoring Program for Water Quality in San Francisco Bay (Contribution No. 769), with in-kind contributions from San Francisco Baykeeper and the 5 Gyres Institute. The authors would like to acknowledge participants from the following wastewater treatment plants: Palo Alto Regional Water Quality Control Plant, Fairfield Suisun Sewer District, Central

Contra Costa Sanitary District, East Bay Municipal Utility District, East Bay Dischargers Authority, City of San Mateo Wastewater Treatment Plant, San José/Santa Clara Regional Wastewater Facility, and San Francisco International Airport Sanitary Waste Treatment Plant and Reclaimed Water Facility. The authors acknowledge Pete Kauhanen for preparation of the GIS image, Rusty Holleman for Bay wastewater flow data, and Claire Boudreaux for assistance with visualization. Additional thanks to Oberlin College, Career Center for providing intern support. Comments from Derek Muir, Luisa Valiela, Arleen Feng, Bridgette DeShields, Jackson Vanfleet Brown, and an anonymous reviewer greatly strengthened the resulting publication.

Appendix A. Supplementary content

Supplementary content to this article can be found online at <http://dx.doi.org/10.1016/j.marpolbul.2016.05.077>.

References

- Bond, A.L., Provencher, J.F., Daoist, P.-Y., Lucas, Z.N., 2014. Plastic ingestion by fulmars and shearwaters at Sable Island, Nova Scotia, Canada. *Mar. Pollut. Bull.* 87, 68–75.
- Browne, M.A., Dissanayake, A., Galloway, T.S., Lowe, D.M., Thompson, R.C., 2008. Ingested microscopic plastic translocates to the circulatory system of the mussel, *Mytilus edulis* (L.). *Environ. Sci. Technol.* 42, 5026–5031.
- Browne, M.A., Crump, P., Niven, S.J., Teuten, E., Tonkin, A., Galloway, T., Thompson, R., 2011. Accumulation of microplastic on shorelines worldwide: sources and sinks. *Environ. Sci. Technol.* 45, 9175–9179.
- Carr, S.A., Liu, J., Tesoro, A.G., 2016. Transport and fate of microplastic particles in wastewater treatment plants. *Water Res.* 91, 174–182.
- Castañeda, R.A., Avlijas, S., Simard, M.A., Ricciardi, A., 2014. Microplastic pollution in St. Lawrence River sediments. *Can. J. Fish. Aquat. Sci.* 71, 1767–1771.
- Cole, M., Lindeque, P., Fileman, E., Halsband, C., Goodhead, R., Moger, J., Galloway, T.S., 2013. Microplastic ingestion by zooplankton. *Environ. Sci. Technol.* 47, 6646–6655.
- Davis 3rd, W., Murphy, A.G., 2015. Plastic in surface waters of the inside passage and beaches of the Salish Sea in Washington state. *Mar. Pollut. Bull.* 97, 169–177.
- Devriese, L.I., van der Meulen, M.D., Maes, T., Bekaert, K., Paul-Pont, I., Frère, L., Robbens, J., Vethaak, A.D., 2015. Microplastic contamination in brown shrimp (*Crangon crangon*, Linnaeus 1758) from coastal waters of the southern North Sea and channel area. *Mar. Pollut. Bull.* 98, 179–187.
- Dubai, F., Liebezeit, G., 2013. Suspended microplastics and black carbon particles in the jade system, southern North Sea. *Water Air Soil Pollut.* 224, 1352–1359.
- Eriksen, M., Mason, S., Wilson, S., Box, C., Zellers, A., Edwards, W., Farley, H., Amato, S., 2013. Microplastic pollution in the surface waters of the Laurentian Great Lakes. *Mar. Pollut. Bull.* 77, 177–182.
- Fendall, L.S., Sewell, M.A., 2009. Contributing to marine pollution by washing your face: Microplastics in facial cleansers. *Mar. Pollut. Bull.* 58, 1225–1228.
- Fossia, M.C., Marsilia, L., Bainia, M., Giannettia, M., Coppola, D., Guerrantia, C., Calania, I., Minutolic, R., Laurianod, G., Finoiad, M.G., Rubegnig, F., Panigadaf, S., Bérubég, M., Ramírez, J.U., Panti, C., 2016. Fin whales and microplastics: the Mediterranean Sea and the Sea of Cortez scenarios. *Environ. Pollut.* 209, 68–78.
- Free, C.M., Jensen, O.P., Mason, S.A., Eriksen, M., Williamson, N.J., Boldgiv, B., 2014. High-levels of microplastic pollution in a large, remote, mountain lake. *Mar. Pollut. Bull.* 85, 156–163.
- Hammer, S., Nager, R.G., Johnson, P.C.D., Furness, R.W., Provencher, J.F., 2016. Plastic debris in great skua (*Stercorarius skua*) pellets corresponds to seabird prey species. *Mar. Pollut. Bull.* 103, 206–210.
- Hidalgo-Ruz, V., Gutow, L., Thompson, R.C., Thiel, M., 2012. Microplastics in the marine environment: a review of the methods used for identification and quantification. *Environ. Sci. Technol.* 46, 3060–3075.
- Klosterhaus, S.L., Grace, R., Hamilton, M.C., Yee, D., 2013. Method validation and reconnaissance of pharmaceuticals, personal care products, and alkylphenols in surface waters, sediments, and mussels in an urban estuary. *Environ. Int.* 54, 92–99.
- Lavers, J.L., Bond, A.L., Hutton, I., 2014. Plastic ingestion by flesh-footed shearwaters (*Puffinus carneipes*): implications for fledgling body condition and the accumulation of plastic-derived chemicals. *Environ. Pollut.* 187, 124–129.
- Lenz, R., Enders, K., Stedmon, C.A., Mackenzie, D.M., Nielsen, T.G., 2015. A critical assessment of visual identification of marine microplastic using Raman spectroscopy for analysis improvement. *Mar. Pollut. Bull.* 100, 82–91.
- Mason, S.A., Garneau, D., Sutton, R., Chu, Y., Ehmman, K., Barnes, J., Fink, P., Papazissimos, D., Rogers, D., 2016. Municipal Wastewater Treatment Plant Effluent as a Pathway of Microplastic Pollution (in review).
- Masura, J., Baker, J., Foster, G., Arthur, C., Herring, C., 2015. Laboratory methods for the analysis of microplastics in the marine environment: recommendations for quantifying synthetic particles in waters and sediments. NOAA Technical Memorandum NOS-OR&R-48. National Oceanic and Atmospheric Administration, Silver Springs, MD.
- McCormick, A., Hoellein, T.J., Mason, S.A., Schluep, J., Kelly, J.J., 2014. Microplastic is an abundant and distinct microbial habitat in an urban river. *Environ. Sci. Technol.* 48, 11863–11871.
- Miranda, D.A., Carvalho-Souza, G.F., 2016. Are we eating plastic ingesting fish? *Mar. Pollut. Bull.* 103, 109–114.

- Napper, I.E., Bakir, A., Rowland, S.J., Thompson, R.C., 2015. Characterisation, quantity and sorptive properties of microplastics extracted from cosmetics. *Mar. Pollut. Bull.* 99, 178–185.
- Nicolau, L., Marçalo, A., Ferreira, M., Sá, S., Vingada, J., Eira, C., 2016. Ingestion of marine litter by loggerhead sea turtles, *Caretta caretta*, in Portuguese continental waters. *Mar. Pollut. Bull.* 103, 179–185.
- NYS OAG, 2015. Discharging Microbeads to our Waters: An Examination of Wastewater Treatment Plants in New York. New York State Office of the Attorney General, Environmental Protection Bureau, Albany, NY.
- Peters, C.A., Bratton, S.P., 2016. Urbanization is a major influence on microplastic ingestion by sunfish in the Brazos River Basin, Central Texas, USA. *Environ. Pollut.* 210, 380–387.
- Remy, F., Collard, F., Gilbert, B., Compere, P., Eppe, G., Lepoint, G., 2015. When microplastic is not plastic: the ingestion of artificial cellulose fibers by macrofauna living in seagrass Macrophytodebris. *Environ. Sci. Technol.* 49, 11158–11166.
- Rocha-Santos, T., Duarte, A.C., 2014. A critical overview of the analytical approaches to the occurrence, the fate and the behavior of microplastics in the environment. *Trends Anal. Chem.* 65, 47–53.
- Rochman, C.M., Tahir, A., Williams, S.L., Baxa, D.V., Lam, R., Miller, J.T., Teh, F.C., Werorilangi, S., Teh, S.J., 2015. Anthropogenic debris in seafood: plastic debris and fibers from textiles in fish and bivalves sold for human consumption. *Sci. Rep.* 5, 14340.
- Romeo, T., Pietro, B., Pedà, C., Consoli, P., Andaloro, F., Fossi, M.C., 2015. First evidence of presence of plastic debris in stomach of large pelagic fish in the Mediterranean Sea. *Mar. Pollut. Bull.* 95, 358–361.
- Rubissow-Okamoto, A., 2014. Stormwater: beyond the bag ban. *Estuary News*, December 2014, pp. 6–8.
- Teuten, E.L., Rowland, S.J., Galloway, T.S., Thompson, R.C., 2007. Potential for plastics to transport hydrophobic contaminants. *Environ. Sci. Technol.* 41, 7759–7764.
- Thompson, R.C., Moore, C.J., vom Saal, F.S., Swan, S.H., 2009. Plastics, the environment and human health: current consensus and future trends. *Philos. Trans. R. Soc. B* 364, 2153–2166.
- van Franeker, J.A., Law, K.L., 2015. Seabirds, gyres and global trends in plastic pollution. *Environ. Pollut.* 203, 89–96.
- Wright, S.L., Thompson, R.C., Galloway, T.S., 2013. The physical impacts of microplastics on marine organisms: a review. *Environ. Pollut.* 178, 483–492.
- Yonkos, L.T., Friedel, E.A., Perez-Reyes, A.C., Ghosal, S., Arthur, C.D., 2014. Microplastics in four estuarine rivers in the Chesapeake Bay, U.S.A. *Environ. Sci. Technol.* 48, 14195–14202.