



ELSEVIER

Contents lists available at ScienceDirect

Marine Pollution Bulletin

journal homepage: www.elsevier.com/locate/marpolbul

Development of a reef fish biological condition gradient model with quantitative decision rules for the protection and restoration of coral reef ecosystems

Patricia Bradley^a, Ben Jessup^a, Simon J. Pittman^b, Christopher F.G. Jeffrey^{c,o}, Jerald S. Ault^d, Lisamarie Carrubba^e, Craig Lilyestrom^f, Richard S. Appeldoorn^g, Michelle T. Schärer^g, Brian K. Walker^h, Melanie McFieldⁱ, Deborah L. Santavy^{j,*}, Tyler B. Smith^k, Graciela García-Moliner^l, Steven G. Smith^d, Evelyn Huertas^m, Jeroen Gerritsen^a, Leah M. Oliver^j, Christina Horstmann^{i,p}, Susan K. Jacksonⁿ

^a Tetra Tech, Inc., Owing Mills, MD, USA

^b Seascope Analytics Ltd, Plymouth, UK

^c CSS-Inc., Fairfax, VA, USA

^d University of Miami, Rosenstiel School of Marine and Atmospheric Science, Miami, FL, USA

^e NOAA, National Marine Fisheries Service, Silver Spring, MD, USA

^f Puerto Rico Department of Natural and Environmental Resources, San Juan, PR, USA

^g University of Puerto Rico, Mayagüez, PR, USA

^h Nova Southeastern University, Halmos College of Natural Sciences and Oceanography, Fort Lauderdale, FL, USA

ⁱ Smithsonian Marine Station, Fort Pierce, FL, USA

^j U.S. Environmental Protection Agency (US EPA), Office of Research and Development (ORD), Center for Environmental, Measurement and Modeling (CEMM), Gulf Ecosystem Measurement and Modeling Division (GEMMD), Gulf Breeze, FL, USA

^k University of the Virgin Islands, St. Thomas, VI, USA

^l NOAA, Caribbean Fisheries Management Council, San Juan, PR, USA

^m U.S. Environmental Protection Agency, Region 2, Guaynabo, PR, USA

ⁿ U.S. Environmental Protection Agency, Office of Water, Washington, DC, USA

^o Under Contract to NOAA, National Centers for Coastal Ocean Science, Marine Spatial Ecology Division, Biogeography Branch, Silver Spring, MD, USA

^p Oak Ridge Institute for Science Education Fellow, US EPA, ORD, CEMM, GEMMD, Gulf Breeze, FL, USA

ARTICLE INFO

Keywords:

Biological attributes
Biocriteria
Thresholds
Sustainability

ABSTRACT

Coral reef ecosystems are declining due to multiple interacting stressors. A bioassessment framework focused on stressor-response associations was developed to help organize and communicate complex ecological information to support coral reef conservation. This study applied the Biological Condition Gradient (BCG), initially developed for freshwater ecosystems, to fish assemblages of U.S. Caribbean coral reef ecosystems. The reef fish BCG describes how biological conditions changed incrementally along a gradient of increasing anthropogenic stress. Coupled with physical and chemical water quality data, the BCG forms a scientifically defensible basis to prioritize, protect and restore water bodies containing coral reefs. Through an iterative process, scientists from across the U.S. Caribbean used fishery-independent survey data and expert knowledge to develop quantitative decision rules to describe six levels of coral reef ecosystem condition. The resultant reef fish BCG provides an effective tool for identifying healthy and degraded coral reef ecosystems and has potential for global application.

1. Introduction

While climate changes are affecting reefs globally (Hughes et al., 2003; Hoegh-Guldberg et al., 2007, 2011, 2017; Carpenter et al., 2008; Knowlton and Jackson, 2008), local anthropogenic stressors contribute

directly to reef declines and can exacerbate climate change impacts (Rogers, 1990; Edinger et al., 1998; Jackson et al., 2001; Fabricius et al., 2005; Mora, 2008; Bejarno and Appeldoorn, 2013; Vega Thurber et al., 2014; Ennis et al., 2016; Robinson et al., 2017; Moustaka et al., 2018). Fishes represent a diverse taxonomic group providing ecological

* Corresponding author at: US EPA, ORD, CEMM, GEMMD, Gulf Breeze, FL, USA.

E-mail address: santavy.debbie@epa.gov (D.L. Santavy).

<https://doi.org/10.1016/j.marpolbul.2020.111387>

Received 23 February 2019; Received in revised form 11 June 2020; Accepted 16 June 2020

Available online 19 August 2020

0025-326X/ Published by Elsevier Ltd.

functions that are critical to the ecological integrity of coral reef ecosystems (Pratchett et al., 2014; Lefcheck et al., 2019). As such, fishes are important measures of the biological condition of coral reef ecosystems. For example, herbivores provide top-down control of algae that may otherwise replace living corals (Hughes, 1994; Burkepille and Hay, 2008), large predators provide top-down control on the fishes that prey on herbivores (Mumby et al., 2006; Stallings, 2008, 2009), and invertivores aid in controlling the abundance of coral feeders and bioeroders. Reef fish also provide economic and cultural value, such as food provisioning via subsistence and commercial fishing, and support tourism and recreational activities (Pendleton, 1995; Hawkins and Roberts, 2004; Principe et al., 2012; Ault et al., 2008, 2014; Brander and van Beukering, 2013; Spalding et al., 2017). Given their diverse functional roles in the ecosystem, using reef fish as indicators of coral reef ecosystem condition can help managers set targets for protection and restoration of coral reefs. For coral reefs, biological assessments using underwater survey techniques are commonly employed to directly measure the status of one or more taxonomic assemblage (e.g., corals, fish) and the chemical and physical attributes that support those assemblages (Jameson et al., 2001; Hill and Wilkinson, 2004; Jokiel et al., 2004; Brandt et al., 2009; Smith et al., 2011; Santavy et al., 2012; Jackson et al., 2014). These assessments are routinely used by states and territories to evaluate coral reef status and trends (Turgeon and Asch, 2002; Waddell, 2005; Waddell and Clarke, 2008). In the U.S. Gulf of Mexico and Caribbean, over-fishing and habitat degradation, including loss of nursery areas, have dramatically altered fish community composition across coral reef ecosystems (Claro, 1991; Paddock et al., 2009; Graham et al., 2017; Kadison et al., 2017). Reef fish species at all trophic levels have been subjected to intense fishing pressure (Munro, 1983; Hughes, 1994; Jackson et al., 2001; Pandolfi et al., 2003; Newman et al., 2006; Ault et al., 2005). Large groupers, snappers, hogfishes, and parrotfishes are now rare, with a resultant loss of predation and herbivory (Pittman et al., 2010; Appeldoorn, 2011; Ault et al., 2005, 2013). Sedimentation from development along tropical shorelines and runoff from agricultural land use is widely considered to have adversely impacted fish communities, particularly through suppressed feeding capability, poor water quality, and changes to benthic habitat (Rogers, 1990; Bejarno and Appeldoorn, 2013; Wenger et al., 2015; Neves et al., 2016; Brown et al., 2017).

Coral reef managers have little control over global or continental scale changes in climate and other environmental conditions; however, they may be able to substantially reduce local anthropogenic stresses by developing and enforcing laws, regulations and policies for waterbody activities and watershed land use. The U.S. Clean Water Act (CWA) (33 USC § 1251 et seq., 1972) can be used to protect coral reef ecosystems (Bradley et al., 2008, 2009, 2010). The CWA long-term objective is to restore and maintain the chemical, physical and biological integrity of the nation's waters. To help achieve this visionary objective, the CWA directs jurisdictions (states, territories and tribes) to adopt water quality standards (WQS) as provisions of their laws or regulations. A key component of WQS is water quality criteria (physical, chemical and biological criteria). Water quality criteria are scientifically defensible thresholds established to protect the goals, or designated uses, for a waterbody. When the WQS are not attained, the waterbody is determined to be impaired and management response is needed to address the impairment.

As part of the WQS process, the Environmental Protection Agency's (EPA) guidance recommends jurisdictions to develop and adopt into their water quality standards biological criteria (henceforth "biocriteria") (EPA, 1990, 2002, 2011b, 2013b, 2016) to protect aquatic life. Biological monitoring surveys provide the foundational information for bioassessments and establishing biocriteria (EPA, 1990, 2002, 2011a, 2013b, 2016; Ault et al., 1999; Davis and Simon, 2004; Bradley et al., 2010; Smith et al., 2011; Bryan et al., 2016). Biological assessment programs and, in some places, biocriteria have been implemented nationwide for streams and rivers (EPA, 2016).

The Biological Condition Gradient (BCG) is an approach to assess the biological condition of a waterbody relative to natural expectations comparable to the concept of biological integrity. *Biological integrity* has been defined as a balanced, integrated, adaptive community of organisms having a composition and diversity comparable to that of the natural habitats (Frey, 1977). This definition includes the ecosystem functions and processes that generate and maintain the community (Karr and Chu, 2000). As such, *biological integrity* is integral to concept of *ecological integrity* which embodies four main components: nativeness, pristineness, diversity, and resilience (Schallenberg et al., 2011). *Resilience* refers to the ability of an ecosystem to maintain key functions and processes in the face of stresses or pressures by resisting and then adapting to change (Holling, 1973; Nystrom and Folke, 2001). Coral reef resilience has been linked to certain physical and ecological characteristics that provide some reefs with a greater likelihood of resisting and/or recovering from disturbance (Salm et al., 2001; West and Salm, 2003).

Originally developed and applied for freshwater ecosystems, the BCG is part of EPA's biological assessment and criteria "toolbox" that includes biological indices, models, statistical methods, and practical guidance (Davies and Jackson, 2006; EPA, 2011a, 2013b, 2016). The BCG (Fig. 1) provides a common language to describe how biological attributes of an aquatic ecosystem (ordinate, y-axis) are expected to change along a gradient of increasing anthropogenic stress (abscissa, x-axis) ranging from observable biological condition found at undisturbed or minimally-disturbed reference sites (i.e., sites with high environmental quality, pristine, or intact conditions) to sites with high anthropogenic stress (i.e., partially to completely degraded). Biological attributes include aspects of size-structured population abundance, community composition, inter-habitat connectivity and ecosystem function. Although the theoretical stressor-response curve is continuous, condition levels are discrete intervals defined along the stress gradient by a consistent, cogent narrative for each level.

A set of resilience indicators has been developed and applied to various coral reef ecosystems (Obura and Grimsditch, 2009; McClanahan et al., 2012; Maynard et al., 2015). The BCG method enables incorporation of indicators of ecological resilience directly into the model. The BCG ordinate can include indicators of biological resilience, while indicators of physical and chemical resilience can be incorporated in the BCG abscissa.

Freshwater BCG developments have typically separated fishes and benthic invertebrates, leading to BCG models for each (e.g. EPA, 2016; Gerritsen et al., 2017). Benthic invertebrates are an obvious choice because they have high site fidelity (limited mobility), primarily integrate stressors at much smaller scales, and are not immediately impacted by stressors not directly related to water quality (e.g., fishing pressure). Fishes can be highly mobile and wide-ranging, but their absence in a given location may be a reflection of acute or chronic stressors. Coral reef fish site fidelity varies in scale depending on the species, growth stage, and local habitat availability (Walker et al., 2009; Grober-Dunsmore et al., 2007; Kuffner et al., 2007; Pittman et al., 2007), however, assemblages are specific to certain habitat combinations and ecosystem regions (Ames, 2017). Their stressors vary from those of benthic invertebrates, especially due to the many species targeted for food, sport, and the aquarium trade. Loss of assemblages has been related to decreases in water quality and habitat degradation (Knowlton and Jackson, 2008).

Some jurisdictions have used the BCG to support various aspects of water quality management, including: (1) more precise definitions of designated aquatic life uses; (2) setting goals for protection or restoration of aquatic life; (3) identification and protection of high quality waters; (4) assessing condition and identifying degraded waterbodies; (5) tracking progress in restoration and protection; and, (6) development of biological criteria. The BCG is an effective tool for clear communication with the public and stakeholders of the biological condition of their waters in the context of the CWA biological integrity objectives,




Fig. 1. Conceptual model of the biological condition gradient (Davies and Jackson, 2006).

and the likely outcomes of water quality management decisions (EPA, 2011a, 2011b, 2016).

In 2004 the U.S. Ocean Action Plan recommended EPA develop biological assessment methods and biological criteria methods for states and territories to evaluate the health of coral reefs and associated water quality (The White House, 2004; Bradley et al., 2008, 2010). EPA developed a rapid bioassessment protocol (RBP) for stony coral demographics (Fisher, 2007) that was successfully tested in the Florida Keys (Fisher et al., 2007). EPA subsequently tested the stony coral indicators derived from the RBP to determine their responsiveness to anthropogenic disturbance (Fisher et al., 2008) and developed a probabilistic survey design with spatially-balanced random site selection for St. Croix in 2007 and St. Thomas and St. John in 2009 (Fisher et al., 2014). EPA expanded their survey methodology to include fish, gorgonians and sponges (Santavy et al., 2012). In 2009 the U.S. Coral Reef Task Force (USCRTF) selected the Guánica Bay watershed as the location for its first multi-agency initiative to reduce watershed impacts on coral reefs in the coastal zone, leading EPA to focus its bioassessment development efforts on southern Puerto Rico.

2. Materials and methods

A quantitative reef fish BCG model was developed for coral reef ecosystems. Development of the reef fish BCG model included five steps:

- (1) collect statistically robust data that are fully representative of the study domain where the BCG is to be applied, and encompass most, if not all, of the possible BCG levels;
- (2) conduct preliminary data assimilation and analysis, putting data into formats readily used in the development process, and examine stressor-response relationships for individual taxa and community assemblages relative to proposed gradients;
- (3) convene an expert panel familiar with the local and regional environment and species, including expected species and assemblage responses to stressors in the region of concern;
- (4) develop the quantitative decision rules for the reef fish BCG model; and
- (5) test the model, adjust, iterate and recalibrate.

2.1. Step 1: collect and organize bioassessment data

Two underwater coral reef fish surveys were conducted by EPA in 2010 and 2011 along the south coast of Puerto Rico that support development of the coral reef fish BCG. The 2010 survey was designed to reflect coral reef impacts due to increased sediment exposure resulting from land-based human disturbances at 76 stations (Oliver et al., 2014; Bradley et al., 2014) (Fig. 2). Stations were selected to represent a range of potential land-based sediment and pollution threats as modeled by WRI and NOAA (2006) “Reefs at Risk” and NOAA’s “Summit to Sea” respective approaches. This project analyzed sediment production on land using soil type and relative erodibility, precipitation data and slope, coupled with an inverse distance weighting function to simulate reduced sediment threat to coastal habitats located further from shore. Puerto Rico 2010 stations were selected from the WRI and NOAA (2006) geospatial dataset, which assigned relative sedimentation threat to mapped coastal habitats (Kendall et al., 2001). The 2011 survey used a probability-based design (Fig. 3) to determine status and biological condition of reef fish and coral communities (Fisher et al., 2019). Both surveys were conducted on coral reefs within 4.8 km of shore (including shores of small islands) at depths ≤ 12 m as characterized in NOAA’s benthic habitat map (Kendall et al., 2001) to reduce depth effects on assemblage structure and to reflect exposure to land-based stressors in nearshore waters. The surveys included visual assessments of all reef fishes (species size-structured abundance), stony corals (taxa,




Fig. 2. Location and distribution of 2010 EPA sampling stations. Seventy-six targeted coral survey locations (red circles) at regular intervals across human disturbance gradients (low, medium and high sediment threat levels) were distributed across linear reefs within 1.5 km of shore (including cays) and between 2 and 12 m depth. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)




Fig. 3. Location and distribution of 2011 EPA sampling stations (Fisher et al., 2019). Sixty randomly selected coral survey locations (red circles) were distributed across linear reefs within 1.5 km of shore (including cays in the target substrate) and between 2 and 12 m depth. Coral reef and colonized hardbottom substrate shown in gray shading. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

individual 3D colony sizes, amount of live tissue on coral colonies, and the occurrence of adverse health conditions such as bleaching, disease or overgrowth by boring sponges), reef rugosity, selected macro-invertebrates (e.g., queen conch, spiny lobster, reef crabs, sea urchins and long-spined urchins), and morphometric data for gorgonians and sponges (colony height, diameter, and morphology) (Santavy et al., 2012).

2.2. Step 2: conduct preliminary data analysis and data preparation

Survey data were subjected to a thorough QA/QC procedure to eliminate uncorrectable unmatched or conflicting data, sites deemed to be in non-target habitat types, and to correct older taxonomic names or synonyms. The data were then put into an Excel workbook for use by the experts. Except for fishing pressure, literature on stressor/response relationships that ties individual stressors to reef fish community metrics was limited (Bradley et al., 2014; Bradley et al., 2016), so limited stressor information was provided in the experts' workbooks.

For each site surveyed, information included depth, distance from shore and shelf edge, reef type, habitat type, and rugosity. Roberts and Ormond (1987) stated that depth alone can be a good indicator of fish species richness; however, depth is also a defining variable for reef type (Walker et al., 2009). Distance from the shore was included because certain fish species are more likely to migrate from nearby near-shore nursery habitats to adult reef habitats (Appeldoorn et al., 1997, 2003; Lindeman et al., 2000; Nagelkerken et al., 2015; Dahlgren and Eggleston, 2000; Cocheret de la Morinière et al., 2002; Christensen et al., 2003; Aguilar-Perera, 2004; Mumby et al., 2004, 2008; Aguilar-Perera and Appeldoorn, 2007; McField and Kramer, 2007; Meynecke et al., 2008; Schärer-Umpierre, 2009; Sale et al., 2010). Shelf breaks are areas of unique habitats and physical properties (Shcherbina et al., 2008) that support equally unique fish assemblages (Kimmel, 1985; Cerveny, 2006; Pittman et al., 2010). Additionally, they are an important spawning habitat for a variety of species (Thompson and Munro, 1974; Johannes, 1978; Colin et al., 1987; Shapiro et al., 1993; Sadovy et al., 1994a, 1994b; Sala et al., 2001; Claro and Lindeman, 2003; Nemeth et al., 2006; Ojeda-Serrano et al., 2007a, 2007b; Heyman and Kjerfve, 2008; Sanchez et al., 2017).

To account for confounding effect of habitat complexity on species richness, we used several measures or indicators. These included the rugosity index: the ratio of the length of a chain over the distance covered along a transect by the chain when draped over stony corals and non-coral substrates (Risk, 1972; Hobson, 1972; Talbot and Goldman, 1972; McCormick, 1994; Rogers et al., 1994; Lang, 2003; Santavy et al., 2012). While the rugosity index accounts for important vertical dimensions, it does not fully reflect the three-dimensional availability of fish habitat. Therefore, the data also included additional indicators of habitat complexity, including colony surface area estimates for the three major sessile benthic populations, stony corals,

sponges and gorgonians (Courtney et al., 2007; Santavy et al., 2012; Fisher et al., 2007, 2014).

Commonly used metrics that characterize the fish community metrics were calculated. These included: species richness, density, mean length and standard deviation, total fish biomass, number of fish schools, percent of fish in various families (i.e., *Acanthuridae*, *Scaridae*, *Chaetodontidae*, *Haemulidae*, *Pomacentridae*, *Labridae*, *Lutjanidae* and *Carangidae* and *Epinephelidae*), and relative biomass of herbivores and piscivores (Caldow et al., 2009; Santavy et al., 2012). Additionally, the list of fish species observed at the site was provided, including density and biomass by species.

2.3. Step 3: convene an expert panel

A panel of coral reef and reef fish experts was assembled in 2012 (Bradley et al., 2014; Santavy et al., 2016). The experts were chosen based on their scientific expertise in Caribbean coral reef taxonomic groups, as well as community structure, organism condition, ecosystem function and ecosystem inter-habitat connectivity. Experts included research scientists from federal and state organizations, academia, and non-governmental organizations (NGOs), as well as water quality managers and natural resource managers from Puerto Rico and the U.S. Virgin Islands (USVI). A list of the BCG experts is available in Bradley et al. (2016).

2.4. Step 4: develop BCG model decision rules

Four expert workshops (August 2012, April 2014, October 2015 and March 2019) were held in Puerto Rico to develop, test, and calibrate the BCG model for coral reef ecosystems. The first workshop provided proof-of-concept that the BCG framework developed for freshwater aquatic ecosystems could be adapted for coral reef ecosystems (Bradley et al., 2014; Santavy et al., 2016). The coral reef experts examined video footage from panoramic and linear transect views and supporting photographs of 12 sites from the bioassessments conducted in 2010 and 2011. The sites were selected to represent a range of biological condition. A blind identification system masked all site locations from the coral reef experts. Each expert was asked to draw upon their personal experience and expertise to rate the reef condition for each site. Workshop materials were organized by site and included a photo diary of key representative photos, so coral reef experts would rate the biological condition of each site to document the traits or characteristics used to support their ratings. The experts were asked to consider all aspects of the reef and specifically instructed to consider the characteristics of the condition of corals, sponges, gorgonians, fish, algae, reef rugosity, and topographical heterogeneity (Bradley et al., 2014).

While our working group simultaneously examined multiple assemblages for potential development of the BCG, after the first workshop we decided to separate into fishes and benthic assemblage subgroups for development of the quantitative BGC model, principally because the experts tended to be knowledgeable about either fishes or benthic organisms, and because the two assemblages respond differently to stressors. This paper describes the model development process and results achieved by the fish experts. The benthic process is described in a separate publication (Santavy et al., 2016).

During facilitated discussions in the second, third and fourth workshops and multiple webinars, the fish experts had three subtasks in the development of decision rules: (1) arrive at a common understanding of the target species that make up the database, including community structure, expected occurrences, and sensitivities to stressors; (2) use their understanding of the species to assign a set of individual sites to BCG levels; and, (3) use these results to develop a narrative and ultimately quantitative descriptions of expected species compositions for each BCG level. For subtask 1, all information appropriate to the target species was used: scientific and technical literature; panel members' knowledge and experience; and, empirical

Table 1
BCG attributes and their descriptions (modified from EPA, 2016).

Attribute	Description
I. Historically documented, long-lived, or regionally endemic taxa	<p>Taxa known to have been supported according to historical, museum or archeological records, or taxa with restricted distribution (occurring only in a locale as opposed to a region), often due to unique life history requirements. They may be long-lived and late maturing and have low fecundity, limited mobility, multiple habitat requirements as with diadromous species, or require a mutualistic relationship with other species. They may be among listed Endangered or Threatened (E/T) or special concern species. Predictability of occurrence is often low, and therefore requires documented observation. The taxa that are assigned to this category require expert knowledge of life history and regional occurrence of the taxa to appropriately interpret the significance of their presence or absence. Long-lived species are especially important as they provide evidence of multi-annual persistence of habitat condition.</p> <p>Taxa that are highly sensitive to pollution or anthropogenic disturbance. Tend to occur in low numbers relative to total population density, but they might make up a large relative proportion of richness. In high quality sites, they might be ubiquitous in occurrence or might be restricted to certain micro-habitats. They often have slow growth – long-lived (K-strategists) vs. short-lived—fast growth (r-strategists). In coral reef ecosystems, large-bodied, slow-growing, late-maturing fishes (K-strategists) are generally more sensitive to fishing pressure and environmental stress than faster-growing, shorter-lived species (Beverton and Holt, 1957; Man et al., 1995; Jennings et al., 1998; Coleman et al., 2000; Goodwin et al., 2006; Ault et al., 2008). The distinguishing characteristic for this attribute category was found to be sensitivity and not relative rarity, although some of these taxa might be uncommon in the data set (e.g., very small percent of sample occurrence or sample density), therefore, these are the first to disappear with disturbance or pollution.</p>
III. Intermediate sensitive taxa	<p>Taxa that are abundant in relatively undisturbed conditions but are sensitive to anthropogenic disturbance/pollution. They have a broader range of tolerance than Attribute II taxa and can be found in reduced density and richness in moderately disturbed or polluted stations. These taxa often comprise a substantial portion of natural communities.</p>
IV. Intermediate tolerant taxa	<p>Taxa that commonly comprise a substantial portion of the fish assemblage in undisturbed habitats, as well as in moderately disturbed or polluted habitats. They exhibit physiological or life-history characteristics that enable them to thrive under a broad range of thermal, flow, or oxygen conditions. Many have generalist or facultative feeding strategies enabling utilization of diverse food types. These species have little or no detectable response to moderate stress, and they are often equally abundant in both reference and moderately stressed sites. Some intermediate tolerant taxa may show an “intermediate disturbance” response, where densities and frequency of occurrence are relatively high at intermediate levels of stress, but they are intolerant of excessive pollution loads or habitat alteration.</p>
V. Tolerant taxa	<p>Tolerant taxa are those that typically comprise a low proportion of natural communities. These taxa are more tolerant of a greater degree of disturbance and stress than other organisms and are, thus, resistant to a variety of pollution or habitat induced stress. They may increase in number (sometimes greatly) under severely altered or stressed conditions. They may possess adaptations in response to organic pollution, hypoxia, or toxic substances. These are the last survivors in severely disturbed systems and can prevail in great numbers due to lack of competition or predation by less tolerant organisms, and they are key community components of level 5 and 6 conditions.</p>
VI. Non-native or intentionally introduced species	<p>Any species not native to the ecosystem. Species introduced or spread from one region of the U.S. to another outside their normal ranges are non-native, or non-indigenous. This category also includes species introduced from other continents and referred to as “alien” species. This attribute represents both an effect of human activities and a stressor in the form of biological pollution. The BCG identifies the presence of native taxa expected under undisturbed or minimally disturbed conditions as an essential characteristic of BCG level 1 and 2 conditions. The BCG only allows for the occurrence of non-native taxa in these levels if those taxa do not displace native taxa and do not have a detrimental effect on native structure and function. Condition levels 3 and 4 depict increasing occurrence of non-native taxa. Extensive replacement of native taxa by tolerant or invasive, non-native taxa can occur in levels 5 and 6. Anomalies of the organisms; indicators of individual health (e.g., deformities, lesions, tumors).</p>
VII. Organism condition	<p>Ecosystem function refers to processes required for the performance of a biological system expected under naturally occurring conditions (e.g., primary and secondary production, respiration, nutrient cycling, and decomposition).</p>
VIII. Ecosystem function	<p>Assessing ecosystem function includes consideration of the aggregate performance of dynamic interactions within an ecosystem, such as the interactions among taxa (e.g., food web dynamics) and energy and nutrient processing rates (e.g., energy and nutrient dynamics) (Cairns, 1977). Additionally, ecosystem function includes aspects of all levels of biological organization (e.g., individual, population, and community condition). Altered interactions between individual organisms and their abiotic and biotic environments might generate changes in growth rates, reproductive success, movement, or mortality. These altered interactions are ultimately expressed at ecosystem-levels of organization (e.g., shifts from heterotrophy to autotrophy, onset of eutrophic conditions) and as changes in ecosystem process rates (e.g., photosynthesis, respiration, production, decomposition).</p>
IX. Spatial and temporal extent of detrimental effects	<p>(continued on next page)</p>

Table 1 (continued)

Attribute	Description
X. Ecosystem connectivity	<p>The spatial and temporal extent of stressor effects includes the near-field to far-field range of observable effects of the stressors on a water body. Such information can be conveyed by biological assessments provided the spatial density of sampling sites is sufficient to convey changes along a pollution continuum (USEPA, 2013a). Use of a continuum provides a method for determining the severity (i.e., departure from the desired state) and extent (i.e., distance over which adverse effects are observed) of an impairment from one or more sources. As with attribute VIII above, attribute IX has not yet been developed and applied in BCG models. It is included for future development and application. State scientists involved in the development of the BCG conceptual model stated that this attribute was important to include for future testing and development.</p> <p>Access or linkage (in space/time) to materials, locations and conditions required for maintenance of interacting populations of aquatic life. It is the opposite of fragmentation and is necessary for persistence of metapopulations and natural flows of energy and nutrients across ecosystem boundaries. Ecosystem connectivity can be indirectly expressed by certain species that depend on the connectivity, or lack of connectivity, within an aquatic ecosystem to fully complete their life cycles and thus maintain their populations. There are two commonly recognized categories of connectivity based upon the typical life history (i.e., two-phase life cycle) of most reef associated fishes: (1) pre-settlement connectivity through larval dispersal and (2) post-settlement connectivity (Aguilar-Perera, 2004).</p>

species associations with both natural habitats, as well as anthropogenic stressors. Most coral reef assessment data were collected at the site or reef scale. BCG attributes relevant to this scale included aspects of taxonomic composition and community structure (attributes I–V) and non-native taxa (attribute VI) (EPA, 2016). Taxa differ in their sensitivity or tolerance to stressors, but sensitivity can vary both among species and by stressor (Davies and Jackson, 2006). The fish experts used the BCG attribute definitions (Table 1), their expert knowledge and experience, available literature, and frequency of a species occurring in the data set to assign 357 Caribbean fish species to the taxonomic attributes based on their sensitivities to two anthropogenic stressors (sediments and fishing). For fishing pressure, the fish experts considered whether each species was subject to fishing pressure, the category of fishing pressure (e.g., commercial, recreational or ornamental), and whether that species was regulated under federal or territorial fishing laws (EPA, 2016). For sensitivity to sediment threat, the experts assigned each fish species to a BCG Attribute based on habitat preferences (e.g., ontogenetic shifts from juvenile to adult habitats, as well as observations where the experts regularly observed a species). Fish response to the two stressor categories often differed, and experts took both stressors into consideration in assigning the fish to BCG attributes. If a fish species was sensitive to one of the two stressor categories, it was considered to be sensitive (i.e., the assignment was based upon the more sensitive response).

Shifts in taxa as a function of differing sensitivities to disturbance are well documented (e.g., shifts from K-selected to r-selected strategists following disturbance). Non-native species were identified as BCG Attribute VI, reflecting the detrimental effects of nonnative taxa on native species (Davies and Jackson, 2006; EPA, 2016). Some taxa were not associated with any attribute (assigned to “x”) because the fish experts were unfamiliar or had little supporting information in the literature relative to stressor tolerance or because the survey methodology did not allow an accurate count of the species (e.g., cryptic species).

In subtask 2, assigning sites to BCG levels, the objective was to assign the levels based solely on natural site classification and species composition. Prior to the second workshop, the facilitation team selected a set of 38 sites from the EPA 2010/2011 surveys to span the range and gradient of stress that occurs in southern Puerto Rico. During the 2nd workshop, the facilitator projected the data for each site onto a screen and pointed out the site data and summary indicators. Site-specific information on potential anthropogenic stressors was withheld from the panel to prevent bias in their assessments. We recognized that some natural classification variables might be confounded with anthropogenic stress; e.g., distance from shore is related to distance from land-based discharges and runoff, and rugosity may be reduced because corals have died. The experts then spent several minutes individually considering the data. The facilitator then called on each expert to propose a BCG level for the site, provide the critical or most important information they used to inform the decision, including any confounding or conflicting information, and how they resolved these conflicts (EPA, 2016; Gerritsen et al., 2017). Once all experts had provided their individual ratings, the experts discussed the ratings and rationales, and revised their individual ratings, if desired.

Experts were often unwilling to select a single, discrete BCG level and rated a site as “better than a 4 but not a 3” for reasons such as low total number of taxa or other factors related to site characteristics versus fish community data. To accommodate this, experts rated sites with additional descriptors of “good”, “poor” or “middle” for each BCG level and these were scored as (+) or (–). The quantitative decision model yielded numeric memberships between 0 and 1 for each BCG level, and all memberships summed to 1. This allowed for ties between levels, as well as dominant membership in a single level and smaller memberships in adjacent levels.

Subsequent to the facilitated site rating process, the fish experts provided narrative statements to describe what they expected to see for each BCG level starting from the highest quality condition observed in

Table 2
Preliminary narrative BCG with four distinct levels of condition for all reef assemblages: very good/excellent; good; fair; and poor.
(Source: Bradley et al., 2014)

Level	Physical structure	Corals	Gorgonians	Sponges	Fish	Other Vertebrates	Other Invertebrates	Algae/Plants	Condition
Very Good – Excellent (BCG 1-2)	High rugosity or 3D structure; substantial reef built above bedrock; many irregular surfaces provide habitat for fish; very clear water; no sediment, flocs or films	High species diversity including rare; large old colonies (<i>Orbicella</i>) with high tissue coverage; balanced population structure (old & middle-sized colonies, recruits); <i>Acropora</i> thickets present	Gorgonians present but sub-dominant to corals	Large autotrophic and highly sensitive sponge species abundant	Populations have balanced species abundance, sizes, biomass, and trophic interactions; Large piscivores present (groupers, barracuda, sharks)	Large, long-lived species present and diverse (turtles, dolphins)	<i>Diadema</i> , lobster, small crustaceans and polychaetes abundant; some large sensitive anemone species	Crustose coralline algae abundant; turf algae present but cropped and grazed by <i>Diadema</i> ; low abundance fleshy algae	Low prevalence of disease or tumors; mostly live tissue on colonies
Good (BCG 3)	Moderate to high rugosity; moderate reef built above bedrock; some irregular cover for fish habitat; water slightly turbid; low sediment, flocs or film on substrate	Moderate coral diversity; large old colonies (<i>Orbicella</i>) with some tissue loss; varied population structure (usually old colonies, few middle aged and some recruitment); <i>Acropora</i> thickets maybe present; rare species absent	Gorgonians more abundant than in BCG Levels 1–2	Autotrophic species present but highly sensitive species missing	Decline of large apex predators (e.g., groupers, snappers, etc.) noticeable; small reef fish more abundant than Levels 1–2	Large, long-lived species locally extirpated (turtles, eels)	<i>Diadema</i> , lobster, small crustaceans and polychaetes less abundant than Level 1–2; large sensitive anemones species missing	Crustose coralline algae present but less than Levels 1–2; turf algae present and longer; more fleshy algae present	Disease and tumor prevalence slightly above background level; more colonies have irregular tissue loss
Fair (BCG 4)	Low rugosity, limited reef built above bedrock; erosion of reef structure obvious; water turbid; more sediment accumulation, flocs and films; <i>Acropora</i> usually gone or present as rubble for recruitment substrate	Reduced coral diversity; emergence of tolerant species, few or no large old colonies (<i>Orbicella</i>) mostly dead; <i>Acropora</i> thickets gone; large remnants mostly dead	Gorgonians more abundant than in Levels 1–3; replace sensitive corals and sponges species	Mostly heterotrophic sponges with tolerate species and clonids	Near absence of large piscivores; small reef fish abundant (mostly Damsel fish)	Large, long-lived species locally extirpated (turtles, eels)	<i>Diadema</i> absent, <i>Palythoa</i> overgrowing corals, crustaceans, polychaetes, and sensitive anemones conspicuously absent	Some coralline algae; turf is uncropped covered in sediment; lots fleshy algae with high diversity (e.g., <i>Dictyota</i>); possibly smothering sessile invertebrates; absence of crustose coralline algae	High incidence of diseased coral, sponges, gorgonians; evidence high mortality; usually less tissue than dead portions on colonies
Poor (BCG 5-6)	Very low rugosity, no or low reef built above bedrock ; poor fish habitat; very turbid water; thick sediment film & high flocs covering bottom; no substrate for recruits	Absence of most species, colonies small, only highly tolerant species with little or no tissue	Small & sparse colonies, mostly small sea fans, often diseased	Heterotrophic sponges buried deep in sediment, highly tolerant sponge species	No large fish, few intolerant species, lack of multiple trophic levels	Usually devoid of other vertebrates	Low or no reef invertebrates; high abundance of sediment dwelling organisms such as polychaetes, holothurians	High cover of fleshy algae (<i>Dictyota</i>); possibly smothering sessile invertebrates; no turf or coralline algae; complete absence of crustose coralline algae	High incidence disease on small colonies of corals, sponges gorgonians; if present, low or no tissue

the data set. This narrative became the basis for BCG rule development.

2.5. Step 5: test model and review model's performance

Following the deliberations in the second workshop, quantitative rules were developed using the fish experts' narrative statements and distribution statistics for attribute metrics and other measures of the assemblage in each BCG level. Rules are logic statements that the fish experts used to make their decisions. Once the rules are quantified (Step 5, below), a knowledgeable person can follow them to obtain the same BCG level ratings as the group of fish experts, making the actual decision criteria transparent to water quality managers and stakeholders. Rules and reasoning of the experts, whether quantitative or qualitative, were compared to data summaries of the sites evaluated by the experts. For example, if the panel identified a small to moderate number of sensitive taxa for BCG level 3, then the number of sensitive taxa in samples the panel assigned to BCG level 3 were examined (e.g., sensitive taxa ranged from 4 to 8). The statistical distribution of the data in sites assessed by the panel, including modes and quantiles, were used to establish decision thresholds for classifying sites to each BCG level. Quantiles helped to establish the fuzzy boundaries of the decision rules (see below). This process was repeated for all rules and attributes identified by the panel as being important to their decisions.

The decision rules were tested and refined by the expert panel in webinars following the second workshop. The rules were reviewed at the third workshop (with several new panel members present) to confirm that the rules were consistent with expert logic, observations, and empirical data. Mathematical fuzzy logic that mimicked human reasoning was used to develop an inference model to replicate the fish

experts' decision process (EPA, 2016). Fuzzy logic is “a precise logic of imprecision and approximate reasoning” (Zadeh, 2008) that has been directly applied worldwide during environmental assessments where imprecise and incomplete information is used to make decisions on the quality and sustainability of systems (Castella and Speight, 1996; Ibelings et al., 2003; Ionnidou et al., 2003; EPA, 2016; Gerritsen et al., 2017). The development of BCG inference models is explained specifically in Gerritsen et al. (2017), and a general tutorial on fuzzy logic can be found in Klir (2004).

Membership of a site in a given BCG level was interpreted according to rules applicable to each attribute or metric that the panel deemed important for the BCG level. For example, for BCG level 3, the rule for the metric *total taxa* was: total taxa ≥ 20 (15–25). This meant that the panel agreed that the rule for the metric *total taxa* should be a desired mean value of 20 for that metric, but an absolute minimum of 15, and full membership at a value of 25. Hence, membership of the site in BCG level 3 was 0 (zero) when the metric *total taxa* was less than or equal to 15, 50% when there were exactly 20 of the metric and 1 (100%) when the value equals or exceeded 25. The panel also specified other rules expressed in the same way. Rules for individual metrics were typically combined with logical AND, i.e., the minimum value of all the memberships was taken as the final membership. Some rules were identified as alternates, i.e., either A or B, and these were combined with a logical OR, i.e., the maximum value of the two alternative rules.

The fish experts reviewed the model rules and results and suggested revisions to their ratings or the rules when needed. As rule development and refinement proceeded, it became apparent that differences among classification variables could be encapsulated by two habitat types: patch reef and hard bottom.




Fig. 4. Distribution of fish panelists' BCG level assignments expressed as difference from the group median in 1/3 BCG level steps. Calibration (top) and confirmation (bottom) stations from the Puerto Rico reef fish dataset.

During Workshop 3, the experts were asked to review 11 confirmation sites selected to span the range and gradient of stress that occurs in southern Puerto Rico, to apply the fish rules that had been established in Workshop 2. Experts were requested to assign a BCG level to each site and to state reasons if they disagreed with any given quantitative rule. The experts requested and received the size structure distributions for all stations and for each species. No disagreements with rules were stated and the experts completed the confirmation stations. There were, however, several issues that arose that warrant further investigation (see Discussion).

Performance of the model was described in terms of agreement between model results and the median of expert ratings per site. We assessed the number of sites where the draft BCG decision model's level rating

exactly matched the fish experts' median opinion ("exact match") and the number of sites where the model predicted a BCG level that differed from the median expert opinion ("mismatch" sites). For the mismatched sites, the BCG level rating differences between the fish experts and the model were examined to determine whether there was a bias.

3. Results

During the first workshop, experts rated 12 shallow reef sites from Puerto Rico as either good, fair, or poor based on videos and photos. Using only the 12 sites, the experts developed a preliminary narrative BCG with four distinct levels of condition: very good - excellent; good; fair; and poor

Table 3
Narrative rules for fish BCG model in Puerto Rico coral reefs.

BCG Level 1	Definition: Biological conditions as they existed (or still exist) in the absence of measurable effects of stressors and provides the basis for comparison to the next five levels.
	Fish Narrative Rules: Populations have balanced species abundance, sizes, biomass, and trophic interactions; Large piscivores present (groupers, barracuda, sharks)
BCG Level 2	Definition: Minimal changes in structure of the biotic community and minimal changes in ecosystem function— <i>virtually all native taxa are maintained with some changes in biomass and/or abundance; ecosystem functions are fully maintained within the range of natural variability</i>
	Fish Narrative Rules: Populations have balanced species abundance, sizes, biomass, and trophic interactions; Large piscivores present (groupers and snappers, but not sharks); schools of piscivores present
BCG Level 3	Definition: Evident changes in structure of the biotic community and minimal changes in ecosystem function— <i>Some changes in structure due to loss of some rare native taxa; shifts in relative abundance of taxa, but intermediate sensitive taxa are common and abundant; ecosystem functions are fully maintained through redundant attributes of the system</i>
	Fish Narrative Rules: Decline of large apex predators (e.g., groupers, snappers, etc.) noticeable, however still present; small reef fish more abundant than Levels 1–2; large body parrotfish present; high within-family diversity
BCG Level 4	Definition: Moderate changes in structure of the biotic community with minimal changes in ecosystem function. <i>Moderate changes in structure because of replacement of some sensitive-ubiquitous taxa by more tolerant taxa, but reproducing populations of some sensitive taxa are maintained; overall balanced distribution of all expected major groups; ecosystem functions largely maintained through redundant attributes.</i>
	Fish Narrative Rules: Near absence of large piscivores, however t least one piscivore present; small reef fish abundant (mostly Damsel fish and wrasses); parrotfish present
BCG Level 5	Definition: Major changes in structure of the biotic community and moderate changes in ecosystem function. <i>Sensitive taxa are markedly diminished; conspicuously unbalanced distribution of major groups from distributions expected; organism condition shows signs of physiological stress; ecosystem function shows reduced complexity and redundancy.</i>
	Fish Narrative Rules: No large fish, few intolerant species, lack of multiple trophic levels; more than 4-5 fish species
BCG Level 6	Definition: Severe changes in structure of the biotic community and major loss of ecosystem function. <i>Extreme changes in structure; wholesale changes in taxonomic composition; extreme alterations from normal densities and distributions; organism condition is often poor; ecosystem functions are severely altered.</i>
	Fish Narrative Rules: Does not meet Level 5 rules

(Table 2). The experts agreed that there were no longer any reefs in Puerto Rico that met the BCG level 1 definition corresponding to very good-excellent condition (Bradley et al., 2014).

During the second workshop, the fish experts assigned fish species to BCG Attributes I–VI (Tables Appendix A1–A6) with the following frequency:

- Attribute I: Historically Documented, Long-lived, or Regionally Endemic Taxa – 15 taxa
- Attribute II: Highly Sensitive Taxa - 54 taxa
- Attribute III: Intermediate Sensitive Taxa – 108 taxa
- Attribute IV: Intermediate Tolerant Taxa - 51 taxa
- Attribute V: Tolerant Taxa - 4 taxa
- Attribute VI: Non-native or Intentionally Introduced Taxa - 3 taxa
- X – Taxa not assigned to an attribute – 122 taxa

The fish experts assessed 38 calibration sites from the Puerto Rico surveys during the 2nd and 3rd workshops and several webinars. The fish experts agreed that all the stations had some degree of disturbance, including ubiquitous effects from fishing pressure and reef degradation. No sites were assigned to BCG level 2, so only conceptual rules were developed for level 2. BCG level 1 was not expected to occur in Puerto Rico and was not described conceptually or with model rules by the fish experts. All sites were rated as BCG levels 3–6, and intermediate levels were assigned as ‘+’ (exhibiting characteristic of the next best conditions but not enough to rank site in higher level) and ‘-’ (exhibiting characteristics that suggest somewhat worse conditions but not enough to rank site in lower level). This information was used to help define the

condition thresholds at which experts might assign sites to different BCG levels. The fish experts showed a high degree of agreement in their decisions. The “granularity” of the individual decisions was one third (the difference between level 4 and 4+), and this was used to estimate consistency of the experts. For the calibration sites, 85% of individual assessments were within one third of the BCG level of the group median, and 90% were within two thirds of the BCG level. Fig. 4 (top) shows the distribution of individual panelist scores compared to the group median for each site. Confirmation sites (11) were rated during the third workshop, resulting in fewer very close agreements compared to the calibration ratings: 78% of ratings were within one third of the BCG level of the panel median and 95% were within two thirds of the BCG level (Fig. 4 bottom). The lower agreements were likely due to some “drift” of panel members as a result of the intervening time between workshops. Quantifying the drift would require asking each expert how and why their decision differed from the decision rules. Because the drift was relatively small, we did not pursue this.

The narrative decision rules expressed during deliberations exhibited a general pattern of decreasing richness and biomass, especially of sensitive or specialist fish, as biological condition degrades (Table 3). Most of the narrative rules could be translated to numeric decision rules (Table 4). In BCG deliberations, the experts determined how the rules for each level were to be applied: (1) all rules must be met; (2) some rules have alternate rules (e.g., a very low percentage of tolerant individuals may substitute for a high percentage of sensitive individuals); or, (3) some number of rules for that level must be met (EPA, 2016). For example, the fish experts had higher expectations for fish communities in reef habitat than in hard-

Table 4

BCG reef fish assemblage decision rules. Numbers in parentheses are lower and upper bounds for group membership. Puerto Rico rules are based on 4 × 25 m belt transect data collected during 2010–2011 (Santavy et al., 2012). Florida rules are based on 15 m dia cylinder RVC point count data (Smith et al., 2011) collected during 2014–2016.

BCG metric	Narrative rules	Quantitative rules
BCG Level 2 (No survey samples were identified, rules are conceptual)		
Total taxa	Richness is high – valid taxa only	≥ 20 (15 - 25)
Rare, endemic & special species (Attribute I species)	Present	≥ 1
Highly sensitive taxa (Attribute II species)	Present	≥ 1 (0 - 2)
Proportion of all sensitive taxa (Attribute I, II, and III species)	Sensitive taxa constitute large proportion of species richness	≥ 50% (45 - 55)
Total biomass	High fish biomass – valid taxa only	Puerto Rico: ≥ 65 (50 – 80 g/m ²) Florida: ≥ 11.5 (kg/177 m ²)
Large groupers	Present (<i>Epinephelus</i> and <i>Mycteroperca</i>)	≥ 1 (0 - 1)
Large predators	Present	≥ 1 (0 - 2)
Piscivore individuals	Abundant	≥ 20 individuals
BCG Level 3		
Total taxa	Richness moderate to high – valid taxa only	≥ 15 (10 - 20)
Number of all sensitive taxa (Attribute I, I, and III species)	Sensitive taxa are a small to moderate proportion of fish species richness	≥ 6 (4 - 8)
Total biomass (g/m ²)	Total fish biomass is moderate to high – valid taxa only	Puerto Rico: ≥ 35 (30 – 40 g/m ²) Florida: ≥ 6.5 (5.6 – 7.4 kg/177m ²)
Piscivores	Presence of snappers or other piscivores	≥ 1
Parrotfish	Presence of large parrotfish	≥ 1 (0 - 2)
Damselfish	Don't dominate observed species	< 25% (20 - 30)
Groupers	Groupers present (<i>Dermatolepis</i> , <i>Epinephelus</i> , <i>Mycteroperca</i> , and <i>Cephalopholis</i>)	≥ 1
Rule application:	Reef Habitats: More stringent requirements Hard-bottom Habitats: Less stringent requirements	Require 6 of 7 rules Require 5 of 7 rules
BCG Level 4		
Total taxa	Richness low to moderate – valid taxa only	≥ 9 (4 - 14)
Number of all sensitive taxa (Attribute I, II, and III species)	Some sensitive taxa	≥ 3 (1 - 5)
Total biomass (g/m ²)	Low or higher – valid taxa only	Puerto Rico: ≥ 11 (7 – 15 g/m ²) Florida: ≥ 1.1 (0.7 – 1.5 kg/177 m ²)
BCG Level 5		
Total taxa	Sparse – valid taxa only	≥ 5 (2 - 8)
Total biomass (g/m ²)	Very low – valid taxa only	Islands: ≥ 2 (1 - 3) (g/m ²) Florida: ≥ 0.35 (0.18– 0.52 kg/177 m ²)
BCG Level 6 Does not meet Level 5 rules		

^a See Supplemental Material Appendix A Table A7.

bottom habitats. Therefore, in BCG level 3, seven rules are expressed; however, six rules must be met to assign the BCG level 3 in reef habitat, while only five must be met in hard-bottom habitats.

Model performance is summarized in Table 5, showing number and percent of model assessments compared to expert panel assessments. Model output was expressed as membership of a site in a BCG level. While model output was potentially continuous from zero to one, input data and rules were often expressed as whole numbers (e.g., number of species in a family), and model output often included exact half memberships (0.5) of adjacent BCG levels. To avoid false precision greater than the model or data could support, we interpreted intermediate memberships (0.4 to 0.6) as

half memberships of two adjacent levels. This was used only to compare model results to panel consensus. The panel did not consider a half-level mismatch with their consensus to be a meaningfully different assessment, and a half-level was similar to the spread in ratings among panel members. Accordingly, the panel did not adjust ratings or modify rules for small mismatches. On average, the quantitative model was 92% accurate in replicating the expert panel assessments within one-half BCG level for the calibration datasets, and 82% accurate for the confirmation dataset. There were no mismatches greater than one BCG level.

The 4th workshop focused on potential transferability of the Puerto Rico model to a different jurisdiction (e.g., Florida Keys) and possible

Table 5

Performance of BCG quantitative fish model for calibration and confirmation datasets, by Puerto Rican reef fish panel. “Better” and “worse” indicate model assessment of coral reef condition compared to panel (e.g., “better” if model assessed BCG Level 2, but panel assessed BCG Level 3, and so forth). Percent differences are reported with the number of differences followed in parentheses.

(Source: Bradley et al., 2016)

Dataset	Model Performance Difference					Total
	Model 1 level better	Model 1/2 level better	Exact match	Model 1/2 level worse	Model 1 level worse	
Calibrate	5% (2)	0 (0)	79% (30)	13% (5)	3% (1)	100% (38)
Confirm	9% (1)	9% (1)	73% (8)	0 (0)	9% (1)	100% (11)

management applications of the model. To test transferability of the model, the experts rated 14 stations collected in the Florida Keys and Dry Tortugas at depths shallower than 16 m, which were co-sampled by both the fish and benthic teams (Bohnsack and Bannerot, 1986). The stations were selected by the Reef Visual Census (RVC) leads across a stressor gradient: water quality (low anthropogenic impact – Dry Tortugas, low-moderate impact – Florida Keys forereef, and high impact – Hawk's Channel); and fishing pressure based upon management zones (low – Dry Tortugas National Park, medium – Florida Keys, Marine Protected Areas, high – Florida Keys outside of Marine Protected Areas). The quantitative BCG model developed for Puerto Rico was 79% accurate in replicating the expert panel assessments within one-half BCG level for the Florida Keys calibration. The biomass metric was the rule that was not met in the mis-matched sites. The experts felt that species attribute levels might need to be revisited based on location, particularly because fishing pressure varies significantly by jurisdiction.

The experts were asked to consider how the benthic and fish models could be used together for evaluating sites. They applied the BCG rules for both assemblages to several sites. One example scenario was when the benthic organisms met the benthic level 3 rules, but the fish only met the fish level 5 rules. The panel assessed the site as *degraded but with high potential for recovery of the fish population because important habitat and food for fish present*.

4. Discussion

Since 2005, several U.S. states and other entities (e.g., river basin associations and counties) have either calibrated, or are in the process of calibrating, the BCG for freshwater aquatic ecosystems (EPA, 2016). These methods have been shown to be applicable to several stream and riverine environments and taxa: perennial freshwater streams for benthic macroinvertebrates (primarily insects) throughout the United States; freshwater fishes in streams and lakes (EPA, 2016; Gerritsen et al., 2017); and benthic diatoms (Hausmann et al., 2016). This paper extends that utility to reef fishes in coral reef ecosystems.

A regional panel of experts assigned fish species inhabiting Puerto Rico's near-shore linear coral reefs to attributes of sensitivity to human disturbance, natural prevalence, historic species importance in the Caribbean, and native or exotic origin. The experts developed fish rules for six levels of coral reef condition, with a well-defined narrative for each level.

A quantitative decision model had a high degree of fidelity to the expert decisions: the model replicated the expert consensus within one BCG level for 100% of sites and replicated the expert consensus within a half BCG level for 82% to 92% of the sites. This degree of predictive accuracy is as good or better than the examples described for freshwater systems (Gerritsen et al., 2017; Hausmann et al., 2016).

BCG model development, calibration, and validation were successful for the available data using the expert process described. However, there were several issues that arose during workshop and webinar discussions that could be further investigated for incorporation into future BCG model revisions or model result interpretations. The issues addressed fish characteristics (size-structure expectations, longevity, and reproductive strategies), site condition effects (the undisturbed baseline condition and water quality indicators), fish community variations related to habitat (habitat classifications and inter-habitat connectivity), and data collection methods (consistency and sufficiency). These issues are discussed below with possible approaches for resolution.

4.1. Size-structure expectations

Observations of juvenile and adult fish at a reef site might indicate that a full life cycle is supported at the site, inferring inter-habitat connectivity at the site for certain species. With observation of a single

life stage, experts were uncertain about the propensity of the reef site to support nursery function for juveniles or maintenance of an adult population. Therefore, in the BCG rating process, experts requested information about the size-structured abundance distribution of the fish observed. The experts were familiar with critical sizes that might indicate single or multiple life stages and could relate the size and life-stage information to the biological condition of the reef fish community. Unfortunately, fish sizes were recorded at 5-cm intervals for all species, but association of juvenile and adult stages had not yet been completed for this dataset. A listing of juvenile and adult size ranges for fish species might be available in the literature or could be created by the experts based on expert judgment. Stevens et al.'s (2019) recent synthesis of life history demographic parameters for Florida and Caribbean reef fishes could greatly facilitate these efforts. Enumeration of juvenile and adults for future rating exercises would allow calculation of life-stage metrics for reef fish. The life stage metrics might allow better discrimination of BCG levels and inter-habitat connectivity.

4.2. Longevity and reproductive strategies

In coral reef ecosystems, large-bodied, slow-growing, late-maturing fishes (K-strategists) are generally more sensitive to exploitation than faster-growing, shorter-lived species (r-strategists) (Beverton and Holt, 1957; Man et al., 1995; Jennings et al., 1998; Coleman et al., 2000; Goodwin et al., 2006; Ault et al., 1998, 2008). Consideration of K/r strategies informs coral reef fish population responses to environmental stress, which is largely determined by life-history traits with K-strategists being more susceptible to fishing pressure than r-strategists (Musick et al., 2000; Ault et al., 2005, 2008, 2014). The BCG Attribute definitions (Davies and Jackson, 2006) include considerations of these life history traits: Attributes I and II include long-lived, late maturing, low fecundity species; while Attributes IV and V include early colonizers with rapid turn-over times and “boom/bust” population characteristics. However, species-specific life history data was not included in this BCG evaluation and was therefore not considered in the assignment of species to coral reef BCG attributes.

4.3. Undisturbed baseline condition

A challenge in developing the coral reef BCG was the difficulty in determining reference conditions for biological integrity because fish populations in Puerto Rico have been exploited since at least the 15th century and were already decimated by the 1950s (Goreau, 1959; Jackson, 1997; Greenstein et al., 1998; Jackson and Sala, 2001; Jackson et al., 2001; Jameson et al., 2003; Pandolfi et al., 2003), and no anthropogenically unimpacted reference sites were available in Puerto Rico. During the 1st workshop, the BCG experts discussed using the Healthy Reefs Initiative (HRI) data thresholds for the Mesoamerican reef as possible reference conditions for Puerto Rico. HRI based these thresholds on the wider Atlantic and Gulf Rapid Reef Assessment (AGRRA) data for the Wider Caribbean, which the experts considered to be comparable to the U.S. EPA methodology. The fish experts reviewed the HRI thresholds (HRI, 2012), and then, through a methodical, facilitated process, used their expert judgment to define a preliminary set of attributes for reference conditions for Puerto Rico. Calibrating the model with surveys from relatively unimpacted areas elsewhere in the Caribbean may eventually be useful in further testing the reference condition attributes; however, differences in fish observation protocols may present a complication.

4.4. Water quality indicators

The U.S. EPA coral reef research in south Puerto Rico and USVI evaluated potential relationships between reef condition metrics and estimates of coral reef stressors, with each stressor incorporating

relative proximity from reef survey locations to a human disturbance. Methods to designate disturbance ranged in complexity from general sources of disturbance such as towns and industrial centers (Fisher et al., 2008; Oliver et al., 2014), and distance to bays with impaired water quality (Oliver et al., 2018), ST (Oliver et al., 2018), to measures that integrate spatially-explicit land use/land cover such as the Landscape Development Intensity Index (LDI) (Oliver et al., 2011, 2018) and ST (Oliver et al., 2018). Reef survey methods progressed from an initial focus on stony coral communities (Fisher, 2007) to include fish, gorgonians, and sponges (Santavy et al., 2012). The LDI showed the most, and consistently inverse relationships with measures of coral cover, rugosity, colony size and species diversity (Oliver et al., 2011, 2018), consistent with other studies (Rogers, 1990; Fabricius and De'ath, 2001; Fabricius et al., 2005; Cleary et al., 2006) and a general hypothesis that intensifying land-based human activities in watersheds is associated with a decline of adjacent reefs. The BCG expert panelists were concerned about fishing pressure and sedimentation threat to fish communities. Large scale modeling of sediment plumes and potential delivery to Indonesian reefs offers a potential approach to coupling watershed sediment production with an ocean transport model that accounts for current dynamics and particle settling (Rude et al., 2016). The predictive potential for LDI (Oliver et al., 2011) to indicate deteriorated reef condition was demonstrated despite simple assumptions applied to connect St. Croix watershed LDI values to reef survey locations without accounting for ocean currents, wind or bathymetry that undoubtedly influence transport of specific stressors to coral reef communities. Refinements in stressor modeling needed to inform a comprehensive stressor gradient for the BCG require data with appropriate scale to the reef communities of interest. Coral reef stressor gradients cannot be as clearly defined as those in streams. Streams have a distinct catchment and actual flow distance from a source to particular sampling sites can be measured. Coral reefs and all coastal marine ecosystems are not linear systems, and land-based stressors from multiple watersheds may impact a given reef as they become dispersed by wave action, wind and oceanic currents.

The journal's reviewers recommended that we use multivariate statistical techniques to examine indicators of different environmental gradients by linking fish species composition to proxies of exposure to various stressors. Ordination and cluster permutation analyses (i.e., PRIMER-e Ver. 7; Clarke and Gorley, 2015; Clarke et al., 2014) were used to identify how the BCG Level groups were related within and among the four BCG Levels 3–6 (Table 6), and how Levels were related to the environmental gradients used in our study (Fig. 5). The environmental variables tested were distance to shore, distance to shelf, distance to disturbance, sediment threat (ST), reef rugosity, coral colony density, coral species richness, and percent two-dimensional coral cover. A cluster analysis identified similarly patterns among the evaluated sites to find major fish species associated with each BCG Level (Table 7). Ordination techniques (non-metric Multiple-dimensional Scaling) showed how environmental variables mapped in relationship to the fish BCG Levels.

Table 6

BCG Level relationship to each other Level (BCG Levels 3–6). Dissimilarity calculated using Euclidian Distance Coefficient. Test results from ANOSIM and SIMPER calculated among group dissimilarities.

BCG Levels	R statistic	Significance Level	Among Group Dissimilarity
3 vs 4	0.013	33.9	58.7
3 vs 5	0.681	0.1 ***	74.2
3 vs 6	0.991	0.7 ***	91
4 vs 5	0.439	1.3 **	70.7
4 vs 6	0.934	0.6 ***	80
5 vs 6	0.607	6.7	90

Three environmental variables were seen to explain some of the fish species composition (i.e., ST, percent live coral cover (2-D) and rugosity). Increased sediment threat was related to decreasing biological condition (BCG Levels 5 and 6), whereas increased rugosity and live 2D coral cover were correlated to good and fair BCG Levels (3 and 4). As rugosity increased so did live coral cover (2D) and coral density, all variables related to increased ecosystem complexity through higher topography and amount of live coral on reefs essential for the habitats, foraging and refugia needs of many reef fish species. The distance to shore, distance to shelf and distance to disturbance as defined did not provide any explanatory information for the biotic patterns. This might imply the need to consider a different approach for estimating distance from disturbance.

Based on fish density and biomass, BCG Levels 3 and 4 were not significantly different from one another, and BCG Levels 5 and 6 were not significantly different from one another (Table 6). The primary fish species within BCG Levels 3 and 4 were *Thalassoma bifasciatum* and *Sparisoma aurofrenatum*, with eight other fish species also present in both BCG Levels 3 and 4. This similarity caused BCG Levels 3 and 4 to be not different. This is an example where expert judgment considering the importance of the biology and ecology of different species and their functional contributions could outweigh the importance of just density or biomass of fish species determined by a statistical analysis. These statistical analyses corroborate the conclusion that expert judgment is essential in this process to provide the knowledge of nuanced ecological functions for which data were not available.

4.5. Habitat classifications

In coral reef ecosystems, there is a strong positive correlation of habitat complexity with fish species richness (Luckhurst and Luckhurst, 1978; Carpenter et al., 1981; Roberts and Ormond, 1987; McClanahan, 1994; McCormick, 1994; Green, 1996; Friedlander and Parrish, 1998; Sale, 1991; Friedlander et al., 2003; Gratwicke and Speight, 2005a, 2005b; Kuffner et al., 2007; Pittman et al., 2007; Aguilar-Perera and Appeldoorn, 2008; Walker et al., 2009; Smith et al., 2011). Reef fish data can be associated with the NOAA benthic habitat maps to help determine the expected assemblages in different habitats throughout a mapped space (Pittman et al., 2007). For example, the main factors used to determine reef fish assemblages in biogeographic regions on the Southeast Florida reef tract were reef vs. hardbottom substrates, depth, relief, and geographic space (Smith et al., 2011; Fisco, 2016; Ames, 2017). Important species traits might show patterns only found at in-shore or offshore survey sites, exhibiting a distribution restricted by water depth, or geographically widespread across depth, which might influence their potential role as indicators in the BCG model. For example, the absence of a fish species from a nearshore site may not be indicative of the condition of the coral reef ecosystem if that species' range does not occur in nearshore reefs. Similarly, the frequent occurrence of a species in waters known to be impaired due to the influx of land-based pollutants may mean the species is more pollution-tolerant than a species found only in waters that do not contain influxes of land-based pollutants, assuming benthic variables are similar in both locations. The combination of the depth distribution, distance to shore, and the frequency of occurrence provide an indication of relative abundance for each fish species and a simplified geographical habitat width for each species. Improved information on species and functional traits for Caribbean fish could aid in improving and interpreting results when applying the BCG fish model to other Caribbean locations.

4.6. Inter-habitat connectivity

As mentioned, the distance to shore was recognized as a possible site variable because it represented inter-habitat connectivity among larval, juvenile, and adult fish habitats. Beneficial off-reef habitats for reef fish are not all near the shore. These off-reef habitats must be




Fig. 5. Non-metric multidimensional scaling (nMDS) plot of the fish density composition at the 38 sites that were used in development of the BCG model clustered by Bray-Curtis similarity (Stress = 0.19). a) Vector plot overlay shows the direction of linear increase of environmental variable concentrations, and the multiple correlation of each (transformed) variable on the 2D ordination points. The significant gradient vectors are bolded. b-d) Bubble plots overlay the same nMDS plot but with circles of increasing size representing the environmental variable at those sites.

Table 7
Preliminary narrative BCG with four distinct levels of condition for all reef assemblages: very good/excellent; good; fair; and poor.

BCG Level	Species	Avg. Similarity	Similarity SD	% Individual Contribution	% Cumulative Contribution
BCG Level 3					
42.34% average within group similarity	<i>Thalassoma bifasciatum</i>	6.43	2.06	15.19	15.19
	<i>Sparisoma aurofrenatum</i>	4.44	3.3	10.49	25.68
	<i>Scarus iseri</i>	2.87	1.24	6.78	32.46
	<i>Stegastes partitus</i>	2.83	1.19	6.68	39.14
	<i>Ocyurus chrysurus</i>	2.48	1	5.86	45
	<i>Acanthurus bahianus</i>	2.26	0.97	5.35	50.35
	<i>Acanthurus coeruleus</i>	2.17	1.01	5.11	55.46
	<i>Microspathodon chrysurus</i>	2.05	1.02	4.85	60.31
	<i>Stegastes adustus</i>	1.96	0.68	4.64	64.95
	<i>Acanthurus chirurgus</i>	1.3	0.57	3.08	68.03
<i>Sparisoma viride</i>	1.29	0.69	3.05	71.09	
BCG Level 4					
40.69% average within group similarity	<i>Thalassoma bifasciatum</i>	9.81	4.21	24.11	24.11
	<i>Sparisoma aurofrenatum</i>	4.1	1.27	10.07	34.19
	<i>Stegastes partitus</i>	3.19	0.88	7.84	42.03
	<i>Acanthurus bahianus</i>	2.45	0.77	6.02	48.05
	<i>Ocyurus chrysurus</i>	2.32	0.77	5.71	53.75
	<i>Acanthurus coeruleus</i>	2.14	0.79	5.25	59.01
	<i>Microspathodon chrysurus</i>	2.08	0.78	5.12	64.13
	<i>Halichoeres bivittatus</i>	2.04	0.66	5.01	69.14
	<i>Scarus iseri</i>	1.91	0.65	4.68	73.82
BCG Level 5					
26.35% average within group similarity	<i>Halichoeres bivittatus</i>	7.26	0.91	27.54	27.54
	<i>Stegastes adustus</i>	7.06	0.88	26.8	54.34
	<i>Thalassoma bifasciatum</i>	6.48	0.84	24.61	78.95
BCG Level 6					
65.45% average within group similarity	<i>Stegastes diencaeus</i>	39.93	SD=0!	61.01	61.01
	<i>Stegastes leucostictus</i>	25.51	SD=0!	38.99	100

accessible to benefit certain reef fish species. Knowledge of the inter-habitat connectivity between sampling locations and off-reef habitats and the necessity of such habitats for each fish species would improve assessments and interpretation of assessments for reef fish samples and sites. The experts recommended that high-resolution reef bottom topography (e.g., LIDAR) would allow for better estimation of inter-habitat connectivity. With high-resolution topography, features related to inter-habitat connectivity would be recognizable and quantifiable. High-resolution topography would also indicate elements of rugosity as well as inter-habitat connectivity, allowing characterization of broad-scale relief and a possible basis for classification of reefs.

4.7. Data collection methods

Some experts felt that the data collection methods (Santavy et al., 2012; NOAA, 2013a) were not optimal for assessing fish assemblages. Transect methods for reef fish are biased in that some species disperse before they are counted (Ydenberg and Dill, 1986; Januchowski-Hartley et al., 2011; Lindfield et al., 2014; Emslie et al., 2018), significantly impacting estimates of reef fish richness and density (Chapman et al., 1974; Kulbicki, 1998; Kulbicki and Sarraména, 1999). Some experts recommended using the stationary point count (SPC) fish survey method (Bohnsack and Bannerot, 1986; Ault et al., 1998, 2005, 2014; Brandt et al., 2009; Richards et al., 2011; Smith et al., 2011), which allows the fishes time to adjust to the presence of a diver before counting begins. In this method, pairs of divers record the number, size and species of all fishes observed within cylinders visually estimated at 15 m in diameter. Other fish experts preferred the transect method (Santavy et al., 2012; NOAA, 2013a), suggesting that the fish counter has more opportunity to observe cryptic species than when using the point-count method. They also suggested that the belt transect is also much better for low-visibility environments, which are very common in Puerto Rico and the U.S. Virgin Islands. The fish experts reached consensus that a single commonly used method of counting fish would be much better than using different methods and would reduce variance in getting density estimates. The fish experts also recommended revising the field method for measuring topographic complexity (e.g., rugosity) for each reef station where fish were counted. They felt that methods which measure vertical relief along the entirety of a transect (e.g., Dustan et al., 2013; NOAA, 2013b) could provide more information about reef rugosity.

4.8. Application

To facilitate use by water quality managers, the BCG rule application will be automated once finalized. Additionally, clear instructions will be provided for each BCG fish rule. For example, the fish rule “at least one large-bodied parrotfish present” requires clarification of what scientists mean by “large-bodied parrotfish” (Appendix A Table A7). A precise definition is being documented for each rule, and guidance material is being developed so the tool can be easily applied and interpreted.

The BCG provides a powerful framework for an operational monitoring and assessment program, for communicating resource condition to the public, and for assisting in management decisions to protect or remediate water resources. The levels of the BCG are biologically recognizable, measurable stages in condition of coral reef ecosystems. As such, the BCG can be used to inform biological assessments of Caribbean coral reefs. The BCG is a defensible means to translate scientific understanding to support both regulatory and non-regulatory water quality and natural resource programs and inform biocriteria development. Biologically based aquatic life uses coupled with numeric biological criteria provide a direct measure of the aquatic resource that is being protected (e.g., coral reefs), complimenting the stressor and

exposure criteria which are comprised of chemical, toxicological, and physical parameters. The BCG provides a framework that can help relate chemical, physical and biological assessments and criteria for a more integrated, comprehensive evaluation of the condition of a waterbody. Additionally, the fish and benthic BCG models can be combined for a robust interpretation since these communities can respond differently to stressors.

While the BCG model was developed using data from Puerto Rico, it is important to note that the BCG is a general framework that could potentially be applied to other coral reef ecosystems, as demonstrated by the proof-of-concept work done using sites from Florida Keys and Dry Tortugas. In order to use the BCG, states and territories would need to adapt it to their own coral reef habitat and monitoring data and develop a numeric model scheme specific to their jurisdiction. In summary, (1) the BCG conceptual framework is applicable to other coral reef ecosystems; (2) the methods used to develop the BCG in Puerto Rico are likely applicable to other coral reef ecosystems (e.g., the process to elicit expert judgment); (3) the qualitative rules may be applicable, but will require vetting by regional experts, using regional datasets to test and refine the rules; and, (4) quantitative rules are jurisdiction-specific.

Many organizations produce periodic report cards that provide information on coral reef status and trends, as well as management strategies being employed to improve these ecosystems (e.g., The Healthy Reefs Initiative, The Atlantic and Gulf Rapid Reef Assessment, The Florida Keys National Marine Sanctuary, Australia Institute of Marine Science (AIMS) Long Term Monitoring Program for The Great Barrier Reef, etc.). The BCG is complementary to these report cards and can be used to help communicate monitoring results.

5. Conclusion

Coral reef resources have historically been managed by natural resource agencies employing a variety of approaches, including fisheries regulations, marine protected areas, and endangered species protection (Bradley et al., 2010). Regulations, including those for fisheries and protected species may be enhanced by an integrated and ecologically broad systems science approach (e.g., Ault et al., 2005). Such an integrated approach is the coral reef fish BCG model. For example, water quality managers could use the BCG to distinguish high-quality coral habitats for greater protection, or to gauge the effectiveness of management actions to meet restoration goals for coral reefs adjacent to urban and agricultural areas. Successful BCG development for Puerto Rico reef fish communities provides a common framework that can be used by other jurisdictions. However, quantitative calibrations and validations may likely be region specific. Therefore, broader application in the Caribbean or the Pacific will require additional focused study.

CRediT authorship contribution statement

Patricia Bradley: Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Resources, Data curation, Writing - original draft, Writing - review & editing, Visualization, Supervision, Project administration, Funding acquisition. **Ben Jessup:** Conceptualization, Methodology, Software, Validation, Formal analysis, Investigation, Resources, Data curation, Writing - original draft, Writing - review & editing, Visualization, Project administration. **Simon J. Pittman:** Formal analysis, Investigation, Writing - review & editing. **Christopher F.G. Jeffrey:** Methodology, Formal analysis, Investigation, Data curation, Writing - original draft, Writing - review & editing. **Jerald S. Ault:** Methodology, Formal analysis, Investigation, Data curation, Writing - review & editing. **Lisamarie Carrubba:** Methodology,

Formal analysis, Investigation, Writing - review & editing. **Craig Lilyestrom**: Methodology, Investigation, Writing - review & editing. **Richard S. Appeldoorn**: Methodology, Investigation, Writing - review & editing. **Michelle T. Schärer**: Methodology, Investigation, Writing - review & editing. **Brian K. Walker**: Methodology, Investigation, Writing - review & editing. **Melanie McField**: Methodology, Investigation, Writing - review & editing. **Deborah L. Santavy**: Conceptualization, Methodology, Investigation, Writing - review & editing. **Tyler B. Smith**: Methodology, Investigation, Writing - review & editing. **Graciela García-Moliner**: Methodology, Investigation, Writing - review & editing. **Steven G. Smith**: Formal analysis, Investigation, Data curation, Writing - review & editing. **Evelyn Huertas**: Methodology, Investigation, Writing - review & editing. **Jeroen Gerritsen**: Conceptualization, Methodology, Software, Validation, Investigation, Data curation, Writing - review & editing, Project administration. **Steven G. Smith**: Conceptualization, Methodology, Formal analysis, Investigation, Writing - review & editing, Visualization. **Christina Horstmann**: Formal analysis, Investigation, Writing - review & editing. **Susan K. Jackson**: Conceptualization, Methodology, Validation, Formal analysis, Investigation, Data curation, Writing - review & editing, Visualization, Project administration, Funding acquisition.

Declaration of competing interest

The authors declare that they have no known competing financial

interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgments

We would like to thank the Caribbean Coral Reef Institute for hosting the 1st and 4th workshops on Magueyes Island; the El Yunque National Forest Headquarter for providing the conference facilities for the 2nd workshop; and the Caribbean Landscape Cooperative (CLCC) and the International Institute of Tropical Forestry (IITF) for providing the conference facilities for the 3rd workshop. Also, we wish to acknowledge the continuing support received from William Fisher and Buddy LoBue (U.S. EPA).

The U.S. Environmental Protection Agency (EPA) through its Office of Research (ORD) and Development and Office of Water (OW) funded and collaborated in the research described here under two contracts to Great Lakes Environmental Center: Contract EP-C-09-001, Work Assignment #3-01, and Contract #EP-C-14-016, Work Assignment 1-1. The views expressed in this paper are those of the authors and do not necessarily reflect the views or policies of the U.S. EPA. This document has been reviewed by the U.S. EPA, Office of Research and Development, Office of Water, and Region 2, and has been approved for publication. Any mention of trade names, products, or services does not imply an endorsement by the U.S. Government or the U.S. EPA. The U.S. EPA does not endorse any commercial products, services, or enterprises.

Appendix A. Assignment of Fish Species to BCG Attribute Levels I-VI and Large bodied fish species

Fish Species Attribute Assignments made by the expert during multiple workshops and webinars. Assigned BCG attributes for fish species are based upon sensitivity to fishing pressure and sediment stress across the US Caribbean and South Florida. Assignment of fish species to BCG Attribute Levels I-VI: Level I species are historically documented, long lived, or regionally endemic taxa (Table Appendix A1); Level II species are highly sensitive to fishing pressures and sediment threats (Table Appendix A2); Level III species are intermediately sensitive taxa to fishing pressures and sediment threats (Table Appendix A3); Level IV species are intermediately tolerant taxa to fishing pressures and sediment threats (Table Appendix A4); Level V species are tolerant taxa to fishing pressures and sediment threats; and Level VI species are non-native or intentionally introduced species (Table Appendix A5). Abbreviations for the trophic guilds are: H=herbivore, P=piscivores, I=invertivore, and Z=zoo-planktivore (from Caldow et al. 2009). Piscivore size indicated as either large (P-L) or small (P-S).

Table Appendix A1

BCG Attribute I species are historically documented, long-lived, or regionally endemic taxa in the US Caribbean and Florida.

Species Name	Common Name	Trophic Guild
<i>Acanthostracion polygonius</i>	Honeycomb cowfish	I
<i>Acanthostracion quadricornis</i>	Scrawled cowfish	I
<i>Carcharhinus limbatus</i>	Blacktip shark	P-L
<i>Carcharhinus perezii</i>	Caribbean reef shark	P-L
<i>Epinephelus itajara</i>	Atlantic goliath grouper	P-L
<i>Epinephelus morio</i>	Red grouper	I
<i>Epinephelus striatus</i>	Nassau grouper	P-L
<i>Mycteroperca bonaci</i>	Black grouper	P-L
<i>Mycteroperca interstitialis</i>	Yellowmouth grouper	P-S
<i>Mycteroperca tigris</i>	Tiger grouper	P-L
<i>Mycteroperca venenosa</i>	Yellowfin grouper	P-L
<i>Scarus coelestinus</i>	Midnight parrotfish	H
<i>Scarus coeruleus</i>	Blue parrotfish	H
<i>Scarus guacamaia</i>	Rainbow parrotfish	H
<i>Sphyrna mokarran</i>	Great hammerhead shark	P-L

Table Appendix A2

BCG Attribute II species are highly sensitive taxa to fishing pressures and sediment threats in the US Caribbean and Florida.

Species Name	Common Name	Trophic Guild
<i>Aetobatus narinari</i>	Spotted eagle ray	I
<i>Aluterus scriptus</i>	Scrawled filefish	I
<i>Amblycirrhitis pinos</i>	Red-spotted hawkfish	Z

(continued on next page)

Table Appendix A2 (continued)

Species Name	Common Name	Trophic Guild
<i>Anisotremus surinamensis</i>	Black margate	I
<i>Astrapogon stellatus</i>	Conchfish	I
<i>Aulostomus maculatus</i>	Trumpetfish	P-S
<i>Cantherhines macrocerus</i>	America white-spotted filefish	I
<i>Cantherhines pullus</i>	Orange-spotted filefish	H
<i>Caranx crysos</i>	Blue runner	P-S
<i>Caranx hippos</i>	Crevalle jack	P-L
<i>Cephalophilus furcifer</i>	Atlantic creolefish	Z
<i>Chaenopsis limbaughi</i>	Yellowface pikeblenny	I
<i>Chaetodipterus faber</i>	Atlantic spadefish	I
<i>Chromis cyanea</i>	Blue chromis	Z
<i>Chromis multilineata</i>	Brown chromis	Z
<i>Clepticus parrae</i>	Creole wrasse	Z
<i>Dactylopterus volitans</i>	Flying gurnard	I
<i>Dasyatis americana</i>	Southern stingray	I
<i>Elacatinus genie</i>	Cleaner goby	H
<i>Elacatinus multifasciatus</i>	Green-banded goby	I
<i>Elacatinus oceanops</i>	Neon goby	I
<i>Elacatinus prochilos</i>	Broad stripe goby	I
<i>Elacatinus saucrum</i>	Leopard goby	I
<i>Enchelycore nigricans</i>	Viper moray	P-S
<i>Fistularia tabacaria</i>	Blue-spotted cornet fish	P-S
<i>Galeocerdo cuvier</i>	Tiger shark	P-L
<i>Ginglymostoma cirratum</i>	Nurse shark	P-L
<i>Gramma loreto</i>	Fairy basslet	I
<i>Haemulon chrysargyreum</i>	Smallmouth grunt	I
<i>Halichoeres radiatus</i>	Puddingwife	I
<i>Heteropriacanthus cruentatus</i>	Glasseye snapper	Z
<i>Holacanthus ciliaris</i>	Queen angelfish	I
<i>Holacanthus tricolor</i>	Rock beauty	I
<i>Hypoplectrus gemma</i>	Blue hamlet	I
<i>Hypoplectrus hybrid</i>	Hybrid hamlet	I
<i>Lachnolaimus maximus</i>	Hogfish	I
<i>Lactophrys triqueter</i>	Smooth trunkfish	I
<i>Lactophrys bicaudalis</i>	Spotted trunkfish	I
<i>Lactophrys trigonus</i>	Trunkfish	I
<i>Lutjanus analis</i>	Mutton snapper	I
<i>Lutjanus cyanopterus</i>	Cubera snapper	P-L
<i>Lutjanus jocu</i>	Dog snapper	P-L
<i>Melichthys niger</i>	Black durgon	H
<i>Negaprion brevirostris</i>	Lemon Shark	P-L
<i>Pareques acuminatus</i>	Highhat	I
<i>Priacanthus arenatus</i>	Bigeye	I
<i>Priolepis hipoliti</i>	Rusty goby	I
<i>Prognathodes aculeatus</i>	Longsnout butterflyfish	I
<i>Scomberomorus regalis</i>	Cero	P-S
<i>Seriola dumerili</i>	Greater amberjack	P-L
<i>Seriola rivoliana</i>	Almaco jack	P-L
<i>Serranus tigrinus</i>	Harlequin bass	I
<i>Thalassoma bifasciatum</i>	Bluehead	I
<i>Trachinotus falcatus</i>	Permit	I
<i>Trachinotus goodei</i>	Palometa	P-S
<i>Xanthichthys ringens</i>	Sargassum triggerfish	Z

Table Appendix A3

BCG Attribute III fish species are intermediately sensitive taxa to fishing pressure and sediment threats in the US Caribbean and Florida.

Species Name	Common Name	Trophic Guild
<i>Abudefduf taurus</i>	Night sergeant	H
<i>Acanthemblemaria aspera</i>	Roughhead blenny	I
<i>Acanthemblemaria maria</i>	Secretary blenny	I
<i>Acanthemblemaria spinosa</i>	Spinyhead blenny	I
<i>Acanthurus chirurgus</i>	Doctorfish	H
<i>Acanthurus coeruleus</i>	Blue tang	H
<i>Acanthurus tractus</i>	Ocean surgeonfish	H
<i>Apogon aurolineatus</i>	Bridle cardinalfish	Z
<i>Apogon binotatus</i>	Barred cardinalfish	Z
<i>Apogon lachneri</i>	Whitestar cardinalfish	Z
<i>Apogon quadrisquamatus</i>	Sawcheek cardinalfish	Z
<i>Astrapogon puncticulatus</i>	Blackfin cardinalfish	I

(continued on next page)

Table Appendix A3 (continued)

Species Name	Common Name	Trophic Guild
<i>Balistes vetula</i>	Queen triggerfish	I
<i>Bodianus pulchellus</i>	Spotfin hogfish	I
<i>Bodianus rufus</i>	Spanish hogfish	I
<i>Canthidermis sufflamen</i>	Ocean triggerfish	I
<i>Caranx latus</i>	Horse-eye jack	P-S
<i>Caranx lugubris</i>	Black jack	P-L
<i>Centropomus undecimalis</i>	Common snook	P-S
<i>Centropyge aurantonotus</i>	Flameback angelfish	H
<i>Cephalopholis cruentata</i>	Graysby	P-S
<i>Cephalopholis fulva</i>	Coney	P-S
<i>Chaetodon capistratus</i>	Four-eye butterflyfish	I
<i>Chaetodon ocellatus</i>	Spotfin butterflyfish	I
<i>Chaetodon striatus</i>	Banded butterflyfish	I
<i>Chilomycterus antennatus</i>	Bridled burrfish	I
<i>Chromis insolata</i>	Sunshine fish	Z
<i>Coryphopterus dicrus</i> *	Colon goby	I
<i>Coryphopterus eidolon</i> *	Pallid goby	I
<i>Coryphopterus lipernes</i>	Peppermint goby	I
<i>Cosmocampus elucens</i>	Shortfin pipefish	I
<i>Diodon holocanthus</i>	Balloonfish	I
<i>Echidna catenata</i>	Chain moray	I
<i>Elacatinus chancei</i>	Shortstripe goby	I
<i>Elacatinus louisae</i>	Spotlight goby	I
<i>Emmelichthys atlanticus</i>	Bonnetmouth	P-S
<i>Epinephelus adscensionis</i>	Rock hind	I
<i>Epinephelus guttatus</i>	Red hind	P-S
<i>Equetus lanceolatus</i>	Jackknife fish	I
<i>Equetus punctatus</i>	Spotted drum	I
<i>Gymnothorax miliaris</i>	Goldentail moray	P-S
<i>Gymnothorax vicinus</i>	Purplemouth moray	P-S
<i>Haemulon album</i>	Margate (white)	I
<i>Haemulon carbonarium</i>	Caesar grunt	I
<i>Haemulon flavolineatum</i>	French grunt	I
<i>Haemulon macrostomum</i>	Spanish grunt	I
<i>Haemulon parra</i>	Sailors choice	I
<i>Halichoeres garnoti</i>	Yellowhead wrasse	I
<i>Halichoeres maculipinna</i>	Clown wrasse	I
<i>Halichoeres pictus</i>	Rainbow wrasse	I
<i>Hippocampus reidi</i>	Longsnout seahorse	I
<i>Holocentrus adscensionis</i>	Squirrelfish	I
<i>Holocentrus rufus</i>	Longspine squirrelfish	I
<i>Hypoplectrus aberrans</i>	Yellowbelly hamlet	I
<i>Hypoplectrus chlorurus</i>	Yellowtail hamlet	I
<i>Hypoplectrus guttavarius</i>	Shy hamlet	I
<i>Hypoplectrus indigo</i>	Indigo hamlet	I
<i>Hypoplectrus nigricans</i>	Black hamlet	P-S
<i>Hypoplectrus puella</i>	Barred hamlet	I
<i>Hypoplectrus randallorum</i>	Tan hamlet	I
<i>Hypoplectrus unicolor</i>	Butter hamlet	P-S
<i>Kyphosus sectator</i>	Chub (Bermuda/yellow)	H
<i>Labrisomus nuchipinnis</i>	Hairy blenny	I
<i>Liopropoma rubre</i>	Peppermint basslet	I
<i>Lutjanus buccanella</i>	Blackfin snapper	P-S
<i>Lutjanus mahogoni</i>	Mahogany snapper	P-S
<i>Lutjanus synagris</i>	Lane snapper	P-S
<i>Malacanthus plumieri</i>	Sand tilefish	I
<i>Malacoctenus aurolineatus</i> *	Goldline blenny	I
<i>Malacoctenus macropus</i> *	Rosy blenny	I
<i>Malacoctenus versicolor</i>	Barfin blenny	I
<i>Megalops atlanticus</i>	Tarpon	P-L
<i>Microspathodon chrysurus</i>	Yellowtail damselfish	H
<i>Monacanthus ciliatus</i>	Fringed filefish	H
<i>Monacanthus tuckeri</i>	Slender filefish	Z
<i>Mulloidichthys martinicus</i>	Yellow goatfish	I
<i>Myrichthys breviceps</i>	Sharptail eel	I
<i>Myrichthys ocellatus</i>	Gold-spotted eel	I
<i>Myripristis jacobus</i>	Blackbar soldierfish	I
<i>Neonifon marianus</i>	Longjaw squirrelfish	I
<i>Odontoscion dentex</i>	Reef croaker	Z
<i>Ophichthus ophis</i>	Spotted snake eel	P-S
<i>Opistognathus aurifrons</i>	Yellowhead jawfish	Z
<i>Opistognathus macrogathus</i>	Banded jawfish	I
<i>Opistognathus whitehursti</i>	Dusky jawfish	I
<i>Parablennius marmoratus</i> *	Seaweed blenny	Z

(continued on next page)

Table Appendix A3 (continued)

Species Name	Common Name	Trophic Guild
<i>Pempheris schomburgkii</i>	Glassy sweeper	I
<i>Pomacanthus arcuatus</i>	Gray angelfish	I
<i>Pomacanthus paru</i>	French angelfish	I
<i>Pseudupeneus maculatus</i>	Spotted goatfish	I
<i>Rypticus saponaceus</i>	Greater soapfish	
<i>Sargocentron bullisi</i>	Deepwater squirrelfish	I
<i>Sargocentron coruscum</i>	Reef squirrelfish	I
<i>Scarus iseri</i>	Striped parrotfish	H
<i>Scarus taeniopterus</i>	Princess parrotfish	H
<i>Scarus vetula</i>	Queen parrotfish	H
<i>Scomberomorus cavalla</i>	King mackerel	
<i>Scomberomorus maculatus</i>	Spanish mackerel	
<i>Scorpaena plumieri</i>	Spotted scorpionfish	I
<i>Selar crumenophthalmus</i>	Bigeye scad	P-S
<i>Serranus tabacarius</i>	Tobaccofish	P-S
<i>Sparisoma atomarium</i>	Greenblotch parrotfish	H
<i>Sparisoma chrysopterygum</i>	Redtail parrotfish	H
<i>Sparisoma rubripinne</i>	Yellowtail parrotfish	H
<i>Sparisoma viride</i>	Stoplight parrotfish	H
<i>Sphaeroides spengleri</i>	Bandtail puffer	I
<i>Sphyrna barracuda</i>	Great barracuda	P-L
<i>Sphyrna picudilla</i>	Southern sennet	P-S
<i>Stegastes partitus</i>	Bicolor damselfish	H

* Species not assigned to BCG attribute level in Florida.

Table Appendix A4

BCG Attribute IV species are intermediately tolerant taxa to fishing pressures and sediment threats in the US Caribbean and Florida.

Species Name	Common Name	Trophic Guild
<i>Abudefduf saxatilis</i>	Sergeant major	I
<i>Alphesthes afer</i>	Mutton hamlet	I
<i>Anisotremus virginicus</i>	Porkfish	I
<i>Apogon maculatus</i>	Flame fish	Z
<i>Apogon pseudomaculatus</i>	Two-spot cardinalfish	Z
<i>Apogon townsendi</i>	Belted cardinalfish	Z
<i>Archosargus rhomboidalis</i>	Sea bream	H
<i>Bothus lunatus</i>	Peacock flounder	P-S
<i>Bothus ocellatus</i>	Eyed flounder	P-S
<i>Calamus bajonado</i>	Jolthead porgy	I
<i>Calamus calamus</i>	Saucereye porgy	I
<i>Calamus nodosus</i>	Knobbed porgy	I
<i>Calamus penna</i>	Sheepshead porgy	I
<i>Calamus pennatula</i>	Pluma	I
<i>Calamus proridens</i>	Littlehead porgy	
<i>Calamus UNK</i>	Porgy	I
<i>Canthigaster rostrata</i>	Sharpnose puffer	I
<i>Carangoides bartholomaei</i>	Yellow Jack	P-L
<i>Carangoides ruber</i>	Bar jack	P-S
<i>Chloroscombrus chrysurus</i>	Atlantic bumper	Z
<i>Conger triporiceps</i>	Many tooth conger	P-S
<i>Coryphopterus glaucofraenum</i> *	Bridled goby	I
<i>Coryphopterus personatus/hyalinus</i> *	Masked/Glass goby	I
<i>Cryptotomus roseus</i>	Blue-lip parrotfish	H
<i>Ctenogobius saepepallens</i> *	Dash goby	I
<i>Diodon hystrix</i>	Porcupine fish	I
<i>Eucinostomus argenteus</i>	Spotfin mojarra/Silver mojarra	
<i>Eucinostomus jonesii</i>	Slender mojarra	I
<i>Eucinostomus melanopterus</i>	Flagfin mojarra	I
<i>Gnatholepis thompsoni</i> *	Gold-spot goby	H
<i>Gymnothorax funebris</i>	Green moray	P-S
<i>Gymnothorax moringa</i>	Spotted moray	P-S
<i>Haemulon aurolineatum</i>	Tomtate	I
<i>Haemulon plumieri</i>	White grunt	I
<i>Haemulon sciurus</i>	Blue-striped grunt	I
<i>Haliichoeres bivittatus</i>	Slippery dick	I
<i>Inermia vittata</i>	Boga	Z
<i>Lutjanus apodus</i>	Schoolmaster	P-S
<i>Lutjanus griseus</i>	Gray snapper	P-S
<i>Ocyurus chrysurus</i>	Yellowtail snapper	Z
<i>Ophioblennius macclurei</i> *	Redlip blenny	H
<i>Paradiplogrammus bairdi</i>	Lancer dragonet	I

(continued on next page)

Table Appendix A4 (continued)

Species Name	Common Name	Trophic Guild
<i>Sargocentron vexillarium</i>	Dusky squirrelfish	I
<i>Serranus baldwini</i>	Lantern bass	I
<i>Serranus flaviventris</i>	Twinspot bass	P-S
<i>Serranus tortugarum</i>	Chalk bass	Z
<i>Sparisoma aurofrenatum</i>	Redband parrotfish	H
<i>Sparisoma radians</i>	Bucktooth parrotfish	H
<i>Stegastes adustus</i>	Dusky damselfish	H
<i>Stegastes diencaeus</i>	Longfin damselfish	H
<i>Stegastes leucostictus</i>	Beaugregory	H
<i>Stegastes planifrons</i>	Threespot damselfish	I
<i>Stegastes variabilis</i>	Cocoa damselfish	H
<i>Xyrichtys splendens</i>	Green razorfish	Z

* Species not assigned to BCG attribute level in Florida.

Table Appendix A5

BCG Attribute Level V species are tolerant taxa to fishing pressures and sediment threats and Level VI species are non-native or intentionally introduced species in the US Caribbean and Florida.

BCG Attribute No.	Species Name	Common Name	Trophic Guild
V	<i>Diplodus argenteus</i>	Silver porgy	H
V	<i>Gerres cinereus</i>	Yellowfin mojarra	I
V	<i>Mugil cephalus</i>	Striped mullet	Z
V	<i>Sphoeroides testudineus</i>	Checkered puffer	I
V	<i>Synodus foetens</i>	Inshore lizardfish	P-S
VI	<i>Callogobius clitellus</i>	Saddled goby	I
VI	<i>Pterois volitans</i>	Red lionfish	P

Table Appendix A6

Fish species not assigned to an attribute as the survey methods used were insufficient to detect these species (often cryptic) in the US Caribbean and Florida.

Species Name	Common Name	Trophic Guild
<i>Ablennes hians</i>	Flat needlefish	P-S
<i>Acanthemblemaria UNK</i>	Tube Blenny	I
<i>Acanthocybium solandri</i>	Wahoo	
<i>Acanthurus UNK</i>	Surgeonfish	H
<i>Acentronura dendritica</i>	Pipehorse	I
<i>Albula vulpes</i>	Bonefish	I
<i>Alectis ciliaris</i>	African pompano	P-S
<i>Apogon UNK</i>	Cardinalfish	Z
<i>Archosargus probatocephalus</i>	Sheepshead	I
<i>Atherinomorus stipes</i>	Hardhead silverside	Z
<i>Balistes capriscus</i>	Gray triggerfish	I
<i>Bathygobius soporator</i>	Frillfin goby	I
<i>Belontiidae UNK</i>	Needlefish	P-S
<i>Bollmannia boqueronensis</i>	White-eye goby	I
<i>Bothus UNK</i>	Flounder	P-S
<i>Canthigaster jamestyleri</i>	Goldface toby	I
<i>Canthigaster UNK</i>	Puffer	I
<i>Carcharhinus leucas</i>	Bull shark	
<i>Caranx UNK</i>	Jack	P-S
<i>Centropristis striata</i>	Black sea bass	P-S
<i>Centropyge argi</i>	Cherubfish	H
<i>Chaenopsis ocellata</i>	Bluethroat pikeblenny	I
<i>Chaenopsis UNK</i>	Pike blenny	I
<i>Chaetodon sedentarius</i>	Reef butterflyfish	I
<i>Chromis enchrysur</i>	Yellowtail reeffish	I
<i>Chromis scotti</i>	Purple reeffish	Z
<i>Clupeidae UNK</i>	Herrings	Z
<i>Coryphopterus UNK</i>	Goby	I
<i>Coryphopterus punctipectophorus</i>	Spotted goby	
<i>Ctenogobius stigmaticus</i>	Marked goby	I
<i>Decapterus macarellus</i>	Mackerel scad	Z
<i>Decapterus punctatus</i>	Round scad	
<i>Decapterus UNK</i>	Scad	Z
<i>Dermatolepis inermis</i>	Marbled grouper	P-S
<i>Diplectrum bivittatum</i>	Dwarf sand perch	I

(continued on next page)

Table Appendix A6 (continued)

Species Name	Common Name	Trophic Guild
<i>Diplectrum formosum</i>	Sand perch	P-S
<i>Diplodus holbrooki</i>	Spottail pinfish	H
<i>Doratonotus megalepis</i>	Dwarf wrasse	I
<i>Echeneis naucrates</i>	Sharksucker	Z
<i>Echeneis neucratoides</i>	Whitefin sharksucker	Z
<i>Elacatinus dilepis</i>	Orangesided goby	I
<i>Elacatinus evelynae</i>	Sharknose goby	I
<i>Elacatinus horsti</i>	Yellowline goby	
<i>Elacatinus macrodon</i>	Tiger goby	
<i>Elacatinus UNK</i>	Goby	I
<i>Elacatinus xanthiprora</i>	Yellowprow goby	
<i>Elagatis bipinnulata</i>	Rainbow runner	P-S
<i>Emblemaria pandionis</i>	Sailfin blenny	Z
<i>Emblemaria sp</i>	Tube blenny	Z
<i>Emblemariopsis UNK</i>	Blenny	I
<i>Engraulidae UNK</i>	Anchovies	Z
<i>Enneanectes UNK</i>	Triplefin	H
<i>Eucinostomus gula</i>	Silver jenny	I
<i>Eucinostomus UNK</i>	Mojarra	I
<i>Euthynnus alletteratus</i>	Little tuny	P-S
<i>Gobiidae UNK</i>	Goby	I
<i>Gobiosoma grosvenori</i>	Rockcut goby	I
<i>Gymnothorax UNK</i>	Moray eel	P-S
<i>Haemulon melanurum</i>	Cottonwick	I
<i>Haemulon UNK</i>	Grunt	I
<i>Haemulon striatum</i>	Striped grunt	Z
<i>Halichoeres burekai</i>	Mardi gras wrasse	I
<i>Halichoeres caudalis</i>	Painted wrasse	I
<i>Halichoeres cyanocephalus</i>	Yellowcheek wrasse	I
<i>Halichoeres poeyi</i>	Blackear wrasse	I
<i>Halichoeres UNK</i>	Wrasse	I
<i>Harengula jaguana</i>	Scaled sardine	
<i>Hemiemblemaria simulas</i>	Wrasse blenny	
<i>Hemiramphus brasiliensis</i>	Ballyhoo	
<i>Heteroconger halis</i>	Brown garden eel	Z
<i>Heteroconger longissimus</i>	Brown garden eel	Z
<i>Hippocampus UNK</i>	Pipefish	I
<i>Holacanthus bermudensis</i>	Blue angelfish	I
<i>Holacanthus Townsendi</i>	Townsend angelfish	
<i>Holacanthus UNK</i>	Angelfish	I
<i>Hypleurochilus bermudensis</i>	Barred blenny	I
<i>Hypoplectrus UNK</i>	Hamlet	I
<i>Jenkinsia UNK</i>	Herring	Z
<i>Labrisomus filamentosus</i>	Quillfin blenny	I
<i>Lagodon rhomboides</i>	Pinfish	I
<i>Lonchopisthus micrognathus</i>	Swordtail jawfish	Z
<i>Lophogobius cyprinoides</i>	Crested goby	I
<i>Lutjanus campechanus</i>	Red snapper	P-S
<i>Lutjanus UNK</i>	Snapper	P-S
<i>Malacoctenus boehlkei</i>	Diamond blenny	I
<i>Malacoctenus gilli</i>	Dusky blenny	I
<i>Malacoctenus triangulatus</i>	Saddled blenny	I
<i>Malacoctenus UNK</i>	Scaly blenny	I
<i>Manta birostris</i>	Giant manta	Z
<i>Microgobius carri</i>	Seminole goby	Z
<i>Microgobius signatus</i>	Microgobius signatus	Z
<i>Microgobius UNK</i>	Goby UNK	H
<i>Mullidae UNK</i>	Goatfishes	I
<i>Muraenidae UNK</i>	Moray eel	P-S
<i>Mycteroperca microlepis</i>	Gag	P-S
<i>Mycteroperca phenax</i>	Scamp	P-S
<i>Mycteroperca UNK</i>	Grouper UNK	P-S
<i>Myrichthys UNK</i>	Snake eel	I
<i>Nes longus</i>	Orange-spotted goby	I
<i>Nicholsina usta</i>	Emerald parrotfish	H
<i>Ogcocephalus nasutus</i>	Shortnose batfish	I
<i>Ophichthidae UNK</i>	Snake eel UNK	P-S
<i>Opistognathus UNK</i>	Jawfish	Z
<i>Oxyurichthys stigmaphius</i>	Spotfin goby	I
<i>Pareques umbrosus</i>	Cubby	I
<i>Platybelone argalus</i>	Keeltail needlefish	P-S
<i>Pomacanthus UNK</i>	Angelfish	I
<i>Ptereleotris calliura</i>	Blue dartfish	
<i>Ptereleotris helenae</i>	Hovering dartfish	Z

(continued on next page)

Table Appendix A6 (continued)

Species Name	Common Name	Trophic Guild
<i>Remora remora</i>	Common remora	Z
<i>Rypticus bistrispinus</i>	Freckled soapfish	P-S
<i>Rypticus maculatus</i>	White-spotted soapfish	P-S
<i>Scartella cristata</i>	Molly miller	H
<i>Scarus UNK</i>	Parrotfish	H
<i>Scorpaena UNK</i>	Scorpionfish UNK	I
<i>Scorpaenodes caribbaeus</i>	Reef scorpionfish	
<i>Serraniculus pumilio</i>	Pygmy sea bass	I
<i>Serranus subligarius</i>	Belted sandfish	I
<i>Serranus UNK</i>	Seabass UNK	P-S
<i>Sparisoma UNK</i>	Parrotfish	H
<i>Sphyraena borealis</i>	Northern sennet	P-S
<i>Stephanolepis hispidus</i>	Planehead filefish	H
<i>Stephanolepis setifer</i>	Pygmy filefish	H
<i>Stromateidae UNK</i>	Butterfish	P-S
<i>Syacium UNK</i>	Sand flounder	I
<i>Sygnathus dawsoni</i>	Pipefish	I
<i>Synodus intermedius</i>	Sand diver	P-S
<i>Synodus saurus</i>	Blue-striped lizardfish	P-S
<i>Tigriogobius dilepis</i>	Orange-sided goby	I
<i>Trachinocephalus myops</i>	Snakefish	Z
<i>Triglidae UNK</i>	Searobin Family	I
<i>Tylosurus crocodilus</i>	Houndfish	P-S
<i>Urobatis jamaicensis</i>	Yellow stingray	
<i>Xyrichtys martinicensis</i>	Rosy razorfish	I
<i>Xyrichtys novacula</i>	Pearly razorfish	I
<i>Xyrichtys UNK</i>	Razorfish	I

Table Appendix A7

Large-bodied species for reef fish using 90 cm length threshold to distinguish between large and small body sizes.

Fish Species	Common Name	Piscivore
<i>Aetobatus narinari</i>	Spotted eagle ray	
<i>Carangoides bartholomaei</i>	Yellow jack	X
<i>Caranx hippos</i>	Crevalle jack	X
<i>Caranx lugubris</i>	Black jack	X
<i>Carcharhinus limbatus</i>	Blacktip shark	X
<i>Carcharhinus perezii</i>	Caribbean Reef shark	X
<i>Dasyatis americana</i>	Southern stingray	
<i>Epinephelus striatus</i>	Nassau grouper	X
<i>Epinephelus itajara</i>	Atlantic goliath grouper	X
<i>Epinephelus morio</i>	Red grouper	X
<i>Galeocerdo cuvier</i>	Tiger shark	X
<i>Ginglymostoma cirratum</i>	Nurse shark	X
<i>Lachnolaimus maximus</i>	Hogfish	
<i>Lutjanus analis</i>	Mutton snapper	
<i>Lutjanus cyanopterus</i>	Cubera snapper	X
<i>Lutjanus jocu</i>	Dog snapper	X
<i>Megalops atlanticus</i>	Tarpon	X
<i>Mycteroperca bonaci</i>	Black grouper	X
<i>Mycteroperca tigris</i>	Tiger grouper	X
<i>Mycteroperca venosa</i>	Yellowfin grouper	X
<i>Negaprion brevirostris</i>	Lemon shark	X
<i>Scarus coelestinus</i>	Midnight parrotfish	
<i>Scarus coeruleus</i>	Blue parrotfish	
<i>Scarus guacamaia</i>	Rainbow parrotfish	
<i>Seriola dumerili</i>	Greater amberjack	X
<i>Seriola rivoliana</i>	Almaco jack	X
<i>Sphyraena barracuda</i>	Great barracuda	X
<i>Sphyrna mokarran</i>	Great hammerhead shark	X

Appendix B. Large-bodied reef fish species. Note: 90 cm maximum size cut-off used as threshold between large and small; small predators are still an indicator of good condition and are reflected as a rule in BCG Level 3

Fish species	Common name	Piscivore
<i>Aetobatus narinari</i>	Spotted eagle ray	
<i>Carangoides bartholomaei</i>	Yellow jack	X
<i>Caranx hippos</i>	Crevalle jack	X
<i>Caranx lugubris</i>	Black jack	X

<i>Carcharhinus limbatus</i>	Blacktip shark	X
<i>Carcharhinus perezii</i>	Caribbean reef shark	X
<i>Dasyatis americana</i>	Southern stingray	
<i>Epinephelus striatus</i>	Nassau grouper	X
<i>Epinephelus itajara</i>	Atlantic goliath grouper	X
<i>Epinephelus morio</i>	Red grouper	X
<i>Galeocerdo cuvier</i>	Tiger shark	X
<i>Ginglymostoma cirratum</i>	Nurse shark	X
<i>Lachnolaimus maximus</i>	Hogfish	
<i>Lutjanus analis</i>	Mutton snapper	
<i>Lutjanus cyanopterus</i>	Cubera snapper	X
<i>Lutjanus jocu</i>	Dog snapper	X
<i>Megalops atlanticus</i>	Tarpon	X
<i>Mycteroperca bonaci</i>	Black grouper	X
<i>Mycteroperca tigris</i>	Tiger grouper	X
<i>Mycteroperca venenosa</i>	Yellowfin grouper	X
<i>Negaprion brevirostris</i>	Lemon shark	X
<i>Scarus coelestinus</i>	Midnight parrotfish	
<i>Scarus coeruleus</i>	Blue parrotfish	
<i>Scarus guacamaia</i>	Rainbow parrotfish	
<i>Seriola dumerili</i>	Greater amberjack	X
<i>Seriola rivoliana</i>	Almaco jack	X
<i>Sphyrna barracuda</i>	Great barracuda	X
<i>Sphyrna mokarran</i>	Great hammerhead shark	X

Appendix C. Supplementary data

Supplementary data to this article can be found online at <https://doi.org/10.1016/j.marpolbul.2020.111387>.

References

- 33 USC §1251 et seq, 1972. Federal Water Pollution Control Amendments of 1972 (aka The Clean Water Act).
- Aguilar-Perera, J.A., 2004. Coastal Habitat Connectivity of Reef Fishes From Southwestern Puerto Rico (PhD Thesis). University of Puerto Rico, Mayagüez, Puerto Rico.
- Aguilar-Perera, A., Appeldoorn, R.S., 2007. Variation in juvenile fish density along the mangrove-seagrass-coral reef continuum in SW Puerto Rico. *Mar. Ecol. Prog. Ser.* 348, 139–148.
- Aguilar-Perera, A., Appeldoorn, R.S., 2008. Spatial distribution of marine fishes along a cross-shelf gradient containing a continuum of mangrove-seagrass-coral reefs off southwestern Puerto Rico. *Estuar. Coast. Shelf Sci.* 76, 378–394.
- Ames, C., 2017. Reef Fish Assemblage Biogeography Along the Florida Reef Tract (Master's thesis). Nova Southeastern University Retrieved from NSUWorks. https://nsuworks.nova.edu/cgi/viewcontent.cgi?article=1459&context=occ_stuetd.
- Appeldoorn, R.S., 2011. Can we stop the madness? Managing for resilience in coral reef fisheries. *Proc. Gulf Caribb. Fish. Inst.* 63, 6–9.
- Appeldoorn, R.S., Recksiek, C.W., Hill, R.L., Pagan, F.E., Dennis, G.D., 1997. Marine protected areas and reef fish movements: the role of habitat in controlling ontogenetic migration. In: *Proceedings of the Eighth International Coral Reef Symposium*. 2. pp. 1917–1922.
- Appeldoorn, R.S., Friedlander, A., Sladek Nowlis, J., Ussegilo, P., Mitchell-Chui, A., 2003. Habitat connectivity on the insular platform of Old Providence-Santa Catalina, Colombia: mechanisms, limits and ecological consequences relevant to marine reserve design. *Gulf Caribb. Res.* 14, 61–77.
- Ault, J.S., Bohnsack, J.A., Meester, G.A., 1998. A retrospective (1979–1996) multispecies assessment of coral reef fish stocks in the Florida Keys. *Fish. Bull.* 96 (3), 395–414.
- Ault, J.S., Luo, J., Smith, S.G., Serafy, J.E., Wang, J.D., Diaz, G.A., Humston, R., 1999. A spatial dynamic multistock production model. *Can. J. Fish. Aquat. Sci.* 56 (S1), 4–25.
- Ault, J.S., Smith, S.G., Bohnsack, J.A., 2005. Evaluation of average length as an estimator of exploitation status for the Florida coral reef fish community. *J. Mar. Sci.* 62, 417–423.
- Ault, J.S., Smith, S.G., Luo, J., Monaco, M.E., Appeldoorn, R.S., 2008. Length-based assessment of sustainability benchmarks for coral reef fishes in Puerto Rico. *Environ. Conserv.* 35, 221–231.
- Ault, J.S., Smith, S.G., Bohnsack, J.A., Luo, J., Zurcher, N., McClellan, D.B., Zeigler, T.A., Hallac, D.E., Patterson, M., Feeley, M.W., Ruttenberg, B.I., Hunt, J., Kimball, D., Causey, B., 2013. Assessing coral reef fish population and community changes in response to marine reserves in the Dry Tortugas, Florida, USA. *Fish. Res.* 144, 28–37.
- Ault, J.S., Smith, S.G., Browder, J.A., Nuttle, W., Franklin, E.C., Luo, J., DiNardo, G.T., Bohnsack, J.A., 2014. Indicators for assessing the ecological and sustainability dynamics of southern Florida's coral reef and coastal fisheries. *Ecol. Indic.* 44, 164–172.
- Bejarno, I., Appeldoorn, R.S., 2013. Seawater turbidity and fish communities on coral reefs of Puerto Rico. *Mar. Ecol. Prog. Ser.* 474, 217–226.
- Beverton, R.J.H., Holt, S.J., 1957. On the Dynamics of Exploited Fish Populations. *Fishery Investigations Series II XIX Ministry of Agriculture, Fisheries and Food, Lowestoft* (533 p).
- Bohnsack, J.A., Bannerot, S.P., 1986. A stationary visual census technique for quantitatively assessing community structure of coral reef fishes. In: *NOAA Technical Report NMFS*. 41.
- Bradley, P., Davis, W., Fisher, W., Bell, H., Chan, V., LoBue, C., Wiltse, W., 2008. Biological criteria for protection of U.S. coral reefs. In: *Proceedings of the 11th International Coral Reef Symposium*, Ft. Lauderdale, Florida, 7–11 July 2008.
- Bradley, P., Fisher, W.S., Bell, H., Davis, W.S., Chan, V., LoBue, C., Wiltse, W., 2009. Development and implementation of coral reef biocriteria in U. S. jurisdictions. *Environ. Monit. Assess.* 150, 43–51.
- Bradley, P., Fore, L., Fisher, W., Davis, W., 2010. Coral Reef Biological Criteria: Using the Clean Water Act to Protect a National Treasure. U.S. EPA, ORD, AED, Narragansett, RI (EPA/600/R-10/054).
- Bradley, P., Santavy, D.L., Gerritsen, J., 2014. Workshop on Biological Integrity of Coral Reefs, August 21–22, 2012, Caribbean Coral Reef Institute, Isla Magueyes, La Parguera, Puerto Rico. U.S. EPA, ORD, AED, Narragansett, RI (EPA/600/R-13/350).
- Bradley, P., Jessup, B., Gerritsen, J., Gallindo, A., 2016. The Coral Reef Biological Condition Gradient: Moving Toward Quantitative Criteria From Species-Based Attributes. TetraTech, Inc (Prepared for the U.S. Environmental Protection Agency, Office of Water, Office of Science and Technology, Washington, D.C).
- Brander, L., van Beukering, P., 2013. The Total Economic Value of U.S. Coral Reefs: A Review of the Literature. National Oceanic and Atmospheric Administration's Coral Reef Conservation Program. NOAA, Silver Spring, Maryland.
- Brandt, M.E., Zurcher, N., Acosta, A., Ault, J.S., Bohnsack, J.A., Feeley, M.W., Harper, D.E., Hunt, J.H., Kellison, G.T., McClellan, D.B., Patterson, M.E., Smith, S.G., 2009. A cooperative multi-agency reef fish monitoring protocol for the Florida Keys coral reef ecosystem. In: *Natural Resource Report NPS/SFCN/NRR 2009/150*. National Park Service, Fort Collins, Colorado.
- Brown, C.J., Jupiter, S.D., Lin, H.Y., Albert, S., Klein, C., Maina, J.M., Tulloch, V.J., Wenger, A.S., Mumby, P.J., 2017. Habitat change mediates the response of coral reef fish populations to terrestrial run-off. *Mar. Ecol. Prog. Ser.* 576, 55–68.
- Bryan, D.R., Smith, S.G., Ault, J.S., Feeley, M.W., Menza, C.W., 2016. Feasibility of a regionwide probability survey for coral reef fish in Puerto Rico and the U.S. Virgin Islands. *Mar. Coast. Fish.* 8 (1), 135–146.
- Burkepile, D.E., Hay, M.E., 2008. Herbivore species richness and feeding complementarity affect community structure and function on a coral reef. *Proc. Natl. Acad. Sci.* 105, 16201–16206.
- Cairns, J., 1977. Quantification of Biological Integrity. In: Ballentine, R.K., Guarrain, L.J. (Eds.), *The Integrity of Water*. U.S. Environmental Protection Agency, Office of Water, Washington, DC, USA, pp. 171–187.
- Caldow, C., Clark, R., Edwards, K., Hile, S., Menza, C., Hickerson, E., Schmahl, G.P., 2009. Biogeographic Characterization of Fish Communities and Associated Benthic Habitats Within the Flower Garden Banks National Marine Sanctuary: Sampling Design and Implementation of Scuba Surveys on the Coral Caps. NOAA/National Centers for Coastal Ocean Science, Silver Spring, MD (NOAA Technical Memorandum NOS NCCOS 81).
- Carpenter, K.E., Miclat, R.I., Albaladejo, V.D., Corpuz, V.T., 1981. The influence of substrate structure on the abundance and diversity of Philippine reef fishes. In: *Proceedings of the Fourth International Coral Reef Symposium*. 2. pp. 497–502.
- Carpenter, K.E., Abrar, M., Aeby, G., Aronson, R.B., Banks, S., Bruckner, A., Chiriboga, A., Cortes, J., Delbeek, J.C., DeVantier, L., Edgar, G.J., Edwards, A.J., Fenner, D., Guzman, H.M., Hoeksema, B.W., Hodgson, G., Johan, O., Licuanan, W.Y., Livingstone, S.R., Lovell, E.R., Moore, J.A., Obura, D.O., Ochavillo, D., Polidoro, B.A., Precht, W.F., Quibilan, M.C., Reboton, C., Richards, Z.T., Rogers, A.D., Sanciangco,

- J., Sheppard, A., Sheppard, C., Smith, J., Stuart, S., Turak, E., Veron, J.E.N., Wallace, C., Weil, E., Wood, E., 2008. One-third of reef-building corals face elevated extinction risk from climate change and local impacts. *Science* 321, 560–563.
- Castella, E., Speight, M.C.D., 1996. Knowledge representation using fuzzy coded variables: an example based on the use of Syrphidae (Insecta, Diptera) in the assessment of riverine wetlands. *Ecol. Model.* 85, 13–25.
- Cervený, K., 2006. Distribution Patterns of Reef Fishes in Southwest Puerto Rico, Relative to Structural Habitat, Cross—Shelf Location, and Ontogenetic Stage (Master's Thesis). University of Puerto Rico—Mayagüez.
- Chapman, C.J., Johnstone, A.D.F., Dunn, J.R., Creasey, D.J., 1974. Reactions of fish to sound generated by divers' open-circuit underwater breathing apparatus. *Mar. Biol.* 27, 357–366.
- Christensen, J.D., Jeffrey, C.F.G., Caldow, C., Monaco, M.E., Kendall, M.S., Appeldoorn, R.S., 2003. Cross-shelf habitat utilization patterns of reef fishes in southwestern Puerto Rico. *Gulf Caribb. Res.* 14, 9–27.
- Clarke, K.R., Gorley, R.N., 2015. PRIMER Version 7: User Manual/Tutorial. PRIMER-E, Plymouth, UK.
- Clarke, K.R., Gorley, R.N., Somerfield, P.J., Warwick, R.M., 2014. Change in Marine Communities: An Approach to Statistical Analysis and Interpretation, 3rd edition. PRIMER-E, Plymouth, UK.
- Claro, R., 1991. Changes in fish assemblage structure by the effect of intense fisheries activity. *Trop. Ecol.* 32, 36–46.
- Claro, R., Lindeman, K.C., 2003. Spawning aggregation sites of snapper and grouper species (Lutjanidae and Serranidae) on the insular shelf of Cuba. *Gulf Caribb. Res.* 14, 91–106.
- Cleary, D.F.R., Suharsono, Hoeksema, B.W., 2006. Coral diversity across a disturbance gradient in the Pulau Seribu reef complex off Jakarta, Indonesia. *Biodivers. Conserv.* 15, 3653–3674.
- Cocheret de la Morinière, E., Polluz, B.J.A., Nagelkerken, I., van der Velde, G., 2002. Post-settlement life cycle migration patterns and habitat preference of coral reef fish that use seagrass and mangrove habitats as nurseries. *Estuar. Coast. Shelf Sci.* 55, 309–321.
- Coleman, F.C., Koenig, C.C., Huntsman, G.R., Musick, J.A., Eklund, A.M., McGovern, J.C., Chapman, R.W., Sedberry, G.R., Grimes, C.B., 2000. Long-lived reef fishes: the grouper-snapper complex. *Fisheries* 25, 14–21.
- Colin, P.L., Shapiro, D.Y., Weiler, D., 1987. Preliminary investigations of reproduction of two species of groupers. *Epinephelus guttatus* and *E. striatus* in the West Indies. *Bull. Mar. Sci.* 40, 220–230.
- Courtney, L.A., Fisher, W.S., Raimondo, S., Oliver, L.M., Davis, W.S., 2007. Estimating three-dimensional colony surface area of field corals. *J. Exp. Mar. Biol. Ecol.* 351, 234–242.
- Dahlgren, C.P., Eggleston, D.B., 2000. Ecological processes underlying ontogenetic habitat shifts in a coral reef fish. *Ecology* 81 (8), 2227–2240 2000.
- Davies, S.P., Jackson, S.K., 2006. The biological condition gradient: a descriptive model for interpreting change in aquatic ecosystems. *Ecol. Appl.* 16, 1251–1266.
- Davis, W.S., Simon, T. (Eds.), 2004. Biological Assessment and Criteria: Tools for Water Resource Planning and Decision Making. CRC Press LLC, Boca Raton, Florida, USA (432 pp).
- Dustan, P., Doherty, O., Pardede, S., 2013. Digital reef rugosity estimates coral reef habitat complexity. *PLoS One* 8 (2), e57386.
- Edinger, E.N., Jompa, J., Limmon, G.V., Widjajmoko, W., Risk, M.J., 1998. Reef degradation and coral biodiversity in Indonesia: effects of land-based pollution, destructive fishing practices and changes over time. *Mar. Pollut. Bull.* 36, 617–630.
- Emslie, M.J., Cheal, A.J., MacNeil, M.A., Miller, I.R., Sweatman, H.P.A., 2018. Reef fish communities are spooked by scuba surveys and may take hours to recover. *PeerJ* 6, e4886. <https://doi.org/10.7717/peerj.4886>.
- Ennis, R.S., Brandt, M.E., Wilson, K.R., Smith, T.B., 2016. Coral reef health response to chronic and acute changes in water quality in St. Thomas, United States Virgin Islands. *Mar. Pollut. Bull.* 111, 418–427.
- Fabricius, K., De'ath, G., 2001. Environmental factors associated with the spatial distribution of crustose coralline algae in the Great Barrier Reef. *Coral Reefs* 19, 303–309.
- Fabricius, K., De'ath, G., McCook, L., Turak, E., Williams, D.M., 2005. Changes in algal, coral and fish assemblages along water quality gradients on the inshore Great Barrier Reef. *Mar. Pollut. Bull.* 51, 384–398.
- Fisco, D., 2016. Reef Fish Spatial Distribution and Benthic Habitat Associations on the Southeast Florida Reef Tract (Master's thesis). Nova Southeastern University.
- Fisher, W.S., 2007. Stony Coral Rapid Bioassessment Protocol. US Environmental Protection Agency. EPA/600/R-06/167, July 2007. Office of Research and Development, Washington, DC.
- Fisher, W.S., Davis, W.P., Quarles, R.L., Patrick, J., Campbell, J.G., Harris, P.S., Hemmer, B.L., Parsons, M., 2007. Characterizing coral condition using estimates of three-dimensional colony surface area. *Environ. Monit. Assess.* 125, 347–360.
- Fisher, W.S., Fore, L.S., Hutchins, A., Quarles, R.L., Campbell, J.G., LoBue, C., Davis, W., 2008. Evaluation of stony coral indicators for coral reef management. *Mar. Pollut. Bull.* 56, 1737–1745.
- Fisher, W.S., Fore, L.S., Oliver, L.M., LoBue, C., Quarles, R.L., Campbell, J.G., Harris, P.S., Hemmer, B.L., Vickery, S., Parsons, M., Hutchins, A., Bernier, K., Rodriguez, D., Bradley, P., 2014. Regional status assessment of stony corals in the U.S. Virgin Islands. *Environ. Monit. Assess.* 186, 7165–7181.
- Fisher, W.S., Vivian, D.N., Campbell, J., LoBue, C., Hemmer, R.L., Wilkinson, S., Harris, P., Santavy, D.L., Parsons, M., Bradley, P., Humphrey, A., Oliver, L.M., Harwell, L., 2019. Biological assessment of coral reefs in southern Puerto Rico: supporting coral reef protection under the U.S. Clean Water Act. *Coast. Manag.* 47, 429–452.
- Frey, D.G., 1977. Biological integrity of water: an historical approach. In: *The Integrity of Water*. Proceedings of a Symposium. US Environmental Protection Agency, Washington, DC, pp. 127–140.
- Friedlander, A.M., Parrish, J.D., 1998. Habitat characteristics affecting fish assemblages on a Hawaiian coral reef. *J. Exp. Mar. Biol. Ecol.* 224, 1–30.
- Friedlander, A.M., Brown, E.K., Jokiel, P.L., Smith, W.R., Rodgers, K.S., 2003. Effects of habitat, wave exposure, and marine protected area status on coral reef fish assemblages in the Hawaiian archipelago. *Coral Reefs* 22, 291–305.
- Gerritsen, J., Bouchard Jr., R.W., Zheng, L., Leppo, E.W., Yoder, C.O., 2017. Calibration of the Biological Condition Gradient in Minnesota streams: a quantitative expert-based decision system. *Freshw. Sci.* 36, 427–451.
- Goodwin, N.B., Grant, A., Perry, A.L., Dulvy, N.K., Reynolds, J.D., 2006. Life history correlates of density-dependent recruitment in marine fishes. *Can. J. Fish. Aquat. Sci.* 63, 494–509.
- Goreau, T.F., 1959. The ecology of Jamaican coral reefs. I. Species composition and zonation. *Ecology* 40, 67–90.
- Graham, N.A.J., McClanahan, T.R., MacNeil, M.A., Wilson, S.K., Cinner, J.E., Huchery, C., Holmes, T.H., 2017. Human disruption of coral reef trophic structure. *Curr. Biol.* 27, 231–236.
- Gratwicke, B., Speight, M.R., 2005a. Effects of habitat complexity on Caribbean marine fish assemblages. *Mar. Ecol. Prog. Ser.* 292, 301–310.
- Gratwicke, B., Speight, M.R., 2005b. The relationship between fish species richness, abundance and habitat complexity in a range of shallow tropical marine habitats. *J. Fish Biol.* 66, 650–667.
- Green, A.L., 1996. Spatial, temporal and ontogenetic patterns of habitat use by coral reef fishes (family Labridae). *Mar. Ecol. Prog. Ser.* 133, 1–11.
- Greenstein, B.J., Curran, A., Pandolfi, J.M., 1998. Shifting ecological baselines and the demise of *Acropora cervicornis* in the western North Atlantic and Caribbean province: a Pleistocene perspective. *Coral Reefs* 17, 249–261.
- Grober-Dunsmore, R., Frazer, T., Lindberg, W., Beets, J., 2007. Reef fish and habitat relationships in a Caribbean seascape: the importance of reef context. *Coral Reefs* 26, 201–216.
- Hausmann, S., Charles, D.F., Gerritsen, J., Belton, T.J., 2016. A diatom-based Biological Condition Gradient (BCG) approach for assessing impairment and developing nutrient criteria for streams. *Sci. Total Environ.* 562, 914–927.
- Hawkins, J.P., Roberts, C.M., 2004. Effects of artisanal fishing on Caribbean coral reefs. *Conserv. Biol.* 18, 215–226.
- Healthy Reefs for Healthy People Initiative (HRI), 2012. Report Card for the Mesoamerican Reef 2012. 25 p. URL. <http://www.healthyreefs.org/cms/report-cards>.**
- Heyman, W.D., Kjerfve, B., 2008. Characterization of transient multi-species reef fish spawning aggregations at Gladden Spit, Belize. *Bull. Mar. Sci.* 83, 531–551.
- Hill, J., Wilkinson, C., 2004. Methods for Ecological Monitoring of Coral Reefs. Australian Institute of Marine Science, Townsville.
- Hobson, R.D., 1972. Surface roughness in topography: quantitative approach. In: Chorley, R.J. (Ed.), *Spatial Analysis in Geomorphology*. Methuen and Co., Ltd, London, pp. 221–245.
- Hoegh-Guldberg, O., Mumby, P.J., Hooten, A.J., Steneck, R.S., Greenfield, P., Gomez, E., Harvell, C.D., Sale, P.F., Edwards, A.J., Caldeira, K., Knowlton, N., Eakin, C.M., Iglesias-Prieto, R., Muthiga, N., Bradbury, R.H., Dubi, A., Hatzioiols ME, M.E., 2007. Coral reefs under rapid climate change and ocean acidification. *Science* 318, 1737–1742.
- Hoegh-Guldberg, O., Ortiz, J.C., Dove, S., 2011. The future of coral reefs. *Science* 334, 1494–1495.
- Hoegh-Guldberg, O., Poloczanska, E.S., Skirving, W., Dove, S., 2017. Coral reef ecosystems under climate change and ocean acidification. *Front. Mar. Sci.* 4. <https://doi.org/10.3389/fmars.2017.00158>.
- Holling, C.S., 1973. Resilience and stability in ecological systems. *Annu. Rev. Ecol. Syst.* 4, 1–23.
- Hughes, T.P., 1994. Catastrophes, phase shifts, and large-scale degradation of a Caribbean coral reef. *Science* 265, 1547–1551.
- Hughes, T.P., Baird, A.H., Bellwood, D.R., Card, M., Connolly, S.R., Folke, C., Grosberg, R., Hoegh-Guldberg, O., Jackson, J.B.C., Kleypas, J., Lough, J.M., Marshall, P., Nystrom, M., Palumbi, S.R., Pandolfi, J.M., Rosen, B., Roughgraden, J., 2003. Climate change, human impacts, and the resilience of coral reefs. *Science* 301 (5635), 929–933.
- Ibelings, B.W., Vonk, M., Los, H.F.J., Van Der Molen, D.T., Mooij, W.M., 2003. Fuzzy modeling of cyanobacterial surface waterblooms: validation with NOAA-AVHRR satellite images. *Ecol. Appl.* 13, 1456–1472.
- Ionnidou, I.A., Paraskevopoulos, S., Tzionas, P., 2003. An interactive computer graphics interface for the introduction of fuzzy inference in environmental education. *Interact. Comput.* 18 (4), 683–708.
- Jackson, J.B.C., 1997. Reefs since Columbus. *Coral Reefs* 16, S23–S32.
- Jackson, J.B.C., Sala, E., 2001. Unnatural oceans. *Sci. Mar.* 65, 273–281.
- Jackson, J.B.C., Kirby, M.X., Berger, W.H., Bjorndal, K.A., Botsford, L.W., Bourque, B.J., Bradbury, R.H., Cooke, R., Erlandson, J., Estes, J.A., Hughes, T.P., Kidwell, S., Lange, C.B., Lenihan, H.S., Pandolfi, J.M., Peterson, C.H., Steneck, R.S., Tegner, M.J., Warner, R.R., 2001. Historical overfishing and the recent collapse of coastal ecosystems. *Science* 293, 629–637.
- Jackson, J.B.C., Donovan, M.K., Cramer, K.L., Lam, V.V. (Eds.), 2014. Status and Trends of Caribbean Coral Reefs: 1970–2012. Global Coral Reef Monitoring Network, IUCN, Gland, Switzerland.
- Jameson, S.C., Erdmann, M.V., Karr, J.R., Gibson Jr., G.R., Potts, K.W., 2001. Charting a course toward diagnostic monitoring: a continuing review of coral reef attributes and a research strategy for creating coral reef indexes of biotic integrity. *Bull. Mar. Sci.* 69, 701–744.
- Jameson, S.C., Karr, J.R., Potts, K.W., 2003. Establishing Reference Conditions for the Diagnostic Monitoring and Assessment of Coral Reefs. U.S.EPA, Office of Water,

- Washington, DC (44 p).
- Januchowski-Hartley, F.A., Graham, N.A.J., Feary, D.A., Morove, T., Cinner, J.E., 2011. Fear of fishers: human predation explains behavioural changes in coral reef fishes. *PLoS One* 6 (8), e22761.
- Jennings, S., Reynolds, J.D., Mills, S.C., 1998. Life history correlates of responses to fisheries exploitation. *Proc. R. Soc. Biol. Sci.* 265, 333–339.
- Johannes, R.E., 1978. Reproductive strategies of coastal marine fishes in the tropics. *Environ. Biol. Fish.* 3, 65–84.
- Jokiel, P.L., Brown, E.K., Friedlander, A., Rodgers, S.K., Smith, W.R., 2004. Hawai'i coral reef assessment and monitoring program: spatial patterns and temporal dynamics in reef coral communities. *Pac. Sci.* 58, 159–174.
- Kadison, E., Brandt, M., Nemeth, R., Martens, J., Blondeau, J., Smith, T., 2017. Abundance of commercially important reef fish indicates different levels of over-exploitation across shelves of the U.S. Virgin Islands. *PLoS One* 12, e0180063.
- Karr, J.R., Chu, E.W., 2000. Sustaining living rivers. *Hydrobiologia* 422, 1–14.
- Kendall, M.S., Krueger, C.R., Buja, K.R., Christensen, J.D., Finkbeiner, M., Warner, R.A., Monaco, M.E., 2001. Methods Used to Map the Benthic Habitats of Puerto Rico and the U.S. Virgin Islands. NOAA Technical Memorandum 152. NOS NCCOS CCMA Silver Spring, MD.
- Kimmel, J.J., 1985. A Characterization of Puerto Rican Fish Assemblages (Doctoral Dissertation). University of Puerto Rico—Mayagüez.
- Klir, G.J., 2004. Fuzzy Logic: A Specialized Tutorial. In: Demicco, Robert V., Klir, George J. (Eds.), *Fuzzy Logic in Geology*. Elsevier Academic Press, pp. 352.
- Knowlton, N., Jackson, J.B., 2008. Shifting baselines, local impacts, and global change on coral reefs. *PLoS Biol.* 6, e54. <https://doi.org/10.1371/journal.pbio.0060054>.
- Kuffner, I.B., Brock, J.C., Grober-Dunsmore, R., Bonito, V.E., Hickey, T.D., Wright, C.W., 2007. Relationships between reef fish communities and remotely sensed rugosity measurements in Biscayne National Park, Florida, USA. *Environ. Biol. Fish.* 78, 71–82.
- Kulbicki, M., 1998. How the acquired behaviour of commercial reef fishes may influence the results obtained from visual censuses. *J. Exp. Mar. Biol. Ecol.* 222, 11–30.
- Kulbicki, M., Sarraména, S., 1999. Comparison of density estimates derived from strip transect and distance sampling for underwater visual censuses: a case study of Chaetodontidae and Pomacanthidae. *Aquat. Living Resour.* 12, 315–325.
- Lang, J.C., 2003. Status of coral reefs in the western Atlantic: results of initial surveys, Atlantic and Gulf Rapid Reef Assessment (AGRRA) Program. *Atoll Res. Bull.* 496, 1–630.
- Lefcheck, J.S., Innes-Gold, A.A., Brandl, S.J., Steneck, R.S., Torres, R.E., Rasher, D.B., 2019. Tropical fish diversity enhances coral reef functioning across multiple scales. *Sci. Adv.* 5 (3), 6420.
- Lindeman, K.A., Pugliese, R., Waugh, G.T., Ault, J.S., 2000. Developmental patterns within a multispecies reef fishery: management applications for essential fish habitats and protected areas. *Bull. Mar. Sci.* 66 (3), 929–956.
- Lindfield, S.J., Harvey, E.S., McIlwain, J.L., Halford, A.R., 2014. Silent fish surveys: bubble-free diving highlights inaccuracies associated with SCUBA-based surveys in heavily fished areas. *Methods Ecol. Evol.* 5, 1061–1069.
- Luckhurst, B.E., Luckhurst, K., 1978. Analysis of substrate variables on coral reef fish communities. *Mar. Biol.* 49, 317–323.
- Man, A., Law, R., Polunin, N.V.C., 1995. Role of marine reserves in recruitment to reef fisheries: a metapopulation model. *Biol. Conserv.* 71, 197–204.
- Maynard, J.A., McKagan, S., Raymundo, L., Johnson, S., Ahmadi, G.N., Johnston, L., Houk, P., Williams, G.J., Kendall, M., Heron, S.F., van Hooidek, R., McLeod, E., Tracey, D., Planes, S., 2015. Assessing relative resilience potential of coral reefs to inform management. *Biol. Conserv.* 192, 109–119.
- McClanahan, T.R., 1994. Kenyan coral reef lagoon fish: effects of fishing, substrate complexity and sea urchins. *Coral Reefs* 13, 231–241.
- McClanahan, T.R., Donner, S.D., Maynard, J.A., MacNeil, M.A., Graham, N.A.J., Maina, J., Baker, A.C., Alemu, J.B., Beger, M., Campbell, S.J., Darling, E.S., Eakin, C.M., Heron, S.F., Jupier, S.D., Lundquist, C.J., McLeod, E., Mumby, P.J., Paddock, M.J., Selig, E.R., van Woest, R., 2012. Prioritizing key resilience indicators to support coral reef management in a changing climate. *PLoS One* 7 (8), e42884. <https://doi.org/10.1371/journal.pone.0042884>.
- McCormick, M., 1994. Comparison of field methods for measuring surface topography and their associations with a tropical reef fish assemblage. *Mar. Ecol. Prog. Ser.* 112, 87–96.
- McField, M., Kramer, P.R., 2007. Healthy Reefs for Healthy People: A Guide to Indicators of Reef Health and Social Well-being in the Mesoamerican Reef Region. With Contributions by M Gorrez and M McPherson. (208 pp).
- Meynecke, J.O., Lee, S.Y., Duke, N.C., 2008. Linking spatial metrics and fish catch reveals the importance of coastal wetland connectivity to inshore fisheries in Queensland, Australia. *Biol. Conserv.* 141, 981–996.
- Mora, C., 2008. A clear human footprint in the coral reefs of the Caribbean. *Proc. R. Soc. Lond. B* 275, 767–773.
- Moustaka, M., Langlois, T.J., McLean, D., Bond, T., Fisher, R., Fearn, P., Dorji, P., Evan, R.D., 2018. The effects of suspended sediment on coral reef fish assemblages and feeding guilds of north-west Australia. *Coral Reefs* 37 (3), 659–673.
- Mumby, P.J., Edwards, A.J., Arias-Gonzalez, J.E., Lindeman, K.C., Blackwell, P.G., Gall, A., Gorchynska, M.I., Harborne, A.R., Pescod, C.L., Renken, H., Wabnitz, C.C., Llewellyn, G., 2004. Mangroves enhance the biomass of coral reef fish communities in the Caribbean. *Nature* 427, 533–536.
- Mumby, P.J., Dahlgren, C.P., Harborne, A.R., Kappel, C.V., Micheli, F., Brumbaugh, D.R., Holmes, K.E., Mendes, J.M., Broad, K., Sanchirico, J.N., Buch, K., Box, S., Stoffle, R.W., Gil, A.B., 2006. Fishing, trophic cascades, and the process of grazing on coral reefs. *Science* 311, 98–101.
- Mumby, P.J., Broad, K., Brumbaugh, D.R., Dahlgren, C.P., Harborne, A.R., Hastings, A., Holmes, K.E., Kappel, C.V., Micheli, F., Sanchirico, J.N., 2008. Coral reef habitats as surrogates of species, ecological functions, and ecosystem services. *Conserv. Biol.* 22, 941–951.
- Munro, J.L., 1983. Caribbean Coral Reef Fishery Resources. *WorldFish* (286 pp).
- Musick, J.A., Harbin, M.M., Berkeley, S.A., Burgess, G.H., Eklund, A.M., Findley, L., Gilmore, R.G., Golden, J.T., Ha, D.S., Huntsman, G.R., McGovern, J.C., Parker, S.J., Poss, S.G., Sala, E., Schmidt, T.W., Sedberry, G.R., Weeks, H., Wright, S.G., 2000. Marine, estuarine, and diadromous fish stocks at risk of extinction in North America (exclusive of Pacific salmonids). *Fisheries* 25, 6–30.
- Nagelkerken, I., Sheaves, M., Baker, R., Connolly, R.M., 2015. The seascape nursery: a novel spatial approach to identify and manage nurseries for coastal marine fauna. *Fish. Fish.* 16, 362–371.
- National Oceanic and Atmospheric Administration (NOAA), 2013a. DRAFT Belt Transect Fish Survey Protocol for Atlantic/Caribbean. National Coral Reef Monitoring Program (NCRMP), Silver Spring, MD.
- National Oceanic and Atmospheric Administration (NOAA), 2013b. Draft Line Point-Intercept (LPI) Survey Protocol for the Atlantic/Caribbean. National Coral Reef Monitoring Program (NCRMP). National Coral Reef Monitoring Program (NCRMP), Silver Spring, MD.
- Nemeth, R.S., Kadison, E., Herzlieb, S., Blondeau, J., Whiteman, E., 2006. Status of a yellowfin grouper (*Mycteroperca venenosa*) spawning aggregation in the U.S. Virgin Islands with notes on other species. *Proc. Gulf Caribb. Fish. Inst.* 57, 543–558.
- Neves, L.M., Teixeira-Neves, T.P., Pereira-Filho, G.H., Araújo, F.G., 2016. The farther the better: effects of multiple environmental variables on reef fish assemblages along a distance gradient from river influences. *PLoS One* 11, e0166679.
- Newman, M.J.H., Paredes, G.A., Sala, S., Jackson, J.B.C., 2006. Structure of Caribbean coral reef communities across a large gradient of fish biomass. *Ecol. Lett.* 9, 1216–1227.
- Nystrom, M., Folke, C., 2001. Spatial resilience of coral reefs. *Ecosystems* 4, 406–417.
- Obura, D.O., Grimsditch, G., 2009. Resilience Assessment of Coral Reefs – Assessment protocol for coral reefs, focusing on coral bleaching and thermal stress. IUCN Working Group on Climate Change and Coral Reefs. IUCN, Gland, Switzerland (70 pages).
- Ojeda-Serrano, E., Appeldoorn, R.S., Ruiz-Valentín, I., 2007a. Reef fish spawning aggregations of the Puerto Rican shelf. *Proc. Gulf Caribb. Fish. Inst.* 59, 467–474.
- Ojeda-Serrano, E., Appeldoorn, R.S., Ruiz-Valentín, I., 2007b. Reef fish spawning aggregations of the Puerto Rican shelf. Final report to the Caribbean Coral Reef Institute http://ccri.uprm.edu/researcher/Ojeda/Ojeda_Final_Report_CCRI_SPAG%27s.pdf.
- Oliver, L.M., Lehrter, J.C., Fisher, W.S., 2011. Relating landscape development intensity to coral reef condition in the watersheds of St. Croix, US Virgin Islands. *Mar. Ecol. Prog. Ser.* 427, 293–302.
- Oliver, L.M., Fisher, W.S., Dittmar, J., Hallock, P., Campbell, J., Quarles, R.L., Harris, P., LoBue, C., 2014. Contrasting responses of coral reef fauna and foraminiferal assemblages to human influence in La Parguera, Puerto Rico. *Mar. Environ. Res.* 99, 95–105. <https://doi.org/10.1016/j.marenvres.2014.04.005>.
- Oliver, L.M., Fisher, W.S., Fore, L., Smith, A., Bradley, P., 2018. Assessing land use, sedimentation, and water quality stressors as predictors of coral reef condition in St. Thomas, U.S. Virgin Islands. *Environ. Monit. Assess.* 190, 213–228.
- Paddock, M.J., Reynolds, J.D., Aguilar, C., Appeldoorn, R.S., Beets, J., Burkett, E.W., Chittaro, P.M., Clarke, K., Esteves, R., Fonseca, A.C., Forrester, G.E., 2009. Recent region-wide declines in Caribbean reef fish abundance. *Curr. Biol.* 19, 590–595.
- Pandolfi, J.M., Bradbury, R.H., Sala, E., Hughes, T.P., Bjorndal, K.A., Cooke, R.G., McArdle, D., McClanahan, L., Newman, M.J.H., Paredes, G., Warner, R.R., Jackson, J.B.C., 2003. Global trajectories of the long-term decline of coral reef ecosystems. *Science* 301, 955–958.
- Pendleton, L.H., 1995. Valuing coral reef protection. *Ocean Coast. Manag.* 26, 119–131.
- Pittman, S.J., Christensen, J.D., Caldwell, C., Menza, C., Monaco, M.E., 2007. Predictive mapping of fish species richness across shallow-water seascapes in the Caribbean. *Ecol. Model.* 204, 9–21.
- Pittman, S.J., Hile, S.D., Jeffrey, C.F.G., Clark, R., Woody, K., Herlach, B.D., Caldwell, C., Monaco, M.E., Appeldoorn, R., 2010. Coral Reef Ecosystems of Reserva Natural de La Parguera (Puerto Rico): Spatial and Temporal Patterns in Fish and Benthic Communities (2001–2007). NOAA Technical Memorandum NOS NCCOS 107. NOS NCCOS CCMA Silver Spring, MD.
- Pratchett, M.S., Hoey, A.S., Wilson, S.K., 2014. Reef degradation and the loss of critical ecosystem goods and services provided by coral reef fishes. *Curr. Opin. Environ. Sustain.* 7, 37–43.
- Principe, P., Bradley, P., Yee, S., Fisher, W., Johnson, E., Allen, P., Campbell, D., 2012. Quantifying Coral Reef Ecosystem Services. U.S. Environmental Protection Agency, Office of Research and Development, Research Triangle Park, NC (EPA/600/R-11/206, January 2012).
- Richards, B.L., Williams, I.D., Nadon, M.O., Zgliczynski, B.J., 2011. A towed-diver survey method for mesoscale fishery-independent assessment of large-bodied reef fishes. *Bull. Mar. Sci.* 87, 55–74.
- Risk, M.J., 1972. Fish diversity on a coral reef in the Virgin Islands. *Atoll Res. Bull.* 152, 1–6.
- Roberts, C.M., Ormond, R.F.G., 1987. Habitat complexity and coral reef fish diversity and abundance on Red Sea fringing reefs. *Mar. Ecol. Prog. Ser.* 41, 1–8.
- Robinson, J.P.W., Williams, I.D., Edwards, A.M., McPherson, J., Yeager, L., Vigliola, L., Baum, J.K., 2017. Fishing degrades size structure of coral reef fish communities. *Glob. Chang. Biol.* 23 (3), 1009–1022.
- Rogers, C.S., 1990. Responses of coral reefs and reef organisms to sedimentation. *Mar. Ecol. Prog. Ser.* 62, 185–202.
- Rogers, C.S., Garrison, G., Grober, R., Hillis, Z.M., Franke, M.A., 1994. Coral Reef Monitoring Manual for the Caribbean and Western Atlantic. U. S. National Park Service, St. John, U. S. Virgin Islands.
- Rude, J., Minks, A., Doheny, B., Tyner, M., Maher, K., Huffard, C., Hidayat, N.I.,

- Grantham, H., 2016. Ridge to reef modelling for use within land–sea planning under data-limited conditions. *Aquat. Conserv. Mar. Freshwat. Ecosyst.* 26 (2), 251–264.
- Sadovy, Y., Colin, P.L., Domeier, M.L., 1994a. Aggregation and spawning in the tiger grouper, *Mycteroperca tigris* (Pisces: Serranidae). *Copeia* 1994, 511–516.
- Sadovy, Y., Rosario, A., Román, A., 1994b. Reproduction in an aggregating grouper, the red hind, *Epinephelus guttatus*. *Environ. Biol. Fish* 41, 269–286.
- Sala, E., Ballesteros, E., Starr, R.M., 2001. Rapid decline of Nassau grouper spawning aggregations in Belize: fishery management and conservation needs. *Fisheries* 26, 23–30.
- Sale, P.F., 1991. Habitat structure and recruitment in coral reef fishes. In: Bell, S.S., McCoy, E.D., Mushinsky, H.R. (Eds.), *Habitat Structure: The Physical Arrangement of Objects in Space*. Chapman and Hall, pp. 197–210.
- Sale, P.F., Van Lavieren, H., Ablan Lagman, M.C., Atema, J., Butler, M., Fauvelot, C., Hogan, J.D., Jones, G.P., Lindeman, K.C., Paris, C.B., Steneck, R., Stewart, H.L., 2010. *Preserving Reef Connectivity: A Handbook for Marine Protected Area Managers*. Connectivity Working Group, Coral Reef Targeted Research & Capacity Building for Management Program, UNU-INWEH.
- Salm, R.V., Smith, S.E., Llewellyn, G., 2001. Mitigating the impact of coral bleaching through marine protected area design. In: *Coral Bleaching: Causes, Consequences and Response*. University of Rhode Island, USA, pp. 81–88.
- Sanchez, P.J., Appeldoorn, R.S., Schärer-Umpierre, M.T., Locascio, J.V., 2017. Patterns of courtship acoustics and geophysical features at spawning sites of black grouper (*Mycteroperca bonaci*). *Fish. Bull.* 115, 186–195. <https://doi.org/10.7755/FB.115.2.5>.
- Santavy, D.L., Fisher, W.S., Campbell, J.G., Quarles, R.L., 2012. *Field Manual for Coral Reef Assessments*. U.S. Environmental Protection Agency, Office of Research and Development, Gulf Ecology Division, Gulf Breeze, FL (EPA/600/R-12/029).
- Santavy, D.L., Bradley, P., Gerritsen, J., Oliver, L., 2016. The biological condition gradient, a tool used for describing the condition of US coral reef ecosystems. In: *Proceedings of the 13th International Coral Reef Symposium*, pp. 557–568.
- Schallenberg, M., Kelly, D., Clapcott, J., Death, R., C., MacNeil, Young, R., Sorrell, B., Scarsbrook, M., 2011. *Approaches to Assessing Ecological Integrity of New Zealand Freshwaters*. Department of Conservation, Wellington 6143, New Zealand (40 pp).
- Schärer-Umpierre, M.T., 2009. *Using Landscape Ecology to Describe Habitat Connectivity for Coral Reef Fishes* (PhD Thesis). University of Puerto Rico, Mayagüez, Puerto Rico (216 pp).
- Shapiro, D.Y., Sadovy, Y., McGehee, M.A., 1993. Size, composition, and spatial structure of the annual spawning aggregation of the Red hind, *Epinephelus guttatus* (Pisces: Serranidae). *Copeia* 1993, 399–406.
- Shcherbina, A.Y., Gawarkiewicz, G.G., Linder, C.A., Thorrold, S.R., 2008. Mapping bathymetric and hydrographic features of Glover's Reef, Belize, with a REMUS autonomous underwater vehicle. *Limnol. Oceanogr.* 53, 2264–2272.
- Smith, S.G., Ault, J.S., Bohnsack, J.A., Harper, D.E., Luo, J., McClellan, D.B., 2011. Multispecies survey design for assessing reef-fish stocks, spatially-explicit management performance, and ecosystem condition. *Fish. Res.* 109, 25–41.
- Spalding, M., Burke, L., Wood, S.A., Ashpole, J., Hutchison, J., Ermgassen, P., 2017. Mapping the global value and distribution of coral reef tourism. *Mar. Policy* 82, 104–113.
- Stallings, C.D., 2008. Indirect effects of an exploited predator on recruitment of coral-reef fishes. *Ecology* 89, 2090–2095.
- Stallings, C.D., 2009. Predator identity and recruitment of coral-reef fishes: indirect effects of fishing. *Mar. Ecol. Prog. Ser.* 383, 251–259.
- Stevens, M.H., Smith, S.G., Ault, J.S., 2019. Life history demographic parameter synthesis for Florida and Caribbean reef fishes. *Fish. Fish.* 20, 1196–1217. <https://doi.org/10.1111/faf.12405>.
- Talbot, F.H., Goldman, B., 1972. A preliminary report on the diversity and feeding relationships of reef fishes of One Tree Island, Great Barrier Reef system. In: *Proceedings of the First International Symposium on Corals and Coral Reefs*. 1. pp. 425–443.
- The White House, 2004. *U.S. Ocean Action Plan: The Bush Administration's Response to the U.S. Commission on Ocean Policy*. The White House, Washington, DC.
- Thompson, R., Munro, J.L., 1974. *The Biology, Ecology and Exploitation and Management of the Caribbean Reef Fishes. Part V. Carangidae (Jacks)*. (Research report from the Zoology Department, University of the West Indies).
- Turgeon, D.D., Asch, R.G. (Eds.), 2002. *The State of Coral Reef Ecosystems of the United States and Pacific Freely Associated States: 2002*. National Oceanic and Atmospheric Administration/National Ocean Service/National Centers for Coastal Ocean Science, Silver Spring, MD.
- U.S. Environmental Protection Agency (EPA), 1990. *Biological Criteria: National Program for Surface Waters*. U.S. Environmental Protection Agency, Office of Water, Washington, DC (EPA-440/5-90-004).
- U.S. Environmental Protection Agency (EPA), 2002. *Biological Assessments and Criteria: Crucial Components of Water Quality Programs*. EPA-822-F-02-006.
- U.S. Environmental Protection Agency (EPA), 2011a. *A Primer on Using Biological Assessments to Support Water Quality Management*. EPA 810-R-11-01. U.S. Environmental Protection Agency, Washington, DC.
- U.S. Environmental Protection Agency (EPA), 2011b. *A Field-based Aquatic Life Benchmark for Conductivity in Central Appalachian Streams*. EPA/600/R-10/023F. U.S. Environmental Protection Agency, Office of Research and Development, National Center for Environmental Assessment, Washington, DC. <http://cfpub.epa.gov/ncea/cfm/recordisplay.cfm?deid=233809> (Accessed February 2016).
- U.S. Environmental Protection Agency (EPA), 2013a. *Biological Assessment Program Review: Assessing Level of Technical Rigor to Support Water Quality Management*. EPA 820-R-13-001. U.S. Environmental Protection Agency, Washington, DC.
- U.S. Environmental Protection Agency (EPA), 2013b. *Biological Assessment Program Review: Assessing Level of Technical Rigor to Support Water Quality Management*. EPA 820-R-13-001. U.S. Environmental Protection Agency, Washington, DC.
- U.S. Environmental Protection Agency (EPA), 2016. *A Practitioner's Guide to the Biological Condition Gradient: A Framework to Describe Incremental Change in Aquatic Ecosystems*. EPA 842-R-16-001, Wash. DC.
- Vega Thurber, R.L., Burkepile, D.E., Fuchs, C., Schantz, A.A., McMinds, R., Zaneveld, J.R., 2014. Chronic nutrient enrichment increases prevalence and severity of coral disease and bleaching. *Glob. Chang. Biol.* 20, 544–554.
- Waddell, J.E. (Ed.), 2005. *The State of Coral Reef Ecosystems of the United States and Pacific Freely Associated States: 2005*. National Oceanic and Atmospheric Administration, Silver Spring, MD.
- Waddell, J.E., Clarke, A.M. (Eds.), 2008. *The State of Coral Reef Ecosystems of the United States and Pacific Freely Associated States: 2008*. NOAA Technical Memorandum NOS CCOS 73. NOAA/NCCOS Center for Coastal Monitoring and Assessment's Biogeography Team, Silver Spring, MD.
- Walker, B.K., Jordan, L.K.B., Spieler, R.E., 2009. Relationship of reef fish assemblages and topographic complexity on southeastern Florida coral reef habitats. *J. Coast. Res.* 53, 39–48.
- Wenger, A.S., Fabricius, K.E., Jones, G.P., Brodie, J.E., 2015. Effects of sedimentation, eutrophication, and chemical pollution on coral reef fishes, chap 15. In: Mora, C. (Ed.), *Ecology of Fishes on Coral Reefs*. Cambridge University Press, Cambridge, pp. p145–p153.
- West, J.M., Salm, R.V., 2003. Resistance and resilience to coral bleaching: implications for coral reef conservation and management. *Conserv. Biol.* 17 (4), 956–967.
- World Resources Institute (WRI), U.S. National Oceanographic and Atmospheric Administration (NOAA), 2006. *Land-based Sources of Threat to Coral Reefs in the U.S. Virgin Islands*. (Washington, DC. 14 pp).
- Ydenberg, R.C., Dill, L.M., 1986. The economics of fleeing from predators. *Adv. Study Behav.* 16, 229–249.
- Zadeh, L.A., 2008. Is there a need for fuzzy logic? *Inf. Sci.* 178, 2751–2799.