

## Clinical paper

# The continuous quality improvement project for telephone-assisted instruction of cardiopulmonary resuscitation increased the incidence of bystander CPR and improved the outcomes of out-of-hospital cardiac arrests<sup>☆</sup>

Yoshio Tanaka<sup>a</sup>, Junro Taniguchi<sup>b</sup>, Yukihiro Wato<sup>c</sup>, Yutaka Yoshida<sup>d</sup>, Hideo Inaba<sup>a,\*</sup>

<sup>a</sup> Department of Emergency Medical Science, Kanazawa University Graduate School of Medicine, Kanazawa, Japan

<sup>b</sup> Emergency Medical Center, Ishikawa Prefecture Central Hospital, Kanazawa, Japan

<sup>c</sup> Department of Emergency Medicine, Kanazawa Medical University, Kanazawa, Japan

<sup>d</sup> Section of Anaesthesia, Komatsu Citizens Hospital, Komatsu, Japan

## ARTICLE INFO

## Article history:

Received 29 July 2011

Received in revised form 13 February 2012

Accepted 18 February 2012

## Keywords:

Telephone-CPR

Continuous quality improvement

Cardiopulmonary resuscitation

Dispatcher

Emergency medical service

Out-of-hospital cardiac arrest

## ABSTRACT

**Review:** In 2007, the Ishikawa Medical Control Council initiated the continuous quality improvement (CQI) project for telephone-assisted cardiopulmonary resuscitation (telephone-CPR), which included instruction on chest-compression-only CPR, education on how to recognise out-of-hospital cardiac arrests (OHCAs) with agonal breathing, emesis and convulsion, recommendations for on-line or redialling instructions and feedback from emergency physicians. This study aimed to investigate the effect of this project on the incidence of bystander CPR and the outcomes of OHCAs.

**Materials and methods:** The baseline data were prospectively collected on 4995 resuscitation-attempted OHCAs, which were recognised or witnessed by citizens rather than emergency medical technicians during the period of February 2004 to March 2010. The incidence of telephone-CPR and bystander CPR, as well as the outcomes of the OHCAs, was compared before and after the project.

**Results:** The incidence of telephone-CPR and bystander CPR significantly increased after the project (from 42% to 62% and from 41% to 56%, respectively). The incidence of failed telephone-CPR due to human factors significantly decreased from 30% to 16%. The outcomes of OHCAs significantly improved after the projects. A multiple logistic regression analysis revealed that the CQI project is one of the independent factors associated with one-year (1-Y) survival with favourable neurological outcomes (odds ratio = 1.81, 95% confidence interval = 1.20–2.76).

**Conclusions:** The CQI project for telephone-CPR increased the incidence of bystander CPR and improved the outcome of OHCAs. A CQI project appeared to be essential to augment the effects of telephone-CPR.

© 2012 Elsevier Ireland Ltd. All rights reserved.

## 1. Introduction

Currently, medical control (MC) for the emergency medical service system (EMS) is active in Japan. However, MC, including education, is mainly targeted to emergency medical technicians (EMTs) and is rarely targeted to dispatchers in Japan. Since February 2004, the Ishikawa MC Council has requested that fire departments collect the reasons why dispatchers failed to provide telephone-assisted instruction on cardiopulmonary resuscitation

(telephone-CPR). Telephone-CPR has been shown to increase the incidence of bystander CPR and is expected to improve the outcomes of out-of-hospital cardiac arrests (OHCAs).<sup>1–5</sup>

There are no special qualifications or authorisations required for fire department staff to be dispatchers in Japan. Some dispatchers have no experience as EMTs. An educational programme for dispatchers has not been established. In some fire departments, the actual condition of the OHCA patient, discovered by EMTs during a post-arrival interview, is not relayed back to the dispatchers.

In March of 2007, the Ishikawa MC Council initiated the continuous quality improvement (CQI) project for telephone-CPR. The project included the following: (1) a standardised manual for instruction on chest-compression-only CPR (CC-only CPR), (2) education on how to recognise OHCAs with agonal breathing,<sup>6,7</sup> emesis<sup>8</sup> and convulsion,<sup>9,10</sup> (3) recognition of an impending cardiac arrest,<sup>11</sup> (4) recommendations for on-line or redialling instructions for patients with impending cardiac arrest,<sup>12,13</sup> (5) instructions on

<sup>☆</sup> A Spanish translated version of the abstract of this article appears as Appendix in the final online version at doi:10.1016/j.resuscitation.2012.02.013.

\* Corresponding author at: Department of Emergency Medical Science, Kanazawa University Graduate School of Medicine, 13-1 Takara-machi, Kanazawa 920-8641, Japan. Tel.: +81 76 265 2825; fax: +81 76 234 4243.

E-mail address: [hidinaba@med.kanazawa-u.ac.jp](mailto:hidinaba@med.kanazawa-u.ac.jp) (H. Inaba).

the use of a cellular phone as a communication device at the scene<sup>14</sup> and (6) feedback from emergency physicians and EMT supervisors to dispatchers.

The purpose of this study was to investigate the effects of the CQI project on the incidences of telephone-CPR and CPR before EMT arrival. This study also investigated the effects of the project on the time factors and outcomes of individuals experiencing an OHCA that was recognised or witnessed by citizens and not witnessed by EMTs.

## 2. Methods

The data were collected in accordance with the national guidelines of ethics for epidemiological surveys.<sup>15</sup> This study has been approved by the review board of the Ishikawa MC Council.


### 2.1. Populations and setting

The Ishikawa prefecture encompasses a snowfall area of 4186 km<sup>2</sup> and has a resident population of 1,170,000. The prefecture is divided into four administrative regions: one central or urban region and three semi-rural or rural regions. Sixty-two percent of the residents are located in the central (urban) region, which has an area of 1432 km<sup>2</sup>. The age of the population is more advanced in rural regions (28.5% vs. 20.3% over the age of 65).

EMTs resuscitate patients experiencing an OHCA according to the protocol developed by the Ishikawa MC Council, which is based on the guidelines of the American Heart Association and the Japan Resuscitation Council. The paramedics are authorised to perform the following procedures during the resuscitation: (1) use a semi-automated external defibrillator, (2) use a supra-pharyngeal airway and (3) the infusion of Ringer's lactate. Since July 2004, specially trained paramedics have been permitted to insert tracheal tubes under limited indication criteria,<sup>16</sup> and since April 2006, they have been permitted to administer intravenous adrenaline. The EMTs are not permitted to terminate resuscitation in the field. All resuscitation-attempted victims are transported to hospitals. EMS changes over time are summarised in Fig. 1.

### 2.2. CQI project for telephone-CPR

Since February 2004, telephone-CPR has been conducted by all fire departments (Fig. 1), and the reasons why dispatchers failed to attempt the telephone-CPR have been recorded prospectively (Fig. 3). Our CQI project was based on the prior analysis of the reasons for failure followed by planning and preparing for the project's implementation. Our project was implemented in March 2007 and is presently being continued. The project consists of the following: (1) enforcement of a uniform telephone-CPR manual for CC-only CPR, (2) a standard educational approach on how to detect and recognise OHCA with agonal breathing,<sup>6,7</sup> emesis<sup>8</sup> and anoxic seizure<sup>9,10</sup> as well as an impending cardiac arrest,<sup>11</sup> (3) behaviour modification for dispatchers and (4) conferences between EMTs and dispatchers to resolve discrepancies in their information. The telephone-CPR manual states the following. (1) Telephone-CPR may be initiated by obtaining other information about the situation and events suggestive of cardiac arrest, even when a caller is not able to judge whether an OHCA victim is responsive or breathing normally. (2) When snoring, vomiting and abnormal motions are preferentially described by the caller, a dispatcher should ask for evidence of agonal breathing, emesis or regurgitation and anoxic seizures associated with cardiac arrest. For example, "Is the victim unresponsive, unconscious, or breathing abnormally?" and "Has the victim vomited (or moved abnormally) before becoming unresponsive (or pale)?" or "Are you only seeing the vomitus on the




**Fig. 1.** Changes in the EMS protocol and the incidence of CPR following telephone-CPR before and after the CQI project. "CPR performed without telephone-CPR" means that CPR was performed by citizens on their own initiative despite the absence of telephone-CPR either before or after an emergency call. Percentage values for CC-only CPR are expressed as a fraction of the total number of CPR cases for the "CPR performed without telephone-CPR" and "CPR performed following telephone-CPR" groups.

victim's face or floor?" (3) A dispatcher should request that the caller redial them via a mobile or wireless phone when they are not beside the OHCA victim. (4) A dispatcher should request that the caller redial them if they notice any changes in the patient with impending cardiac arrest, or the dispatcher should call back the caller periodically, when there are sufficient resources.

New programmes were introduced in prefectural school, and simulation trainings for dispatchers in fire departments were included in the educational approach. Telephone-CPR quality was evaluated in all OHCA by emergency physicians and EMT supervisors, and their evaluations, including an appreciative comment, were given as feedback to the dispatchers.

### 2.3. Patient data

Baseline data were collected prospectively by fire departments for 4995 resuscitation-attempted OHCA that were witnessed or recognised by citizens rather than EMTs from 1 February 2004 to 31 March 2010. The collected data were based on the Utstein template<sup>17,18</sup> and included the patient's backgrounds, arrest witness, causes of arrest (presumably cardiac or not), bystander CPR, initial cardiac rhythm, estimated time of collapse, times of the initiation of CPR by bystanders and EMTs, EMT arrival on the scene, sustained return of spontaneous circulation (SROSC), one-month




**Fig. 2.** Annual changes in the incidence of CPR prior to EMT arrival and patient outcomes. The number of OHCA dispatches is shown in numeric values. The interval between the call and EMT arrival on the scene (response time) is shown as a median (25–75%) value.

(1-M) survival, one-year (1-Y) survival and 1-Y survival with a favourable neurological outcome, as determined by the Pittsburgh cerebral performance category (CPC).<sup>19</sup> The SROSC is defined as the continuous presence of palpable pulses for more than 20 min.<sup>17,18</sup> The survival rate at 1-Y was defined as the patient being alive in a hospital at 1-Y or as the patient being alive and discharged from the hospital to a home or to a care or rehabilitation facility within 1-Y. One-year survival with a favourable neurological outcome was defined as a CPC of one (good recovery) or two (moderate

disability) in patients without any neurological disturbance before the arrest event and when the best CPC was equal to the pre-arrest category in patients with neurological disturbance. The primary end point was 1-Y survival with a favourable neurological outcome. The secondary end points were SROSC, 1-M survival and 1-Y survival.

When telephone-CPR was not attempted during an OHCA, the causes of failure were analysed by reviewing voice recordings and/or dispatch reports as well as the EMT's interview records



**Fig. 3.** Incidences of telephone-CPR and failure due to preventable factors before and after the CQI project. All percentage values are expressed as a fraction of all OHCA's.

for callers. Supervisors and emergency physicians involved in the project classified the reasons for failure into two categories: unpreventable factors related to the callers and the scenes and preventable human factors related to the dispatchers.

We compared the incidences of telephone-CPR and CPR before EMT arrival as well as the outcome of OHCA between two time periods: before (1 February 2004 to 28 February 2007) and after the initiation of the project (1 March 2007 to 31 March 2010).

#### 2.4. Statistical analysis

We analysed the data using JMP version 7 for Windows (SAS institute, Cary, NC, USA). The chi-squared test with and without a Pearson's correction was applied for the univariate analyses. The Kruskal–Wallis test was used for non-parametric comparisons. We used a multiple logistic regression analysis to identify the factors associated with outcome. In all analyses,  $p < 0.05$  was considered significant.

### 3. Results

#### 3.1. Incidence of CPR following telephone-CPR before and after the project

As shown in Fig. 1, the CQI project significantly increased the incidence of CPR following telephone-CPR from 24% to 42%. Regardless of whether CPR was initiated either under one's own initiative or following telephone-CPR, the incidence of CC-only CPR as a CPR method significantly increased after the project. Annual changes in the incidence of CPR before EMT arrival are shown in Fig. 2. Although the incidence of CPR before arrival has been continuously increasing, the increase appeared to be augmented by the project. As demonstrated in Fig. 3, the project significantly increased the incidence of telephone-CPR from 42% to 62%. Telephone-CPR was not attempted in some cases with impending cardiac arrest. The project decreased the incidence of failed attempts due to preventable human factors from 30% to 16% when the impending cardiac arrest was included in the preventable cause and from 25% to 12% when the impending arrest was independently categorised.

#### 3.2. Background and management of OHCA patients before and after the CQI project (Table 1 and Fig. 1)

As shown in Table 1, there were differences in the background and characteristics of OHCA patients between the two time periods. The incidence of a presumed cardiac aetiology was significantly higher before the project (52% vs. 48%), and the patient's age was higher after the project (76 vs. 78). Witnessed OHCA of a presumed cardiac aetiology and with a shockable initial rhythm more frequently occurred at the patient's home after the project (47% vs. 68%). Advanced CPR procedures including tracheal intubation (6% vs. 14%) and adrenaline administration (0.3% vs. 6%) were more frequently performed after the project.

As shown in Fig. 1, the incidence of CPR before EMT arrival was 41% before the project and increased to 56% after the CQI (odds ratio = 1.75, 95% confidence interval = 1.57–1.96). Citizens more frequently used CC-only CPR after the project (67% vs. 83%).

#### 3.3. Time factors and prognoses of OHCA before and after the CQI project (Fig. 2 and Table 1)

As shown in Fig. 2, the incidence of SROSC and 1-Y survival with favourable neurological outcomes appeared to be augmented after the project, though a small drop in the incidence of SROSC was observed in the fiscal year term of 2009 having a larger number of OHCA.




Fig. 4. Subcategorised comparisons of the outcomes between the two time periods for OHCA managed without prehospital ACLS procedures. ★ significant difference between the two times periods (using a chi-squared test with and without Person's correction).

As shown in Table 1, the interval between the call and the initiation of CPR (by citizens or EMTs, whichever was performed first) significantly decreased after the project when analysed for all OHCA. The interval between the witnessing or recognition of cardiac arrest and the call significantly increased after the project. For all OHCA given CPR before EMT arrival, the interval between the call and CPR initiation by citizens was significantly reduced. The interval between the call and EMT arrival at the scene significantly increased after the project when analysed for all OHCA. The incidence of 1-Y survival with a favourable neurological outcome was significantly improved after the project (1.9% vs. 2.8%,  $p = 0.0443$ ) when analysed for all OHCA. For the witnessed OHCA with a presumed cardiac aetiology, having an initial shockable rhythm, the incidence of 1-Y survival (22.8% vs. 35.7%,  $p = 0.0243$ ) and 1-Y survival with a favourable neurological outcome (16.3% vs. 27.8%,  $p = 0.0276$ ) were significantly improved after the project.

#### 3.4. Subcategorised comparisons of outcomes between the two time periods for OHCA managed without prehospital advanced cardiac life support (ACLS) procedures (Fig. 4)

Because tracheal intubation and adrenaline administration have been shown to affect the outcomes of OHCA,<sup>16,20–23</sup> we analysed the effect of the project on the outcomes of OHCA managed prior to hospital arrival without tracheal intubation or adrenalin administration. As shown in Fig. 4, we found a significant improvement in the outcomes of OHCA after the project.

#### 3.5. Elucidation of factors related to 1-Y survival with favourable neurological outcomes in all OHCA (Table 2)

A multiple logistic regression analysis following a univariate analysis revealed that the CQI project is one of the independent factors related to 1-Y survival with favourable neurological outcomes. Decreased age, a witnessed cardiac arrest, cardiac arrests with a presumed cardiac aetiology, a shockable initial rhythm, a shorter response time and a shorter interval between the witnessing or recognition of cardiac arrest and a telephone call were other independent factors associated with favourable 1-Y survival. Adrenaline administration was an independent factor related to unfavourable 1-Y survival, whilst tracheal intubation was an independent factor that was not associated with favourable 1-Y survival.

**Table 1**  
Background, time factors and management of OHCA before and after the CQI project.

Background, characteristics and management	All OHCA			Witnessed OHCA with a presumed cardiac aetiology with a shockable initial rhythm		
	Before the project (N=2325)	After the project (N=2670)	p	Before the project (N=23)	After the project (N=126)	p
Arrest – witnessed, % (N)	37% (852)	38% (1004)	p=0.4846	–	–	Undefined
Aetiology – presumably cardiac, % (N)	52% (1199)	48% (1269)	p=0.0044	–	–	Undefined
Location – Home, % (N)	66% (1537)	67% (1796)	p=0.3862	47% (58)	68% (86)	p=0.0007
Sex – male, % (N)	60% (1402)	59% (1583)	p=0.4666	81% (99)	74% (93)	p=0.2090
Age, y median (25–75%)	76 (63–84)	78 (65–85)	p<0.0001	63 (55–72)	64 (54–75)	p=0.4808
Region – central (urban), % (N)	47% (1096)	50% (1326)	p=0.0751	60% (74)	54% (68)	p=0.3233
Tracheal intubation, % (N)	6% (143)	14% (367)	p<0.0001	4% (5)	8% (10)	p=0.1949
Adrenalin, % (N)	0.3% (8)	6% (166)	p<0.0001	0% (0)	6% (7)	p=0.0018
Call to CPR by citizens, min median (25–75%) N	1 (–1–2) N=973	0 (–1–1) N=1489	p<0.0001	0.5 (–1–2) N=66	0.5 (0–2) N=80	p=0.4707
Call to first CPR <sup>a</sup> , min median (25–75%)	5 (1–8)	2 (0–8)	p<0.0001	3 (0–7)	2 (0–7)	p=0.8577
Call to arrival at patient, min median (25–75%)	7 (6–10)	8 (6–10)	p=0.0069	7 (5–9)	8 (6–9)	p=0.2356
Witness/recognition to call, min median (25–75%)	2 (1–5)	3 (1–6)	p<0.0001	1 (0–2)	2 (1–3)	p=0.0125
Duration of transportation, min median (25–75%)	8 (5–12)	9 (6–13)	p=0.0014	9 (6–13)	9 (5–12)	p=0.9538
1-Y survival, % (N)	3.2% (74)	4.2% (112)	p=0.0595	22.8% (28)	35.7% (45)	p=0.0243
1-Y survival with favourable neurological outcome, % (N)	1.9% (45)	2.8% (75)	p=0.0443	16.3% (20)	27.8% (35)	p=0.0276

<sup>a</sup> First CPR: CPR by citizens or EMTs, whichever was performed first.

**4. Discussion**

Telephone-CPR has the potential to improve the frequency of bystander CPR.<sup>1–5</sup> To enhance the beneficial effect of telephone-CPR and reduce the ‘barriers’ to telephone-CPR, educational approaches aimed at dispatchers and citizens are necessary.<sup>24–29</sup> Although teaching dispatchers to understand and recognise bystander descriptions of agonal respiration in patients with OHCA have been reported to result in a significant increase in the incidence of telephone-CPR,<sup>24,29,30</sup> the effects of education on survival after OHCA have not been reported. To the best of our knowledge, this study is the first study to demonstrate the efficacy of the educational approach with the CQI project.

Our CQI project was based on the analysis of reasons why dispatchers failed to initiate telephone-CPR; presumably due to ‘cognitive bias’, some callers interpret ‘snoring’ breathing sounds, abnormal body motion and vomitus on the face or the floor as a ‘vital’ sign of OHCA victims and preferentially describe them to a dispatcher. For example, “She/he is unhealthy but is snoring, moving or vomiting”. This may interfere with a dispatcher’s proper recognition of cardiac arrest and may lead to ‘no further

exploration’ by the dispatcher because snoring, abnormal motion or convulsions and emesis are common symptoms of acute illnesses other than cardiac arrest.<sup>31,32</sup> These failures may be minimised by teaching dispatchers that agonal breathing, anoxic convulsions and emesis or regurgitation often occur in OHCA patients.<sup>8–10,27–29</sup> We also realised that dispatchers frequently fail to attempt telephone-CPR for OHCA that are witnessed after the emergency call. Although the telephone-CPR in these cases may be difficult, a dispatcher in fire departments with sufficient resources may attempt it by recognising prearrest signs and symptoms<sup>11</sup> and by redialling the caller periodically.

The incidence of CPR before EMT arrival was significantly increased after the project (Fig. 1). The increased frequency of CPR before EMT arrival is mainly due to the increased incidence of CPR performed following telephone-CPR (Fig. 2) and the decreased failure of telephone-CPR attempts due to preventable human factors related to the dispatchers (Fig. 3).

The time factors related to citizen-performed CPR were also improved by this project. The interval between the call and the initiation of CPR significantly decreased after the project. For OHCA given CPR before EMT arrival, the interval between the call and CPR

**Table 2**  
Factors related to 1-Y survival with a favourable neurological outcome for all OHCA.

Factors	Survivors (N=120)	Non-survivors (N=4875)	Unadjusted odds ratio (95% C.I.) for survival or p value	Adjusted odds ratio (95% C.I.) for survival by multiple logistic regression
Patient’s age, median (25–75%)	63 (52–76)	77 (64–85)	p<0.0001	0.97 (0.96–0.98)
Sex – male, % (N)	66% (79)	60% (2906)	1.31 (0.89–1.91)	1.24 (0.80–1.57)
Location – home, % (N)	58% (70)	67% (3263)	0.69 (0.48–1.00)	0.96 (0.64–1.46)
Region – central, % (N)	58% (69)	48% (2353)	1.45 (1.01–2.09)	1.05 (0.69–1.59)
Arrest – witnessed, % (N)	78% (94)	36% (1762)	6.38 (4.12–9.90)	3.36 (2.10–5.53)
Aetiology – presumably cardiac, % (N)	73% (87)	49% (2381)	2.76 (1.84–4.14)	1.32 (0.81–2.17)
Initial rhythm – shockable, % (N)	58% (69)	6.5% (317)	19.45 (13.31–28.43)	9.32 (5.84–15.00)
CPR before EMT arrival, % (N)	59% (71)	49% (2391)	1.51 (1.04–2.18)	1.24 (0.69–2.12)
Call to arrival at patient, min median (25–75%)	6 (5–8)	8 (6–10)	p<0.0001	0.88 (0.83–0.93)
Witness/recognition to call, min median (25–75%)	2 (1–3)	2 (1–5)	p<0.0001	0.93 (0.90–0.98)
Call to first CPR <sup>a</sup> , min median (25–75%)	2 (0–6)	4 (0–8)	p=0.0016	0.98 (0.92–1.00)
Duration of transportation, min median (25–75%)	9 (5–14)	8 (5–13)	p=0.5973	1.00 (1.00–1.02)
CQI project, % (N)	63% (75)	53% (2595)	1.46 (1.01–2.13)	1.81 (1.20–2.76)

<sup>a</sup> First CPR: CPR by citizens or EMTs, whichever was performed first.

by citizens was significantly reduced. These favourable changes in time factors appear to be due not only to the increased incidence of citizen-performed CPR but also to the increased incidence of CC-only CPR because instruction on CC-only CPR accelerates the initiation of CC.<sup>12,33</sup>

Some of the other time factors related to survival, including the intervals between the recognition or witnessing of cardiac arrest and the call and between the call and EMT arrival on the scene were significantly prolonged after the project. These unfavourable changes were not related to the CQJ project and may be attributed to the increased age of the patients<sup>34</sup> and an increased number of dispatches in our region. Despite these unfavourable changes, a univariate analysis revealed that the outcomes of OHCAs were significantly improved after the project, even when subgroups of OHCAs and witnessed OHCAs with a presumed cardiac aetiology with a shockable initial rhythm were analysed independently (Table 1). A small decline of SROSC incidence was observed in the fiscal year beginning in February 2009 (Fig. 2). In this term, we had a larger number of OHCAs, and the interval between the call and the arrival on the scene was significantly prolonged, presumably due to unusually heavy snowfall in this winter.

The use of ACLS, including tracheal intubation and adrenaline administration, has been shown to affect the outcome of OHCAs.<sup>16,20–23</sup> We compared the outcomes between the two time periods for OHCAs managed prior to hospital arrival without pre-hospital ACLS procedures because the incidence of ACLS was higher after the project. We showed that the CQJ project improved the outcome of OHCAs. We further confirmed the effect of the CQJ project in all OHCAs using a multiple logistic regression analysis, which included all prehospital factors related to 1-Y survival. Whilst tracheal intubation was not an independent factor associated with a favourable 1-Y survival, adrenaline administration was a potent independent factor related to unfavourable 1-Y survival. These effects of ACLS procedures were carefully interpreted because the procedures were performed only by specially trained paramedics in Japan.

The risk of bystander CPR for patients not in cardiac arrest needs to be considered.<sup>35–38</sup> In this study, we did not determine how frequently telephone-CPR was attempted on patients who were not in cardiac arrest.<sup>39</sup> At present, no complications associated with telephone-CPR have been reported to the Ishikawa MC Council. We are now conducting surveillance of our telephone-CPR system and have reported the preliminary results of this analysis.<sup>40</sup>

#### 4.1. Limitations

Because this study was not randomised and not conducted in an on-off manner, the results of the outcome analysis might therefore be a result of coincidence. The CQJ project was initiated shortly after the EMT protocol changed, according to the 2005 Consensus (C2005), and the improvement of patient management by EMTs, particularly the quality of chest compressions, may have influenced the outcome of OHCAs. However, starting in April 2003, the quality assurance of patient care by EMTs has been a main issue addressed by our MC Council.

## 5. Conclusions

The CQJ project for telephone-CPR initiated shortly after the implementation of C2005 increased the incidence of CPR before EMT arrival and improved the outcomes of OHCAs in our community having the standard Japanese EMS system. These observations suggest that a CQJ project is essential to improve the outcome of telephone-CPR.

## Conflict of interest statement

We have no conflict of interest to disclose.

## Acknowledgements

We would like to thank all of the fire departments in Ishikawa, Japan for their help with data collection.

## References

- Eisenberg MS, Hallstorm AP, Carter WB, et al. Emergency CPR instruction via telephone. *Am J Public Health* 1985;75:47–50.
- O'Nwill JF, Deakin CD. Evaluation of telephone CPR advice for adult cardiac arrest patient. *Resuscitation* 2007;74:63–7.
- Bohm K, Rosenqvist M, Hollenberg J, et al. Dispatcher-assisted telephone-guided cardiopulmonary resuscitation: an underused life saving systems. *Eur J Emerg Med* 2007;14:256–9.
- Rea TD, Eisenberg MS, Culley LL, et al. Dispatcher-assisted cardiopulmonary resuscitation and survival in cardiac arrest. *Circulation* 2001;104:2513–6.
- Bohm K, Vaillancourt C, Charette ML, et al. In patients with out-of cardiac arrest, dose the provision of dispatch cardiopulmonary resuscitation instruction as opposed to no instructions improve outcome: a systematic review of the literature. *Resuscitation* 2011;82:1490–5.
- Vaillancourt C, Jensen JL, Grimshaw J, et al. A survey of factors associated with the successful of agonal breathing and cardiac arrest by 9-1-1 call takers: design and methodology. *BMC Emerg Med* 2009;9:14.
- Bohm K, Stålhandske B, Rosenqvist M, et al. Tuition of emergency medical dispatchers in the recognition of agonal respiration increases the use of telephone assisted CPR. *Resuscitation* 2009;80:1025–8.
- Simons RW, Rea TD, Becker LJ, et al. The incidence and significance of emesis associated with out-of-hospital cardiac arrest. *Resuscitation* 2007;74:427–31.
- Clawson J, Olola C, Heward A, et al. Cardiac arrest predictability in seizure patients based on emergency medical dispatcher identification of previous seizure or epilepsy history. *Resuscitation* 2007;75:298–304.
- Clawson J, Olola C, Scott G, et al. Effect of a medical priority dispatch system key question addition in the seizure/convulsion/fitting protocol to improve recognition of ineffective (agonal) breathing. *Resuscitation* 2008;79:257–64.
- Müller D, Agrawal R, Arntz H-R. How sudden is sudden cardiac death? *Circulation* 2006;114:1146–50.
- 2005 International consensus on cardiopulmonary resuscitation (CPR) and emergency cardiovascular care (ECC) science with treatment recommendation. Section 1, part 2: adult basic life support. *Circulation* 2005;112:III-5–16.
- Berg RA, Hemphill R, Abella BS, et al. Part 5: adult basic life support: 2010 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2010;122:S685–705.
- Koster RW, Baubin MA, Bossaert LL, et al. European Resuscitation Council guidelines for resuscitation 2010 Section 2: adult basic life support and use of automated external defibrillators. *Resuscitation* 2010;81:1277–92.
- The Ministry of Health, Labor and Welfare in Japan [accessed 04.12.11, at <http://www.mhlw.go.jp/general/seido/kousei/i-kenkyu/index.html>].
- Takei Y, Enami M, Yachida T, et al. Tracheal intubation by paramedics under limited intubation criteria may improve the short-term outcome of out-of-hospital cardiac arrests with noncardiac origin. *J Anesth* 2010;24:716–25.
- Recommended guidelines for uniform reporting of data from out-of-hospital cardiac arrest: the 'Utstein style'. Prepared by a Task Force of Representatives from the European Resuscitation Council, American Heart Association, Heart and Stroke Foundation of Canada, Australian Resuscitation Council. *Resuscitation* 1991;22:1–26.
- Jacob I, Nadkarni V, Bahr J, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update and simplification of the Utstein templates for resuscitation registries: a statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian Resuscitation Council, New Zealand Resuscitation Council, Heart and Stroke Foundation of Canada, Inter American Heart Foundation, Resuscitation Councils of Southern Africa). *Resuscitation* 2004;63:233–49.
- Safar P. Resuscitation after brain ischemia. In: Grenvik A, Safar P, editors. *Brain Failure and Resuscitation*. New York: Churchill Livingstone; 1981. p. 155–84.
- Stiell IG, Wells GA, Field B, et al. Advanced cardiac life support in out-of-hospital cardiac arrest. *N Engl J Med* 2004;351:647–56.
- Olasveengen TM, Sunde K, Brunborg C, et al. Intravenous drug administration during out-of-hospital cardiac arrest: a randomized trial. *JAMA* 2009;302:2222–9.
- Studnek JR, Thestrup L, Vandevender S, et al. The association between prehospital endotracheal intubation attempts and survival to hospital discharge among out-of-hospital cardiac arrest patient. *Acad Emerg Med* 2010;17:918–25.
- Jacobs IG, Finn JC, Jelinek GA, et al. Effect of adrenalin on survival in out-of-hospital cardiac arrest: a randomised double-blind placebo-controlled trial. *Resuscitation* 2011;82:1138–43.
- Vaillancourt C, Charette ML, Stiell IG, et al. An evaluation of 9-1-1 calls to assess the effectiveness of dispatch-assisted cardiopulmonary resuscitation (CPR) instructions: design and methodology. *BMC Emerg Med* 2008;8:12.

25. Bohm K, Stålhandske B, Rosenqvist M, et al. Tuiton of emergency medical dispatchers in the recognition of agonal respiration increases the use of telephone assisted CPR. *Resuscitation* 2009;80:1025–8.
26. Vaillancourt C, Verma A, Trickett J, et al. Evaluating the effectiveness of dispatch-assisted cardiopulmonary resuscitation instructions. *Acad Emerg Med* 2007;14:877–83.
27. Nurmi J, Pettila V, Biber B, et al. Effect of protocol compliance to cardiac arrest identification by emergency medical dispatchers. *Resuscitation* 2006;70:463–9.
28. Vaillancourt C, Charette ML, Bohm K, et al. In out-of-hospital cardiac arrest patient, dose the description of any specific symptoms to the emergency medical dispatcher improve the accuracy of the diagnosis of cardiac arrest: a systematic review of the literature. *Resuscitation* 2011;82:1483–9.
29. Berdowski J, Beekhuis F, Zwinderman AH, et al. Importance of the first link: description and recognition of an out-of-hospital cardiac arrest in an emergency call. *Circulation* 2009;119:2096–102.
30. Roppolo LP, Westfall A, Pepe PE, et al. Dispatcher assesments for agonal breathing improved detection of cardiac arrest. *Resuscitation* 2009;80:769–72.
31. Ingram DA, Fulton RA, Portal RW, et al. Vomiting as a diagnostic aid in acute ischaemic cardiac pain. *BMJ* 1980;281:636–7.
32. Krieger DW, Demchuk Am, Kasner SE, et al. Early clinical and radiological predictors of fatal brain swelling in ischemic atroke. *Stroke* 1999;30:287–92.
33. International Guidelines 2000 for CPR and ECC. Part 3: adult basic life support. *Circulation* 2000;102:1-22–59.
34. Takei Y, Inaba H, Yachida T, et al. Analysis of reasons for emergency call delays in Japan in relation to location: high incidence of correctable causes and the impact of delays on patient outcomes. *Resuscitation* 2010;81:1492–8.
35. Krischer JP, Fine EG, Davis JH, et al. Complications of cardiac resuscitation. *Chest* 1987;92:287–91.
36. Corbett SW, O'Callaghan T. Detection of traumatic complications of cardiopulmonary resuscitation by ultrasound. *Ann Emerg Med* 1997;29:317–21.
37. Machii M, Inaba H, Nakae H, et al. Cardiac rupture by penetration of fractured sternum: a rare complication of cardiopulmonary resuscitation. *Resuscitation* 2000;43:151–3.
38. Spoormans I, Van Hoorenbeeck K, Balliu L, et al. Gastric perforation after cardiopulmonary resuscitation: review of the literature. *Resuscitation* 2010;81:272–80.
39. White L, Rogers J, Bloomingdale M, et al. Dispatcher-assisted cardiopulmonary resuscitation: risks for patients not in cardiac arrest. *Circulation* 2010;121:91–7.
40. Inaba H, Enami M, Taniguchi J, et al. One-year survey of telephone-assisted instruction (TAI) of chest compression only cardiopulmonary resuscitation (CC only-CPR) for victims presumed to be in cardiac arrest. *Resuscitation* 2010;81:S10.