
Journal Pre-proof



Improved Outcomes for Out-of-Hospital Cardiac Arrest Patients Treated by Emergency Life-Saving Technicians Compared with Basic Emergency Medical Technicians: A JCS-ReSS Study Report

Hiromichi Naito MD, PhD Tetsuya Yumoto MD, PhD Takashi Yorifuji MD, PhD Yoshio Tahara MD, PhD Naohiro Yonemoto MPH, PhD Hiroshi Nonogi MD, PhD Ken Nagao MD, PhD Takanori Ikeda MD, PhD Naoki Sato MD, PhD Hiroyuki Tsutsui MD, PhD

PII: S0300-9572(20)30186-6

DOI: https://doi.org/doi:10.1016/j.resuscitation.2020.05.007

Reference: RESUS 8525

To appear in: *Resuscitation*

Received Date: 18 November 2019

Revised Date: 18 April 2020

Accepted Date: 3 May 2020

Please cite this article as: Naito H, Yumoto T, Yorifuji T, Tahara Y, Yonemoto N, Nonogi H, Nagao K, Ikeda T, Sato N, Tsutsui H, Improved Outcomes for Out-of-Hospital Cardiac Arrest Patients Treated by Emergency Life-Saving Technicians Compared with Basic Emergency Medical Technicians: A JCS-ReSS Study Report, *Resuscitation* (2020), doi: https://doi.org/10.1016/j.resuscitation.2020.05.007

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2020 Published by Elsevier.

Journal Pre-proof

Title

Improved Outcomes for Out-of-Hospital Cardiac Arrest Patients Treated by Emergency Life-Saving Technicians Compared with Basic Emergency Medical Technicians: A JCS-ReSS Study Report

Authors:

Hiromichi Naito MD PhD[1,2], Tetsuya Yumoto MD PhD[1,2], Takashi Yorifuji MD PhD[1,3], Yoshio Tahara MD PhD[1,4], Naohiro Yonemoto MPH PhD[1,5], Hiroshi Nonogi MD PhD[1,6], Ken Nagao MD PhD[1,7], Takanori Ikeda MD PhD[1,8], Naoki Sato MD PhD[1,9], Hiroyuki Tsutsui MD PhD[1,10]

Author Affiliations:

1. JCS-ReSS Group

2. Okayama University Hospital, Advanced Emergency and Critical Care Medical Center

3. Department of Epidemiology, Okayama University Graduate School of Medicine, Dentistry and Pharmaceutical Sciences

4. National Cerebral and Cardiovascular Center, Department of Cardiovascular

Journal Pre-proof

Journal Pre-proof

Medicine

5. National Center of Neurology and Psychiatry

6. Shizuoka General Hospital, Intensive Care Center

7. Nihon University Hospital, Cardiovascular Center

8. Toho University Faculty of Medicine, Department of Cardiovascular Medicine

9. Kawaguchi Cardiovascular and Respiratory Hospital, Cardiovascular Medicine

10. Kyushu University Faculty of Medical Sciences, Department of Cardiovascular Medicine

Corresponding Author:

Hiromichi Naito MD PhD, Okayama University Hospital, Advanced Emergency and Critical Care Medical Center, 2-5-1 Shikata, Okayama, Japan 700-8558; e-mail address: naito.hiromichi@gmail.com; telephone number: +81-86-235-7427

Keywords:

Paramedic; Prehospital; Emergency Medical Services; Cardiopulmonary Resuscitation; Advanced Life Support

Journal Pre-proof

Journal Pre-proof

Word Count: 2,850 words, Number of References: 39, Number of Tables: 3, Number of Figures: 1

Abstract

Background: Emergency life-saving technicians (ELSTs) are specially trained prehospital medical providers believed to provide better care than basic emergency medical technicians (BEMTs). ELSTs are certified to perform techniques such as administration of advanced airways or adrenaline and are considered to have more knowledge; nevertheless, ELSTs' effectiveness over BEMTs regarding out-of-hospital cardiac arrest (OHCA) remains unclear. We investigated whether the presence of an ELST improves OHCA patient outcomes.

Methods: In a retrospective study of adult OHCA patients treated in Japan from 2011 to 2015, we compared two OHCA patient groups: patients transported with at least one ELST and patients transported by only BEMTs. The primary outcome measure was one-month favorable neurological outcomes, defined as Cerebral Performance Category ≤ 2. A multivariable logistic regression model was used to calculate odds ratios (ORs) and their confidence intervals (CIs) to evaluate the effect of ELSTs.

Results: Included were 552,337 OHCA patients, with 538,222 patients in the ELST group

Journal Pre-proof

and 14,115 in the BEMT group. The ELST group had a significantly higher odds of favorable neurological outcomes (2.5% vs. 2.1%, adjusted OR 1.39, 95% CI 1.17–1.66), one-month survival (4.9% vs. 4.1%, adjusted OR 1.37, 95% CI 1.22–1.54), and return of spontaneous circulation (8.1% vs. 5.1%, adjusted OR 1.90, 95% CI 1.72–2.11) compared with the BEMT group. However, ELSTs ' limited procedure range (adrenaline administration or advanced airway management) did not promote favorable neurological outcomes.

Conclusions: Compared with the BEMT group, transport by the ELST group was associated with better neurological outcomes in OHCA.

Introduction

Out-of-hospital cardiac arrest (OHCA) is a leading public health concern [1][2][3]; favorable neurological outcomes remain low. Improvement in the health care system [4], public education [5], and advanced care in the receiving hospital [6] can ameliorate survival and neurological outcomes in OHCA. Prehospital medical providers play important roles worldwide [7][8]. In Japan, the Fire and Disaster Management Agency (FDMA) employs two levels of public prehospital medical providers, emergency life-saving

Journal Pre-proof

technicians (ELSTs) and basic emergency medical technicians (BEMTs). ELSTs are certified and receive more prehospital emergency care education than BEMTs [9].

Japanese ambulances typically have three providers, including at least one ELST; however, approximately 3% of cases are transported without ELSTs (only BEMTs). Legally, only ELSTs may administer advanced airways using tools like supraglottic devices in Japan; specially-trained ELSTs can perform endotracheal intubation (ETI) [9]. Another important potentially life-saving skill only specially trained ELSTs may perform is adrenaline administration. Considering their skills and knowledge, treatment and transport by advanced prehospital medical providers seem to be associated with improved survival [10][11][12]; however, the survival or neurological benefit of advanced prehospital medical providers remains controversial globally [13][14][15]. In addition, ELST skill maintenance requires costly efforts and training [9][16].

ELSTs' effectiveness over BEMTs in OHCA treatment and transport has not been fully elucidated. We conducted a large-scale, observational, population-based cohort study on OHCA patients treated by BEMTs and ELSTs and investigated whether presence of ELSTs improved OHCA patient outcomes.

Journal Pre-proof

Journal Pre-proof

Methods

Emergency Medical System (EMS) and Providers

Japan is a 378,000 km^2 area with 127 million people. In 2015, there were 750 local fire departments with 1,709 fire stations and 3,145 dispatch centers [17]; the EMS is available 24 hours daily. Ambulances respond to scenes from the nearest fire stations. Each ambulance commonly has three emergency personnel, including at least one ELST nationally certified and trained to provide prehospital emergency care. Since April 1991, the number of ELSTs has been increasing. However, only BEMTs are present for some cases.

Two levels of Japanese prehospital medical providers (BEMTs and ELSTs) have been previously described [12]. BEMTs are accredited by local EMS after 250 hours of medical education, perform basic life support (BLS), including bag-valve-mask (BVM) ventilation, and are encouraged to apply automated defibrillators. ELSTs are accredited in two ways: ① They must work as a BEMT in prehospital care and transportation for approximately five years, complete a half year of ELST education and training, and pass a national examination, or ② After two years of ELST education at an EMT school/college, they must pass a national examination. ELSTs perform Advanced Life Support, applying semi-automated defibrillators. When ELSTs have difficulty performing BVM ventilation, they may secure the airway using supraglottic devices (laryngeal tube, i-gel, laryngeal mask,

Journal Pre-proof

109 combi-tube, etc.) for OHCA patients under direct telephone orders of a local medical
110 consultant (emergency physician). Additionally, ELSTs who complete the additional
111 training and perform 30 successful live intubations in the operating room are certified to
112 perform ETI. ETI criteria include OHCA patients with asphyxia due to foreign-body airway
113 obstruction without successful airway through a supraglottic device or judgement by a
114 medical consultant on direct order. Similarly, ELSTs with additional training may establish
115 intravenous (IV) lines (additional training, 10 successful IV placements, one adrenaline
116 administration to a cardiac arrest patient under supervision). If return of spontaneous
117 circulation (ROSC) doesn't occur after initial defibrillation, advanced ELSTs can administer
118 adrenaline to pulseless electrical activity, asystole, or refractory ventricular fibrillation
119 (VF)/ventricular tachycardia (VT) rhythms after giving shock. Adrenaline is administered
120 via prefilled syringes every three to five minutes until ROSC or hospital arrival.
121 Synchronized radio-controlled watches are used to record all procedures. ELSTs and
122 BEMTs are legally banned from terminating resuscitation in the field. They may not attempt
123 resuscitation in cases with definite deaths like incineration, decapitation, rigor mortis,
124 decomposition, or "do not attempt resuscitation" orders [18].
125
126 *Study Design and Data Collection*

Journal Pre-proof

The Okayama University Ethics Committee approved the study (1806-012) and waived the requirement for written informed consent. This nationwide, retrospective, population-based, observational study used the All-Japan Utstein Registry database for OHCA patients run by the FDMA [5].

OHCA patient data were prospectively collected based on Utstein-style guidelines [19][20] by the local EMS and subsequently integrated into the national registry. Cardiac arrest was defined as stoppage of mechanical cardiac activity substantiated by lack of circulation signs. Cardiac arrest was assumed of cardiac origin unless caused by respiratory or cerebrovascular diseases; external causes, including hanging, trauma, drowning, asphyxia, and drug overdose; malignant tumors; or other non-cardiac causes. Diagnoses of cardiac or non-cardiac causes were clinically decided by in charge physicians in collaboration with EMS providers. Variables included gender, age, witness status, bystander-initiated cardiopulmonary resuscitation (CPR), initial electrocardiogram rhythm, resuscitation time-course, cardiac arrest region, cardiac or non-cardiac origin, ELST status, use of advanced airway, establishment of IV line, adrenaline administration, hospital admission, ROSC, one-month neurological status, and one-month survival. ROSC was defined as recovery of any spontaneous palpable pulse with any time duration confirmed with cardiac rhythm monitoring [18]. Cerebral Performance Category (CPC) scale was used to evaluate

Journal Pre-proof

neurological outcomes [19][20].

Patient Selection and Endpoint

All 18-year-old or older patients with OHCA of cardiac and non-cardiac causes transported by the FDMA from January 1, 2011 through December 31, 2015 were included. Cases where resuscitation was not attempted and cases involving patients under 18, the presence of doctor during transport, and treatment by doctor before transport were excluded. We compared two OHCA patient groups, the ELST group (patients transported by EMS, including at least one ELST) and the BEMT group (patients transported only by BEMTs). The primary outcome measure was one-month favorable neurological outcome defined as CPC 1 or 2. Secondary outcome measures included one-month survival and ROSC. Additionally, advanced airway efficacy and adrenaline use were evaluated in the ELST group.

Data Analysis

Continuous variables were described using median with interquartile ranges (IQR). Categorical variables were described using percentages. The Mann-Whitney U-test was used to compare EMS contact to initial defibrillation times between the ELST and BEMT groups.

Journal Pre-proof

For the primary outcome, univariable logistic regression was used to determine association between favorable neurological outcome and groups. Multivariable logistic regression was performed to adjust covariates (year of cardiac arrest, age, gender, initial recorded cardiac rhythm, estimated cardiac or non-cardiac origin, bystander CPR, dispatcher instruction for CPR, time from EMS call to hospital arrival, cardiac arrest region). Logistic regression analysis results were expressed using odds ratios (ORs) and 95% confidence intervals (CIs). For the secondary outcome, including effect of ELST on one-month survival and ROSC, univariable and multivariable logistic regression were used. Effectiveness of advanced treatments (airway management, adrenaline administration) only allowed by ELSTs is still conflicting; to identify advantages of ELSTs' technical procedure skills, univariable and multivariable logistic regression were also used in the ELST group to evaluate whether these advanced treatments contribute to the outcomes. Same variables were used for adjustment in multivariable logistic regression. Statistical analysis was performed using Stata version 15 (StataCorp LP, College Station, TX). *P*-values below 0.05 were considered statistically significant.

Results

Journal Pre-proof

Figure 1 shows an OHCA patient flow chart. The registry documented 629,471 OHCA patients from 2011 to 2015 in Japan. Patients meeting exclusion criteria (16,210 resuscitations not attempted; 7,687 under 18 years old; 18,951 transported with doctors; 34,286 treated by doctors) were excluded. Finally, 552,337 were included, 538,222 in the ELST group and 14,115 in the BEMT group.

Table 1 shows baseline ELST and BEMT group characteristics, prehospital procedure characteristics, and outcomes. The proportion of males (ELST group vs. BEMT group: 56.6 vs. 56.9%), age (median [IQRs]: 79 [67-86] vs. 79 [68-86] years), witnessed collapse (40.3 vs. 40.0%), and time from EMS call to hospital arrival (32 [26-39] vs. 31 [26-39] min) did not differ between the groups. Proportions of initial rhythm VF/VT (7.1 vs. 6.6%) and dispatcher instruction for CPR (53.0 vs. 50.0%) were higher in the ELST group, while estimated cardiac origin (59.5 vs. 62.2%) and bystander CPR (50.9 vs. 56.0%) were lower. The proportion of patients transported by BEMTs tended to decrease as the years progressed. The proportion of BEMTs tended to be higher in low population density regions. Time from EMS arrival to initial defibrillation did not differ between the ELST and BEMT groups (2 [1-3] vs. 2 [1-4] min, $p = 0.25$). Advanced airway management was conducted in 42.2% of ELST group patients (34.0% supraglottic airways; 7.3% ETI); adrenaline administration was conducted in 16.0% of the ELST group. Overall, there was a 2.5%

Journal Pre-proof

199 incidence of favorable neurological outcomes, 4.9% of one-month survival, and 8.0% of
200 ROSC.

201 In the univariable logistic regression (**Table 2**), all outcomes, including favorable
202 neurological outcomes (2.5 vs. 2.1%, OR 1.19, 95% CI 1.06-1.34), one-month survival (4.9
203 vs. 4.1%, OR 1.21, 95% CI 1.12-1.32), and ROSC (8.1 vs. 5.1%, OR 1.65, 95% CI
204 1.53-1.78), were higher in the ELST group. Multivariable logistic regression was conducted
205 to adjust for potential confounders (**Table 2**). ELST presence was significantly associated
206 with favorable neurological outcomes (adjusted OR 1.39, 95%, CI 1.17-1.66), one-month
207 survival (adjusted OR 1.37, 95% CI 1.22-1.54), and ROSC (adjusted OR 1.90, 95% CI
208 1.73-2.11).

209 We explored whether technical procedures (advanced airway management;
210 adrenaline administration) limited to ELSTs promoted outcomes with univariable and
211 multivariable logistic regression (**Table 3**). Use of advanced airways did not promote
212 favorable neurological outcomes (adjusted OR 0.34, 95% CI 0.32-0.36) or one-month
213 survival (adjusted OR 0.72, 95% CI 0.70-0.75). Administration of adrenaline did not
214 promote favorable neurological outcomes (adjusted OR 0.33, 95% CI 0.31-0.35) or
215 one-month survival (adjusted OR 0.84, 95% CI 0.80-0.87). However, both advanced airways
216 (adjusted OR 1.08, 95% CI 1.05-1.11) and adrenaline (adjusted OR 3.73, 95% CI

Journal Pre-proof

217 3.64-3.83) increased ROSC.

218

219

220 Discussion

221 We examined the efficacy of ELST presence during treatment and transport for
222 OHCA patient outcomes using a large national registry in Japan and found that patients
223 treated by ELSTs had better neurological outcomes (adjusted OR of 1.39) than patients
224 treated only by BEMTs. Adjusted one-month survival and ROSC were also better in the
225 ELST group compared with the BEMT group. However, intervention skills such as advanced
226 airway use or adrenaline administration did not fully explain ELSTs' contribution to
227 favorable neurological outcomes. ELSTs' greater knowledge of pathophysiology and
228 appropriate transport to receiving hospitals to treat OHCA patients may be their advantages.

229 Prehospital medical providers contribute to treating and transporting patients,
230 including OHCA patients, all over the world [21][22]; their techniques and knowledge may
231 vary widely by region. Since prehospital medical providers (in Japan, BEMTs or ELSTs)
232 commonly play important roles in advising/helping first-contacted public BLS providers or
233 treating and transporting OHCA patients to hospitals, their skills or knowledge can be
234 decisive factors in OHCA patient outcomes. OHCA experience (case volume) of prehospital

Journal Pre-proof

medical provider has been reported to predict ROSC [11]. Similarly, association between OHCA exposure of prehospital medical provider and patient survival has been reported. OHCA exposure during the preceding three years had a positive effect on patient survival; moreover, outcomes improved when it was less than six months since prehospital medical providers last treated OHCA patients [23]. In Osaka, Japan, Kajino et al. compared the efficacy of supraglottic devices vs. ETI; in their sub-analysis, the presence of an ETI-certified ELST was significantly associated with favorable neurological outcomes (adjusted OR of 1.86, 95% CI 1.04 to 3.34), similar to our results [24]. Our study did not address the number of EMTs or ELSTs; however, in the Osaka study, the presence of three ELSTs was associated with improved one-month survival with favorable neurological outcomes in OHCA compared with presence of one ELST [12]. In addition, an over 50% ratio of on-scene advanced EMTs was associated with improved survival in OHCA patients with witnessed, non-shockable rhythm in Taipei [10]. In their study, the presence of four EMTs with an advanced EMT ratio of over 50% was associated with the best outcomes. However, number of EMTs alone does not seem to improve outcomes. More EMTs in ambulances did not affect rates of ROSC, survival, or neurological favorable outcomes of OHCA patients in Tokyo [25]. Similarly, Eschmann et al. reported that in their two-tiered system, the presence of three or advanced EMTs at the scene was not associated with

Journal Pre-proof

253 improved survival compared with one basic EMT and two advanced EMTs [13]. At least one
254 well-trained and experienced prehospital medical provider such as an ELST seems to be
255 needed on scene for decision-making to improve outcomes.

256 Whether advanced airways contribute to OHCA outcomes remains debatable;
257 however, these skills are essential in some cases with a limited number of crews. However,
258 the use of advanced airways did not promote ELST favorable neurological outcomes,
259 implying that ELSTs' advantages extend beyond the technique. The results provided
260 additional evidence of the limited impact of advanced techniques like advanced airways or
261 adrenaline administration on neurological outcomes. Our results were consistent with those
262 from previous studies showing no improved outcomes with advanced airway device use in
263 Japan [14]. However, these skills may still potentially improve outcomes [26][27][28].
264 Proper indications for advanced airways are needed; moreover, outcomes may vary
265 according to the prehospital medical provider's proficiency in performing the procedure.
266 Adrenaline administration was conducted in fewer patients (16%) compared to the US [29]
267 or UK [30]. Regional protocols for administering adrenaline have variations; some districts
268 do not recommend adrenaline administration to unwitnessed asystole [31]. Moreover, the
269 need for direct phone orders from a medical consultant physician to establish an intravenous
270 access and a subsequent phone call to administer adrenaline may have resulted in decreased

Journal Pre-proof

271 incidence or delayed adrenaline administration. Adrenaline administration increased ROSC;
272 however, our data did not show efficacy of adrenaline administration for favorable
273 neurological outcomes. Past studies have conflicting results: in one trial, prehospital
274 epinephrine use was associated with higher chance of ROSC, but lower chance of survival
275 and good neurological outcomes [15]. Contrarily, after accounting for time-dependent
276 patient imbalance, Nakahara et al. reported higher proportions of overall and
277 neurologically-intact survival among those with non-shockable rhythms in adrenaline
278 administration [32]. In the recent randomized controlled trial, adrenaline use resulted in
279 significantly higher survival, but a lower rate of good neurological outcome in those who
280 survived [30]. Adrenaline timing [33][34] or dosing [35][36][37] and appropriate patients
281 [38] should be further evaluated in prehospital settings.

282 BLS quality is crucial in OHCA outcomes; more educated prehospital medical
283 providers may have performed better quality BLS. Other possible explanations for better
284 neurological survival with ELSTs is that they performed better quality defibrillation. Shorter
285 interruptions in chest compressions before and after defibrillatory shock is an independent
286 predictor of survival from shockable OHCA [39]. Although EMS contact time to initial
287 defibrillation time did not differ between groups, ELSTs with more experience/education
288 and semi-automated defibrillator use may have facilitated faster defibrillation.

Journal Pre-proof

289 Results of multivariable logistic regression showed evidence supporting ELST
290 presence. ELSTs' knowledge of factors other than skills/procedures like defibrillation or
291 administration of advanced airways or adrenaline may have had more impact on outcomes.
292 Receiving hospital characteristics may be associated with OHCA outcomes. Transport to
293 Japanese nationally-certified critical care medical centers assumed to be high-volume
294 independently predicted good neurological outcomes in OHCA without field ROSC [6].
295 Receiving hospital choice may have been better with more educated or experienced
296 prehospital medical providers like ELSTs. In addition, ELSTs may have provided hospitals
297 more precise meaningful information from the field to prepare for definitive treatment.
298 Future research should determine these factors affecting the association between advanced
299 prehospital medical providers' ability and improved outcomes.
300
301
302 Limitations
303 Our study has several limitations. First, we couldn't obtain information on several
304 factors associated with outcomes in BEMT or ELST transport. We didn't address ELST
305 education level (whether or not they are certified in ETI/adrenaline administration),
306 number of BEMTs and ELSTs, or BEMTs/ELST experience. Second, the number of

Journal Pre-proof

BEMTs was much smaller than the number of ELSTs. Moreover, there was some difference in characteristics between groups. For instance, ELST group showed higher rates of patients with VF/VT and lower rates of patients with estimated cardiac origin. This may attribute to regional variation for BEMT-only transport: (i.e., more frequent BEMT transport in rural areas). We tried to adjust the difference between the two groups using known factors potentially related to outcomes using factors obtained in the database. Third, other unmeasured confounding factors may have influenced outcomes, i.e. patient comorbidity, location of arrest, cardiac arrest etiology, receiving hospital information, hospital post-cardiac arrest management, CPR quality information. Finally, like with all epidemiological studies, data validity, integrity, and ascertainment bias were possible limitations. However, large sample size, population-based design to cover all OHCA in Japan, and uniform data collection based on Utstein-style guidelines for reporting cardiac arrest were intended to lessen these potential biases.

Conclusions

Compared with the BEMT group, treatment and transport by the ELST group was associated with improved favorable neurological outcomes for OHCA. EMS response to

Journal Pre-proof

OHCA should include at least one advanced level prehospital provider.

Conflict of interest

Conflicts of interest: none.

Acknowledgements

Thank you to all EMS providers and concerned physicians.

Figure Legends

Figure 1

Flow diagram of patients analyzed. The ELST group comprised patients transported by emergency medical service, including at least one ELST. The BEMT group comprised patients transported only by BEMTs. BEMT: basic emergency medical technician; CPR: cardiopulmonary resuscitation; ELST: emergency life-saving technician; OHCA: out-of-hospital cardiac arrest.

Journal Pre-proof

Journal Pre-proof

References

[1] Kitamura T, Iwami T, Kawamura T, Nitta M, Nagao K, Nonogi H, et al. Nationwide improvements in survival from out-of-hospital cardiac arrest in Japan. Circulation 2012;126:2834–43.

[2] Gräsner JT, Lefering R, Koster RW, Masterson S, Böttiger BW, Herlitz J, et al. EuReCa ONE—27 Nations, ONE Europe, ONE Registry: A prospective one month analysis of out-of-hospital cardiac arrest outcomes in 27 countries in Europe. Resuscitation 2016;105:188–95.

[3] Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, Cushman M, et al. Heart Disease and Stroke Statistics—2016 Update. Circulation 2016;133:e38-360.

[4] Japanese Circulation Society Resuscitation Science Study Group. Chest-compression-only bystander cardiopulmonary resuscitation in the 30:2 compression-to-ventilation ratio era. Nationwide observational study. Circ J 2013;77:2742–50.

[5] Kitamura T, Kiyohara K, Sakai T, Matsuyama T, Hatakeyama T, Shimamoto T, et al. Public-Access Defibrillation and Out-of-Hospital Cardiac Arrest in Japan. N Engl J Med 2016;375:1649–59.

[6] Kajino K, Iwami T, Daya M, Nishiuchi T, Hayashi Y, Kitamura T, et al. Impact of

Journal Pre-proof

transport to critical care medical centers on outcomes after out-of-hospital cardiac arrest. Resuscitation 2010;81:549–54.

[7] Spaite DW, Stiell IG, Bobrow BJ, de Boer M, Maloney J, Denninghoff K, et al. Effect of transport interval on out-of-hospital cardiac arrest survival in the OPALS study: implications for triaging patients to specialized cardiac arrest centers. Ann Emerg Med 2009;54:248–55.

[8] Drennan IR, Verbeek PR. The Role of EMS in Regionalized Systems of Care. CJEM 2015;17:468–74.

[9] Takahashi H, Sagisaka R, Natsume Y, Tanaka S, Takyu H, Tanaka H. Does dispatcher-assisted CPR generate the same outcomes as spontaneously delivered bystander CPR in Japan? Am J Emerg Med 2018;36:384–91.

[10] Sun J, Chiang W, Hsieh M, Huang EP, Yang W-S, Chien Y-C, et al. The effect of the number and level of emergency medical technicians on patient outcomes following out of hospital cardiac arrest in Taipei. Resuscitation 2018;122:48–53.

[11] Tuttle J, Hubble M. Paramedic Out-of-hospital Cardiac Arrest Case Volume Is a Predictor of Return of Spontaneous Circulation. West J Emerg Med 2018;19:654–9.

[12] Kajino K, Kitamura T, Iwami T, Daya M, Ong MEH, Nishiyama C, et al. Impact of the number of on-scene emergency life-saving technicians and outcomes from out-of-hospital

Journal Pre-proof

cardiac arrest in Osaka City. Resuscitation 2014;85:59–64.

[13] Eschmann NM, Pirrallo RG, Aufderheide TP, Lerner EB. The association between emergency medical services staffing patterns and out-of-hospital cardiac arrest survival. Prehosp Emerg Care 2010;14:71–7.

[14] Hasegawa K, Hiraide A, Chang Y, Brown DFM. Association of prehospital advanced airway management with neurologic outcome and survival in patients with out-of-hospital cardiac arrest. JAMA 2013;309:257–66.

[15] Hagihara A. Prehospital Epinephrine Use and Survival among Patients with Out-of-Hospital Cardiac Arrest. JAMA - J Am Med Assoc 2013;43:205–6.

[16] Wang HE, Mann NC, Mears G, Jacobson K, Yealy DM. Out-of-hospital airway management in the United States. Resuscitation 2011;82:378–85.

[17] Ambulance Service Planning Office of Fire and Disaster Management Agency of Japan. Statistics of fire department and fire fighters, 2017 (in Japanese). Available at: https://www.fdma.go.jp/publication/hakusho/h29/data/2180.html. Accessed November 1, 2019.

[18] Nagao K, Nonogi H, Yonemoto N, Gaieski DF, Ito N, Takayama M, et al. Duration of prehospital resuscitation efforts after out-of-hospital cardiac arrest. Circulation 2016;133:1386–96.

Journal Pre-proof

Journal Pre-proof

[19] Cummins RO, Chamberlain DA, Abramson NS, Allen M, Baskett PJ, Becker L, et al. Recommended guidelines for uniform reporting of data from out-of-hospital cardiac arrest: the Utstein Style. A statement for health professionals from a task force of the American Heart Association, the European Resuscitation Council, the Heart and Stroke. Circulation 1991;84:960–75.

[20] Jacobs I, Nadkarni V, Bahr J, Berg RA, Billi JE, Bossaert L, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: update and simplification of the Utstein templates for resuscitation registries. A statement for healthcare professionals from a task force of the international liaison committee on resusci. Resuscitation 2004;63:233–49.

[21] Martin AC, O'Meara P. Perspectives from the frontline of two North American community paramedicine programs: an observational, ethnographic study. Rural Remote Health 2019;19:4888.

[22] O'Meara P, Duthie S. Paramedicine in Australia and New Zealand: A comparative overview. Aust J Rural Health 2018;26:363–8.

[23] Dyson K, Bray JE, Smith K, Bernard S, Straney L, Finn J. Paramedic Exposure to Out-of-Hospital Cardiac Arrest Resuscitation Is Associated With Patient Survival. Circ Cardiovasc Qual Outcomes 2016;9:154–60.

[24] Kajino K, Iwami T, Kitamura T, Daya M, Ong MEH, Nishiuchi T, et al. Comparison of

Journal Pre-proof

supraglottic airway versus endotracheal intubation for the pre-hospital treatment of out-of-hospital cardiac arrest. Crit Care 2011;15:R236.

[25] Hagiwara S, Oshima K, Aoki M, Miyazaki D, Sakurai A. American Journal of Emergency Medicine Does the number of emergency medical technicians affect the neurological outcome of patients with out-of-hospital cardiac arrest ? Am J Emerg Med 2017;35:391–6.

[26] Jabre P, Penaloza A, Pinero D, Duchateau F-X, Borron SW, Javaudin F, et al. Effect of Bag-Mask Ventilation vs Endotracheal Intubation During Cardiopulmonary Resuscitation on Neurological Outcome After Out-of-Hospital Cardiorespiratory Arrest: A Randomized Clinical Trial. JAMA 2018;319:779–87.

[27] Izawa J, Komukai S, Gibo K, Okubo M, Kiyohara K, Nishiyama C, et al. Pre-hospital advanced airway management for adults with out-of-hospital cardiac arrest: nationwide cohort study. BMJ 2019;364:l430.

[28] Izawa J, Iwami T, Gibo K, Okubo M, Kajino K, Kiyohara K, et al. Timing of advanced airway management by emergency medical services personnel following out-of-hospital cardiac arrest: A population-based cohort study. Resuscitation 2018;128:16–23.

[29] Hansen M, Schmicker RH, Newgard CD, Grunau B, Scheuermeyer F, Cheskes S, et al. Time to Epinephrine Administration and Survival From Nonshockable Out-of-Hospital

Journal Pre-proof

433 Cardiac Arrest Among Children and Adults. Circulation 2018;137:2032–40.

434 [30] Perkins G, Ji C, Deakin C, Quinn T, et al. A Randomized Trial of Epinephrine in

435 Out-of-Hospital Cardiac Arrest. N Engl J Med 2018:1–11.

436 [31] Tomio J, Nakahara S, Takahashi H, Ichikawa M, Nishida M, Morimura N, et al.

437 Effectiveness of Prehospital Epinephrine Administration in Improving Long-term

438 Outcomes of Witnessed Out-of-hospital Cardiac Arrest Patients with Initial

439 Non-shockable Rhythms. Prehosp Emerg Care 2017;21:432–41.

440 [32] Nakahara S, Tomio J, Takahashi H, Ichikawa M, Nishida M, Morimura N, et al.

441 Evaluation of pre-hospital administration of adrenaline (epinephrine) by emergency

442 medical services for patients with out of hospital cardiac arrest in Japan: Controlled

443 propensity matched retrospective cohort study. BMJ 2013;347:1–10.

444 [33] Tanaka H, Takyu H, Sagisaka R, Ueta H, Shirakawa T, Kinoshi T, et al. Favorable

445 neurological outcomes by early epinephrine administration within 19 minutes after EMS

446 call for out-of-hospital cardiac arrest patients. Am J Emerg Med 2016;34:2284–90.

447 [34] Koscik C, Pinawin A, McGovern H, Allen D, Media DE, Ferguson T, et al. Rapid

448 epinephrine administration improves early outcomes in out-of-hospital cardiac arrest.

449 Resuscitation 2013;84:915–20.

450 [35] Fisk CA, Olsufka M, Yin L, McCoy AM, Latimer AJ, Maynard C, et al. Lower-dose

Journal Pre-proof

epinephrine administration and out-of-hospital cardiac arrest outcomes. Resuscitation 2018;124:43–8.

[36] Callaway CW, Donnino MW. Testing Epinephrine for Out-of-Hospital Cardiac Arrest. N Engl J Med 2018:1–2.

[37] Wang CH, Huang CH, Chang WT, Tsai MS, Yu PH, Wu YW, et al. The influences of adrenaline dosing frequency and dosage on outcomes of adult in-hospital cardiac arrest: A retrospective cohort study. Resuscitation 2016;103:125–30.

[38] Goto Y, Maeda T, Goto YN. Effects of prehospital epinephrine during out-of-hospital cardiac arrest with initial non-shockable rhythm: An observational cohort study. Crit Care 2013;17:1–11.

[39] Cheskes S, Schmicker RH, Christenson J, Salcido DD, Rea T, Powell J, et al. Perishock pause: an independent predictor of survival from out-of-hospital shockable cardiac arrest. Circulation 2011;124:58–66.

Journal Pre-proof

Journal Pre-proof

Table 1. Patient Characteristics and Outcomes of ELST and BEMT Groups

	ELST	BEMT	All
Patient characteristics			
Male sex			312,615/552,337 (56.6%)
	304,590/538,222 (56.6%)	8,025/14,115 (56.9%)	
Age – median [IQR]			79 [67-86]
	79 [67-86]	79 [68-86]	
Initial rhythm VF/VT			37,779/531,586 (7.1%)
	36,879/517,989 (7.1%)	900/13,597 (6.6%)	
	PEA/Asystole		493,807/531,586 (92.9%)
	481,110/517,989 (92.9%)	12,697/13,597 (93.4%)	
Estimated cardiac origin			329,021/552,337 (59.6%)
	320,239/538,222 (59.5%)	8,782/14,115 (62.2%)	
Witnessed collapse			222,329/553,337 (40.3%)
	216,685/538,222 (40.3%)	5,644/14,115 (40.0%)	
Bystander CPR			246,326/482,684 (51.0%)
	240,091/471,611 (50.9%)	6,235/11,073 (56.3%)	
Dispatcher instruction for CPR			288,006/544,419 (52.9%)
	281,129/530,670 (53.0%)	6,877/13,749 (50.0%)	
Time from EMS call to hospital arrival – median [IQR]			32 [26-39] min
	32 [26-39] min	31 [26-39] min	
Year of cardiac arrest			
	2011		116,911
	113,687/116,911 (97.2%)	3,224/116,911 (2.8%)	
	2012		112,923
	110,244/112,923 (97.6%)	2,679/112,923 (2.4%)	
	2013		106,714
	104,267/106,714 (97.7%)	2,447/106,714 (2.3%)	
	2014		108,741
	105,104/108,741 (96.7%)	3,637/108,741 (3.3%)	

Journal Pre-proof

Journal Pre-proof

	2015		107,048
	104,920/107,048 (98.0%)	2,128/107,048 (2.0%)	
Region			
	Hokkaido		22,056/552,337 (4.0%)
	21,532/538,222 (4.0%)	524/14,115 (3.7%)	
	Tohoku		47,322/552,337 (8.6%)
	44,747/538,222 (8.3%)	2,575/14,115 (18.2%)	
	Kanto		186,808/552,337 (33.8%)
	184,580/538,222 (34.3%)	2,228/14,115 (15.8%)	
	Chubu		102,970/552,337 (18.6%)
	101,195/538,222 (18.8%)	1,775/14,115 (12.6%)	
	Kinki		88,715/552,337 (16.1%)
	85,054/538,222 (15.8%)	3,661/14,115 (25.9%)	
	Chugoku		32,481/552,337 (5.9%)
	31,683/538,222 (5.9%)	798/14,115 (5.7%)	
	Shikoku		18,518/552,337 (3.4%)
	17,957/538,222 (3.3%)	561/14,115 (4.0%)	
	Kyushu		53,467/552,337 (9.7%)
	51,474/538,222 (9.6%)	1,993/14,115 (14.1%)	
Interventions			
	Time from EMS arrival to initial defibrillation (VF/VT) – median [IQR]		2 [1-3] min
2 [1-3] min	2 [1-4] min		
	Use of advanced airways		N/A
	220,135/522,001 (42.2%)	N/A	
	Supraglottic airways		N/A
	183,098/538,222 (34.0%)	N/A	
	Endotracheal intubation		N/A
	39,339/538,222 (7.3%)	N/A	
	Use of intravenous line and adrenaline		
	Intravenous line establishment		N/A
	172,371/538,998 (32.0%)	N/A	
	Administration of adrenaline		N/A
	85,432/535,420 (16.0%)	N/A	
Outcomes			

Journal Pre-proof

One-month favorable neurological outcome (overall)			13,536/552,334 (2.5%)
	13,244/538,219 (2.5%)	292/14,115 (2.1%)	
	Initial rhythm VF/VT		6,877/37,779 (18.2%)
	6,726/36,879 (18.2%)	151/900 (16.8%)	
	Initial rhythm PEA/Asystole		3,834/493,804 (0.78%)
	3,746/481,107 (0.78%)	88/12,697 (0.69%)	
One-month survival (overall)			26,829/552,337 (4.9%)
	26,257/538,222 (4.9%)	572/14,115 (4.1%)	
	Initial rhythm VF/VT		10,036/37,779 (26.6%)
	9,828/36,879 (26.7%)	208/900 (23.1%)	
	Initial rhythm PEA/Asystole		12,643/493,807 (2.6%)
	12,360/481,110 (2.6%)	283/12,697 (2.2%)	
ROSC			44,340/552,337 (8.0%)
	43,625/538,222 (8.1%)	715/14,115 (5.1%)	

Patient characteristics, interventions, and outcomes are shown. All patient characteristics variables were used to adjust for the outcomes in the multivariable logistic regression. BEMT: basic emergency technician, CPR: cardiopulmonary resuscitation, ELST: emergency life-saving technician, EMS: emergency medical service, IQR: interquartile range, N/A: not applicable, PEA: pulseless electrical activity, ROSC: return of spontaneous circulation, VF/VT: ventricular fibrillation/ventricular tachycardia

Journal Pre-proof

Journal Pre-proof

Table 2. Univariable and Multivariable Logistic Regression Comparing Favorable Neurological Outcomes, One-Month Survival, and ROSC

	Crude OR (95% CI)	Adjusted OR (95% CI)
One-month favorable neurological outcome		
Overall		
BEMT	1 (ref)	1 (ref)
ELST	1.19 (1.06-1.34)	1.39 (1.17-1.66)
VF/VT		
BEMT	1 (ref)	1 (ref)
ELST	1.11 (0.93-1.32)	1.41 (1.13-1.77)
PEA/Asystole		
BEMT	1 (ref)	1 (ref)
ELST	1.12 (0.91-1.39)	1.31 (1.01-1.73)
One-month survival		
BEMT	1 (ref)	1 (ref)
ELST	1.2 (1.12-1.32)	1.37 (1.22-1.54)
ROSC		
BEMT	1 (ref)	1 (ref)
ELST	1.65 (1.53-1.78)	1.90 (1.72-2.11)

Variables, including gender, age, initial shockable rhythm, estimated cardiac origin, witnessed collapse, bystander CPR, dispatcher instruction for CPR, time from EMS call to hospital arrival, cardiac arrest year, and region, were used to adjust for the outcomes in the multivariable logistic regression. CI: confidence interval, CPR: cardiopulmonary resuscitation, BEMT: basic emergency medical technician, ELST: emergency life-saving technician, EMS: emergency medical service, OR: odds ratio, PEA: pulseless electrical activity, ROSC: return of spontaneous circulation, VF/VT: ventricular fibrillation/ventricular tachycardia

Journal Pre-proof


Journal Pre-proof

Table 3. Univariable and Multivariable Logistic Regression Comparing the Presence or Absence of Technical Intervention on Favorable Neurological Outcomes, One-Month Survival, and ROSC in the ELST group

	Crude OR (95% CI)	Adjusted OR (95% CI)
Favorable neurological outcome		
Advanced airways	0.30 (0.29-0.32)	0.34 (0.32-0.36)
Adrenaline	0.50 (0.47-0.53)	0.33 (0.31-0.35)
One-month survival		
Advanced airways	0.60 (0.59-0.62)	0.72 (0.70-0.75)
Adrenaline	1.02(0.99-1.06)	0.84 (0.80-0.87)
ROSC		
Advanced airways	0.97 (0.95-0.99)	1.08 (1.05-1.11)
Adrenaline	3.89 (3.81-3.97)	3.73 (3.64-3.83)

Variables, including gender, age, initial shockable rhythm, estimated cardiac origin, witnessed collapse, bystander CPR, dispatcher instruction for CPR, time from EMS call to hospital arrival, cardiac arrest year, and region, were used to adjust for the outcomes in the multivariable logistic regression. CI: confidence interval, CPR: cardiopulmonary resuscitation, ELST: emergency life-saving technician, EMS: emergency medical service system, OR: odds ratio, ROSC: return of spontaneous circulation

Journal Pre-proof


OHCA occurred from
2011 to 2015
n = 613,261
Resuscitation not
attempted
n = 29,238
Patients under 18
n = 7,687
Transported with doctor
on board
n = 18,951
Treatment by doctor
n = 34,286
Patients included in
the analysis
n = 552,337
ELST group
n = 538,222
BEMT group
n = 14,115