

# Socioeconomic Disparities in the Use of Cardioprotective Medications Among Patients With Peripheral Artery Disease

An Analysis of the American College of Cardiology's NCDR PINNACLE Registry

Sumeet Subherwal, MD, MBA,\* Manesh R. Patel, MD,\* Fengming Tang, MS,†  
Kim G. Smolderen, PhD,†‡ W. Schuyler Jones, MD,\* Thomas T. Tsai, MD, MSc,§  
Henry H. Ting, MD, MBA,|| Deepak L. Bhatt, MD, MPH,¶ John A. Spertus, MD, MPH,†  
Paul S. Chan, MD, MSc†

*Durham, North Carolina; Kansas City, Missouri; Aurora, Colorado; Rochester, Minnesota;  
Boston, Massachusetts; and Tilburg, the Netherlands*

<b>Objectives</b>	The aim of this paper was to examine disparities in the use of cardioprotective medications in the treatment of peripheral artery disease (PAD) by socioeconomic status (SES).
<b>Background</b>	PAD is associated with increased cardiovascular risk and is more prevalent among those of lower SES. However, the use of guideline-recommended secondary preventive measures for the treatment of PAD across diverse income subgroups and the influence of practice site on potential treatment disparities by SES are unknown.
<b>Methods</b>	Within the National Cardiovascular Disease Registry (NCDR) PINNACLE Registry, 62,690 patients with PAD were categorized into quintiles of SES, as defined by the median income of each patient's zip code. The association between SES and secondary preventive treatment with antiplatelet and statin medications was evaluated using sequential hierarchical modified Poisson models, adjusting first for practice site and then for clinical variables.
<b>Results</b>	Compared with the highest SES quintile (median income: >\$60,868), PAD patients in the lowest SES quintile (median income: <\$34,486) were treated less often with statins (72.5% vs. 85.8%; RR: 0.84; 95% CI: 0.83 to 0.86; $p < 0.001$ ) and antiplatelet therapy (79.0% vs. 84.6%; RR: 0.93; 95% CI: 0.91 to 0.94; $p < 0.001$ ). These differences were markedly attenuated after controlling for practice site variation: statins (adjusted RR: 0.97; 95% CI: 0.95 to 0.99; $p = 0.003$ ) and antiplatelet therapy (adjusted RR: 0.98; 95% CI: 0.97 to 1.00; $p = 0.012$ ). Additional adjustment for patients' clinical characteristics had minimal impact, with slight further attenuation with statins (adjusted RR: 1.00; 95% CI: 0.99 to 1.01; $p = 0.772$ ) and antiplatelet therapy (adjusted RR: 1.00; 95% CI: 0.99 to 1.01; $p = 0.878$ ).
<b>Conclusions</b>	Among PAD patients, the practice site at which patients received care largely explained the observed SES differences in treatment with guideline-recommended secondary preventive medications. Future efforts to reduce treatment disparities in these vulnerable populations should target systems improvement at practices serving high proportions of patients with low SES. (J Am Coll Cardiol 2013;62:51-7) © 2013 by the American College of Cardiology Foundation

The Institute of Medicine has challenged the U.S. health care system to minimize disparities in treatment and to provide equitable access to evidence-based therapies to all

patients (1). Although there have been numerous studies investigating disparities in the care of cardiac patients (2-4), research on disparities in peripheral arterial disease (PAD)

From the \*Duke University, Durham, North Carolina; †Mid America Heart Institute, Saint Luke's Hospital, Kansas City, Missouri; ‡Tilburg University, Tilburg, the Netherlands; §University of Colorado, Aurora, Colorado; ||Mayo Clinic, Rochester, Minnesota; and the ¶Veterans Affairs Boston Healthcare System, Brigham and Women's Hospital, and Harvard Medical School, Boston, Massachusetts. Dr. Subherwal's work on this paper was supported by a National Institutes of Health Training Grant (no. T32 HL 69749-9). Dr. Smolderen's work on this paper was

supported by the Outcomes Research post-doctoral fellowship awarded by the American Heart Association Pharmaceutical Roundtable and David and Stevie Spina (grant no. AHA: 0875149N), by the Netherlands Organization for Scientific Research (VENI grant no. 916.11.179), and by an unrestricted grant from W.L. Gore & Associates, Inc. Dr. Chan's work on this paper was supported by a Career Development Grant Award (no. K23HL102224) from the NHLBI. This research was also supported by the American College of Cardiology Foundation's National

## Abbreviations and Acronyms

**PAD** = peripheral artery disease

**SES** = socioeconomic status

have been limited (5–9), even though PAD affects over 7 million Americans (10) and disproportionately affects those of lower socioeconomic status (SES) (11,12). Equitable access to inexpensive, guideline-recommended secondary preventive therapies has the potential to improve cardiovascular outcomes in this vulnerable population (13–17), but whether treatment rates differ across income groups remains unknown. Given that PAD is associated with increased cardiovascular risk and mortality (18–22), illuminating current practice patterns by SES for evidence-based secondary preventive strategies is particularly important in defining opportunities to better improve care.

Accordingly, we examined PAD treatment rates by SES within the American College of Cardiology's (ACC) National Cardiovascular Disease Registry (NCDR) PINNACLE Registry, which prospectively captures information on the clinical care of outpatients, including the use of guideline-recommended secondary preventive therapies. Given the potential variability in care across clinics, we explicitly sought to examine both the variations in secondary preventive treatment of PAD by SES and whether treatment differences by SES were explained at the site level, with the hope that our findings would not only identify potential disparities by income but also define targets for future interventions to reduce disparities in PAD care.

## Methods

**Study population.** The PINNACLE Registry was launched in 2008 and represents the first national, prospective, office-based, quality-improvement registry of cardiovascular

patients in the United States (23,24). Among participating practices, patient data were collected at the point of care for a variety of cardiovascular conditions, including coronary artery disease, heart failure, atrial fibrillation, and PAD. Participation in this quality-improvement initiative is voluntary.

For the purposes of this study, we identified 66,282 patients with a diagnosis of PAD enrolled from 61 practices between July 1, 2010, and June 30, 2011. Within the PINNACLE Registry, PAD was defined by one of the following self-identified criteria by patients: 1) claudication, either with exertion or at rest; 2) amputation for arterial vascular insufficiency; 3) vascular reconstruction, bypass surgery, or percutaneous intervention to the extremities (excluding dialysis fistulas and vein stripping); 4) documented aortic aneurysm with or without repair; or 5) positive noninvasive test (e.g., ankle brachial index  $\leq 0.9$ , ultrasound, magnetic resonance, computed tomography) or diagnostic angiographic stenosis of  $>50\%$  in any major peripheral artery (e.g., renal, subclavian, femoral, iliac). We excluded 2,945 patients from whom information on SES was missing (zip code data were not available). Our final study cohort comprised 62,690 PAD patients from 61 sites. For the analyses in this study, as patients may have had multiple visits in the PINNACLE Registry, we used information from the first visit to represent each patient only once.

**SES and processes of care.** The key independent variable was patients' SES, which was defined by the median income of the patient's zip code of residence. This approach to categorize levels of socioeconomic status has been used in previous reports of various disease conditions (25–27). The primary study outcome was treatment with two secondary preventive medications: an antiplatelet agent (aspirin or clopidogrel) and a statin, both of which are Class I indications for PAD by the ACC/American Heart Association (AHA) PAD guidelines and PAD performance measures (28,29). Patients with documented contraindications to antiplatelet therapy (e.g., history of gastrointestinal bleeding) or statins were excluded from the analysis of each treatment. Moreover, for the analyses of antiplatelet therapy, we further excluded 9,295 patients already on warfarin therapy given that warfarin may influence the use of antiplatelet therapy.

**Statistical analysis.** Patients were categorized into quintiles of SES, with quintile 1 representing the lowest SES and quintile 5 the highest. Baseline differences across quintiles of SES were then compared using the Mantel-Haenszel trend test for categorical variables and the linear trend test for continuous variables.

Separate multivariable hierarchical modified Poisson models were used to assess the relationship between SES and treatment with antiplatelet therapy and statins. We employed two-level hierarchical models to adjust for clustering of patients within practices, with individual practices modeled as random effects and other patient characteristics modeled as fixed effects within each practice (30). This approach allowed us to control for measured and unmeasured

Cardiovascular Data Registry (NCDR). The views expressed in this manuscript represent those of the author(s), and do not necessarily represent the official views of the NCDR or its associated professional societies, identified at [www.ncdr.com](http://www.ncdr.com). The PINNACLE Registry is an initiative of the American College of Cardiology Foundation. Bristol-Myers Squibb and Pfizer Inc. are Founding Sponsors of the PINNACLE Registry. Dr. Patel has received research grants from Johnson & Johnson, the National Heart, Lung and Blood Institute (NHLBI), AstraZeneca, Medtronic, and Maquet; is a consultant/advisory board member for Bayer Healthcare, Genzyme, Ortho-McNeil-Janssen, Pleuristem, and Baxter. Dr. Jones has received research grants from AstraZeneca and Bristol-Myers Squibb. Dr. Bhatt is on the advisory board of Elsevier Practice Update Cardiology, Medscape Cardiology; is on the Board of Directors of the Boston VA Research Institute, Society of Chest Pain Centers; is the Chair of the American Heart Association Get With The Guidelines Science Subcommittee; has received honoraria from American College of Cardiology (Editor, Clinical Trials, Cardiosource), Duke Clinical Research Institute (clinical trial steering committees), Slack Publications (Chief Medical Editor, Cardiology Today's Intervention), WebMD (CME steering committees); is the Senior Associate Editor, *Journal of Invasive Cardiology*; is on the Data Monitoring Committees of Population Health Research Institute; has received research grants from Amarin, AstraZeneca, Bristol-Myers Squibb, Eisai, Ethicon, Medtronic, Sanofi Aventis, The Medicines Company; and has received unfunded research from FlowCo, PLx Pharma, and Takeda. Dr. Spertus has grants and contracts with the ACCF, NIH, NHLBI, ACCF, Lilly, Gilead, Amoryte, Genentech; consulting with Janssen, Genentech, United Healthcare, Amgen; copyrights of the SAQ, KCCQ, PAQ; and equity in Health Outcomes Sciences. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

Manuscript received March 7, 2013; accepted April 17, 2013.

between-practice confounding, as the use of hierarchical models ensured that patients with similar SESs were compared with each other from the same practice.

To better understand the extent of the practice-site variation in accounting for treatment differences by SES, we performed a 2-step sequential adjustment. First, we adjusted for practice site only, to assess the extent to which differences by SES were attenuated. This step allowed us to understand whether treatment differences by SES persisted when patients of different SESs within the same site were compared. Next, we additionally controlled for clinical characteristics, including age, sex, insurance status, diabetes, dyslipidemia, history of MI, history of revascularization in the previous 12 months, history of congestive heart failure, and history of stroke.

For each analysis, the null hypothesis was evaluated at a 2-sided significance level of 0.05, with 95% confidence intervals (CIs) calculated using robust SEs. All analyses were performed with SAS version 9.3 (SAS Institute, Cary, North Carolina) and R version 2.10.0.

## Results

The baseline characteristics of the 62,690 PAD patients by SES quintiles are summarized in Table 1. There was no difference in baseline characteristics of patients among practices categorized by tertiles of mean patient income (Table 2). The median number of patients per site was 151 (interquartile range [IQR]: 16 to 1,328) patients per site. Patients in the lowest SES quintile were from zip codes with median household incomes of <\$34,486 annually, while those in the

highest quintiles were from zip codes with annual median household incomes of >\$60,868.

The median age of the overall cohort was 70.0 (IQR: 62.0–78.0) years, and 62.4% were men. Nearly 60% of patients had private insurance, and only 4.2% were uninsured. There were high prevalences of coronary artery disease (85.4%), dyslipidemia (81.5%), and hypertension (83.3%) in the cohort. Nearly one third of patients were diabetic and one quarter were active smokers. Finally, 30.3% of patients had undergone coronary artery bypass surgery, whereas 41.4% had undergone percutaneous coronary intervention within the previous year.

Compared with patients in the highest SES quintiles, patients in the lower SES quintiles were slightly younger, more frequently female, and less likely to have private health insurance. Patients in lower SES quintiles were also more likely to have undergone percutaneous coronary intervention in the previous year and to be active smokers, and were less likely to have undergone coronary artery bypass surgery in the previous year and to have had a prior stroke. Lastly, rates of dyslipidemia, diabetes, hypertension, congestive heart failure, and prior myocardial infarction were clinically similar across quintiles.

**Use of cardioprotective therapy.** Treatment rates with statins decreased in a graded fashion going from higher to lower SES: 85.8% in quintile 5 to 72.5% in quintile 1 (Table 3). Compared with patients in the highest SES quintile, PAD patients in the lowest SES quintile were 16% less likely to be treated with statins (unadjusted rate ratio [RR]: 0.84; 95% CI: 0.83 to 0.86;  $p < 0.0001$ ). Notably, sites with higher mean incomes among its patients had greater

**Table 1** Baseline Demographic and Clinical Characteristics of the Study Cohort, Stratified by Socioeconomic Status

Characteristic	Socioeconomic Status					All Patients (N = 62,690)	p Value*
	Quintile 1 (n = 12,521)	Quintile 2 (n = 12,513)	Quintile 3 (n = 12,565)	Quintile 4 (n = 12,511)	Quintile 5 (n = 12,580)		
Patient income, range,† US\$	4,583–34,486	34,486–41,117	41,118–50,371	50,372–60,868	60,869–200,001	4,583–200,001	<0.001
Age, median (IQR), yrs	69.0 (61.0–78.0)	71.0 (62.0–79.0)	70.0 (62.0–78.0)	71.0 (63.0–79.0)	71.0 (63.0–79.0)	70.0 (62.0–78.0)	<0.001
Male, %	60.6	62.8	62.5	61.7	64.2	62.4	<0.001
Insurance category,‡ n (%)							
Private	54.5	63.3	59.7	58.7	61.0	59.4	<0.001
Public	41.1	33.0	37.2	37.9	35.2	36.9	<0.001
None	4.5	3.7	3.1	3.4	3.8	3.7	<0.001
Comorbidity, n (%)							
CAD	86.9	86.2	86.6	83.6	83.5	85.4	<0.001
Dyslipidemia	77.2	80.8	82.1	82.7	84.5	81.5	<0.001
Diabetes mellitus	33.4	34.0	32.2	31.5	32.2	32.7	<0.001
Hypertension	82.3	85.2	84.4	82.3	82.4	83.3	0.010
Stroke or TIA	11.5	24.1	25.2	16.4	18.0	19.0	<0.001
MI	30.9	29.9	31.8	28.7	31.5	30.6	0.982
CABG in previous 12 months	23.29	25.4	33.8	33.2	35.8	30.3	<0.001
PCI in previous 12 months	47.0	48.8	43.0	33.7	34.7	41.4	<0.001

\*Continuous variables compared using linear trend test; categorical variables compared using Mantel-Haenszel trend test. †Calculated as the median household income per year in each patient's zip code.

‡Public insurance refers to Medicare, Medicaid, military, and state insurance.

CABG = coronary artery bypass grafting; CAD = coronary artery disease; IQR = interquartile range; MI = myocardial infarction; PCI = percutaneous coronary intervention; TIA = transient ischemic attack; USD = US dollars.

**Table 2** Baseline Demographic and Clinical Characteristics Within Practices, by Tertile of Mean Patient Income

Characteristic	Mean Income			All Patients (N = 61)	p Value*
	Tertile 1 (n = 20)	Tertile 2 (n = 20)	Tertile 3 (n = 21)		
Age, yrs	68.7 (66.3–71.7)	71.1 (68.4–72.5)	70.6 (69.5–72.0)	70.3 (68.3–72.2)	0.16
Male, %	57.8 (53.1–62.9)	62.8 (55.4–69.1)	60.6 (55.4–65.2)	60.6 (55.4–66.1)	0.81
Insurance category,† %					
Private	46.9 (26.0–69.4)	45.6 (34.5–74.9)	44.8 (24.7–68.2)	45.5 (25.0–72.9)	0.69
Public	50.0 (23.7–69.3)	32.8 (21.3–65.1)	47.1 (28.4–68.8)	45.9 (23.0–68.8)	0.48
None	2.9 (0.0–9.7)	2.4 (0.9–7.6)	3.3 (1.0–5.9)	2.6 (0.8–7.5)	0.47
Comorbidity, n (%)					
CAD	84.9 (70.5–95.8)	83.9 (76.6–98.4)	77.3 (60.4–91.1)	83.2 (70.0–94.6)	0.05
Dyslipidemia	75.0 (60.4–88.7)	86.4 (74.2–92.3)	78.8 (74.9–92.0)	79.8 (71.4–90.9)	0.17
Diabetes mellitus	40.0 (30.7–50.0)	30.3 (17.4–37.5)	33.9 (29.2–41.4)	33.9 (27.0–43.6)	0.58
Hypertension	90.2 (78.4–100)	87.6 (75.8–98.3)	82.1 (78.4–92.1)	85.3 (78.3–97.0)	0.19
Stroke or TIA	7.8 (2.7–11.7)	7.0 (1.6–11.8)	7.0 (3.6–9.0)	7.2 (3.1–11.0)	0.88
MI	27.1 (9.6–51.7)	22.4 (10.6–39.2)	26.0 (11.2–30.4)	26.0 (10.0–40.0)	0.14
CABG in previous 12 months	8.0 (4.4–21.7)	2.4 (0.021.4)	12.4 (1.5–31.4)	6.8 (1.2–29.1)	0.56
PCI in previous 12 months	24.3 (8.3–50.0)	14.3 (3.4–51.4)	18.8 (7.2–47.2)	19.3 (7.1–50.0)	0.92

Values are median (IQR). \*Continuous variables compared using linear trend test; categorical variables compared using Mantel-Haenszel trend test.

†Public insurance refers to Medicare, Medicaid, military, and state insurance.

Abbreviations as in Table 1.

percentages of patients prescribed a statin medication (weighted correlation coefficient: 0.48) (Fig. 1). After adjustment for the practice at which a patient received care, treatment differences by SES were markedly attenuated (adjusted RR for quintile 1 vs. 5: 0.97; 95% CI: 0.97 to 0.99;  $p = 0.003$ ). Further adjustment for clinical variables was associated with only a small attenuation of differences by SES (fully adjusted RR: 1.00; 95% CI: 0.99 to 1.01;  $p = 0.772$ ). A similar pattern was observed for quintiles 2, 3, and 4 (Table 4).

Rates of any antiplatelet treatment were lowest among those in quintile 1 (79.0%, vs 84.6% in quintile 5) (Table 3). Compared with patients in the highest SES quintile,



patients in the lowest SES quintile were 7% less likely to be treated with any antiplatelet medication (unadjusted RR: 0.93; 95% CI: 0.91 to 0.94;  $p < 0.0001$ ). As with statin treatment, sites with a higher median income among its patients had greater percentages of patients prescribed an antiplatelet agent (weighted correlation coefficient: 0.30) (Fig. 2). After adjustment for the practice at which a patient received care, treatment differences by SES were nearly eliminated (adjusted RR: 0.98; 95% CI: 0.97 to 1.00;  $p = 0.012$ ). Further adjustment for the clinical characteristics of patients had minimal effect on attenuation of effect between the quintiles of SES (fully adjusted RR: 1.00; 95% CI: 0.99 to 1.01;  $p = 0.878$ ) (Table 4).

**Table 3** Medication\* Treatment Rates, Stratified by Socioeconomic Status†

Medication	Socioeconomic Status					All Patients (N = 62,690)	p Value‡
	Quintile 1 (n = 12,521)	Quintile 2 (n = 12,513)	Quintile 3 (n = 12,565)	Quintile 4 (n = 12,511)	Quintile 5 (n = 12,580)		
Any antiplatelet: clopidogrel and/or aspirin	79.0	83.0	84.8	83.1	84.6	82.9	<0.001
Statin	72.5	77.6	83.2	81.5	85.8	80.1	<0.001

\*Medications described among those eligible (without contraindication). †Stratified by range of patient income (in USD) (calculated as the median household income per year in each patient's zip code), as follows: quintile 1 = 4,583–34,486; quintile 2 = 34,486–41,117; quintile 3 = 41,118–50,371; quintile 4 = 50,372–60,868; and quintile 5 = 60,869–200,001. ‡Continuous variables compared using linear trend test; categorical variables compared using Mantel-Haenszel trend test unless otherwise noted.

KW = Kruskal-Wallis test; other abbreviations as in Table 1.



Discussion

Among outpatients with PAD, we found that treatment with antiplatelet and statin therapies differed by SES. These differences, however, were largely explained by the clinical practice at which patients received care. Our findings suggest

that initiatives to reduce disparities in medication treatment for PAD should target practices with high proportions of low-SES patients.

Studies from different populations have demonstrated that cardiovascular risk factors and disease disproportionately affect those of lower SES (31–36), and among patients with

Table 4      Association of Socioeconomic Status* With PAD Treatment						
Treatment/Quintile	Unadjusted		Adjusted for Practice Site		Further Adjusted for Clinical Characteristics†	
	RR (95% CI)	p Value	RR (95% CI)	p Value	RR (95% CI)	p Value
Statin therapy						
Quintile 5	[Reference]	–	[Reference]	–	[Reference]	–
Quintile 4	0.95 (0.94–0.96)	<0.001	0.98 (0.97–1.00)	0.0134	0.98 (0.97–1.00)	0.007
Quintile 3	0.97 (0.96–0.98)	<0.001	0.98 (0.97–1.00)	0.0124	0.99 (0.98–1.00)	0.172
Quintile 2	0.90 (0.89–0.91)	<0.001	0.98 (0.95–1.00)	0.0341	0.99 (0.97–1.01)	0.327
Quintile 1	0.84 (0.83–0.86)	<0.001	0.97 (0.95–0.99)	0.0029	1.00 (0.99–1.01)	0.772
Any antiplatelet‡						
Quintile 5	[Reference]	–	[Reference]	–	[Reference]	–
Quintile 4	0.99 (0.98–1.00)	0.022	1.00 (0.99–1.01)	0.556	1.01 (1.00–1.01)	0.237
Quintile 3	1.00 (0.99–1.01)	0.946	0.99 (0.98–1.01)	0.347	1.00 (0.99–1.01)	0.788
Quintile 2	0.98 (0.97–0.99)	<0.0001	1.00 (0.99–1.01)	0.794	1.01 (1.00–1.02)	0.071
Quintile 1	0.93 (0.91–0.94)	<0.0001	0.98 (0.97–1.00)	0.012	1.00 (0.99–1.01)	0.878

\*Stratified by range of patient income (in USD) (calculated as the median household income per year in each patient's zip code), as follows: quintile 1 = 4,583–34,486; quintile 2 = 34,486–41,117; quintile 3 = 41,118–50,371; quintile 4 = 50,372–60,868; and quintile 5 = 60,869–200,001. †Clinical characteristics adjusted model: model adjusted for practice, age, sex, history of myocardial infarction, revascularization in the previous 12 months, insurance, congestive heart failure, diabetes, stroke, dyslipidemia, and tobacco use. ‡Excluding patients on warfarin.

CI = confidence interval; PAD = peripheral artery disease; RR = relative risk.



cardiac disease, those with lower SES experience higher morbidity and mortality (4,37–40). Patient-level risk factors, such as increased health-risk behaviors, may account for some of the increased morbidity and mortality seen in patients of low SES (4,41). However, identifying factors beyond patient risk factors and behaviors (36), which are not easily modifiable, is crucial to quality initiatives that address the Institute of Medicine mandates to improve outcomes and reduce disparities. For instance, several studies have reported variation in compliance with evidence-based therapies for other cardiac conditions by SES, and may partly explain the association between lower SES and worse outcomes (37,42,43). However, these prior studies have not examined differences in treatments of PAD by SES. Moreover, they have not examined the extent to which the site at which a patient receives his or her care influences treatment rates.

The present study expands on findings of other disparities research and focuses on PAD, in which outcomes research has been limited. Prior studies of disparities in PAD have focused on rates of utilization of lower extremity revascularization by SES or reported rates of optimal medical therapy by insurance status (5–9). This present study confirms findings similar to those in other cardiac disease states, in that the use of secondary preventive treatments for PAD was lower among those of low SES. This finding highlights an important gap in the quality of care of patients with PAD, especially because the cost of aspirin and generic statins is low and should not pose significant barriers to patient access to these evidence-based therapies.

The present analysis further contributes to disparities research by highlighting the central role of the practice at which patients receive their care in explaining treatment differences by SES. We found that disparities in medication use between the highest and lowest SESs were markedly attenuated after adjustment for site-level variation; this finding suggests that differences in medication use were predominately explained by differences in sites that largely treat patients of low SES compared with sites caring for largely higher-SES patients. We believe these findings serve as an important paradigm for future efforts to reduce disparities in care, which will need to target clinical practices as intervention units and go beyond patient-level interventions. For PAD, future studies are needed to determine whether system-wide improvements at the practice level (e.g., identification of patients with PAD, initiation of secondary prevention medications, and physician education) or resource interventions at practices with high proportions of low-SES patients (e.g., electronic medical systems, decision aids) will reduce disparities in treatment by SES. Quality-improvement initiatives that provide feedback to sites by providing reports benchmarking performance of the site in relation to pre-specified goals or national averages for select performance metrics may help change behavior at a site.

**Study limitations.** First, we relied on patient diagnoses for PAD, which were self-reported by practices. It is possible that some patients were not classified as having PAD;

however, we believe any misclassification would have been nondifferential and are unlikely to have influenced our findings. Moreover, we were unable to examine severity of PAD, as we did not have physiological (e.g., ankle-brachial index) or angiographic data on all patients. Similarly, we were unable to analyze the variability in use of medication by the specific PAD diagnosis (i.e., surgery vs. noninvasive test). Second, we defined SES by median income in each patient's residential zip code, which is a common strategy in previous studies (25–27), and did not examine other socioeconomic variables, such as educational level, which were not available in the PINNACLE registry. Third, our study was conducted among cardiology practices participating in PINNACLE, a quality-improvement registry; therefore, treatment rates by SES may differ in nonparticipating practices, including primary care centers. Given voluntary enrollment in this quality-improvement initiative, it is possible that the rates of medication use are higher than expected in nonparticipating sites. We did not adjust for race in the clinical model given that it was frequently missing (nearly 50% of patients had missing data on this variable); however, most of the variation in treatment between the SES groups could be accounted for by practice-level variation. Furthermore, although it is possible that the differences in therapies by SES could be mediated by some, but not all, physicians within a practice, the current PINNACLE registry does not provide information on provider characteristics sufficient to allow us to currently examine this possibility. Regardless, further education and resources geared at practices with high proportions of low-SES patients have the potential to improve compliance to therapies indicated for PAD. Finally, we were unable to examine longitudinal outcomes in this study.

## Conclusions

Among patients with PAD, treatment with evidence-based antiplatelet and statin therapies differed by patients' SES. These differences, however, were largely explained by the clinical practice at which patients received care, suggesting variation in treatment patterns across centers. Future efforts to reduce treatment disparities by SES in PAD and to improve outcomes in these vulnerable populations should target practices serving high proportions of patients of low SES.

---

**Reprint requests and correspondence:** Dr. Sumeet Subherwal, Duke Clinical Research Institute, 2400 Pratt Street, Room 0311 Terrace Level, Durham, North Carolina 27705. E-mail: [sumeet.subherwal@duke.edu](mailto:sumeet.subherwal@duke.edu).

---

## REFERENCES

1. Quality of Health Care in America, Institute of Medicine. Crossing the Quality Chasm: A New Health System for the 21st Century. New York, NY: The National Academies Press, 2001.
2. Rathore SS, Berger AK, Weinfurt KP, et al. Race, sex, poverty, and the medical treatment of acute myocardial infarction in the elderly. *Circulation* 2000;102:642–8.

3. Rathore SS, Foody JM, Wang Y, et al. Race, quality of care, and outcomes of elderly patients hospitalized with heart failure. *JAMA* 2003;289:2517-24.
4. Alter DA, Chong A, Austin PC, et al. Socioeconomic status and mortality after acute myocardial infarction. *Ann Intern Med* 2006;144:82-93.
5. LaMorte WW, Scott TE, Menzoian JO. Racial differences in the incidence of femoral bypass and abdominal aortic aneurysmectomy in Massachusetts: relationship to cardiovascular risk factors. *J Vasc Surg* 1995;21:422-31.
6. Huber TS, Wang JG, Wheeler KG, et al. Impact of race on the treatment for peripheral arterial occlusive disease. *J Vasc Surg* 1999;30:417-26.
7. Eslami MH, Zayaruzny M, Fitzgerald GA. The adverse effects of race, insurance status, and low income on the rate of amputation in patients presenting with lower extremity ischemia. *J Vasc Surg* 2007;45:55-9.
8. Meadows TA, Bhatt DL, Hirsch AT, et al. Ethnic differences in the prevalence and treatment of cardiovascular risk factors in US outpatients with peripheral arterial disease: insights from the Reduction of Atherothrombosis for Continued Health (REACH) Registry. *Am Heart J* 2009;158:1038-45.
9. Yeh D, Jones M, Schulman C, Karmacharya J, Velazquez OC. Uninsured South Florida vascular surgery patients are less likely to receive optimal medical management than their insured counterparts. *J Vasc Surg* 2010;51 Suppl:S4-8.
10. Pande RL, Perlstein TS, Beckman JA, Creager MA. Secondary prevention and mortality in peripheral artery disease: National Health and Nutrition Examination Study, 1999 to 2004. *Circulation* 2011;124:17-23.
11. Kröger K, Dragano N, Stang A, et al. An unequal social distribution of peripheral arterial disease and the possible explanations: results from a population-based study. *Vasc Med* 2009;14:289-96.
12. Carson AP, Rose KM, Catellier DJ, et al. Cumulative socioeconomic status across the life course and subclinical atherosclerosis. *Ann Epidemiol* 2007;17:296-303.
13. Yusuf S, Sleight P, Pogue J, Bosch J, Davies R, Dagenais G, for the Heart Outcomes Prevention Evaluation Study Investigators. Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. *N Engl J Med* 2000;342:145-53.
14. A randomised, blinded, trial of clopidogrel versus aspirin in patients at risk of ischaemic events (CAPRIE). *Lancet* 1996;34:1329-39.
15. Lindholm LH. MRC/BHF Heart Protection Study of cholesterol-lowering with simvastatin in 5963 people with diabetes: a randomised placebo-controlled trial. *Commentary. Lancet* 2003;36:2005-16.
16. Antithrombotic Trialists' Collaboration. Collaborative meta-analysis of randomised trials of antiplatelet therapy for prevention of death, myocardial infarction, and stroke in high risk patients. *BMJ* 2002;324(7329):71-86.
17. Feringa HHH, van Waning VH, Bax JJ, et al. Cardioprotective medication is associated with improved survival in patients with peripheral arterial disease. *J Am Coll Cardiol* 2006;47:1182-7.
18. Leng GC, Lee AJ, Fowkes FGR, et al. Incidence, natural history and cardiovascular events in symptomatic and asymptomatic peripheral arterial disease in the general population. *Int J Epidemiol* 1996;25:1172-81.
19. Steg PG, Bhatt DL, Wilson PWF, et al. One-year cardiovascular event rates in outpatients with atherothrombosis. *JAMA* 2007;297:1197-206.
20. Resnick HE, Lindsay RS, McDermott MM, et al. Relationship of high and low ankle brachial index to all-cause and cardiovascular disease mortality. *Circulation* 2004;109:733-9.
21. Criqui MH, Langer RD, Fronck A, et al. Mortality over a period of 10 years in patients with peripheral arterial disease. *N Engl J Med* 1992;326:381-6.
22. Criqui MH, Ninomiya JK, Wingard DL, Ji M, Fronck A. Progression of peripheral arterial disease predicts cardiovascular disease morbidity and mortality. *J Am Coll Cardiol* 2008;52:1736-42.
23. Chan PS, Oetgen WJ, Buchanan D, et al. Cardiac performance measure compliance in outpatients: the American College of Cardiology and National Cardiovascular Data Registry's PINNACLE (Practice Innovation And Clinical Excellence) program. *J Am Coll Cardiol* 2010;56:8-14.
24. Chan PS, Oetgen WJ, Spertus JA. The Improving Continuous Cardiac Care (IC3) Program and outpatient quality improvement. *Am J Med* 2010;123:217-9.
25. Burstin HR, Lipsitz SR, Brennan TA. Socioeconomic status and risk for substandard medical care. *JAMA* 1992;268:2383-7.
26. Santry HP, Gillen DL, Lauderdale DS. Trends in bariatric surgical procedures. *JAMA* 2005;294:1909-17.
27. Rao SV, Schulman KA, Curtis LH, Gersh BJ, Jollis JG. Socioeconomic status and outcome following acute myocardial infarction in elderly patients. *Arch Intern Med* 2004;164:1128-33.
28. Rooke TW, Hirsch AT, Misra S, et al. 2011 ACCF/AHA focused update of the guideline for the management of patients with peripheral artery disease (updating the 2005 guideline). *Circulation* 2011;124:2020-45.
29. Olin JW, Allie DE, Belkin M, et al. ACCF/AHA/ACR/SCAI/SIR/SVM/SVN/SVS 2010 performance measures for adults with peripheral artery disease: a report of the American College of Cardiology Foundation/American Heart Association Task Force on Performance Measures, the American College of Radiology, the Society for Cardiac Angiography and Interventions, the Society for Interventional Radiology, the Society for Vascular Medicine, the Society for Vascular Nursing, and the Society for Vascular Surgery (Writing Committee to Develop Clinical Performance Measures for Peripheral Artery Disease) Developed in Collaboration With the American Association of Cardiovascular and Pulmonary Rehabilitation; the American Diabetes Association; the Society for Atherosclerosis Imaging and Prevention; the Society for Cardiovascular Magnetic Resonance; the Society of Cardiovascular Computed Tomography; and the PAD Coalition Endorsed by the American Academy of Podiatric Practice Management. *J Am Coll Cardiol* 2010;56:2147-81.
30. Goldstein H. Multilevel statistical models. Volume 847. 4th edition. London, UK: Wiley, 2010.
31. Cooper R, Cutler J, Desvigne-Nickens P, et al. Trends and disparities in coronary heart disease, stroke, and other cardiovascular diseases in the United States: findings of the National Conference on Cardiovascular Disease Prevention. *Circulation* 2000;102:3137-47.
32. Smith SC, Clark LT, Cooper RS, et al. Discovering the full spectrum of cardiovascular disease. *Circulation* 2005;1110:e134-9.
33. Lynch J, Kaplan GA, Salonen R, Salonen JT. Socioeconomic status and progression of carotid atherosclerosis: prospective evidence from the Kuopio Ischemic Heart Disease Risk Factor Study. *Arterioscler Thromb Vasc Biol* 1997;17:513-9.
34. Deans KA, Bezlyak V, Ford I, et al. Differences in atherosclerosis according to area level socioeconomic deprivation: cross sectional, population based study. *BMJ* 2009;339:b4170.
35. Winkleby MA, Robinson TN, Sundquist J, Kraemer HC. Ethnic variation in cardiovascular disease risk factors among children and young adults. *JAMA* 1999;281:1006-13.
36. Albert MA, Glynn RJ, Buring J, Ridker PM. Impact of traditional and novel risk factors on the relationship between socioeconomic status and incident cardiovascular events. *Circulation* 2006;1144:2619-26.
37. Kapral MK, Wang H, Mamdani M, Tu JV. Effect of socioeconomic status on treatment and mortality after stroke. *Stroke* 2002;33:268-75.
38. Cox AM, McKevitt C, Rudd AG, Wolfe CDA. Socioeconomic status and stroke. *Lancet Neurol* 2006;5:181-8.
39. Rathore SS, Masoudi FA, Wang Y, et al. Socioeconomic status, treatment, and outcomes among elderly patients hospitalized with heart failure: findings from the National Heart Failure Project. *Am Heart J* 2006;152:371-8.
40. Koch CG, Li L, Kaplan GA, et al. Socioeconomic position, not race, is linked to death after cardiac surgery. *Circ Cardiovasc Qual Outcomes* 2010;3:267-76.
41. Lantz PM, House JS, Lepkowski JM, Williams DR, Mero RP, Chen J. Socioeconomic factors, health behaviors, and mortality. *JAMA* 1998;279:1703-8.
42. Philbin EF, McCullough PA, DiSalvo TG, Dec GW, Jenkins PL, Weaver WD. Socioeconomic status is an important determinant of the use of invasive procedures after acute myocardial infarction in New York State. *Circulation* 2000;102 Suppl 3:III107-15.
43. Hernandez AF, Fonarow GC, Liang L, et al. Sex and racial differences in the use of implantable cardioverter-defibrillators among patients hospitalized with heart failure. *JAMA* 2007;298:1525-32.

**Key Words:** disparities ■ medications ■ peripheral artery disease ■ secondary prevention.