

CLINICAL RESEARCH

Cardiovascular Risk

## Apolipoprotein(a) Genetic Sequence Variants Associated With Systemic Atherosclerosis and Coronary Atherosclerotic Burden But Not With Venous Thromboembolism

Anna Helgadóttir, MD, PhD,<sup>1,2,3</sup> Solveig Gretarsdóttir, PhD,<sup>1</sup> Gudmar Thorleifsson, PhD,<sup>1</sup> Hilma Holm, MD,<sup>1</sup> Riyaz S. Patel, MD,<sup>4,5</sup> Thorarinn Gudnason, MD, PhD,<sup>6</sup> Gregory T. Jones, PhD,<sup>7</sup> Andre M. van Rij, MD,<sup>7</sup> Danny J. Eapen, MD,<sup>4</sup> Annette F. Baas, PhD,<sup>8</sup> David-Alexandre Tregouet, PhD,<sup>9</sup> Pierre-Emmanuel Morange, MD,<sup>10</sup> Joseph Emmerich, MD, PhD,<sup>11</sup> Bengt Lindblad, MD, PhD,<sup>12</sup> Anders Gottsäter, MD, PhD,<sup>12</sup> Lambertus A. Kiemeny, PhD,<sup>13</sup> Jes S. Lindholt, MD, PhD,<sup>14</sup> Natzi Sakalihasan, MD, PhD,<sup>15</sup> Robert E. Ferrell, PhD,<sup>16</sup> David J. Carey, PhD,<sup>17</sup> James R. Elmore, MD,<sup>18</sup> Philip S. Tsao, PhD,<sup>19</sup> Niels Garup, MD, PhD,<sup>20</sup> Torben Jørgensen, MD, DMSci,<sup>21,22</sup> Daniel R. Witte, MD, PhD,<sup>23</sup> Torben Hansen, MD, PhD,<sup>20,24</sup> Oluf Pedersen, MD, DMSci,<sup>20,25,26,27</sup> Roberto Pola, MD, PhD,<sup>28,29</sup> Eleonora Gaetani, MD,<sup>28</sup> Hulda B. Magnadóttir, MD,<sup>30</sup> Cisca Wijmenga, PhD,<sup>31</sup> Gerard Tromp, PhD,<sup>17</sup> Antti Ronkainen, MD, PhD,<sup>32</sup> Ynte M. Ruigrok, MD,<sup>33</sup> Jan D. Blankensteijn,<sup>34,35</sup> Thomas Mueller, MD,<sup>36</sup> Philip S. Wells, MD,<sup>37</sup> Javier Corral, PhD,<sup>38</sup> Jose Manuel Soria, PhD,<sup>39</sup> Juan Carlos Souto, MD, PhD,<sup>40</sup> John F. Peden, PhD,<sup>41</sup> Shapour Jalilzadeh, MD, PhD,<sup>41</sup> Bongani M. Mayosi, DPhil,<sup>42</sup> Bernard Keavney, MD,<sup>43</sup> Rona J. Strawbridge PhD,<sup>44</sup> Maria Sabater-Lleal, PhD,<sup>44</sup> Karl Gertow, PhD,<sup>44</sup> Damiano Baldassarre, PhD,<sup>45</sup> Kristiina Nyyssönen, PhD,<sup>46</sup> Rainer Rauramaa, MD, PhD,<sup>47</sup> Andries J. Smit, MD, PhD,<sup>48</sup> Elmo Mannarino, MD,<sup>49</sup> Philippe Giral, MD,<sup>50</sup> Elena Tremoli, PhD,<sup>45</sup> Ulf de Faire, MD, PhD,<sup>51</sup> Steve E. Humphries, PhD,<sup>52</sup> Anders Hamsten, MD, PhD,<sup>44</sup> Vilhelmina Haraldsdóttir, MD,<sup>53</sup> Isleifur Olafsson, MD, PhD,<sup>53</sup> Magnus K. Magnusson, MD,<sup>3,53</sup> Nilesh J. Samani, MD,<sup>54,55</sup> Allan I. Levey, MD, PhD,<sup>4</sup> Hugh S. Markus, MD,<sup>56</sup> Konstantinos Kostulas, MD, PhD,<sup>57</sup> Martin Dichgans, MD,<sup>58</sup> Klaus Berger, MD,<sup>59</sup> Gregor Kuhlenbäumer, MD,<sup>60</sup> E. Bernd Ringelstein, MD,<sup>61</sup> Monika Stoll, PhD,<sup>62</sup> Udo Seedorf, PhD,<sup>62</sup> Peter M. Rothwell, MD, PhD,<sup>63</sup> Janet T. Powell, MD,<sup>64</sup> Helena Kuivaniemi, MD, PhD,<sup>17</sup> Pall T. Onundarson, MD,<sup>3,53</sup> Einar Valdimarsson, MD,<sup>65</sup> Stefan E. Matthiasson, MD, PhD,<sup>66</sup> Daniel F. Gudbjartsson, PhD,<sup>1</sup> Guðmundur Thorgeirsson, MD, PhD,<sup>3,6</sup> Arshed A. Quyyumi, MD,<sup>4</sup> Hugh Watkins, MD, PhD,<sup>41</sup> Martin Farrall, MD,<sup>41</sup> Unnur Thorsteinsdóttir, PhD,<sup>1,3</sup> Kari Stefansson, MD, PhD<sup>1,3</sup>

Reykjavik, Iceland; Atlanta, Georgia; Pittsburgh and Danville, Pennsylvania; Stanford, California; Boston, Massachusetts; Lebanon, New Hampshire; Cardiff, Oxford, Newcastle, London, and Leicester, United Kingdom; Otago, New Zealand; Utrecht, Nijmegen, Groningen, and Amsterdam, the Netherlands; Paris and Marseille, France; Malmö, Stockholm, and Huddinge, Sweden; Viborg, Copenhagen, Gentofte, Glostrup, Odense, and Aarhus, Denmark; Liège, Belgium; Rome, Milan, and Perugia, Italy; Kuopio, Finland; Linz, Austria; Ottawa, Ontario, Canada; Murcia and Barcelona, Spain; Cape Town, South Africa; and Munich, Kiel, and Münster, Germany

## Objectives

The purpose of this study is investigate the effects of variants in the apolipoprotein(a) gene (*LPA*) on vascular diseases with different atherosclerotic and thrombotic components.

## Background

It is unclear whether the *LPA* variants rs10455872 and rs3798220, which correlate with lipoprotein(a) levels and coronary artery disease (CAD), confer susceptibility predominantly via atherosclerosis or thrombosis.

## Methods

The 2 *LPA* variants were combined and examined as *LPA* scores for the association with ischemic stroke (and TOAST [Trial of Org 10172 in Acute Stroke Treatment] subtypes) (effective sample size [ $n_e$ ] = 9,396); peripheral arterial disease ( $n_e$  = 5,215); abdominal aortic aneurysm ( $n_e$  = 4,572); venous thromboembolism ( $n_e$  = 4,607); intracranial aneurysm ( $n_e$  = 1,328); CAD ( $n_e$  = 12,716), carotid intima-media thickness ( $n$  = 3,714), and angiographic CAD severity ( $n$  = 5,588).

## Results

*LPA* score was associated with ischemic stroke subtype large artery atherosclerosis (odds ratio [OR]: 1.27;  $p = 6.7 \times 10^{-4}$ ), peripheral artery disease (OR: 1.47;  $p = 2.9 \times 10^{-14}$ ), and abdominal aortic aneurysm (OR: 1.23;  $p = 6.0 \times 10^{-5}$ ), but not with the ischemic stroke subtypes cardioembolism (OR: 1.03;  $p = 0.69$ ) or small vessel disease (OR: 1.06;  $p = 0.52$ ). Although the *LPA* variants were not associated with carotid intima-media thickness, they were associated with the number of obstructed coronary vessels ( $p = 4.8 \times 10^{-12}$ ). Furthermore, CAD cases carrying *LPA* risk variants had increased susceptibility to atherosclerotic manifestations outside of the coronary tree (OR: 1.26;  $p = 0.0010$ ) and had earlier onset of CAD (−1.58 years/allele;  $p = 8.2 \times 10^{-8}$ ) than CAD cases not carrying the risk variants. There was no association of *LPA* score with venous thromboembolism (OR: 0.97;  $p = 0.63$ ) or intracranial aneurysm (OR: 0.85;  $p = 0.15$ ).

## Conclusions

*LPA* sequence variants were associated with atherosclerotic burden, but not with primarily thrombotic phenotypes. (J Am Coll Cardiol 2012;60:722–9) © 2012 by the American College of Cardiology Foundation

From <sup>1</sup>Population Genomics, deCODE Genetics, Reykjavik, Iceland; <sup>2</sup>Institute of Human Genetics, University of Iceland, Reykjavik, Iceland; <sup>3</sup>Faculty of Medicine, University of Iceland, Reykjavik, Iceland; <sup>4</sup>Emory University School of Medicine, Atlanta, Georgia; <sup>5</sup>Cardiff University, Cardiff, United Kingdom; <sup>6</sup>Department of Cardiology and Cardiovascular Research Institute, Landspítali University Hospital and University of Iceland, Reykjavik, Iceland; <sup>7</sup>Department of Surgery, Dunedin School of Medicine, University of Otago, Otago, New Zealand; <sup>8</sup>Department of Medical Genetics, University Medical Center Utrecht, Utrecht, the Netherlands; <sup>9</sup>INSERM UMRS 937, ICAN Institute, and Université Pierre et Marie Curie, Paris, France; <sup>10</sup>INSERM UMRS 1062 and Aix-Marseille Université, Marseille, France; <sup>11</sup>INSERM UMRS 765, Médecine Vasculaire–HTA, Hôpital Européen Georges-Pompidou, Université Paris-Descartes, Paris, France; <sup>12</sup>University of Lund, Department of Vascular Diseases, Skåne University Hospital, Malmö, Sweden; <sup>13</sup>Department of Epidemiology, Biostatistics and HTA, Radboud University Nijmegen Medical Centre, Nijmegen, the Netherlands; <sup>14</sup>Vascular Research Unit, Viborg Hospital, Viborg, Denmark; <sup>15</sup>University Hospital of Liège, Liège, Belgium; <sup>16</sup>Department of Human Genetics, School of Public Health, University of Pittsburgh, Pittsburgh, Pennsylvania; <sup>17</sup>Weis Center for Research, Geisinger Clinic, Danville, Pennsylvania; <sup>18</sup>Department of Vascular Surgery, Geisinger Clinic, Danville, Pennsylvania; <sup>19</sup>Division of Cardiovascular Medicine, Stanford University School of Medicine, Stanford, California; <sup>20</sup>Novo Nordisk Foundation Center for Basic Metabolic Research, Faculty of Health Sciences, University of Copenhagen, Copenhagen, Denmark; <sup>21</sup>Research Centre for Prevention and Health, Glostrup Hospital, Glostrup, Denmark; <sup>22</sup>Faculty of Health Sciences, University of Copenhagen, Copenhagen, Denmark; <sup>23</sup>Steno Diabetes Center, Gentofte, Denmark; <sup>24</sup>University of Southern Denmark, Odense, Denmark; <sup>25</sup>Hagedorn Research Institute, Gentofte, Denmark; <sup>26</sup>Institute of Biomedical Sciences, Faculty of Health Sciences, University of Copenhagen, Copenhagen, Denmark; <sup>27</sup>Faculty of Health Sciences, University of Aarhus, Aarhus, Denmark; <sup>28</sup>Laboratory of Vascular Biology and Genetics, Department of Medicine, A. Gemelli University Hospital, Catholic University School of Medicine, Rome, Italy; <sup>29</sup>Division of Cardiovascular Research, Department of Medicine, Caritas St. Elizabeth's Medical Center, Tufts University School of Medicine, Boston, Massachusetts; <sup>30</sup>Upper Valley Neurology Neurosurgery, Lebanon, New Hampshire; <sup>31</sup>University of Groningen, University Medical Center Groningen, Department of Genetics, Groningen, the Netherlands; <sup>32</sup>Department of Neurosurgery, Kuopio University Hospital, Kuopio, Finland; <sup>33</sup>Utrecht Stroke Center, Department of Neurology and Neurosurgery, Rudolf Magnus Institute of Neuroscience, University Medical Center Utrecht, Utrecht, the Netherlands; <sup>34</sup>Division of Surgery, VU Medical Center, Amsterdam, the Netherlands; <sup>35</sup>Department of Vascular Surgery, Radboud University Medical Center, Nijmegen, the Netherlands; <sup>36</sup>Department of Laboratory Medicine, Konventhospital Barmherzige Brüder, Linz, Austria; <sup>37</sup>Department of

Medicine, Division of Hematology, University of Ottawa/Ottawa Hospital Research Institute, Ottawa, Ontario, Canada; <sup>38</sup>Department of Medicine, University of Murcia, Centro Regional de Hemodonación, Murcia, Spain; <sup>39</sup>Unit of Genomics of Complex Diseases, Research Institute, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain; <sup>40</sup>Haemostasis and Thrombosis Unit, Department of Haematology, Hospital de la Santa Creu i Sant Pau, Barcelona, Spain; <sup>41</sup>Department of Cardiovascular Medicine, The Wellcome Trust Centre of Human Genetics, University of Oxford, Oxford, United Kingdom; <sup>42</sup>Faculty of Health Sciences, University of Cape Town, Cape Town, South Africa; <sup>43</sup>Institute of Genetic Medicine, Newcastle University, United Kingdom; <sup>44</sup>Atherosclerosis Research Unit, Department of Medicine Solna, Karolinska Institutet, Stockholm, Sweden; <sup>45</sup>Dipartimento di Scienze Farmacologiche e Biomolecolari, Università di Milano, Centro Cardiologico Monzino, IRCCS, Milan, Italy; <sup>46</sup>Institute of Public Health and Clinical Nutrition, University of Eastern Finland, Kuopio, Finland; <sup>47</sup>Foundation for Research in Health, Exercise and Nutrition, Kuopio Research Institute of Exercise Medicine, Kuopio, Finland; <sup>48</sup>Department of Medicine, University Medical Center, Groningen, the Netherlands; <sup>49</sup>Internal Medicine, Angiology and Arteriosclerosis Diseases, Department of Clinical and Experimental Medicine, University of Perugia, Perugia, Italy; <sup>50</sup>Assistance Publique–Hôpitaux de Paris; Service Endocrinologie–Métabolisme, Groupe Hôpitalier Pitie-Salpêtrière, Unités de Prévention Cardiovasculaire, Paris, France; <sup>51</sup>Division of Cardiovascular Epidemiology, Institute of Environmental Medicine, Karolinska Institutet, Stockholm, Sweden; <sup>52</sup>Centre for Cardiovascular Genetics, BHF Laboratories, Institute Cardiovascular Science, University College London, London, United Kingdom; <sup>53</sup>Landspítali University Hospital, Reykjavik, Iceland; <sup>54</sup>Department of Cardiovascular Sciences, University of Leicester, Clinical Sciences Wing, Glenfield Hospital, Leicester, United Kingdom; <sup>55</sup>Leicester NIHR Biomedical Research Unit in Cardiovascular Disease and <sup>56</sup>Stroke and Dementia Research Centre, St George's University of London, London, United Kingdom; <sup>57</sup>Department of Neurology, Karolinska Institutet at Karolinska University Hospital, Huddinge, Sweden; <sup>58</sup>Institute for Stroke and Dementia Research, Klinikum der Universität München, Ludwig Maximilians University Munich, Munich, Germany; <sup>59</sup>Institute of Epidemiology and Social Medicine, University of Münster, Münster, Germany; <sup>60</sup>Institute of Experimental Medicine, University of Kiel, Kiel, Germany; <sup>61</sup>Department of Neurology, University of Münster, Münster, Germany; <sup>62</sup>Leibniz-Institute of Arteriosclerosis Research at the University of Münster, Germany; <sup>63</sup>Nuffield Department of Clinical Neurosciences, John Radcliffe Hospital, Oxford, United Kingdom; <sup>64</sup>Vascular Surgery Research Group, Imperial College at Charing Cross, London, United Kingdom; <sup>65</sup>Department of Neurology, Landspítali University Hospital, Reykjavik, Iceland; and <sup>66</sup>Lackning, Medical Clinics, Reykjavik, Iceland. Work performed at deCODE was funded in part through grants from the U.S. National Heart, Lung and Blood Institute (NHLBI), National Institutes of Health

Recent studies strongly support a causal relationship between circulating lipoprotein (a) (Lp[a]) and coronary artery disease (CAD) (1-3). However, the pathophysiologic mechanism of this relationship is unclear. Substantial experimental evidence supports the notion that Lp(a) has both atherogenic and thrombogenic properties (4-7), suggesting that Lp(a) may have a role in both atherosclerotic and thrombotic aspects of arterial disease and potentially also in venous thrombosis.

Alleles of 2 single nucleotide polymorphisms (SNPs) in the *LPA* gene, *rs10455872* and *rs3798220*, have been shown to be associated with high plasma levels of Lp(a) and CAD (1,8). The variants, which are poorly correlated ( $r^2 < 0.001$ ), together explain about 36% of the variance in Lp(a) levels (1). Each minor allele of *rs10455872* (G; 6.2% frequency) and *rs3798220* (C; 1.4% frequency) increased log Lp(a) by 1.08 and 1.15 standard deviation units and the risk for CAD by 47% and 68%, respectively (1). Evidence for the association between plasma Lp(a) and other atherosclerotic and thrombotic diseases is weaker; some studies have supported such an association, but others have not (2,3,9-18). Furthermore, it has not been established whether Lp(a), or Lp(a) correlated variants, confer the same risks to all of the ischemic stroke TOAST (Trial of Org 10172 in Acute Stroke Treatment) subtypes (19,20). Therefore, additional information about the role of Lp(a) at different vascular sites obtained from relatively large datasets that supports or refutes previously found associations is needed.

The objective of this study was to investigate the effects of the *LPA* variants on the risks for vascular diseases with different atherosclerotic and thrombotic components, to

provide insight into the mechanism by which Lp(a) confers susceptibility. We investigated the association of the aforementioned Lp(a) correlated SNPs, with vascular diseases related to atherosclerosis and/or thrombosis: ischemic stroke (and its TOAST subtypes), peripheral arterial disease (PAD), abdominal aortic aneurysm (AAA), venous thromboembolism (VTE), and CAD in many case-control series. We also tested for association with intracranial aneurysm (IA), an arterial disease without arteriosclerotic or thrombotic etiology. Additionally, the effects of *LPA* SNPs on atherosclerotic burden, as well as on carotid intima-media thickness (IMT), an early marker of atherosclerosis, were examined.

## Methods

**Study populations.** Samples from 35 case-control series that included patients with ischemic stroke (effective sample size [ $n_e$ ] = 9,396), PAD ( $n_e$  = 5,215), AAA ( $n_e$  = 4,572), VTE ( $n_e$  = 4,607), IA ( $n_e$  = 1,328), and CAD ( $n_e$  = 12,716), as well as from 3,714 subjects with carotid IMT measurements, were analyzed. Samples from 2 cross-sectional studies, including 5,588 subjects who had undergone coronary angiography, were used to assess the association with CAD severity, as well as the association with myocardial infarction (MI), among those with angiographic CAD. The association with age at onset of CAD was assessed in 9,608 cases. See further description in the Online Appendix.

**Ethical approval.** All studies involved in the current analyses were approved by local research ethics committees and informed consent was obtained from all participants.

(R01HL089650-02) and the European Community's Seventh Framework Programme FAD (Fighting Aneurysmal Diseases) project grant agreement HEALTH-F2-2008-200647 and ENGAGE (European Network for Genetic and Genomic Epidemiology) project, grant agreement HEALTH-F4-2007-201413. Funding for the New Zealand sample recruitment was provided by the Health Research Council of New Zealand. The MARTHA (MARseille THrombosis Association study) project was supported by a grant from the Program Hospitalier de Recherche Clinique and the FARIVE (Facteurs de Risque et de Récidives de la Maladie Thromboembolique Veineuse) study by grants from the Fondation pour la Recherche Médicale, the Program Hospitalier de recherche Clinique (PHRC 20002; PHRC2009 RENOVA-TV [REcherche de Nouveaux VAriants de susceptibilité à la Thrombose Veineuse]), the Fondation de France, and the Leducq Foundation. The Spanish VTE study was funded by RECAVA (Red Temática de Investigación Cooperativa en Enfermedades Cardiovasculares)-RD06/0039. The recruitment of patients with abdominal aortic aneurysm and controls from Belgium, Canada, and Pittsburgh, Pennsylvania, was funded in part by grants from the NHLBI (HL064310 to H.K. and HL044682 to R.E.F.). The sample collection at Geisinger Clinic was funded by a grant from the Pennsylvania Commonwealth Universal Research Enhancement program (to D.J.C.), a grant from the Geisinger Clinical Research Fund (to J.R.E.), and a Grant-In-Aid from the American Heart Association (to D.J.C.). The Geisinger MyCode Project was funded in part by a grant from the Ben Franklin Technology Development Fund of PA. AAA-STOP (Simple Treatment Or Prevention) was supported by a grant from the NIH-NHLBI (1P50HL08300 and 1R01HL101388). The Danish study was in part supported by The Lundbeck Foundation Centre for Applied Medical Genomics in Personalised Disease Prediction, Prevention and Care (LuCamp). Collection of data from the Finnish sample with intracranial aneurysm was funded in part by the NINDS (National Institute of Neurological Disease and Stroke) (NS034395 to G.T.), the American Heart Association, Michigan Affiliate (to G.T.), and the University of Kuopio (to A.R.). The PROCARDIS (PreOcioous Coronary ARtery DISease) study was supported by the European Community Sixth Framework Program (LSHM-CT-2007-037273). Drs. Helgadottir, Watkins, and Farrall acknowledge support from the BHF (British Heart Foundation), Centre of Research Excellence, RE/08/004, as does SEH (BHF PG2008/08).

Dr. Samani is supported by BHF and is also an NIHR (National Institute for Health Research) Senior Investigator. Dr. Keavney is supported by a BHF Personal Chair. Drs. Watkins and Farrall acknowledge support from the Wellcome Trust core award to the Wellcome Trust Centre for Human Genetics (090532/Z/09/Z). Dr. Baas was supported by the Dr. E. Dekker program of the Netherlands Heart Foundation (2009T001). Dr. Ruigrok was supported by the NWO (Netherlands Organization for Scientific Research) VENI (Innovational Research Incentives Scheme) grant 916.10.016. Dr. Soria was supported by "Programa d'Estabilització d'Investigadors de la Direcció d'Estratègia i Coordinació del Departament de Salut" (Generalitat de Catalunya). The GAIT (Genetic Analysis of Idiopathic Thrombophilia) project was supported by PI-08/0420, PI-08/0756, SAF2008/01859, RECAVA-RD06/0014 and SGR 01068 from Agència de Gestió d'ajuts Universitaris i de Recerca. The IMPROVE (Carotid Intima Media Thickness (IMT) and IMT-Progression as Predictors of Vascular Events in a High Risk European Population) study was supported by the European Commission (Contract number: QL-G1-CT-2002-00896), the Swedish Heart-Lung Foundation, the Swedish Research Council (projects 8691 and 0593), the Knut and Alice Wallenberg Foundation, the Stockholm County Council (project 562183), Academy of Finland (grant #110413), and the British Heart Foundation (RG2008/014). Drs. Helgadottir, Gretarsdottir, Gudbjartsson, Holm, Thorsteindottir, and Thorleifsson are employees of deCODE genetics, a biotechnology company. deCODE genetics intends to incorporate the variants described in this paper into its genetic testing services. Dr. Thorsteindottir reports that she receives stock options from deCODE genetics. Dr. Witte is an employee of Steno Diabetes Center A/S, owned by Novo Nordisk A/S, and reports that he owns shares in Novo Nordisk A/S. Dr. Wells is a member of the advisory boards of Boehringer-Ingelheim, Pfizer, and Bayer. Dr. Ringelstein is a consultant to Boehringer Ingelheim, Sygnis, Neurobiological Technologies, Novartis, Non Nordisk A/S, Sanofi-Aventis, Bayer Vital, Ma Science, Servier, UCB, and Trommsdorff, and has received travel expenses and honoraria. All other authors have reported that they have no relationships relevant to the contents of this paper to disclose.

Manuscript received August 23, 2011; revised manuscript received December 5, 2011, accepted January 3, 2012.



**Genotyping.** DNA extraction and single SNP genotyping was carried out with various high-throughput methods with laboratory-specific quality-control procedures as detailed in Online Table 1. As the *LPA* SNPs were not represented on available genome-wide chips, genotypes for the Icelandic samples were imputed using methods previously described (21).

**Statistical analysis.** For the main analyses, we assessed the association between the outcome variables and the total number of minor alleles of either SNP (*rs3798220*[C] or *rs10455872*[G]), assuming that both minor alleles have the same effect on the outcome. The effect of each SNP on the binary outcomes, as reported in Online Table 2, was also assessed by comparing it with the effect of the major alleles of both SNPs.

Logistic and linear regression models were used for binary and quantitative outcomes, respectively, to test for linear trend with allele score (*LPA* score defined as the number of minor alleles of *rs3798220* and *rs10455872*) to derive effect estimates, standard errors, and 2-sided p values, assuming additive allelic effects (on the linear or log scale as appropriate). These tests were performed as implemented in NEMO (22), the R-software, and STATA software version 10.

The IMT trait was logarithmically transformed before analysis, and adjusted for age, sex, body mass index, physical exercise, and study center (country of origin). Age at angiography and sex were included as covariates in the linear regression model when fitting the number of affected coronary vessels to the *LPA* score. In the combined analysis, additional adjustments were made for study site and ethnicity.

To allow for the relatedness between subjects in the GAIT (Genetic Analysis of Idiopathic Thrombophilia) studies, the method of generalized estimating equations was used as implemented in the generalized estimating equations package (23) for the R-software. For the Icelandic studies, p values and 95% confidence interval (CI) of ORs are given after adjustment for the relatedness of the subjects and other possible population stratification using the method of genomic control. As the association tests for IMT in the Oxfordshire Family Blood Pressure studies were not significant, no adjustments for relatedness were made.

A fixed-effects meta-analysis approach was used to estimate an inverse-variance weighted average of the ORs from all studies of each outcome category. Tests of whether summary ORs were equal to 1, and whether the true effects in all studies were the same (likelihood ratio chi-square heterogeneity test) were performed. Allelic frequencies of *rs3798220*(C) and *rs10455872*(G), provided in the introduction, are the average frequency of the 15 populations of European origin examined in this study.

Given the number of traits tested, we considered p values <0.0017 significant, assuming a Bonferroni correction factor of 30. The power of the study to detect significant association ( $p \leq 0.0017$ , assuming an OR of 1.3) between *LPA* score and ischemic stroke, PAD, AAA, or VTE, was >97%. The power was slightly less for the ischemic stroke subtypes large artery atherosclerosis (LAA), cardioembo-

lism (CE), or small vessel disease (SVD), or 74%, 78%, and 52%, respectively, and for IA the power was 34%. The power to detect an association between *LPA* score and IMT with effect size  $\geq 0.1$  mm was >99%, at a p value even as low as 0.0005. Prospective studies have suggested that a 0.1-mm increase in IMT is associated with a 20% to 30% increase in risk for developing CAD (24,25), and each *LPA* minor allele increases the risk for CAD by 20% to 30%. An effect size of 0.1 mm per variant allele could therefore be expected if the effect of *LPA* variants on CAD (or other atherosclerotic diseases) was mediated entirely through early atherogenesis and reflected in changes in the carotid artery IMT.

As the case-control ratios differed between studies, the effective sample size ( $n_e$ ), representing the sample size for cases in which the case-control ratio was 1, for each study was calculated as the harmonic mean of the number of cases and controls and is reported in Figures 1 and 2. For the Icelandic studies, the effect sample size was further adjusted for relatedness by dividing it by the genomic control inflation factor. The combined effective sample sizes were used in the power calculations.


## Results

***LPA* SNP association with different vascular traits.** The associations between each SNP and the vascular diseases are shown for all study groups in Online Tables 2A to 2D. As the estimated effects of the 2 SNPs were not significantly different, and the mechanisms by which both SNPs affect Lp(a) have been suggested to be similar (1), for reasons of simplicity (and to increase power) the main analyses (Figs. 1 and 2, Tables 1, 2, and 3) report the effect estimates for the 2 SNPs combined (*LPA* score), which assumes that their effects are equal. *LPA* score results for each study group are detailed in Online Tables 3A to 3C.

Results from all studies combined show nominally significant association between the *LPA* score and ischemic stroke overall, with an estimated OR of 1.10 (95% CI: 1.02 to 1.18;  $p = 0.016$ ) (Fig. 1). Ischemic stroke subtype analysis revealed that the association is restricted to LAA (OR: 1.27; 95% CI: 1.11 to 1.46;  $p = 6.7 \times 10^{-4}$ ), as the effects of *LPA* variants on the CE and SVD subtypes were nonsignificant ( $p > 0.50$ ). The *LPA* score was associated

### Abbreviations and Acronyms

<b>AAA</b>	= abdominal aortic aneurysm
<b>CAD</b>	= coronary artery disease
<b>CE</b>	= cardioembolism
<b>CI</b>	= confidence interval
<b>IA</b>	= intracranial aneurysm
<b>IMT</b>	= intima-media thickness
<b>LAA</b>	= large artery atherosclerosis
<b>Lp(a)</b>	= apolipoprotein (a)
<b>LPA</b>	= apolipoprotein(a) gene
<b>MI</b>	= myocardial infarction
<b><math>n_e</math></b>	= effective sample size
<b>OR</b>	= odds ratio
<b>PAD</b>	= peripheral arterial disease
<b>SNP</b>	= single nucleotide polymorphism
<b>SVD</b>	= small vessel disease
<b>TOAST</b>	= Trial of Org 10172 in Acute Stroke Treatment
<b>VTE</b>	= venous thromboembolism




**Figure 1** Association of LPA Score With Vascular Diseases

Forest plot of the associations of apolipoprotein(a) gene (*LPA*) score with ischemic stroke, and TOAST (Trial of Org 10172 in Acute Stroke Treatment) subtypes (large artery atherosclerosis [LAA], cardioembolism [CE], and small vessel disease [SVD]), peripheral arterial disease (PAD), abdominal aortic aneurysm (AAA), venous thromboembolism (VTE), coronary artery disease (CAD), and intracranial aneurysm (IA) in individuals of European origin. Combined odds ratios (ORs) from multiple studies are indicated by squares, with an area in proportion to the sample size, and with horizontal lines representing the 95% confidence intervals (CIs). The plot shows the effective sample sizes ( $n_e$ ) and p values for the associations.

with both PAD (OR: 1.47; 95% CI: 1.33 to 1.62;  $p = 2.9 \times 10^{-14}$ ) and AAA (OR: 1.23; 95% CI: 1.11 to 1.36;  $p = 6.0 \times 10^{-5}$ ). The previously reported association between CAD and *LPA* score (1) in individuals of European ancestry was confirmed (OR: 1.32; 95% CI: 1.24 to 1.42;  $p = 1.1 \times 10^{-15}$ ). In addition, the *LPA* score was associated with CAD in African Americans (OR: 2.49; 95% CI: 1.08 to 5.72;  $p = 0.032$ ) (Online Table 4). In contrast, the *LPA* score was not associated with VTE ( $p = 0.63$ ), IA ( $p = 0.15$ ) (Fig. 1), or mean common carotid IMT ( $p = 0.083$ ) (Table 1).

**The association of *LPA* score with PAD, LAA, and AAA among patients with and without concomitant CAD.** Given the substantial overlap between CAD and the other atherosclerotic diseases, it is conceivable that the associations of the *LPA* variants with LAA stroke, PAD, and AAA are mediated through (or inflated by) the association of these diseases with CAD. Therefore, we reanalyzed the associations between the *LPA* variants and the atherosclerotic phenotypes in cases with and without a history of CAD in studies in which this information was available



**Figure 2** Association of LPA Score With CAD, With and Without Atherosclerosis in Other Vascular Territories, and With Age-Group at Onset of CAD

**(A)** Forest plot of the association of *LPA* score with CAD in Iceland, for cases with concomitant cardiovascular disease (CVD; including LAA, PAD, or AAA) and for CAD after excluding cases with known CVD.  $n_e$  Values, ORs, 95% CIs and p values are shown. p Values contrasting the 2 groups of CAD cases are shown (\*). **(B)** Forest plot of the association of *LPA* score with early onset CAD (diagnosed before the age of 50 years in men and 60 years in women), intermediate onset CAD (diagnosed at the age of 50 to <70 years in men and 60 to <75 years in women), and late onset CAD (diagnosed after age 70 years in men and 75 years in women). Individuals of European origin from Iceland and Atlanta, Georgia, were combined using an inverse variance-weighted meta-analysis approach (fixed-effects model). Abbreviations as in Figure 1.

**Table 1 Association of LPA Score With Carotid IMT\***

Study Group	N	IMT <sub>log</sub> (mm)		p Value	p <sub>het</sub> Value
		β	SE		
IMPROVE	2,984	-0.0064	0.0036	0.076	—
Oxfordshire Family Blood Pressure Study	730	-0.0002	0.0224	0.99	—
Combined	3,714	-0.0060	0.0036	0.083	0.70

\*Linear regression coefficients (β) with corresponding standard errors (SE) and p values, assuming additive allelic effects. The model included age, sex, body mass index, and physical exercise as covariates. Additional adjustments were made for study site (country) for the IMPROVE study.

IMPROVE = Carotid Intima Media Thickness and IMT-Progression as Predictors of Vascular Events in a High Risk European Population; IMT = intima-media thickness; LPA = apolipoprotein(a) gene; p<sub>het</sub> = p value for heterogeneity test.

(Online Tables 5A and 5B). After the exclusion of data from patients with CAD, the effect estimates for LPA score became lower for PAD (OR: 1.17; p = 0.12) and AAA (OR: 1.11; p = 0.16), suggesting a stronger association of the LPA score with atherosclerotic disease manifested in more than 1 vascular bed, although the effect estimate was not lowered for LAA (OR: 1.30; p = 0.013) (Online Table 5A).

**Association with CAD, with and without atherosclerosis in other vascular territories.** To investigate whether LPA score increased the risk for a widespread atherosclerotic disease, we divided the Icelandic CAD cases into those with confirmed atherosclerosis related disease in other vascular beds (PAD, AAA, or LAA), and those without a known history of these diseases. While, for CAD with concomitant PAD, AAA, and/or LAA the OR was 1.54 (p = 8.9 × 10<sup>-9</sup>), the OR for CAD without known PAD, AAA, and/or LAA was 1.22 (p = 2.4 × 10<sup>-7</sup>). On direct comparison of the 2 groups of CAD cases, each LPA risk allele increased the risk for atherosclerosis in other vascular beds by 26% (p = 0.0010) (Fig. 2).

**Association with angiographic extent of CAD and with MI among patients with angiographic CAD.** Online Table 6 shows the distribution of the number of affected (>50% stenosis) coronary index vessels, and the proportions with MI among 5,588 subjects who had undergone coronary angiography. After adjusting for sex, age at angiogra-

**Table 2 Association of LPA Score With Angiographic CAD Severity\***

Study Group	N	No. of Affected Coronary Index Vessels		p Value
		β	SE	
Iceland	2,330	0.245	0.061	5.8 × 10 <sup>-5</sup>
Atlanta-European American	2,718	0.281	0.052	6.3 × 10 <sup>-8</sup>
Atlanta-African American	540	0.396	0.219	0.071
Combined	5,588	0.267	0.038	4.8 × 10 <sup>-12</sup>

\*Linear regression coefficients (β) with corresponding standard errors (SE) and p values, assuming additive allelic effects. Age at angiography and sex were included as covariates in the model for each study group. In the combined analysis additional adjustments were made for study group (site/ethnicity).  
CAD = coronary artery disease; LPA = apolipoprotein(a) gene.

**Table 3 Association Between LPA Score and the Age at Diagnosis of CAD\***

Parameter	n	Age at Diagnosis		p Value
		$\beta$	SE	
European ancestry				
Age at first CAD diagnosis	9,275	-1.581	0.295	$8.2 \times 10^{-8}$
Age at first MI	5,148	-0.918	0.416	0.028
Age at first CAD diagnosis for MI cases	5,148	-1.334	0.400	0.00085
African Americans				
Age at first CAD diagnosis	333	-3.541	2.521	0.16
Age at first MI	185	-2.221	3.249	0.50
Age at first CAD diagnosis for MI cases	185	-6.009	3.159	0.059

\*Linear regression coefficients (β) with corresponding standard errors (SE) and p values, assuming additive allelic effects. The model included sex as a covariate. In addition, study site (Iceland/Atlanta) was a covariate in the model when analyzing those of European ancestry.

MI = myocardial infarction. Other abbreviations as in Table 2.

phy, study site, and ethnicity, in a linear regression model, each LPA risk allele increased the number of diseased vessels by a mean of 0.267 (p = 4.8 × 10<sup>-12</sup>). Effect estimates for African Americans were similar to those of patients of European origin (Table 2). In a logistic regression model comparing patients with angiographic CAD with MI (n = 1,817) or without MI (n = 1,908), the LPA score was not associated with MI after adjusting for the same variables and the age at first CAD diagnosis (OR: 0.99; p = 0.90).

**Association with age of onset of CAD and MI.** The correlation between LPA score and the age at diagnosis of CAD was tested, with adjustments for sex and study site (Table 3, Fig. 2). Each LPA risk allele was associated with a mean of 1.58 years' earlier diagnosis of CAD (p = 8.2 × 10<sup>-8</sup>) among 9,276 cases of European origin from Iceland and Atlanta, Georgia (Online Table 4). Restricting data to those from cases with MI diagnoses, the corresponding reduction in age at diagnosis of first MI was 0.92 years (p = 0.028). Effect estimates for age at first CAD diagnosis among 333 African Americans were consistent with those observed for whites, but did not reach nominal significance levels.

## Discussion

We report an association between 2 Lp(a)-related SNPs (combined and measured as LPA score), and the PAD, AAA, and LAA subtype of ischemic stroke, with each variant allele increasing the risk by 47%, 23%, and 27% for the 3 diseases, respectively. In contrast, our analyses show no association between the LPA variants and VTE or CE and SVD stroke subtypes. We show that patients with CAD carrying LPA risk alleles have increased susceptibility to atherosclerotic manifestations outside of the coronary tree and are more likely to be diagnosed earlier with CAD than are CAD cases not carrying these variants. Further, our study provides evidence for an association between the LPA variants and the number of coronary arteries (out of 4 index

vessels) with >50% stenosis. Thus the *LPA* variants, and by implication, Lp(a) levels, have an effect on the atherosclerotic burden of large vessels throughout the arterial tree. To the best of our knowledge, relationships between sequence variants affecting Lp(a) levels and either systemic distribution of atherosclerosis or angiographic CAD severity, have not been reported before. However, several studies, that in general have been limited by small sample sizes, have assessed the association between Lp(a) levels and severity of CAD and provided conflicting results (26–30).

In apparent contrast to the correlation between *LPA* score and earlier age at onset of CAD, *LPA* score was not associated with IMT, a marker for subclinical atherosclerosis (31,32). This is consistent with results from a recent population-based cohort study showing that Lp(a) levels (and SNPs correlated with Lp[a]), were not associated with IMT (33), and in line with a recent meta-analysis of genome-wide association studies from the CHARGE (Cohorts for Heart and Aging Research in Genomic Epidemiology) consortium (34) that found little evidence for genetic overlap across carotid IMT and CAD. We suggest that the effects of the *LPA* variants, and thus Lp(a) levels, on the risk for atherosclerotic manifestations are mediated at later stages in the pathogenic pathway and are thus not reflected by variation in carotid IMT.

Our study was limited by the fact that measurements of Lp(a) levels were not available, rendering it impossible to show directly that the association of the *LPA* risk variants with atherosclerotic phenotypes is mediated through Lp(a) levels. Therefore, the possibility that the risk conferred by the *LPA* score is mediated through mechanisms other than increased Lp(a) levels, cannot be disregarded. However, previous studies have shown a strong and consistent association between the 2 *LPA* variants and Lp(a) levels (1,35), and the association between the *LPA* risk variants and CAD has been shown to disappear with Lp(a) level adjustments (1), strongly supporting the view that the effect on atherosclerotic phenotypes is mediated through Lp(a) levels.

Lp(a) has been implicated in both atherogenesis and thrombosis. We show that *LPA* risk alleles that correlate with high plasma Lp(a) levels were associated with vascular diseases with a strong atherosclerotic component, such as CAD, PAD and LAA, and AAA, but we failed to demonstrate an association between *LPA* alleles and diseases without an atherosclerotic etiology, such as VTE and IA, or diseases less related to atherosclerosis, such as the CE and SVD stroke subtypes. Furthermore, among patients with angiographic CAD, after adjustments for age at onset of CAD, the *LPA* variants were not associated with MI, indicating that the risk is mediated through atherosclerotic plaque deposition, rather than plaque rupture, or subsequent thrombosis. Given the assumption that the risk for *LPA* variants on vascular diseases is mediated through Lp(a) levels, our results suggest that the implicit thrombogenic properties of Lp(a) (4,5) are insufficient to increase the risks

for the thrombotic stroke subtypes CE and SVD, or to promote VTE.

## Conclusions

We report an association between 2 variants in the *LPA* gene and 3 atherosclerosis-related diseases (i.e., the LAA subtype of ischemic stroke, PAD, and AAA). In addition to replicating the previously reported association between the *LPA* variants and CAD, we report a correlation between the *LPA* variants and the number of obstructed coronary arteries. Furthermore, the risk variants were associated with an earlier diagnosis of CAD. Evaluating *LPA* risk alleles may therefore contribute to more effective primary and secondary prevention of atherosclerotic disease. In contrast, we found no association with the CE and SVD ischemic stroke subtypes, which have less obvious atherosclerotic components, or VTE, a primarily thrombotic disease. These findings imply that the risk conferred by high Lp(a) levels is mediated through the atherosclerotic, rather than the thrombotic, aspects of vascular disease.

## Acknowledgments

The authors thank the individuals who participated in the study and whose contribution made this work possible.

**Reprint requests and correspondence:** Dr. Anna Helgadottir, DeCODE Genetics, Sturlugata 8, IS-101 Reykjavik, Iceland. E-mail: annah@decode.is.

## REFERENCES

- Clarke R, Peden JF, Hopewell JC, et al. Genetic variants associated with Lp(a) lipoprotein level and coronary disease. *N Engl J Med* 2009;361:2518–28.
- Erqou S, Kaptoge S, Perry PL, et al. Lipoprotein(a) concentration and the risk of coronary heart disease, stroke, and nonvascular mortality. *JAMA* 2009;302:412–23.
- Erqou S, Thompson A, Di Angelantonio E, et al. Apolipoprotein(a) isoforms and the risk of vascular disease: systematic review of 40 studies involving 58,000 participants. *J Am Coll Cardiol* 2010;55:2160–7.
- Caplice NM, Panetta C, Peterson TE, et al. Lipoprotein(a) binds and inactivates tissue factor pathway inhibitor: a novel link between lipoproteins and thrombosis. *Blood* 2001;98:2980–7.
- Miles LA, Fless GM, Levin EG, Scanu AM, Plow EF. A potential basis for the thrombotic risks associated with lipoprotein(a). *Nature* 1989;339:301–3.
- Schreiner PJ, Morrisett JD, Sharrett AR, et al. Lipoprotein[a] as a risk factor for preclinical atherosclerosis. *Arterioscler Thromb* 1993;13:826–33.
- Sotiriou SN, Orlova VV, Al-Fakhri N, et al. Lipoprotein(a) in atherosclerotic plaques recruits inflammatory cells through interaction with Mac-1 integrin. *Faseb J* 2006;20:559–61.
- Luke MM, Kane JP, Liu DM, et al. A polymorphism in the protease-like domain of apolipoprotein(a) is associated with severe coronary artery disease. *Arterioscler Thromb Vasc Biol* 2007;27:2030–6.
- Hopewell JC, Clarke R, Parish S, et al. Lipoprotein(a) genetic variants associated with coronary and peripheral vascular disease, but not with stroke risk in the Heart Protection Study. *Circ Cardiovasc Genet* 2011;4:68–73.
- Cheng SW, Ting AC, Wong J. Lipoprotein(a) and its relationship to risk factors and severity of atherosclerotic peripheral vascular disease. *Eur J Vasc Endovasc Surg* 1997;14:17–23.



11. Volpato S, Vigna GB, McDermott MM, et al. Lipoprotein(a), inflammation, and peripheral arterial disease in a community-based sample of older men and women (the InCHIANTI study). *Am J Cardiol* 105:1825–30.
12. Jones GT, van Rij AM, Cole J, et al. Plasma lipoprotein(a) indicates risk for 4 distinct forms of vascular disease. *Clin Chem* 2007;53:679–85.
13. Takagi H, Manabe H, Kawai N, Goto SN, Umemoto T. Circulating lipoprotein(a) concentrations and abdominal aortic aneurysm presence. *Interact Cardiovasc Thorac Surg* 2009;9:467–70.
14. Pradhan AD, Shrivastava S, Cook NR, Rifai N, Creager MA, Ridker PM. Symptomatic peripheral arterial disease in women: nontraditional biomarkers of elevated risk. *Circulation* 2008;117:823–31.
15. Ridker PM, Stampfer MJ, Rifai N. Novel risk factors for systemic atherosclerosis: a comparison of C-reactive protein, fibrinogen, homocysteine, lipoprotein(a), and standard cholesterol screening as predictors of peripheral arterial disease. *JAMA* 2001;285:2481–5.
16. Aboyans V, Criqui MH, Denenberg JO, Knoke JD, Ridker PM, Fronck A. Risk factors for progression of peripheral arterial disease in large and small vessels. *Circulation* 2006;113:2623–9.
17. Sofi F, Marcucci R, Abbate R, Gensini GF, Prisco D. Lipoprotein(a) and venous thromboembolism in adults: a meta-analysis. *Am J Med* 2007;120:728–33.
18. Tsai AW, Cushman M, Rosamond WD, Heckbert SR, Polak JF, Folsom AR. Cardiovascular risk factors and venous thromboembolism incidence: the longitudinal investigation of thromboembolism etiology. *Arch Intern Med* 2002;162:1182–9.
19. Jones GT, Deng M, Hammond-Tooke GD, McCormick SP, van Rij AM. Increased plasma lipoprotein(a) found in large-artery atherosclerotic, but not small-artery occlusive, stroke. *Clin Chem* 2009;55:1888–90.
20. Ohira T, Shahar E, Chambless LE, Rosamond WD, Mosley TH Jr., Folsom AR. Risk factors for ischemic stroke subtypes: the Atherosclerosis Risk in Communities study. *Stroke* 2006;37:2493–8.
21. Kong A, Masson G, Frigge ML, et al. Detection of sharing by descent, long-range phasing and haplotype imputation. *Nat Genet* 2008;40:1068–75.
22. Gretarsdottir S, Thorleifsson G, Reynisdottir ST, et al. The gene encoding phosphodiesterase 4D confers risk of ischemic stroke. *Nat Genet* 2003;35:131–8.
23. Chen MH, Yang Q. GWAF: an R package for genome-wide association analyses with family data. *Bioinformatics* 26:580–1.
24. Bots ML, Hoes AW, Koudstaal PJ, Hofman A, Grobbee DE. Common carotid intima-media thickness and risk of stroke and myocardial infarction: the Rotterdam Study. *Circulation* 1997;96:1432–7.
25. Hodis HN, Mack WJ, LaBree L, et al. The role of carotid arterial intima-media thickness in predicting clinical coronary events. *Ann Intern Med* 1998;128:262–9.
26. Dahlen GH, Guyton JR, Attar M, Farmer JA, Kautz JA, Gotto AM Jr. Association of levels of lipoprotein Lp(a), plasma lipids, and other lipoproteins with coronary artery disease documented by angiography. *Circulation* 1986;74:758–65.
27. Korhonen T, Savolainen MJ, Koistinen MJ, et al. Association of lipoprotein cholesterol and triglycerides with the severity of coronary artery disease in men and women. *Atherosclerosis* 1996;127:213–20.
28. Zampoulakis JD, Kyriakousi AA, Poralis KA, et al. Lipoprotein(a) is related to the extent of lesions in the coronary vasculature and to unstable coronary syndromes. *Clin Cardiol* 2000;23:895–900.
29. Wang XL, Tam C, McCredie RM, Wilcken DE. Determinants of severity of coronary artery disease in Australian men and women. *Circulation* 1994;89:1974–81.
30. Gazzaruso C, Geroldi D, Garzaniti A, et al. Apolipoprotein(a) phenotypes as genetic markers of coronary atherosclerosis severity. *Int J Cardiol* 1998;64:277–84.
31. O'Leary DH, Polak JF, Kronmal RA, Manolio TA, Burke GL, Wolfson SK Jr. Carotid-artery intima and media thickness as a risk factor for myocardial infarction and stroke in older adults. Cardiovascular Health Study Collaborative Research Group. *N Engl J Med* 1999;340:14–22.
32. Rohani M, Jogestrand T, Ekberg M, et al. Interrelation between the extent of atherosclerosis in the thoracic aorta, carotid intima-media thickness and the extent of coronary artery disease. *Atherosclerosis* 2005;179:311–6.
33. Kivimaki M, Magnussen CG, Juonala M, et al. Conventional and Mendelian randomization analyses suggest no association between lipoprotein(a) and early atherosclerosis: the Young Finns Study. *Int J Epidemiol* 2011;40:470–8.
34. Bis JC, Kavousi M, Franceschini N, et al. Meta-analysis of genome-wide association studies from the CHARGE consortium identifies common variants associated with carotid intima media thickness and plaque. *Nat Genet* 2011;43:940–7.
35. Ronald J, Rajagopalan R, Cerrato F, et al. Genetic variation in LPAL2, LPA, and PLG predicts plasma lipoprotein(a) level and carotid artery disease risk. *Stroke* 2011;42:2–9.

**Key Words:** association ■ atherosclerosis ■ genetic ■ lipoprotein(a) ■ thrombosis.

#### APPENDIX:

For supplementary tables and text, please see the online version of this article.