



Stochastic shell models driven by a multiplicative fractional Brownian-motion



Hakima Bessaih^{a,*}, María J. Garrido-Atienza^b, Björn Schmalfuss^c

^a Department of Mathematics, University of Wyoming, Laramie 82071, USA

^b Dpto. Ecuaciones Diferenciales y Análisis Numérico, Universidad de Sevilla, Apdo. de Correos 1160, 41080-Sevilla, Spain

^c Institut für Stochastik, Friedrich Schiller Universität Jena, Ernst Abbe Platz 2, 77043, Jena, Germany

HIGHLIGHTS

- We study a shell model driven by a fractional Brownian motion.
- We prove the existence and uniqueness of a global mild solution.
- The existence method is based on a compactness argument.
- Fractional calculus is used throughout the paper.

ARTICLE INFO

Article history:

Received 28 October 2014

Received in revised form

20 January 2016

Accepted 26 January 2016

Available online 4 February 2016

Communicated by Edriss S. Titi

Keywords:

Stochastic PDEs

Fractional Brownian-motion

Pathwise solutions

Fractional calculus

ABSTRACT

We prove existence and uniqueness of the solution of a stochastic shell-model. The equation is driven by an infinite dimensional fractional Brownian-motion with Hurst-parameter $H \in (1/2, 1)$, and contains a non-trivial coefficient in front of the noise which satisfies special regularity conditions. The appearing stochastic integrals are defined in a fractional sense. First, we prove the existence and uniqueness of variational solutions to approximating equations driven by piecewise linear continuous noise, for which we are able to derive important uniform estimates in some functional spaces. Then, thanks to a compactness argument and these estimates, we prove that these variational solutions converge to a limit solution, which turns out to be the unique pathwise mild solution associated to the shell-model with fractional noise as driving process.

© 2016 Elsevier B.V. All rights reserved.

1. Introduction

In this paper we consider some shell-models under the influence of a noise. Shell-models of turbulence describe the evolution of complex Fourier-like components of a scalar velocity field $u_n(t) \in \mathbb{C}$ and the associated wavenumbers k_n , where the discrete index n is referred as the shell-index. The evolution of the infinite sequence $(u_n)_{n \in \mathbb{N}}$ is given by

$$\dot{u}_n(t) + \nu k_n^2 u_n(t) + b_n(u(t), u(t)) = g_n(t, u(t)) \dot{\omega}(t), \quad n \in \mathbb{N} \quad (1.1)$$

with the constraints $u_{-1}(t) = u_0(t) = 0$ and $u_n(t) \in \mathbb{C}$ for $n \in \mathbb{N}$. $\dot{\omega}$ denotes a noise path that will be described below. Here $\nu \geq 0$ and, in analogy with Navier–Stokes-equations, ν represents a kinematic viscosity; $k_n = k_0 \lambda^n$ ($k_0 > 0$ and $\lambda > 1$) and g_n is a forcing term. The exact form of $b_n(u, v) \in \mathbb{C}$ varies from one model to another. However in all various models, it is assumed that $b_n(u, v)$ is chosen in such a way that

$$\Re \sum_{n=1}^{\infty} b_n(u, v) \bar{v}_n = 0, \quad (1.2)$$

* Corresponding author.

E-mail addresses: bessaih@uwyo.edu (H. Bessaih), mgarrido@us.es (M.J. Garrido-Atienza), bjoern.schmalfuss@uni-jena.de (B. Schmalfuss).

where \Re denotes the real part and \bar{x} the complex conjugate of x . Eq. (1.2) implies a formal law of conservation of energy in the inviscid ($\nu = 0$) and unforced form of (1.1). In particular, we define the bilinear terms b_n as

$$b_n(u, v) = i(ak_{n+1}\bar{u}_{n+1}\bar{v}_{n+2} + bk_n\bar{u}_{n-1}\bar{v}_{n+1} - ak_{n-1}\bar{u}_{n-1}\bar{v}_{n-2} - bk_{n-1}\bar{u}_{n-2}\bar{v}_{n-1})$$

in the GOY-model (see [1,2]) and by

$$b_n(u, v) = -i(ak_{n+1}\bar{u}_{n+1}v_{n+2} + bk_n\bar{u}_{n-1}v_{n+1} + ak_{n-1}u_{n-1}v_{n-2} + bk_{n-1}u_{n-2}v_{n-1})$$

in the SABRA-model (see [3]). The two parameters a, b are real numbers and $u = (u_n)_{n \in \mathbb{N}}, v = (v_n)_{n \in \mathbb{N}}$. There are several shell-models in literature, the GOY- and SABRA-models defined above have been introduced in [1–3]. The viscous version of the GOY- and SABRA-models, well posedness, global regularity of solutions and smooth dependence on the initial data can be found in [4].

In recent years, shell-models of turbulence have attracted a lot of interest for their ability to capture some of the statistical properties of the three-dimensional turbulence while presenting a structure much simpler than the Navier–Stokes-equations. The stochastic version of the GOY-model under the influence of an additive white noise has been studied in [5] where some statistical properties in terms of the invariant measure have been shown. For the same model in [6,7] a Gaussian invariant measure is associated and a flow constructed.

The introduction of randomness in the Shell model arises from a need to understand the velocity fluctuations observed under identical experimental conditions. Random body forces also arise as control terms, or from random disturbances such as structural vibrations that act on the fluid. In the case of the Navier–Stokes equations, Kolmogorov was the first to introduce white noise in the Navier–Stokes equations in order to obtain an invariant measure for the system. In this article we consider a long term multiplicative noise allowing us to model some memory effects. Such a noise is given by a trace-class fractional Brownian-motion in our state space with Hurst-parameter $H \in (1/2, 1)$. Our motivation in studying this problem comes from the fact that the Hurst parameter might be used in studying or modeling the exponent of the associated structure functions. A first attempt can be found in [5] on a dyadic model by using a white noise. In addition to noise memory length, fractional Brownian-motion is self-similar with parameter H , and when combining this with possible scaling in the space parameter, other regularity properties can be realized, see [8] for details. We hope to perform in the near future some numerical simulations on this model around this question. On the mathematical side, in contrast to white noise, a fractional-Brownian motion is not a martingale, and therefore the multiplicative noise term cannot be presented by an Itô-integral. However, to deal with stochastic integration where the integrator is only Hölder-continuous with an exponent larger than $1/2$, one can use the Young-integration, see Young [9] or the adaptation to a stochastic set up by Zähle [10]. Since the definition of these integrals is based on fractional derivatives (see Samko et al. [11] for a general presentation), this theory is often called fractional calculus. An advantage of this theory is, and in contrast to the Itô-integral which is given in general by a limit in probability of Darboux-sums derived from an adapted integrand, that we can define our integral pathwise which means that for any sufficiently regular integrand and integrator the integral is well defined. Or in other words, the exceptional sets of measure zero which appear in the classical Itô-integration does not depend on the integrand. Moreover, integrals can be defined for non-adapted integrands.

The main issue of our work is to prove existence and uniqueness of a pathwise solution of the stochastic shell-model driven by a fractional multiplicative noise. Applying an infinite dimensional version of the fractional integration theory we are able to present (1.1) in a mild sense where the last term of this equation defines a fractional integral. In particular, the properties of the nonlinear term B generated by the sequence $(b_i(u, v))_{i \in \mathbb{N}}$ allow to present such a solution in a mild form. Nevertheless, in a first step we replace the fractional noise path by a piecewise linear continuous approximation. Considering (1.1) with such a noise path, we are able to construct global and unique mild solutions. It is important to emphasize that the classical contraction method cannot be used alone since the bilinear term $(b_i(u, v))_{i \in \mathbb{N}}$ causes to have estimates that do not close with the right norms. This is why we have first to construct weak solutions and get some a priori estimates. These weak solutions have to be constructed with a smoother path noise in order to define the corresponding stochastic integral. The a priori estimates combined with the estimates obtained from the mild form are then used to pass to the limit by means of a compactness argument, and the limit will turn out to be a mild solution of the original problem. The uniqueness of solutions is proved by an argument that uses the balance of suitable norms. As we mentioned before, just using the mild form in its usual norm does not allow to close the estimates. For that reason we combine the a priori estimates and the norms obtained from the mild form to solve an algebraic system of two inequalities where the unknown is given in terms of the difference of two mild solutions starting from the same initial condition but in two different norms. The solution of this system is zero and this is what allows us to conclude the uniqueness of solutions. We believe that our result on existence of solutions can be generalized to the Navier–Stokes equations although careful calculations have to be performed on the nonlinear term which is the main difference with the current result. We might have to work in slightly different spaces, and this will be done in the forthcoming paper [12].

Articles dealing with pathwise solutions for quite general stochastic ordinary differential equations driven by a multiplicative fractional-Brownian motion are, e.g., [13] and [14]. In the infinite dimensional context, there are also articles studying the existence of pathwise solutions, like [15] (dealing with variational solutions) and [16–18] and [19], for the mild solution. In these papers the Hurst-parameter $H \in (1/2, 1)$, the diffusion and the drift are assumed to be Lipschitz-continuous and the existence of solutions is proved using pathwise arguments through the fractional integrals. On the other hand, there is an extensive literature for fluid flows driven by a Brownian-motion but only a few papers where the noise is a fractional Brownian-motion. For instance, in [20] the considered fractional Brownian-motion has Hurst-parameter bigger than $1/2$, and the authors find a local solution of the 3D Navier–Stokes-equation by using the Young-integral. In [8] the 2D Navier–Stokes-equation driven by a fractional Brownian-motion with no restriction on the values of the Hurst-parameter is studied, but the considered noise is additive.

An interesting advantage of considering the existence of pathwise solutions for the stochastic shell-model is that they will generate a random dynamical system, which gives us the possibility to an intensive asymptotic analysis of (1.1). In particular, this is the foundation to show the existence of random attractors and the analysis of their structure. In the forthcoming paper [12] the dynamics of the stochastic shell-model is investigated by using the random dynamical system theory. The generation of a random dynamical system as well as the study of the corresponding random attractor for another kind of stochastic evolution equations with multiplicative fractional noise have been very recently investigated in the papers [19,21], and [14].

We also would like to point out that, despite the fact that there are similarities between the 2D Navier–Stokes-equation and the shell-model, more effort and more involved techniques will be necessary to obtain similar results for the stochastic 2D Navier–Stokes-equation than the ones proved in this paper.

The paper is organized as follows: in Section 2 we introduce the functional analytical framework. In Section 3, we define the fractional derivatives and the stochastic integral using some type of generalized Young-integrals. In Section 4, we introduce the different assumptions on the diffusion and give the definitions of the different solutions. In Section 5, we prove that the system driven by a smoother path has a unique weak solution, that it is also a mild solution. Furthermore, we obtain some fundamental uniform estimates for the solution of the system driven by such a kind of smooth path. In Section 6, thanks to these uniform estimates and a compactness reasoning we construct a unique pathwise mild solution to the shell-model having a fractional Brownian-motion as driving path. Section 7 is devoted to an example of a particular diffusion fitting the assumptions required for developing the abstract framework. Finally, Appendix contains the proofs of some results that have been used in different sections of the paper.

As usual we denote by c a positive constant that can change their value from line to line.

2. Preliminaries

2.1. Spaces and operators

For any $\alpha \in \mathbb{R}$, let us introduce the following spaces, see Constantin et al. [4] for the details,

$$V_\alpha = \left\{ u = (u_1, u_2, \dots) \in \mathbb{C}^\infty : \sum_{n=1}^{\infty} k_n^{4\alpha} |u_n|^2 < \infty \right\}^1,$$

where we remind that $(k_n)_{n \in \mathbb{N}}$ are the associated wavenumbers to $(u_n)_{n \in \mathbb{N}}$. The space V_α is a separable Hilbert-space with scalar product $(u, v)_{V_\alpha} = \sum_{n=1}^{\infty} k_n^{4\alpha} u_n \bar{v}_n$. Denote by $\|\cdot\|_{V_\alpha}$ its norm. We have the compact embedding

$$V_{\alpha_1} \subset V_{\alpha_2} \quad \text{if } \alpha_1 > \alpha_2.$$

Let us denote by $V := V_0$ and its norm simply by $\|\cdot\|$ and its scalar product by $(\cdot, \cdot)_V$.

Let $A : D(A) = V_1 \rightarrow V$ be the linear unbounded operator defined as

$$A : (u_1, u_2, \dots) \mapsto (-\nu k_1^2 u_1, -\nu k_2^2 u_2, \dots).$$

For simplicity let us set $\nu = 1$. It is known that A generates an analytic semigroup $S(\cdot)$ which follows from the Lax–Milgram lemma, see Sell and You [23, Theorem 36.6], and this semigroup is exponentially stable. Furthermore, $V_\alpha = D(A^\alpha)$ and $(u, v)_{V_\alpha} = (A^\alpha u, A^\alpha v)_V$, $u, v \in V_\alpha$.

Let $L(V_\delta, V_\gamma)$ denote the space of linear continuous operators from V_δ into V_γ . As usual, $L(V)$ denotes $L(V, V)$. The following properties are well known for analytic semigroups and their generators: for $\zeta \geq \alpha$ there exists a constant $c > 0$ such that

$$|S(t)|_{L(V_\alpha, V_\zeta)} = |A^\zeta S(t)|_{L(V_\alpha, V)} \leq \frac{c}{t^{\zeta-\alpha}} e^{-\lambda t}, \quad t > 0, \quad (2.1)$$

$$|S(t) - \text{id}|_{L(V_{\sigma+\nu}, V_{\theta+\nu})} \leq ct^{\sigma-\theta}, \quad \text{for } \sigma \in [\theta, 1 + \theta], \nu \in \mathbb{R}, \quad (2.2)$$

where λ in (2.1) is a positive constant, see for instance Pazy [24, Theorem 2.6.13]. From these inequalities, for $\nu, \eta \in [0, 1], \zeta, \delta \in \mathbb{R}$ such that $\delta \leq \zeta + \nu$, there exists a $c > 0$ such that for $0 \leq q \leq r \leq s \leq t$,

$$\begin{aligned} |S(t-r) - S(t-q)|_{L(V_\delta, V_\zeta)} &\leq c(r-q)^\nu (t-r)^{-\nu-\zeta+\delta}, \\ |S(t-r) - S(s-r) - S(t-q) + S(s-q)|_{L(V)} &\leq c(t-s)^\nu (r-q)^\eta (s-r)^{-(\nu+\eta)}. \end{aligned} \quad (2.3)$$

Define the bilinear operator $B : \mathbb{C}^\infty \times \mathbb{C}^\infty \rightarrow \mathbb{C}^\infty$ as

$$B(u, v) = -(b_1(u, v), b_2(u, v), \dots)$$

where the components b_n satisfy (1.2).

B is well defined when its domain is $V_{1/2} \times V$ or $V \times V_{1/2}$ (see [4]), that is, $B : V_{1/2} \times V \rightarrow V$ and $B : V \times V_{1/2} \rightarrow V$ are bounded operators. The operator B enjoys the following properties

$$\begin{aligned} \Re(B(u, v), w)_V &= -\Re(B(u, w), v)_V, \quad u \in V_{1/2}, v, w \in V, \\ \Re(B(u, v), w)_V &= -\Re(B(u, w), v)_V, \quad u \in V, v, w \in V_{1/2}. \end{aligned}$$

As a consequence, we also have that

$$\Re(B(u, v), v)_V = 0, \quad u \in V, v \in V_{1/2}. \quad (2.4)$$

Moreover, we extend the result of Constantin et al. [4] to more general spaces:

Lemma 2.1. For any $\alpha_1, \alpha_2, \alpha_3 \in \mathbb{R}$

$$B : V_{\alpha_1} \times V_{\alpha_2} \rightarrow V_{-\alpha_3} \quad \text{with } \alpha_1 + \alpha_2 + \alpha_3 \geq \frac{1}{2}$$

and there exists a constant c depending on the α_j 's such that

$$\|B(u, v)\|_{V_{-\alpha_3}} \leq c \|u\|_{V_{\alpha_1}} \|v\|_{V_{\alpha_2}}, \quad u \in V_{\alpha_1}, v \in V_{\alpha_2}.$$

The proof of this result follows by Proposition 1 of Constantin et al. [22], and Bessaih and Ferrario [6] and hence we omit it here.

¹ Here there is an important difference w.r.t. the notation of spaces in [4] and [22].

Let $C([s, t]; V_\mu)$ be the space of continuous functions on $[s, t]$ with values in V_μ and with the usual norm $\|\cdot\|_{C,\mu}$ (or $\|\cdot\|_{C,s,t,\mu}$ when we want to stress the interval). In the particular case that $\mu = 0$, we simply write $\|\cdot\|_C$ (or $\|\cdot\|_{C,s,t}$ respectively). For $\beta \in (0, 1]$ we denote by $C^\beta([s, t]; V_\mu)$ the space of Hölder-continuous functions on $[s, t]$ and with values in V_μ , equipped with the norm

$$\|u\|_{\beta,\mu} = \|u\|_{C,\mu} + \|u\|_{\beta,\mu}, \quad \|u\|_{\beta,\mu} := \sup_{s \leq p < q \leq t} \frac{\|u(q) - u(p)\|_{V_\mu}}{(q - p)^\beta}.$$

In particular, for the case $\beta = 1$ this is the space of Lipschitz-continuous functions.

The spaces $L^p(s, t; V_\mu)$, $p \in [1, \infty]$ have the standard meaning with the usual norms.

As we have mentioned above, sometimes it is important to consider the above norms on different time intervals $[s, t]$, thus in those cases the time interval will be indicated in the index of the norm.

For the previous spaces the following compactness theorem holds true:

Theorem 2.2. (i) For $\alpha > 0$ and $\delta \in \mathbb{R}$, $L^2(s, t; V_\alpha) \cap C^\beta([s, t]; V_{-\delta})$ is compactly embedded into $L^2(s, t; V) \cap C([s, t]; V_{-\delta})$.

(ii) For $0 \leq \delta_1 < \delta_2$ and $0 \leq \beta_1 < \beta_2 \leq 1$ the space $C^{\beta_2}([s, t]; V_{-\delta_1})$ is compactly embedded into $C^{\beta_1}([s, t]; V_{-\delta_2})$.

For the first part see Vishik and Fursikov [25, Chapter IV Theorem 4.1]. For the second part we refer to Maslowski and Nualart [16, Lemma 4.5]. Indeed, we have the compact embedding $V_{-\delta_1} \subset V_{-\delta_2}$.

For $t \in [0, T]$, we now rewrite Eq. (1.1) in an abstract form,

$$du(t) = (Au(t) + B(u(t), u(t))) dt + G(u(t))d\omega(t), \quad u(0) = u_0, \tag{2.5}$$

where G is a nontrivial diffusion term representing the external force, that satisfies certain assumptions which are given in Section 4. Here ω represents a path in $C^{\beta'}([0, T]; V)$, with $\beta' > 1/2$, or in particular, a fractional Brownian-motion with Hurst-parameter $H \in (1/2, 1)$, see the definition in Section 3. The corresponding stochastic evolution equation has therefore a multiplicative noise. In what follows we will describe the type of stochastic integral we are going to consider, which will allow us to give an appropriate meaning to (2.5).

3. Integrals in Hilbert-spaces for Hölder-continuous integrators with Hölder exponents greater than 1/2

In this section we are concerned with the definition of the following infinite dimensional integral

$$\int_{T_1}^{T_2} Z d\omega,$$

where ω is a Hölder-continuous function with Hölder exponent $\beta' > 1/2$ and Z is an appropriate integrand. We follow the recent definition given by Chen et al. [19], and for the sake of completeness, next we shall borrow the main steps of their construction.

We start by considering an abstract separable Hilbert-space \tilde{V} , then for $0 < \alpha < 1$ and general measurable functions $Z : [T_1, T_2] \rightarrow \tilde{V}$ and $\omega : [T_1, T_2] \rightarrow V$, we define the following fractional derivatives

$$D_{T_1+}^\alpha Z[r] = \frac{1}{\Gamma(1-\alpha)} \left(\frac{Z(r)}{(r-T_1)^\alpha} + \alpha \int_{T_1}^r \frac{Z(r) - Z(q)}{(r-q)^{1+\alpha}} dq \right) \in \tilde{V},$$

$$D_{T_2-}^{1-\alpha} \omega_{T_2-}[r] = \frac{(-1)^{1-\alpha}}{\Gamma(\alpha)} \left(\frac{\omega(r) - \omega(T_2-)}{(T_2-r)^{1-\alpha}} + (1-\alpha) \int_r^{T_2} \frac{\omega(r) - \omega(q)}{(q-r)^{2-\alpha}} dq \right) \in V,$$

where $\omega_{T_2-}(r) = \omega(r) - \omega(T_2-)$, being $\omega(T_2-)$ the left-sided limit of ω at T_2 . Here $\Gamma(\cdot)$ denotes the Gamma function.

Let us start with the case when the integrand z and the integrator ζ are one-dimensional. Suppose that $z(T_1+)$, $\zeta(T_1+)$, $\zeta(T_2-)$ exist, being respectively the right-sided limit of z at T_1 and the right- and left-sided limits of ζ at T_1, T_2 , and that $z \in I_{T_1+}^\alpha(L^p(T_1, T_2; \mathbb{R}))$, $\zeta_{T_2-} \in I_{T_2-}^{1-\alpha}(L^{p'}(T_1, T_2; \mathbb{R}))$ with $1/p + 1/p' \leq 1$ and $\alpha p < 1$ (the definition of these spaces can be found, for instance, in Samko et al. [11]). Then following Zähle [10] we define

$$\int_{T_1}^{T_2} z d\zeta = (-1)^\alpha \int_{T_1}^{T_2} D_{T_1+}^\alpha z[r] D_{T_2-}^{1-\alpha} \zeta_{T_2-}[r] dr. \tag{3.1}$$

Suppose now that ζ is Lipschitz-continuous. Then ζ generates a signed measure $d\zeta$ and $\zeta \in I_{T_2-}^\alpha(L^{p'}(T_1, T_2; \mathbb{R}))$. Therefore, in this situation the integral

$$\int_{T_1}^{T_2} z d\zeta$$

can be expressed by (3.1).

Let \hat{V} be a separable Hilbert-space endowed with the norm $\|\cdot\|_{\hat{V}}$ and consider the separable Hilbert-space $L_2(V, \hat{V})$ of Hilbert-Schmidt-operators from V into \hat{V} with the norm $\|\cdot\|_{L_2(V, \hat{V})}$ and inner product $(\cdot, \cdot)_{L_2(V, \hat{V})}$. Let $(e_i)_{i \in \mathbb{N}}$ and $(\hat{f}_i)_{i \in \mathbb{N}}$ be a complete orthonormal basis of V and \hat{V} , resp. A base in $L_2(V, \hat{V})$ is given by

$$E_{ij} e_k = \begin{cases} 0 & j \neq k \\ \hat{f}_i & j = k. \end{cases}$$

Let us consider now mappings $Z : [T_1, T_2] \rightarrow L_2(V, \hat{V})$ and $\omega : [T_1, T_2] \rightarrow V$. Suppose that $z_{ji} = (Z, E_{ji})_{L_2(V, \hat{V})} \in I_{T_1+}^\alpha(L^p(T_1, T_2; \mathbb{R}))$ and $z_{ji}(T_1+)$ exists and $\alpha p < 1$. Moreover, let us also assume that $\zeta_{iT_2-} = (\omega_{T_2-}(t), e_i)_V \in I_{T_2-}^{1-\alpha}(L^{p'}(T_1, T_2; \mathbb{R}))$ such that $1/p + 1/p' \leq 1$, and the mapping

$$[T_1, T_2] \ni r \mapsto \|D_{T_1+}^\alpha Z[r]\|_{L_2(V, \hat{V})} \|D_{T_2-}^{1-\alpha} \omega_{T_2-}[r]\| \in L^1(T_1, T_2; \mathbb{R}).$$

We introduce

$$\begin{aligned} \int_{T_1}^{T_2} Z d\omega &:= (-1)^\alpha \int_{T_1}^{T_2} D_{T_1+}^\alpha Z[r] D_{T_2-}^{1-\alpha} \omega_{T_2-}[r] dr \\ &:= (-1)^\alpha \sum_{j=1}^\infty \left(\sum_{i=1}^\infty \int_{T_1}^{T_2} D_{T_1+}^\alpha z_{ji}[r] D_{T_2-}^{1-\alpha} \zeta_{iT_2-}[r] dr \right) f_j. \end{aligned} \tag{3.2}$$

This last equality is well defined due to the fact that Pettis' theorem and the separability of V ensure that the integrand is weakly measurable and hence measurable. Moreover, the norm of the above integral is given by

$$\begin{aligned} \left\| \int_{T_1}^{T_2} Z d\omega \right\|_{\hat{V}} &= \left(\sum_{j=1}^\infty \left| \sum_{i=1}^\infty \int_{T_1}^{T_2} D_{T_1+}^\alpha z_{ji}[r] D_{T_2-}^{1-\alpha} \zeta_{iT_2-}[r] dr \right|^2 \right)^{\frac{1}{2}} \\ &\leq \int_{T_1}^{T_2} \|D_{T_1+}^\alpha Z[r]\|_{L_2(V, \hat{V})} \|D_{T_2-}^{1-\alpha} \omega_{T_2-}[r]\| dr. \end{aligned}$$

The next result, which proof can be found in [19], considers the definition of the above integral when having suitable Hölder continuous integrator and integrand functions:

Lemma 3.1. *Suppose that $0 \leq T_1 < T_2$, $Z \in C^\beta([T_1, T_2]; L_2(V, \hat{V}))$ and $\omega \in C^{\beta'}([T_1, T_2]; V)$ with $1 - \beta' < \alpha < \beta$. Then*

$$\int_{T_1}^{T_2} Z d\omega \in \hat{V}$$

is well-defined in the sense of (3.2). Also, there exists a constant c depending on $T_2 - T_1$, α , β and β' such that

$$\left\| \int_{T_1}^{T_2} Z d\omega \right\|_{\hat{V}} \leq c \|Z\|_{C^\beta([T_1, T_2]; L_2(V, \hat{V}))} \|\omega\|_{\beta', T_1, T_2} (T_2 - T_1)^{\beta'}.$$

Moreover, the above integral with driving path ω is well-defined even though the integrand is locally Hölder-continuous, which will be the case in the next sections when the semigroup S is part of the integrand, see [19] for the proof of this assertion.

In the following we would like to consider the above integrals when the integrator is a noise given by a fractional Brownian-motion (fBm). An one-dimensional fBm is a centered Gaussian-process given by the auto-covariance

$$R(s, t) = \frac{1}{2}(t^{2H} + s^{2H} - |t - s|^{2H})$$

where $H \in (0, 1)$ is the so-called Hurst-parameter. The value $H = 1/2$ determines a Brownian-motion, which is a martingale and a Markov-process with independent increments. When $H \neq 1/2$ these properties do not hold.

An fBm can be also defined in a separable Hilbert-space. By the following construction we obtain such an infinite-dimensional noise with values in V : let $(\zeta_i)_{i \in \mathbb{N}}$ be a iid-sequence of fBm in \mathbb{R} having the same Hurst-parameter H . Then

$$t \rightarrow \omega(t) := \sum_{i=1}^\infty q_i^{\frac{1}{2}} \zeta_i e_i,$$

where $(q_i)_{i \in \mathbb{N}} \in l_2$, defines an fBm with values in V and with auto-covariance

$$\frac{1}{2} Q(t^{2H} + s^{2H} - |t - s|^{2H}),$$

where the operator Q of diagonal form is defined by

$$(e_i, Qe_j)_V = \delta_{ij} q_i.$$

For simplicity we restrict ourselves to a real fBm. However, taking two one-dimensional independent real fBm ζ^1, ζ^2 then we could construct a one-dimensional complex fBm: $\zeta := 1/\sqrt{2}(\zeta^1 + i\zeta^2)$. Then by the above formula we could construct a complex fBm ω in V .

Remark 3.2. For our further purposes we need the fBm ω to be piecewise linearly approximated. As one can check later, we will use the property that given ω we can find a sequence of piecewise linear continuous functions ω_n converging to ω in a Hölder-continuous space. However, the space of Hölder-continuous functions is not separable, but we can modify it in such a way that the modified space is: for $\gamma \in (0, 1)$, the space

$$C^{0,\gamma}([0, T]; V) := \left\{ \omega \in C^\gamma([0, T]; V) : \lim_{\delta \rightarrow 0} \sup_{|s_1 - s_2| < \delta, [0, T] \ni s_1 \neq s_2} \frac{\|\omega(s_1) - \omega(s_2)\|}{(s_1 - s_2)^\gamma} = 0 \right\},$$

is a separable space since V is itself separable (see [26,27] and [28]). On the other hand, thanks to Kolmogorov’s theorem (see Theorem 1.4.1 in Kunita [29]), it is known that if ω represents an fBm with Hurst parameter H , then there exists a version, that we also call ω , such that $\omega \in C^{\beta'}([0, T]; V)$ for any $\beta' < H$. Let us take $\gamma < \beta' < H$, then it is easy to see that $C^{\beta'}([0, T]; V) \subset C^{0,\gamma}([0, T]; V)$, and therefore this latter space is the one that we should take when the aim is to approximate the fBm. However, for the easy of presentation, in what follows we will assume that ω can be piecewise linearly approximated by a sequence $(\omega_n)_{n \in \mathbb{N}}$ converging in $C^{\beta'}([0, T]; V)$, but this statement must be understood according to the sense given above.

4. Definition of a solution of the stochastic shell-model

In this section we would like to formulate conditions ensuring that (2.5) has a unique global solution.

We emphasize that we have to give a definition of solution which is appropriate in our context: on the one hand, the driving function belongs to $C^{\beta'}([0, T]; V)$ for a $\beta' \in (1/2, 1)$, which means that we cannot define integrals by using the standard integration theory of bounded-variation integrators. In particular our situation here covers the case when the driving function is given by an fBm in V with Hurst-parameter $H > 1/2$. On the other hand, the bilinear operator B is unbounded and not Lipschitz-continuous in this model.

We now give the assumptions for the diffusion operator G of (2.5). In what follows, we choose a constant $\delta > 0$ which will be determined later, and by $L_2(V)$ we denote the space $L_2(V, V)$.

Assumption (G). Assume that the mapping $G : V_{-\delta} \rightarrow L_2(V)$ is bounded and twice continuously Fréchet-differentiable with bounded first and second derivatives $DG(u)$ and $D^2G(u)$, for $u \in V_{-\delta}$. Let us denote, respectively, by c_G, c_{DG} and c_{D^2G} the bounds for G, DG and D^2G . Then, for $u \in V_{-\delta}$

$$\|G(u)\|_{L_2(V)} \leq c_G.$$

Furthermore, for $u_1, u_2 \in V_{-\delta}$,

$$\|G(u_1) - G(u_2)\|_{L_2(V)} \leq c_{DG}\|u_1 - u_2\|_{V_{-\delta}},$$

and for $u_1, u_2, v_1, v_2 \in V_{-\delta}$,

$$\begin{aligned} & \|G(u_1) - G(v_1) - (G(u_2) - G(v_2))\|_{L_2(V)} \\ & \leq c_{DG}\|u_1 - v_1 - (u_2 - v_2)\|_{V_{-\delta}} + c_{D^2G}\|u_1 - u_2\|_{V_{-\delta}}(\|u_1 - v_1\|_{V_{-\delta}} + \|u_2 - v_2\|_{V_{-\delta}}). \end{aligned} \tag{4.1}$$

Notice that $DG : V_{-\delta} \mapsto L_2(V_{-\delta}, L_2(V)) = L_2(V_{-\delta} \times V, V)$ is a bilinear mapping whereas D^2G a trilinear mapping.

In this paper we shall look at the existence and uniqueness of a solution of (2.5) according to the next definition:

Definition 4.1. Let $1/2 < \beta < \beta' \leq 1$ and let $\omega \in C^{\beta'}([0, T]; V), u_0 \in V$ and $\delta \in (\beta, 1)$. A function u is said to be a *mild* solution to (2.5) over the interval $[0, T]$ associated to the initial condition u_0 if

$$u \in C([0, T]; V) \cap L^2(0, T; V_{1/2}) \cap C^\beta([0, T]; V_{-\delta})$$

and such that

$$u(t) = S(t)u_0 + \int_0^t S(t-r)B(u(r), u(r))dr + \int_0^t S(t-r)G(u(r))d\omega(r) \tag{4.2}$$

for every $t \in [0, T]$.

Remark 4.2. Note that the first integral in (4.2) is well defined in V because of the fact that $u \in C([0, T]; V) \cap L^2(0, T; V_{1/2})$ and Lemma 2.1. The stochastic integral in (4.2) must be understood in V according to the definition given in Section 3.

We stress that we are interested in finding a mild solution for (2.5). Following [15] we could also consider weak solutions for our problem. Nevertheless for $u \in L^\infty(0, T; V) \cap L^2(0, T; V_{1/2})$ we have that $B(u, u)$ is sufficiently regular so that we can work with mild solutions.

When ω is regular, we can also interpret the solution in the following weak sense:

Definition 4.3. Assume that ω is piecewise linear continuous in $[0, T]$ with values in V . We say that u is a *weak* solution to (2.5) over the interval $[0, T]$ associated to the initial condition $u_0 \in V$ if

$$u \in C([0, T]; V) \cap L^2(0, T; V_{1/2})$$

and such that

$$(u(t), \varphi)_V + \int_0^t (A^{1/2}u(s), A^{1/2}\varphi)_V ds - \int_0^t (B(u(s), u(s)), \varphi)_V ds = (u_0, \varphi)_V + \int_0^t (G(u(s))\omega'(s), \varphi)_V ds \tag{4.3}$$

holds for every $\varphi \in V_{1/2}$ and $t \in [0, T]$.

5. Solutions of the stochastic shell-model for piecewise linear continuous path noise

In this section we assume that ω is a piecewise linear continuous function. This case is the foundation for studying the more general case which will be treated in the next section. Indeed, in further sections given $\omega \in C^{\beta'}([0, T]; V)$ we shall consider a sequence $(\omega_n)_{n \in \mathbb{N}}$ of piecewise linear continuous paths converging to $\omega \in C^{\beta'}([0, T]; V)$, see [Remark 3.2](#). As we cannot assume that the sequence $(\omega_n)_{n \in \mathbb{N}}$ is uniformly bounded in $L^\infty(0, T; V)$, we will have to construct uniform a priori estimates for the solutions to equations driven by ω_n , which will be based on uniform estimates of $(\omega_n)_{n \in \mathbb{N}}$ with respect to the $C^{\beta'}$ -norm.

We start by studying the existence of solutions for the stochastic Shell-model having this kind of regular driving function:

Proposition 5.1. *Let $u_0 \in V$, ω be a piecewise linear continuous function and assume that G satisfies the assumption **(G)**. Then, there is a unique global weak solution u for Eq. (2.5) in the sense of [Definition 4.3](#).*

Proof. The proof is very classical but, for the sake of completeness, we will sketch it here. Let us denote by P_n the projection operator in V onto the space spanned by e_1, e_2, \dots, e_n . Then, the Galerkin-approximations $(u_n)_{n \in \mathbb{N}}$ to problem (2.5) are solutions of the finite-dimensional systems

$$du_n(t) = (Au_n(t) + P_n B(u_n(t), u_n(t)))dt + P_n G(u_n(t))\omega'(t)dt. \quad (5.1)$$

On the other hand, if G^* denotes the adjoint operator of G , taking the scalar product of (5.1) by u_n , using the property (2.4) and the assumption **(G)**, we get that

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|u_n(t)\|^2 + \|u_n(t)\|_{V_{1/2}}^2 &\leq |(P_n G(u_n(t))\omega'(t), u_n(t))_V| \leq |(\omega'(t), G^*(u_n(t))u_n(t))_V| \\ &\leq \|\omega'(t)\| \|G^*(u_n(t))u_n(t)\| \leq c_G \|\omega'(t)\| \|u_n(t)\| \\ &\leq \frac{c_G^2}{2} \|\omega'(t)\|^2 + \frac{1}{2} \|u_n(t)\|^2. \end{aligned}$$

Hence, using Gronwall's lemma yields that

$$\sup_{t \in [0, T]} \|u_n(t)\|^2 \leq c(\|u_n(0)\|, \|\omega'\|_{L^\infty(0, T; V)}, T)$$

for an appropriate positive constant c , and consequently we also have

$$\int_0^T \|u_n(t)\|_{V_{1/2}}^2 dt \leq c,$$

uniformly in n .

Also, by classical arguments, we get that $(u_n)_{n \in \mathbb{N}}$ is bounded in $C^{\hat{\beta}}([0, T]; V_{-\delta})$ for $\delta > 1/2$ and $\hat{\beta} < 1$. In fact, since $u_n \in L^\infty(0, T; V)$, it follows by [Lemma 2.1](#)

$$\sup_{0 \leq s < t \leq T} \frac{\int_s^t \|A^{-\delta} B(u_n(r), u_n(r))\| dr}{(t-s)^{\hat{\beta}}} \leq c \sup_{0 \leq s < t \leq T} \frac{\int_s^t \|A^{-\frac{1}{2}} B(u_n(r), u_n(r))\| dr}{(t-s)^{\hat{\beta}}} \leq cT^{1-\hat{\beta}} \|u_n\|_{L^\infty(0, T; V)}^2 < \infty,$$

and by **(G)** we arrive at

$$\sup_{0 \leq s < t \leq T} \frac{\int_s^t \|A^{-\delta} G(u_n(r))\omega'(r)\| dr}{(t-s)^{\hat{\beta}}} \leq c \sup_{0 \leq s < t \leq T} \frac{\int_s^t \|G(u_n(r))\|_{L^2(V)} \|\omega'(r)\| dr}{(t-s)^{\hat{\beta}}} \leq cc_G T^{1-\hat{\beta}} \|\omega'\|_{L^\infty(0, T; V)} < \infty.$$

Moreover, applying the interpolation inequality (see [23], Theorem 37.6), we know that there exists a constant $c = c(\delta) \geq 1$ such that

$$\|A^{1-\delta} v\| \leq c \|A^0 v\|^{2\delta-1} \|A^{1/2} v\|^{2-2\delta} \quad \text{for all } v \in V,$$

and therefore, for $\hat{\beta} \in (1/2, 1)$ and $\delta \in (\hat{\beta}, 1)$,

$$\begin{aligned} \sup_{0 \leq s < t \leq T} \frac{\int_s^t \|A^{-\delta} Au_n(r)\| dr}{(t-s)^{\hat{\beta}}} &\leq c \sup_{0 \leq s < t \leq T} \frac{\int_s^t \|u_n(r)\|^{2\delta-1} \|A^{1/2} u_n(r)\|^{2-2\delta} dr}{(t-s)^{\hat{\beta}}} \\ &\leq c \|u_n\|_{L^\infty(0, T; V)}^{2\delta-1} \sup_{0 \leq s < t \leq T} \frac{(\int_s^t dr)^\delta (\int_s^t \|A^{1/2} u_n(r)\|^2 dr)^{1-\delta}}{(t-s)^{\hat{\beta}}} \leq cT^{\delta-\hat{\beta}} \|u_n\|_{L^\infty(0, T; V)}^{2\delta-1} \|u_n\|_{L^2(0, T; V_{1/2})}^{2-2\delta} < \infty. \end{aligned}$$

Hence, by the compactness [Theorem 2.2\(i\)](#) we get a subsequence, still denoted by $(u_n)_{n \in \mathbb{N}}$, that converges strongly in $L^2(0, T; V) \cap C([0, T]; V_{-\delta})$ to some limit u . Since $(u_n)_{n \in \mathbb{N}}$ is bounded in $L^\infty(0, T; V) \cap L^2(0, T; V_{1/2})$ this sequence is relatively weak-star compact in $L^\infty(0, T; V)$ and relatively weak compact in $L^2(0, T; V_{1/2})$. As a consequence, the limit $u \in L^\infty(0, T; V) \cap L^2(0, T; V_{1/2})$. Now, it remains to prove that the limit u is a solution to the system (2.5) according to [Definition 4.3](#). Indeed, assuming that u_n is solution in the sense of [Definition 4.3](#), we can pass to the limit on each term. Furthermore, the regularity of u implies that the right hand side of (4.3) as well as the last two terms of the left hand side of (4.3) are in $C([0, T]; V)$, hence $u \in C([0, T]; V)$. \square

Moreover, we have the following result about mild solutions:

Proposition 5.2. Assume the same hypotheses than in Proposition 5.1 and that $\hat{\beta} \in (1/2, 1)$, $\delta \in (\hat{\beta}, 1)$. Then every weak solution u to (2.5) is a mild solution, that is, $u \in C([0, T]; V) \cap L^2(0, T; V_{1/2}) \cap C^{\hat{\beta}}([0, T]; V_{-\delta})$ and satisfies for every $t \in [0, T]$ the following integral formulation in V :

$$u(t) = S(t)u_0 + \int_0^t S(t-r)B(u(r), u(r))dr + \int_0^t S(t-r)G(u(r))\omega'(r)dr. \tag{5.2}$$

Proof. Suppose that u fulfills (4.3). Then

$$t \mapsto B(u(t), u(t)) + G(u(t))\omega'(t) \in L^2(0, T; V)$$

such that

$$t \mapsto \int_0^t S(t-r)B(u(r), u(r))dr + \int_0^t S(t-r)G(u(r))\omega'(r)dr \in C([0, T]; V),$$

see Pazy [24], proof of Theorem 4.3.1. In addition, every Galerkin-approximation solution of (5.1) satisfies

$$(u_n(t), \varphi)_V = (S(t)P_n u_n(0), \varphi)_V + \int_0^t (S(t-r)P_n B(u_n(r), u_n(r)), \varphi)_V dr + \int_0^t (S(t-r)P_n G(u_n(r))\omega'(r), \varphi)_V dr \tag{5.3}$$

for every $\varphi \in V$ and every $t \in [0, T]$. From the convergence of $(u_n)_{n \in \mathbb{N}}$ in $L^2(0, T; V)$ and the boundedness in $L^2(0, T; V_{1/2})$ it follows that the right hand side of (5.3) converges to

$$(S(t)u_0, \varphi)_V + \int_0^t (S(t-r)B(u(r), u(r)), \varphi)_V dr + \int_0^t (S(t-r)G(u(r))\omega'(r), \varphi)_V dr$$

for every $t \in [0, T]$. On the other hand, from the proof of Proposition 5.1 we know that $(u_n)_{n \in \mathbb{N}}$ converges to u in $C([0, T]; V_{-\delta})$ and hence $u_n(t)$ converges to $u(t)$ in $V_{-\delta}$ for every $t \in [0, T]$. Since the right hand side is in V for every $t \in [0, T]$, $u(t)$ is too. Also, following the same reasoning than in Proposition 5.1, one can prove that $(u_n)_{n \in \mathbb{N}}$ is bounded in $C^\gamma([0, T]; V_{-\hat{\delta}})$ for $\gamma = \hat{\beta} + \varepsilon$ and $\hat{\delta} = \delta - \varepsilon$ for small enough $\varepsilon > 0$ such that $\hat{\delta} > \gamma$. Then it suffices to apply Theorem 2.2(ii) to conclude the proof. \square

From now on, we often use the following property, which is a consequence of the definition of Beta function: for every $0 \leq s < t \leq T$, $a, b > -1$,

$$\int_s^t (r-s)^a (t-r)^b dr = c(t-s)^{a+b+1} \tag{5.4}$$

where c only depends on a and b .

Next we develop a priori estimates that later we need to derive the existence of a solution for a general $\omega \in C^{\beta'}([0, T]; V)$. We cannot use the estimate from Proposition 5.1 because the sequence $(\omega_n)_{n \in \mathbb{N}}$ approximating ω in $C^{\beta'}([0, T]; V)$ is not in general uniformly bounded in $L^\infty(0, T; V)$. That is why in the following estimates the norm $\|\omega\|_{\beta'}$ appears.

Lemma 5.3. Assume that $1/2 < \hat{\beta} < \beta'$, $1 - \beta' < \alpha < \hat{\beta}$, $\delta \in (\hat{\beta}, 1)$, $u_0 \in V$, ω is a piecewise linear continuous function and G satisfies (G). Then, if u is a weak solution to (2.5) in the sense of Definition 4.3, there is a constant $c > 0$ such that for $t \in [0, T]$

$$\|u(t)\|^2 + 2 \int_0^t \|u(r)\|_{V_{1/2}}^2 dr \leq \|u_0\|^2 + c\|\omega\|_{\beta'} t^{\beta'} \|u\|_{C,0,t} + c\|\omega\|_{\beta'} t^{\hat{\beta}+\beta'} (1 + \|u\|_{C,0,t}) \|u\|_{\hat{\beta},-\delta,0,t}. \tag{5.5}$$

Proof. Applying the formula of the square norm, see Teman [30] Lemma III.1.2, using the skew-symmetric property (2.4), and finally integrating over $(0, t)$, this gives us for every $t \in [0, T]$ the following energy inequality

$$\|u(t)\|^2 + 2 \int_0^t \|u(r)\|_{V_{1/2}}^2 dr \leq \|u_0\|^2 + 2 \left| \int_0^t (G^*(u(r))u(r), \omega'(r))_V dr \right|.$$

The integral on the right hand side of the previous expression can be interpreted in the sense of Section 3 using fractional derivatives. Since for any r we have $\|D_{0+}^\alpha G^*(u(r))u(r)\| < \infty$ the expression $D_{0+}^\alpha G^*(u(r))u(r)$ is an element in the space of Hilbert–Schmidt-operators $L_2(V, \mathbb{R}) \simeq V$. Moreover, from the definition of the fractional derivative it is easy to derive that

$$\|D_{t-}^{1-\alpha} \omega_{t-}[r]\| \leq c\|\omega\|_{\beta'} (t-r)^{\alpha+\beta'-1}, \tag{5.6}$$

and therefore we get

$$\left| \int_0^t (G^*(u(r))u(r), \omega'(r))_V dr \right| \leq c\|\omega\|_{\beta'} \int_0^t (t-r)^{\alpha+\beta'-1} \left(\frac{\|G^*(u(r))u(r)\|}{r^\alpha} + \int_0^r \frac{\|G^*(u(r))u(r) - G^*(u(q))u(q)\|}{(r-q)^{1+\alpha}} dq \right) dr.$$

Trivially the boundedness of G implies that $\|G^*(u)u\| \leq c_G \|u\|$ for $u \in V$ and therefore

$$\frac{\|G^*(u(r))u(r)\|}{r^\alpha} \leq \frac{c_G \|u\|_{C,0,t}}{r^\alpha}, \quad r \in [0, t].$$

The boundedness and the Lipschitz-continuity of G imply

$$\begin{aligned} & \int_0^r \frac{\|G^*(u(r))u(r) - G^*(u(q))u(q)\|}{(r-q)^{1+\alpha}} dq \\ & \leq c_G \int_0^r \frac{\|u(r) - u(q)\|_{V_{-\delta}}}{(r-q)^{1+\alpha}} dq + \|u\|_{C,0,t} \int_0^r \frac{\|G^*(u(r)) - G^*(u(q))\|_{L_2(V_{-\delta},V)}}{(r-q)^{1+\alpha}} dq \\ & \leq (c_G + c_{DG}\|u\|_{C,0,t}) \|u\|_{\hat{\beta},-\delta,0,t} \int_0^r (r-q)^{-1-\alpha+\hat{\beta}} dq \\ & = c(c_G + c_{DG}\|u\|_{C,0,t}) \|u\|_{\hat{\beta},-\delta,0,t} r^{\hat{\beta}-\alpha}. \end{aligned}$$

Hence, for an appropriate $c > 0$

$$\left| \int_0^t (G^*(u(r))u(r), \omega'(r))_V dr \right| \leq c \|\omega\|_{\beta'} t^{\beta'} \|u\|_{C,0,t} + c \|\omega\|_{\beta'} t^{\hat{\beta}+\beta'} (1 + \|u\|_{C,0,t}) \|u\|_{\hat{\beta},-\delta,0,t}. \quad \square$$

Lemma 5.4. Under the same conditions of Lemma 5.3, if u is a solution to (2.5) there exist constants $c, \bar{c} > 0$ such that for $t \in [0, T]$

$$\|u\|_{\hat{\beta},-\delta,0,t} \leq \bar{c} t^{\delta-\hat{\beta}} \|u_0\| + c(t^{1-\hat{\beta}} + t^{\delta+1/2-\hat{\beta}}) \|u\|_{C,0,t}^2 + c \|\omega\|_{\beta'} t^{\beta'-\hat{\beta}} (1 + t^{\hat{\beta}} \|u\|_{\hat{\beta},-\delta,0,t}). \quad (5.7)$$

Proof. Consider (5.2) written as

$$u(t) = S(t)u_0 + A^{1/2} \int_0^t S(t-r)A^{-1/2}B(u(r), u(r))dr + \int_0^t S(t-r)G(u(r))\omega'(r)dr.$$

Then the following splitting is considered:

$$\begin{aligned} A^{-\delta}(u(q) - u(p)) &= A^{-\delta}(S(q) - S(p))u_0 + A^{-\delta+1/2} \int_p^q S(q-r)A^{-1/2}B(u(r), u(r))dr \\ &+ A^{-\delta+1/2} \int_0^p (S(q-r) - S(p-r))A^{-1/2}B(u(r), u(r))dr \\ &+ A^{-\delta} \int_p^q S(q-r)G(u(r))\omega'(r)dr + A^{-\delta} \int_0^p (S(q-r) - S(p-r))G(u(r))\omega'(r)dr \\ &=: I_1 + I_2 + I_3 + I_4 + I_5. \end{aligned} \quad (5.8)$$

For the term related to the initial condition, due to the fact that $\delta \in (\hat{\beta}, 1)$ and (2.1), (2.2) we have

$$\sup_{0 \leq p < q \leq t} \frac{\|I_1\|}{(q-p)^{\hat{\beta}}} \leq \sup_{0 \leq p < q \leq t} \frac{\|A^{-\delta}(S(q-p) - \text{Id})S(p)u_0\|}{(q-p)^{\hat{\beta}}} \leq \bar{c} \sup_{0 \leq p < q \leq t} \frac{(q-p)^{\delta} \|u_0\|}{(q-p)^{\hat{\beta}}} \leq \bar{c} t^{\delta-\hat{\beta}} \|u_0\|. \quad (5.9)$$

Moreover, due to Lemma 2.1 and taking into account that $V \subset V_{-\delta+1/2}$,

$$\begin{aligned} \sup_{0 \leq p < q \leq t} \frac{\|I_2\|}{(q-p)^{\hat{\beta}}} &\leq \sup_{0 \leq p < q \leq t} \frac{1}{(q-p)^{\hat{\beta}}} \int_p^q \|A^{-\delta+1/2}S(q-r)A^{-1/2}B(u(r), u(r))\| dr \\ &\leq \sup_{0 \leq p < q \leq t} \frac{c}{(q-p)^{\hat{\beta}}} \int_p^q \|A^{-1/2}B(u(r), u(r))\| dr \\ &\leq \sup_{0 \leq p < q \leq t} \frac{c}{(q-p)^{\hat{\beta}}} (q-p) \|u\|_{C,0,t}^2 \leq ct^{1-\hat{\beta}} \|u\|_{C,0,t}^2. \end{aligned} \quad (5.10)$$

For I_3 , thanks to Lemma 2.1 and (2.3),

$$\begin{aligned} \sup_{0 \leq p < q \leq t} \frac{\|I_3\|}{(q-p)^{\hat{\beta}}} &\leq \sup_{0 \leq p < q \leq t} \frac{c}{(q-p)^{\hat{\beta}}} \int_0^p \|A^{-\delta+1/2}(S(q-p) - \text{Id})S(p-r)A^{-1/2}B(u(r), u(r))\| dr \\ &\leq c \|u\|_{C,0,t}^2 \sup_{0 \leq p \leq t} \int_0^p (p-r)^{\delta-1/2-\hat{\beta}} dr \\ &\leq ct^{\delta+1/2-\hat{\beta}} \|u\|_{C,0,t}^2. \end{aligned} \quad (5.11)$$

Similar estimates to those of I_4, I_5 can be found in [19]. However, and for the completeness of the presentation, we also show these technical estimates in this paper, but we have shifted these calculations into Appendix Section, see Lemma A.1(i) and Corollary A.2(i), where it is proven that

$$\sup_{0 \leq p < q \leq t} \frac{\|I_4\| + \|I_5\|}{(q-p)^{\hat{\beta}}} \leq ct^{\beta'-\hat{\beta}} \|\omega\|_{\beta'} (1 + t^{\hat{\beta}} \|u\|_{\hat{\beta},-\delta,0,t}). \quad (5.12)$$

Hence, collecting all the estimates for the expressions I_j the inequality (5.7) is obtained. \square

Lemma 5.5. Under the assumptions of Lemma 5.3, if u_n is a solution of (4.3) on $[0, T]$ with initial condition $u_0 \in V$ and driven by a piecewise linear continuous path ω_n where $(\omega_n)_{n \in \mathbb{N}}$ is bounded in $C^{\beta'}([0, T]; V)$, then $(u_n)_{n \in \mathbb{N}}$ is uniformly bounded in $C^{\hat{\beta}}([0, T]; V_{-\delta}) \cap C([0, T]; V)$.

The proof of the previous result rests upon the technical Lemmas A.3–A.5 whose proofs are presented into Appendix section.

Remark 5.6. We emphasize that we consider Hölder-continuity with respect to the space $V_{-\delta}$, although all the estimates of the Hölder-norms also make sense for smaller δ . However, the initial condition u_0 is the responsible of having to consider $\delta \in (\hat{\beta}, 1)$, since in (5.7) the exponent in the term $t^{\delta-\hat{\beta}}$ multiplying $\|u_0\|$ must be positive.

6. Construction of solutions

We are now able to construct solutions for the stochastic equation (2.5) and give the main result of this paper. We consider a sequence of solutions $(u_n)_{n \in \mathbb{N}}$ to (2.5) driven by $(\omega_n)_{n \in \mathbb{N}}$, a sequence of piecewise linear continuous approximations of ω converging to ω where ω satisfies Remark 3.2.

First we formulate a general uniqueness theorem.

Theorem 6.1. Suppose that there are two mild solutions u_1, u_2 of (4.2) with $u_1(0) = u_2(0) = u_0 \in V$ and driven by the same path ω . Then, under the given assumptions on A, B and G we have $u_1(t) = u_2(t)$ for $t \in [0, T]$.

Proof. Assume that there exists a maximal interval $[0, t_0]$ contained in $[0, T]$ such that $\Delta u := u_1 - u_2$ is zero on this interval being $t_0 < T$. Then there exists a $0 < \mu < 1$ such that $\Delta u \neq 0$ on $(t_0, t_0 + \mu]$.

We divide the proof in several steps:

(i) First we want to estimate

$$\|\Delta u\|_{\beta, -\delta, t_0, t_0+\mu} = \sup_{t_0 \leq s < t \leq t_0+\mu} \frac{\|\Delta u(t) - \Delta u(s)\|_{V_{-\delta}}}{(t-s)^\beta}.$$

Regarding the non-stochastic integral, we have to estimate

$$\begin{aligned} & \frac{1}{(t-s)^\beta} \left\| \int_s^t S(t-r)A^{-\delta}(B(u_1(r), u_1(r)) - B(u_2(r), u_2(r)))dr \right\| \\ & + \frac{1}{(t-s)^\beta} \left\| \int_{t_0}^s (S(t-r) - S(s-r))A^{-\delta}(B(u_1(r), u_1(r)) - B(u_2(r), u_2(r)))dr \right\| \\ & =: J_1 + J_2. \end{aligned}$$

Since $V_{-1/2} \subset V_{-\delta}$, from Lemma 2.1 we obtain

$$\begin{aligned} \|A^{-\delta}(B(u_1(r), u_1(r)) - B(u_2(r), u_2(r)))\| & \leq c\|B(\Delta u(r), u_1(r))\|_{V_{-1/2}} + c\|B(u_2(r), \Delta u(r))\|_{V_{-1/2}} \\ & \leq c\|\Delta u(r)\|(\|u_1(r)\| + \|u_2(r)\|). \end{aligned}$$

Therefore

$$J_1 \leq \frac{c}{(t-s)^\beta} \int_s^t \|\Delta u(r)\|(\|u_1(r)\| + \|u_2(r)\|)dr \leq c\mu^{1-\beta} \|\Delta u\|_{C, t_0, t_0+\mu} (\|u_1\|_{C, t_0, t_0+\mu} + \|u_2\|_{C, t_0, t_0+\mu}).$$

Notice also that using the properties of the semigroup S

$$\begin{aligned} & \|(S(t-r) - S(s-r))A^{-\delta}(B(u_1(r), u_1(r)) - B(u_2(r), u_2(r)))\| \\ & = \|(S(t-s) - \text{id})S(s-r)(B(u_1(r), u_1(r)) - B(u_2(r), u_2(r)))\|_{V_{-\delta}} \\ & \leq c(t-s)^\delta \|B(u_1(r), u_1(r)) - B(u_2(r), u_2(r))\| \\ & \leq c(t-s)^\delta \|\Delta u(r)\|(\|u_1(r)\|_{V_{1/2}} + \|u_2(r)\|_{V_{1/2}}), \end{aligned}$$

and thus

$$\begin{aligned} J_2 & \leq \frac{1}{(t-s)^\beta} \int_{t_0}^s (t-s)^\delta \|\Delta u(r)\|(\|u_1(r)\|_{V_{1/2}} + \|u_2(r)\|_{V_{1/2}})dr \\ & \leq \mu^{\frac{1}{2}+\delta-\beta} \|\Delta u\|_{C, t_0, t_0+\mu} (\|u_1\|_{L^2(0, T; V_{1/2})} + \|u_2\|_{L^2(0, T; V_{1/2})}). \end{aligned}$$

To analyze the terms corresponding to the stochastic integral, that is,

$$\sup_{t_0 \leq s < t \leq t_0+\mu} \frac{\left\| \int_s^t S(t-r)(G(u_1(r)) - G(u_2(r)))d\omega - \int_{t_0}^s (S(t-r) - S(s-r))(G(u_1(r)) - G(u_2(r)))d\omega \right\|_{V_{-\delta}}}{(t-s)^\beta}$$

we can consider the estimates of I_4, I_5 given in the Appendix, replacing $\|A^{-\delta}(G(u(r)))\|_{L_2(V)}$ by

$$\|A^{-\delta}(G(u_1(r)) - G(u_2(r)))\|_{L_2(V)} \leq c_{DG} \|\Delta u(r)\|_{V_{-\delta}} \tag{6.1}$$

and $\|A^{-\delta}(G(u(r)) - G(u(q)))\|_{L_2(V)}$ by

$$\begin{aligned} & \|A^{-\delta}(G(u_1(r)) - G(u_2(r)) - (G(u_1(q)) - G(u_2(q))))\|_{L_2(V)} \\ & \leq c_{DG} \|\Delta u(r) - \Delta u(q)\|_{V_{-\delta}} + c_{D^2G} (\|\Delta u(r)\|_{V_{-\delta}} \|u_1(r) - u_1(q)\|_{V_{-\delta}} + \|u_2(r) - u_2(q)\|_{V_{-\delta}}), \end{aligned} \quad (6.2)$$

where these two above estimates follow by **(G)**. Then following the steps of [Lemma A.1](#) and taking into account

$$\|\Delta u(r)\|_{V_{-\delta}} = \|\Delta u(r) - \Delta u(t_0)\|_{V_{-\delta}} \leq \|\Delta u\|_{\beta, -\delta, t_0, t_0+\mu} (r - t_0)^\beta,$$

which is true due to the fact that $\Delta u(t_0) = 0$, we obtain the following term as an upper bound of the stochastic part:

$$c \|\omega\|_{\beta'} \mu^{\beta'} \|\Delta u\|_{\beta, -\delta, t_0, t_0+\mu} + c \|\omega\|_{\beta'} \mu^{\beta'} (\|u_1\|_{\beta, -\delta, 0, T} + \|u_2\|_{\beta, -\delta, 0, T}) \|\Delta u\|_{C, t_0, t_0+\mu}.$$

Collecting everything we get

$$\|\Delta u\|_{\beta, -\delta, t_0, t_0+\mu} \leq c_\mu^1 \|\Delta u\|_{\beta, -\delta, t_0, t_0+\mu} + c_\mu^2 \|\Delta u\|_{C, t_0, t_0+\mu}, \quad (6.3)$$

with

$$\begin{aligned} c_\mu^1 &= c \mu^{\beta'} \|\omega\|_{\beta'}, \\ c_\mu^2 &= c (\mu^{\frac{1}{2} + \delta - \beta} (\|u_1\|_{L^2(0, T; V_{1/2})} + \|u_2\|_{L^2(0, T; V_{1/2})}) + \mu^{\beta'} \|\omega\|_{\beta'} (\|u_1\|_{\beta, -\delta, 0, T} + \|u_2\|_{\beta, -\delta, 0, T}) \\ & \quad + \mu^{1-\beta} (\|u_1\|_{C, t_0, t_0+\mu} + \|u_2\|_{C, t_0, t_0+\mu})). \end{aligned} \quad (6.4)$$

(ii) In this second step we are interested in estimating $\|\Delta u\|_{C, t_0, t_0+\mu}$. The non-stochastic part gives us

$$\begin{aligned} & \sup_{t_0 \leq t \leq t_0+\mu} \left\| \int_{t_0}^t S(t-r) (B(u_1(r), u_1(r)) - B(u_2(r), u_2(r))) dr \right\| \\ & \leq c \sup_{t_0 \leq t \leq t_0+\mu} \int_{t_0}^t \|\Delta u(r)\| (\|u_1(r)\|_{V_{1/2}} + \|u_2(r)\|_{V_{1/2}}) dr \\ & \leq c \mu^{\frac{1}{2}} (\|u_1\|_{L^2(0, T; V_{1/2})} + \|u_2\|_{L^2(0, T; V_{1/2})}) \|\Delta u\|_{C, t_0, t_0+\mu}. \end{aligned}$$

To study the norm of the stochastic integral, for $t \in [t_0, t_0 + \mu]$ we split it as follows

$$\begin{aligned} & \|\omega\|_{\beta'} \int_{t_0}^t (t-r)^{\alpha+\beta'-1} \left(\frac{\|S(t-r)(G(u_1(r)) - G(u_2(r)))\|_{L_2(V)}}{(r-t_0)^\alpha} \right. \\ & \quad + \int_{t_0}^r \frac{\|S(t-r) - S(t-\hat{r})\| (G(u_1(r)) - G(u_2(r)))\|_{L_2(V)}}{(r-\hat{r})^{\alpha+1}} d\hat{r} \\ & \quad \left. + \int_{t_0}^r \frac{\|S(t-\hat{r})(G(u_1(r)) - G(u_2(r))) - (G(u_1(\hat{r})) - G(u_2(\hat{r})))\|_{L_2(V)}}{(r-\hat{r})^{\alpha+1}} d\hat{r} \right) dr \\ & =: J_3(t) + J_4(t) + J_5(t). \end{aligned}$$

Following the steps of [Lemma A.1](#), thanks to **(G)** we obtain

$$\sup_{t_0 \leq t \leq t_0+\mu} J_3(t) \leq c \|\omega\|_{\beta'} \mu^{\beta'} \|\Delta u\|_{C, t_0, t_0+\mu},$$

$$\sup_{t_0 \leq t \leq t_0+\mu} J_4(t) \leq c \|\omega\|_{\beta'} \mu^{\beta'} \|\Delta u\|_{C, t_0, t_0+\mu}.$$

Finally, using again **(G)**, since $\|\Delta u(r)\|_{V_{-\delta}} \leq c \|\Delta u(r)\|$,

$$\begin{aligned} \sup_{t_0 \leq t \leq t_0+\mu} J_5(t) & \leq c \|\omega\|_{\beta'} \int_{t_0}^t (t-r)^{\alpha+\beta'-1} \\ & \quad \times \left(\int_{t_0}^r \frac{\|\Delta u(r) - \Delta u(\hat{r})\|_{V_{-\delta}} + \|\Delta u(r)\| (\|u_1(r) - u_1(\hat{r})\|_{V_{-\delta}} + \|u_2(r) - u_2(\hat{r})\|_{V_{-\delta}})}{(r-\hat{r})^{\alpha+1}} d\hat{r} \right) dr \\ & \leq c \|\omega\|_{\beta'} (\|\Delta u\|_{\beta, -\delta, t_0, t_0+\mu} + \|\Delta u\|_{C, t_0, t_0+\mu} (\|u_1\|_{\beta, -\delta, t_0, t_0+\mu} + \|u_2\|_{\beta, -\delta, t_0, t_0+\mu})) \\ & \quad \times \sup_{t_0 \leq t \leq t_0+\mu} \int_{t_0}^t (t-r)^{\alpha+\beta'-1} \left(\int_{t_0}^r (r-\hat{r})^{\beta-\alpha-1} d\hat{r} \right) dr \\ & \leq c \|\omega\|_{\beta'} \mu^{\beta+\beta'} (\|\Delta u\|_{\beta, -\delta, t_0, t_0+\mu} + \|\Delta u\|_{C, t_0, t_0+\mu} (\|u_1\|_{\beta, -\delta, 0, T} + \|u_2\|_{\beta, -\delta, 0, T})). \end{aligned}$$

Hence,

$$\|\Delta u\|_{C, t_0, t_0+\mu} \leq c_\mu^3 \|\Delta u\|_{C, t_0, t_0+\mu} + c_\mu^4 \|\Delta u\|_{\beta, -\delta, t_0, t_0+\mu}, \quad (6.5)$$

with

$$\begin{aligned} c_\mu^3 &= c (\mu^{\frac{1}{2}} (\|u_1\|_{L^2(0, T; V_{1/2})} + \|u_2\|_{L^2(0, T; V_{1/2})}) + \mu^{\beta'} \|\omega\|_{\beta'} + \mu^{\beta'+\beta} \|\omega\|_{\beta'} (\|u_1\|_{\beta, -\delta, 0, T} + \|u_2\|_{\beta, -\delta, 0, T})), \\ c_\mu^4 &= c \mu^{\beta'+\beta} \|\omega\|_{\beta'}. \end{aligned} \quad (6.6)$$

Therefore, solving the system given by (6.3) and (6.5) means that we have to solve a system of inequalities, namely

$$X \leq c_\mu^1 X + c_\mu^2 Y, \quad Y \leq c_\mu^3 Y + c_\mu^4 X$$

with c_μ^i given by (6.4) and (6.6). It is now straightforward to check that for a small enough $\mu \in (0, 1)$ we obtain that $\|\Delta u\|_{C, t_0, t_0+\mu} = 0$, which contradicts the fact that the maximal interval of uniqueness is $[0, t_0]$. Hence the solution of (2.5) is unique. \square

Finally, we can prove the main theorem of the paper:

Theorem 6.2. *Under the assumptions of Lemma 5.3 there exists a mild solution to the stochastic shell-model (2.5) with driving function $\omega \in C^{\beta'}([0, T]; V)$.*

Proof. We divide the proof in several steps:

(i) Let $(\omega_n)_{n \in \mathbb{N}}$ be a sequence of piecewise linear continuous functions converging to ω in $C^{\beta'}([0, T]; V)$, see Remark 3.2, and let $(u_n)_{n \in \mathbb{N}}$ be the sequence of unique solutions driven by $(\omega_n)_{n \in \mathbb{N}}$ with initial condition $u_0 \in V$. From Lemma 5.5 we know that $(u_n)_{n \in \mathbb{N}}$ is uniformly bounded in $C^{\hat{\beta}}([0, T]; V_{-\delta}) \cap C([0, T]; V)$. Then (5.5) implies that $(\|u_n\|_{L^2(0, T; V_{1/2})})_{n \in \mathbb{N}}$ is also bounded and hence $(u_n)_{n \in \mathbb{N}}$ is relatively weak compact in $L^2(0, T; V_{1/2})$. Furthermore, this sequence is relatively compact in $L^2(0, T; V) \cap C([0, T]; V_{-\delta})$ by Theorem 2.2(i).

Moreover, thanks to Corollary A.2(ii), for $\hat{\delta} = \delta - \varepsilon$ with $\varepsilon > 0$ arbitrarily small, we have

$$\|u_n\|_{\hat{\beta}, -\hat{\delta}, 0, t} \leq ct^{\hat{\delta}-\hat{\beta}} \|u_n(0)\| + c(t^{1-\hat{\beta}} + t^{\hat{\delta}-1/2-\hat{\beta}}) \|u_n\|_{C, 0, t}^2 + c\|\omega_n\|_{\beta'} t^{\beta'-\hat{\beta}-\varepsilon} (1 + t^{\hat{\beta}} \|u_n\|_{\hat{\beta}, -\hat{\delta}, 0, t}). \tag{6.7}$$

Following a similar reasoning to the one of Lemma A.5, we can make small the coefficient in front of $\|u_n\|_{\hat{\beta}, -\hat{\delta}, 0, t}$ on the left hand side of (6.7) such that we can solve such an inequality. Moreover, since $(\omega_n)_{n \in \mathbb{N}}$ converges to ω , we obtain that $(u_n)_{n \in \mathbb{N}}$ is uniformly bounded in $C^{\hat{\beta}}([0, T]; V_{-\delta})$. Hence, by Theorem 2.2(ii), this sequence is relatively compact in $C^{\beta}([0, T]; V_{-\delta})$ with $\beta < \hat{\beta}$.

(ii) Let $(u_{n'})_{n' \in \mathbb{N}}$ be a subsequence converging to some limit point $u \in L^2(0, T; V_{1/2}) \cap C^{\beta}([0, T]; V_{-\delta})$. Let us denote this subsequence simply by $(u_n)_{n \in \mathbb{N}}$. Then, since $B : V_{\frac{1}{2}} \times V_{-\delta} \rightarrow V_{-\delta}$ and also $B : V_{-\delta} \times V_{\frac{1}{2}} \rightarrow V_{-\delta}$ and $u_n(0) - u(0) = 0$, applying Lemma 2.1 we have

$$\begin{aligned} & \left\| \int_0^t S(t-r)(B(u_n(r), u_n(r)) - B(u(r), u(r))) dr \right\|_{V_{-\delta}} \\ & \leq \int_0^t (\|B(u_n(r), u_n(r)) - B(u(r), u_n(r))\|_{V_{-\delta}} + \|B(u(r), u_n(r)) - B(u(r), u(r))\|_{V_{-\delta}}) dr \\ & \leq c \int_0^t (\|u_n(r)\|_{V_{1/2}} + \|u(r)\|_{V_{1/2}}) \|u(r) - u_n(r)\|_{V_{-\delta}} dr \\ & \leq c \|u - u_n\|_{\beta, -\delta, 0, T} \int_0^t r^{\beta} (\|u_n(r)\|_{V_{1/2}} + \|u(r)\|_{V_{1/2}}) dr \\ & \leq cT^{\beta+\frac{1}{2}} \|u - u_n\|_{\beta, -\delta, 0, T} (\|u_n\|_{L^2(0, T; V_{1/2})} + \|u\|_{L^2(0, T; V_{1/2})}) \end{aligned}$$

which shows the convergence in $V_{-\delta}$ of the left hand side to zero.

For the stochastic integral we consider the splitting

$$\begin{aligned} & \left\| \int_0^t S(t-r)G(u_n(r))d\omega_n(r) - \int_0^t S(t-r)G(u(r))d\omega(r) \right\|_{V_{-\delta}} \\ & \leq \left\| \int_0^t S(t-r)G(u_n(r))d(\omega_n(r) - \omega(r)) \right\|_{V_{-\delta}} + \left\| \int_0^t S(t-r)(G(u_n(r)) - G(u(r)))d\omega_n(r) \right\|_{V_{-\delta}}. \end{aligned}$$

Similar to (5.12), an upper bound for the first integral on the right hand side is given by

$$cT^{\beta'} \|\omega_n - \omega\|_{\beta'} (1 + T^{\beta} \|u_n\|_{\beta, -\delta, 0, T})$$

and since the set $\{\|u_n\|_{\beta, -\delta, 0, T}\}_{n \in \mathbb{N}}$ is bounded, we obtain the convergence in $V_{-\delta}$ of the first integral on the right hand side. Now using (6.1)-(6.2), setting $u_1 = u_n$, $u_2 = u$ we arrive at

$$\left\| \int_0^t S(t-r)(G(u_n(r)) - G(u(r)))d\omega_n(r) \right\|_{V_{-\delta}} \leq c\|\omega_n\|_{\beta'} T^{\beta'} \|u_n - u\|_{\beta, -\delta, 0, T} (1 + T^{\beta} (1 + \|u_n\|_{\beta, -\delta, 0, T} + \|u\|_{\beta, -\delta, 0, T}))$$

which shows the convergence in $V_{-\delta}$ of the second integral.

Also, since $(u_n)_{n \in \mathbb{N}}$ converges to u in $C([0, T]; V_{-\delta})$, for every $t \in [0, T]$ we have that $u_n(t) \rightarrow u(t)$ in $V_{-\delta}$.

(iii) Since $u \in L^2(0, T; V_{1/2}) \cap L^\infty(0, T; V)$ we have that $t \mapsto B(u(t), u(t)) \in L^2(0, T; V)$ and hence the continuity in V of the first integral of (4.2) with respect to t follows. Moreover, since $u \in C^{\beta}([0, T]; V_{-\delta})$ by (G) we obtain that

$$t \mapsto \int_0^t S(t-r)G(u(r))d\omega \in C([0, T]; V).$$

(iv) Collecting the above properties, on the one hand (i)-(ii) mean that $u \in C^{\beta}([0, T]; V_{-\delta}) \cap L^2(0, T; V_{1/2})$ and u satisfies (4.2) in $V_{-\delta}$. On the other hand, (iii) means that the right hand side of (4.2) belongs to $C([0, T]; V)$, and hence also the left hand side. In conclusion, we have proven the existence of a mild solution u to the stochastic shell-model in the sense of Definition 4.1. \square

7. An example of diffusion term

We define the operator G by a sequence of functions $g_m^n(u) \in \mathbb{C}$ with $u \in V_{-\delta}$, such that for $v \in V$:

$$(G(u)v)_n := \sum_{m=1}^{\infty} g_m^n(u)v_m. \quad (7.1)$$

We now define properties for this sequence such that G satisfies the hypotheses **(G)**. For every $n, m = 1, \dots$, assume that

$$\sup_{u \in V_{-\delta}} \sum_{n,m=1}^{\infty} |g_m^n(u)|^2 =: c_G^2 < \infty. \quad (7.2)$$

In addition, let us assume that the functions g_m^n are twice differentiable having the following properties: For $u, h \in V_{-\delta}$ and $(f_k)_{k \in \mathbb{N}}$ an orthonormal base in $V_{-\delta}$ we have that

$$\begin{aligned} \sum_{n,m=1}^{\infty} (g_m^n(u+h) - g_m^n(u) - Dg_m^n(u)h)^2 &= \sum_{n,m=1}^{\infty} (o_u^{n,m}(\|h\|_{V_{-\delta}}))^2 = o_u(\|h\|_{V_{-\delta}})^2, \\ \sup_{u \in V_{-\delta}} \sum_{n,m,k=1}^{\infty} |Dg_m^n(u)f_k|^2 &=: c_{DG}^2 < \infty. \end{aligned} \quad (7.3)$$

The $o_u^{n,m}$, o_u have the usual properties: $\lim_{h \rightarrow 0} |o_u^{n,m}(\|h\|_{V_{-\delta}})|/\|h\|_{V_{-\delta}} = 0$ and similar for o_u . In addition we assume that for $u, h_1, h_2 \in V_{-\delta}$

$$\begin{aligned} \sum_{n,m=1}^{\infty} (Dg_m^n(u+h_2)h_1 - Dg_m^n(u)h_1 - D^2g_m^n(u)h_1h_2)^2 &= \sum_{n,m=1}^{\infty} (o_{u,h_1}^{n,m}(\|h_2\|_{V_{-\delta}}))^2 =: (o_{u,h_1}(\|h_2\|_{V_{-\delta}}))^2, \\ \sup_{u \in V_{-\delta}} \sum_{n,m,k,l=1}^{\infty} |D^2g_m^n(u)(f_k, f_l)|^2 &=: c_{D^2G}^2 < \infty, \end{aligned} \quad (7.4)$$

where the $o_{u,h_1}^{n,m}$, o_{u,h_1} have the same property as above.

Now we can verify the properties of the operator G formulated in hypothesis **(G)**. It follows from (7.2) that

$$\begin{aligned} \sup_{u \in V_{-\delta}} \|G(u)\|_{L_2(V)}^2 &= \sup_{u \in V_{-\delta}} \sum_{m=1}^{\infty} \|G(u)e_m\|^2 = \sup_{u \in V_{-\delta}} \sum_{n,m=1}^{\infty} |(G(u)e_m)_n|^2 \\ &= \sup_{u \in V_{-\delta}} \sum_{n,m=1}^{\infty} |g_m^n(u)|^2 = c_G^2. \end{aligned}$$

Simple calculations show that (7.3), (7.4) imply that the operators DG and D^2G exist and are bounded. In fact, if $u, h \in V_{-\delta}$, then we have that

$$\|G(u+h) - G(u) - DG(u)h\|_{L_2(V)}^2 = (o(\|h\|_{V_{-\delta}}))^2$$

and

$$\sup_{u \in V_{-\delta}} \|DG(u)\|_{L_2(V \times V_{-\delta}, V)}^2 = \sup_{u \in V_{-\delta}} \sum_{m,k=1}^{\infty} \|DG(u)(e_m, f_k)\|^2 = \sup_{u \in V_{-\delta}} \sum_{n,m,k=1}^{\infty} |Dg_m^n(u)f_k|^2 = c_{DG}^2.$$

Now, using the boundedness of DG we can prove the Lipschitz condition. Similarly, (7.4) implies that the operator D^2G exists and is bounded. Using the boundedness of the second derivative of G standard calculations give (4.1).

Acknowledgment

Hakima Bessaih's research was supported in part by National Science Foundation grant DMS-1418838. Maria Jose Garrido Atienza's research has been supported by MTM2015-63723-P.

Appendix

We start this section by completing the proof of Lemma 5.4, although in the next result (item (i)) we prove a bit more.

Lemma A.1. (i) Let I_4, I_5 be defined in (5.8). Then for any sufficient small $\varepsilon \geq 0$ such that $\beta' - \hat{\beta} > \varepsilon$ we have

$$\sup_{0 \leq p < q \leq t} \frac{\|A^\varepsilon I_4\| + \|A^\varepsilon I_5\|}{(q-p)^{\hat{\beta}}} \leq ct^{\beta' - \hat{\beta} - \varepsilon} \|\omega\|_{\beta'} \left(1 + t^{\hat{\beta}} \|u\|_{\hat{\beta}, -\delta, 0, t}\right).$$

(ii) Let I_2, I_3 be defined in (5.8) and let $0 \leq \varepsilon < \delta - 1/2$, $\beta' - \hat{\beta} > \varepsilon$. Then

$$\sup_{0 \leq p < q \leq t} \frac{\|A^\varepsilon I_2\| + \|A^\varepsilon I_3\|}{(q-p)^{\hat{\beta}}} \leq c(t^{1-\hat{\beta}} + t^{\delta+1/2-\hat{\beta}-\varepsilon}) \|u\|_{\mathbb{C}, 0, t}^2.$$

Proof. Throughout the proof we will use frequently the properties (2.1), (2.2) and (5.4). We choose an α in the same conditions than in Lemma 5.3, that is, $1 - \beta' < \alpha < \hat{\beta}$. We also use the property $L_2(V) \subset L_2(V, V_{-\delta})$.

First, using the definition of the stochastic integral and the estimate (5.6),

$$\begin{aligned} \sup_{0 \leq p < q \leq t} \frac{\|A^\varepsilon I_4\|}{(q-p)^{\hat{\beta}}} &\leq \sup_{0 \leq p < q \leq t} \frac{1}{(q-p)^{\hat{\beta}}} \|\omega\|_{\beta'} \int_p^q (q-r)^{\alpha+\beta'-1} \left(\frac{\|S(q-r)A^\varepsilon A^{-\delta}G(u(r))\|_{L_2(V)}}{(r-p)^\alpha} \right. \\ &+ \left. \int_p^r \frac{\|(S(q-r) - S(q-\hat{r}))A^\varepsilon A^{-\delta}G(u(r))\|_{L_2(V)}}{(r-\hat{r})^{\alpha+1}} d\hat{r} + \int_p^r \frac{\|S(q-\hat{r})A^\varepsilon A^{-\delta}(G(u(r)) - G(u(\hat{r})))\|_{L_2(V)}}{(r-\hat{r})^{\alpha+1}} d\hat{r} \right) dr. \end{aligned}$$

The first term is estimated by

$$\frac{\|S(q-r)A^\varepsilon A^{-\delta}G(u(r))\|_{L_2(V)}}{(r-p)^\alpha} \leq \frac{cC_G}{(r-p)^\alpha (q-r)^\varepsilon}$$

and since $\alpha + \beta' - \varepsilon > 0$, we get

$$\sup_{0 \leq p < q \leq t} \frac{cC_G}{(q-p)^{\hat{\beta}}} \|\omega\|_{\beta'} \int_p^q (q-r)^{\alpha+\beta'-\varepsilon-1} (r-p)^{-\alpha} dr \leq c \|\omega\|_{\beta'} t^{\beta'-\hat{\beta}-\varepsilon}.$$

Concerning the second term, taking an appropriate $\alpha' > \alpha$ such that $\alpha + \beta' > \alpha' + \varepsilon$, we have

$$\int_p^r \frac{\|(S(q-r) - S(q-\hat{r}))A^\varepsilon A^{-\delta}G(u(r))\|_{L_2(V)}}{(r-\hat{r})^{\alpha+1}} d\hat{r} \leq \frac{cC_G}{(q-r)^{\alpha'+\varepsilon}} \int_p^r \frac{(r-\hat{r})^{\alpha'}}{(r-\hat{r})^{\alpha+1}} d\hat{r} \leq \frac{cC_G(r-p)^{\alpha'-\alpha}}{(q-r)^{\alpha'+\varepsilon}},$$

and hence

$$\sup_{0 \leq p < q \leq t} \frac{cC_G}{(q-p)^{\hat{\beta}}} \|\omega\|_{\beta'} \int_p^q \frac{(r-p)^{\alpha'-\alpha}}{(q-r)^{\alpha'+\varepsilon}} (q-r)^{\alpha+\beta'-1} dr \leq c \|\omega\|_{\beta'} t^{\beta'-\hat{\beta}-\varepsilon}.$$

Finally, since $\hat{\beta} > \alpha$

$$\begin{aligned} \int_p^r \frac{\|S(q-\hat{r})A^\varepsilon A^{-\delta}(G(u(r)) - G(u(\hat{r})))\|_{L_2(V)}}{(r-\hat{r})^{\alpha+1}} d\hat{r} &\leq c \int_p^r \frac{\|A^{-\delta}(G(u(r)) - G(u(\hat{r})))\|_{L_2(V)}}{(r-\hat{r})^{\alpha+1} (q-\hat{r})^\varepsilon} d\hat{r} \\ &\leq cC_{DG} \|u\|_{\hat{\beta}, -\delta, 0, t} \frac{1}{(q-r)^\varepsilon} \int_p^r \frac{(r-\hat{r})^{\hat{\beta}}}{(r-\hat{r})^{\alpha+1}} d\hat{r} \leq cC_{DG} \|u\|_{\hat{\beta}, -\delta, 0, t} \frac{(r-p)^{\hat{\beta}-\alpha}}{(q-r)^\varepsilon}, \end{aligned}$$

and since $\beta' + \alpha - \varepsilon > 0$ we have

$$\sup_{0 \leq p < q \leq t} \frac{cC_{DG} \|u\|_{\hat{\beta}, -\delta, 0, t}}{(q-p)^{\hat{\beta}}} \|\omega\|_{\beta'} \int_p^q (q-r)^{\alpha+\beta'-\varepsilon-1} (r-p)^{\hat{\beta}-\alpha} dr \leq c \|\omega\|_{\beta'} \|u\|_{\hat{\beta}, -\delta, 0, t} t^{\beta'-\varepsilon}.$$

Hence, we get that

$$\sup_{0 \leq p < q \leq t} \frac{\|A^\varepsilon I_4\|}{(q-p)^{\hat{\beta}}} \leq ct^{\beta'-\hat{\beta}-\varepsilon} \|\omega\|_{\beta'} \left(1 + t^{\hat{\beta}} \|u\|_{\hat{\beta}, -\delta, 0, t} \right).$$

Thanks to the definition of the stochastic integral and the estimate (5.6) for I_5 we get

$$\begin{aligned} \sup_{0 \leq p < q \leq t} \frac{\|A^\varepsilon I_5\|}{(q-p)^{\hat{\beta}}} &\leq \sup_{0 \leq p < q \leq t} \frac{1}{(q-p)^{\hat{\beta}}} \|\omega\|_{\beta'} \int_0^p (p-r)^{\alpha+\beta'-1} \left(\frac{\|(S(q-r) - S(p-r))A^\varepsilon A^{-\delta}G(u(r))\|_{L_2(V)}}{r^\alpha} \right. \\ &+ \int_0^r \frac{\|(S(q-\hat{r}) - S(p-\hat{r}))A^\varepsilon A^{-\delta}(G(u(r)) - G(u(\hat{r})))\|_{L_2(V)}}{(r-\hat{r})^{\alpha+1}} d\hat{r} \\ &+ \left. \int_0^r \frac{\|(S(q-r) - S(q-\hat{r}) - S(p-r) + S(p-\hat{r}))A^\varepsilon A^{-\delta}G(u(r))\|_{L_2(V)}}{(r-\hat{r})^{\alpha+1}} d\hat{r} \right) dr \\ &=: \sup_{0 \leq p < q \leq t} \frac{1}{(q-p)^{\hat{\beta}}} \|\omega\|_{\beta'} \int_0^p (p-r)^{\alpha+\beta'-1} (I_{5,1} + I_{5,2} + I_{5,3}) dr. \end{aligned}$$

We start with

$$I_{5,1} = \frac{\|(S(q-p) - \text{Id})S(p-r)A^\varepsilon A^{-\delta}G(u(r))\|_{L_2(V)}}{r^\alpha} \leq \frac{cC_G(q-p)^{\hat{\beta}}}{(p-r)^{\hat{\beta}+\varepsilon} r^\alpha}$$

and because $\alpha < 1$ and $\alpha + \beta' - \hat{\beta} - \varepsilon > 0$, the term involving $I_{5,1}$ is estimated by

$$\sup_{0 \leq p < q \leq t} \frac{cC_G}{(q-p)^{\hat{\beta}}} \|\omega\|_{\beta'} (q-p)^{\hat{\beta}} \int_0^p (p-r)^{\alpha+\beta'-1-\hat{\beta}-\varepsilon} r^{-\alpha} dr \leq c \|\omega\|_{\beta'} t^{\beta'-\hat{\beta}-\varepsilon}.$$

On the other hand,

$$\begin{aligned}
 I_{5,2} &= \int_0^r \frac{\|(S(q-p) - \text{Id})S(p-\hat{r})A^\varepsilon A^{-\delta}(G(u(r)) - G(u(\hat{r})))\|_{L_2(V)} d\hat{r}}{(r-\hat{r})^{\alpha+1}} \\
 &\leq c c_{DG} \int_0^r \frac{(p-\hat{r})^{-\hat{\beta}-\varepsilon}(q-p)^{\hat{\beta}}\|u(r) - u(\hat{r})\|_{V_{-\delta}}}{(r-\hat{r})^{\alpha+1}} d\hat{r} \\
 &\leq c c_{DG} \|u\|_{\hat{\beta}, -\delta, 0, t} (p-r)^{-\hat{\beta}-\varepsilon}(q-p)^{\hat{\beta}} \int_0^r \frac{1}{(r-\hat{r})^{\alpha+1-\hat{\beta}}} d\hat{r} \\
 &\leq c c_{DG} \|u\|_{\hat{\beta}, -\delta, 0, t} (p-r)^{-\hat{\beta}-\varepsilon}(q-p)^{\hat{\beta}} r^{\hat{\beta}-\alpha},
 \end{aligned}$$

and thus

$$\begin{aligned}
 &\sup_{0 \leq p < q \leq t} \frac{1}{(q-p)^{\hat{\beta}}} \|\omega\|_{\beta'} \int_0^p (p-r)^{\alpha+\beta'-1} I_{5,2} dr \\
 &\leq c c_{DG} \|\omega\|_{\beta'} \|u\|_{\hat{\beta}, -\delta, 0, t} \sup_{0 \leq p < q \leq t} \int_0^p (p-r)^{\alpha+\beta'-1-\hat{\beta}-\varepsilon} r^{\hat{\beta}-\alpha} dr \\
 &\leq c c_{DG} \|\omega\|_{\beta'} \|u\|_{\hat{\beta}, -\delta, 0, t} t^{\beta'-\varepsilon}.
 \end{aligned}$$

Finally, taking α' close enough to α such that $\alpha' > \alpha$ and $\alpha + \beta' > \alpha' + \hat{\beta} + \varepsilon$ (for a small enough ε), applying the second part of (2.3)

$$\begin{aligned}
 I_{5,3} &\leq c \int_0^r \frac{(q-p)^{\hat{\beta}}(r-\hat{r})^{\alpha'}(p-r)^{-\alpha'-\hat{\beta}-\varepsilon} \|A^{-\delta}G(u(r))\|_{L_2(V)} d\hat{r}}{(r-\hat{r})^{\alpha+1}} \\
 &\leq c c_G (q-p)^{\hat{\beta}} (p-r)^{-\alpha'-\hat{\beta}-\varepsilon} \int_0^r (r-\hat{r})^{\alpha'-\alpha-1} d\hat{r} \\
 &\leq c c_G (q-p)^{\hat{\beta}} (p-r)^{-\alpha'-\hat{\beta}-\varepsilon} r^{\alpha'-\alpha},
 \end{aligned}$$

and hence

$$\begin{aligned}
 &\sup_{0 \leq p < q \leq t} \frac{1}{(q-p)^{\hat{\beta}}} \|\omega\|_{\beta'} \int_0^p (p-r)^{\alpha+\beta'-1} I_{5,3} dr \\
 &\leq c c_G \|\omega\|_{\beta'} \sup_{0 \leq p < q \leq t} \int_0^p (p-r)^{\alpha+\beta'-1-\alpha'-\hat{\beta}-\varepsilon} r^{\alpha'-\alpha} dr \\
 &\leq c c_G \|\omega\|_{\beta'} t^{\beta'-\hat{\beta}-\varepsilon}.
 \end{aligned}$$

Taking into account the previous estimates we finally get

$$\sup_{0 \leq p < q \leq t} \frac{\|A^\varepsilon I_5\|}{(q-p)^{\hat{\beta}}} \leq c t^{\beta'-\hat{\beta}-\varepsilon} \|\omega\|_{\beta'} \left(1 + t^{\hat{\beta}} \|u\|_{\hat{\beta}, -\delta, 0, t}\right).$$

(ii) The proof of this part follows similarly to the estimates (5.10) and (5.11). In particular, for the estimate of $\|A^\varepsilon I_2\|$ we need to use the continuous embedding $V \subset V_{-\delta+\varepsilon+1/2}$, which holds true for small enough $\varepsilon \geq 0$ since $\delta \in (\hat{\beta}, 1)$. \square

Corollary A.2. (i) We derive (5.12) simply taking $\varepsilon = 0$ in Lemma A.1(i).

(ii) As a consequence of Lemma A.1 we can prove the inequality (6.7):

$$\|u_n\|_{\hat{\beta}, -\delta, 0, t} \leq c t^{\hat{\delta}-\hat{\beta}} \|u_n(0)\| + c(t^{1-\hat{\beta}} + t^{\hat{\delta}-1/2-\hat{\beta}}) \|u_n\|_{C, 0, t}^2 + c \|\omega_n\|_{\beta'} t^{\beta'-\hat{\beta}-\varepsilon} (1 + t^{\hat{\beta}} \|u_n\|_{\hat{\beta}, -\delta, 0, t}).$$

Note that, since $\hat{\delta} = \delta - \varepsilon$, now we have the splitting

$$\begin{aligned}
 A^{-\hat{\delta}}(u_n(q) - u_n(p)) &= A^{-\hat{\delta}}(S(q) - S(p))u_n(0) + A^\varepsilon A^{-\delta+1/2} \int_p^q S(q-r)A^{-1/2}B(u_n(r), u_n(r))dr \\
 &\quad + A^\varepsilon A^{-\delta+1/2} \int_0^p (S(q-r) - S(p-r))A^{-1/2}B(u_n(r), u_n(r))dr \\
 &\quad + A^\varepsilon A^{-\delta} \int_p^q S(q-r)G(u_n(r))\omega'_n(r)dr + A^\varepsilon A^{-\delta} \int_0^p (S(q-r) - S(p-r))G(u_n(r))\omega'_n(r)dr \\
 &= A^\varepsilon I_1 + A^\varepsilon I_2 + A^\varepsilon I_3 + A^\varepsilon I_4 + A^\varepsilon I_5,
 \end{aligned}$$

where for $j = 2, 3, 4, 5$ the I_j agree with the integrals previously defined in (5.8). Firstly, similar to (5.9), we obtain that

$$\sup_{0 \leq p < q \leq t} \frac{\|A^\varepsilon I_1\|}{(q-p)^{\hat{\beta}}} \leq \bar{c} t^{\hat{\delta}-\hat{\beta}} \|u_n(0)\|,$$

and by Lemma A.1(ii)

$$\sup_{0 \leq p < q \leq t} \frac{\|A^\varepsilon I_2\| + \|A^\varepsilon I_3\|}{(q - p)^{\hat{\beta}}} \leq c(t^{1-\hat{\beta}} + t^{\hat{\delta}+1/2-\hat{\beta}}) \|u_n\|_{C,0,t}^2.$$

Secondly, on account of the fact that $L_2(V) \subset L_2(V, V_{-\delta})$ and that $V_{-\delta} \subset V_{-\delta}$, from assumption (G) for instance we derive that

$$\|A^{-\delta}(G(u_1) - G(u_2))\|_{L_2(V)} \leq \|G(u_1) - G(u_2)\|_{L_2(V)} \leq c_{DG} \|u_1 - u_2\|_{V_{-\delta}} \leq c \|u_1 - u_2\|_{V_{-\delta}},$$

and therefore, following the same steps than in the proof of Lemma A.1(i) we can deduce

$$\sup_{0 \leq p < q \leq t} \frac{\|A^\varepsilon I_4\| + \|A^\varepsilon I_5\|}{(q - p)^{\hat{\beta}}} \leq ct^{\beta' - \hat{\beta} - \varepsilon} \|\omega_n\|_{\beta'} \left(1 + t^{\hat{\beta}} \|u_n\|_{\hat{\beta}, -\hat{\delta}, 0, t}\right),$$

and this concludes the proof of (6.7).

The rest of the Appendix section is devoted to the proof of Lemma 5.5, which relies upon several results that are proven below.

Lemma A.3. Let $1/2 < \hat{\beta} < \tilde{\beta} < \delta$ and suppose that $u \in C^{\tilde{\beta}}([0, T]; V_{-\delta})$. Then the mapping

$$[s, T] \ni t \mapsto \|u\|_{\hat{\beta}, -\delta, s, t}$$

is continuous and

$$\lim_{t \rightarrow s^+} \|u\|_{\hat{\beta}, -\delta, s, t} = 0.$$

Proof. We only consider here the case $s = 0$. Let us define the following transformation of u given by

$$\hat{u}_{\hat{t}}(r) = \begin{cases} u(r) & : r \leq \hat{t}, \\ u(\hat{t}) & : r \geq \hat{t}. \end{cases}$$

Then for $0 \leq \hat{t} < t \leq T$

$$\|u\|_{\hat{\beta}, -\delta, 0, t} - \|u\|_{\hat{\beta}, -\delta, 0, \hat{t}} = \|u\|_{\hat{\beta}, -\delta, 0, t} - \|\hat{u}_{\hat{t}}\|_{\hat{\beta}, -\delta, 0, t} \leq \|u\|_{\hat{\beta}, -\delta, \hat{t}, t} \leq c(t - \hat{t})^{\tilde{\beta} - \hat{\beta}} \|u\|_{\tilde{\beta}, -\delta, 0, T}$$

from which the desired continuity follows immediately. The convergence to 0 follows in the same way. \square

Lemma A.4. For positive continuous functions $a(t)$, $b(t)$ consider

$$Y = b(t) + a(t)Y^2$$

and assume $4a(t)b(t) < 1$ for every $t \in [0, t_1]$, where $t_1 > 0$ is some positive number. Then there exist two real solutions $Y_1(t) < Y_2(t) \in \mathbb{R}^+$ given by

$$Y_1(t) = \frac{1}{2a(t)} (1 - \sqrt{1 - 4a(t)b(t)}), \quad Y_2(t) = \frac{1}{2a(t)} (1 + \sqrt{1 - 4a(t)b(t)})$$

where $Y_1(t) \leq 2b(t)$. Suppose in addition that $y(t) \geq 0$ is continuous on $[0, t_1]$ such that

$$y(t) \leq b(t) + a(t)y(t)^2, \quad \lim_{t \rightarrow 0^+} y(t) = 0,$$

and that $\lim_{t \rightarrow 0^+} a(t) = 0$. Then we have $y(t) \leq Y_1(t)$ on $[0, t_1]$.

Proof. It follows by Sohr [31, Page 317] that under the conditions of the lemma there exist real solutions Y_1, Y_2 satisfying the above conditions.

On the other hand, y satisfies the above inequality if and only if $y(t) \leq Y_1(t)$ or $y(t) \geq Y_2(t)$. If $y(t) \geq Y_2(t)$ for some $t \in (0, t_1]$ then by the continuity of y , Y_2 and by the fact that $Y_2(t) > Y_1(t)$ on $(0, t_1]$, it follows that $y(t) \geq Y_2(t)$ on $[0, t_1]$. However, under the assumptions we have $\lim_{t \rightarrow 0^+} Y_2(t) = +\infty$ and this is a contradiction with respect to the behavior of y . \square

To simplify the presentation of the following technical result we assume that $T = 1$. Moreover, in what follows, see Lemma A.5, we shall consider inequalities of the type

$$y(t) \leq d(t, x)y(t) + f(t, x) + h(t)y(t)^2, \quad t \in [0, t_1] \tag{A.1}$$

where the increasing functions $d(\cdot, x)$, $f(\cdot, x)$, $h(\cdot)$ are defined by

$$\begin{aligned} d(t, x) &= ct^{\beta'} + 4c^3t^{1+2\beta'} + 2c^2xt^{1+\beta'}, \\ f(t, x) &= \bar{c}xt^{\delta-\hat{\beta}} + cx^2t^{1-\hat{\beta}} + c^2xt^{1+\beta'-\hat{\beta}} + c^3t^{1+2\beta'-\hat{\beta}} + ct^{\beta'-\hat{\beta}}, \\ h(t) &= 4c^3t^{1+2\beta'+\hat{\beta}}. \end{aligned} \tag{A.2}$$

Note that $d(t, x)$ and $f(t, x)$ depend on a positive parameter x . Furthermore, the constants \bar{c} , c , c^2 , c^3 are coming from the estimates of Lemmas 5.3 and 5.4, as we will show in Lemma A.5. In particular, these constants are constricted such that they are including the value $\|\omega\|_{\beta'} = \|\omega\|_{\beta', 0, 1}$. In the following proof we need these constants with norms only for subintervals of $[0, 1]$. However, using $\|\omega\|_{\beta', 0, 1}$

these constants can be chosen independently of the subinterval. In that result, depending on the value of x we shall choose $t_1 > 0$ such that $d(t_1, x) \leq 1/2$. Then, defining $a(t) := 2h(t)$ and $b(t, x) := 2f(t, x)$ we can rewrite (A.1) as

$$y(t) \leq b(t, x) + a(t)y(t)^2, \quad t \in [0, t_1]$$

which looks like the inequality of Lemma A.4. Let us emphasize that with the above choice $\lim_{t \rightarrow 0^+} a(t) = \lim_{t \rightarrow 0^+} 2h(t) = 0$. In the next result we will also choose suitable values of x such that the rest of assumptions of Lemma A.4 also holds.

Lemma A.5. *Let $(\omega_n)_{n \in \mathbb{N}}$ be a sequence of piecewise linear continuous paths converging to ω in the sense of Remark 3.2, and let $(u_n)_{n \in \mathbb{N}}$ be a sequence of solutions for (2.5) on $[0, 1]$ driven by $(\omega_n)_{n \in \mathbb{N}}$ with initial condition $u_0 \in V$. Then, there exist constants $K \geq \hat{K} > 1$ (uniform in n) defining finitely many intervals $(I_i)_{i=1, \dots, i^*}$ by*

$$I_1 = \left[0, \frac{1}{\hat{K}}\right] = [\check{t}_1, \hat{t}_1], \dots, I_i = \left[\hat{t}_{i-1}, \hat{t}_{i-1} + \frac{1}{Ki}\right] = [\check{t}_i, \hat{t}_i]$$

in such a way that on I_i we have uniformly for any $n \in \mathbb{N}$

$$\begin{aligned} \|u_n\|_{\hat{\beta}, -\delta, I_1} &\leq (\hat{K})^{\hat{\beta}}, & \|u_n\|_{\hat{\beta}, -\delta, I_i} &\leq (Ki)^{\hat{\beta}} \\ \|u_n\|_{C, I_1} &\leq \frac{4c\hat{K}^{1-\beta'}}{1-\beta'}, & \|u_n\|_{C, I_i} &\leq \frac{4c(Ki)^{1-\beta'}}{1-\beta'} \end{aligned}$$

for $i = 2, \dots, i^*$. This constant c in particular depends on $\|\omega\|_{\beta', 0, 1}$.

We point out that in the previous result i^* is given by the condition $\check{t}_{i^*} < 1 = T \leq \hat{t}_{i^*}$, and in this case we set $\hat{t}_{i^*} = 1$.

Proof. Let us denote by $x_0 := \max\{1, \|u_0\|\}$, $x_1(t) := \max\{1, \|u_n\|_{C, 0, t}\}$ and $y_1(t) := \|u_n\|_{\hat{\beta}, -\delta, 0, t}$, for $t \in I_1 = [0, \hat{t}_1]$, where \hat{t}_1 will be determined later. The inequality (5.5) implies

$$\begin{aligned} \|u_n(t)\|^2 &\leq \|u_0\|^2 + ct^{\beta'} \|u_n\|_{C, 0, t} + ct^{\hat{\beta}+\beta'} (1 + \|u_n\|_{C, 0, t}) \|u_n\|_{\hat{\beta}, -\delta, 0, t} \\ &\leq x_0^2 + ct^{\beta'} \|u_n\|_{C, 0, t} + ct^{\hat{\beta}+\beta'} (1 + \|u_n\|_{C, 0, t}) \|u_n\|_{\hat{\beta}, -\delta, 0, t}. \end{aligned}$$

Moreover, since $x_0 \geq 1$ we also have

$$1 \leq x_0^2 + ct^{\beta'} \|u_n\|_{C, 0, t} + ct^{\hat{\beta}+\beta'} (1 + \|u_n\|_{C, 0, t}) \|u_n\|_{\hat{\beta}, -\delta, 0, t}.$$

Then

$$\begin{aligned} \max\{\|u_n\|_{C, 0, t}^2, 1\} &\leq x_0^2 + ct^{\beta'} \|u_n\|_{C, 0, t} + ct^{\hat{\beta}+\beta'} (1 + \|u_n\|_{C, 0, t}) \|u_n\|_{\hat{\beta}, -\delta, 0, t} \\ &\leq x_0^2 + ct^{\beta'} \max\{1, \|u_n\|_{C, 0, t}\} + ct^{\hat{\beta}+\beta'} (1 + \max\{1, \|u_n\|_{C, 0, t}\}) \|u_n\|_{\hat{\beta}, -\delta, 0, t} \end{aligned}$$

and therefore

$$x_1^2(t) \leq x_0^2 + c x_1(t) t^{\beta'} + 2c x_1(t) y_1(t) t^{\hat{\beta}+\beta'}. \quad (\text{A.3})$$

Note that in (A.3) we have used that $x_1(t) \geq 1$ and therefore the corresponding last term on the left hand side of (5.5) can be estimated as

$$ct^{\hat{\beta}+\beta'} (1 + x_1(t)) y_1(t) \leq 2c x_1(t) y_1(t) t^{\hat{\beta}+\beta'}.$$

Furthermore, since $t \in [0, 1]$ and $\delta + 1/2 > 1$, we have that $t^{\delta+1/2-\hat{\beta}} < t^{1-\hat{\beta}}$, and thus (5.7) implies

$$y_1(t) \leq \bar{c} x_0 t^{\delta-\hat{\beta}} + c x_1^2(t) t^{1-\hat{\beta}} + ct^{\beta'-\hat{\beta}} + c y_1(t) t^{\beta'}. \quad (\text{A.4})$$

Now combining (A.3) with (A.4) we get

$$y_1(t) \leq \bar{c} x_0 t^{\delta-\hat{\beta}} + c(x_0^2 + c x_1(t) t^{\beta'} + 2c x_1(t) y_1(t) t^{\hat{\beta}+\beta'}) t^{1-\hat{\beta}} + ct^{\beta'-\hat{\beta}} + c y_1(t) t^{\beta'}. \quad (\text{A.5})$$

In addition, from (A.3) the following estimate holds

$$x_1(t) \leq \frac{ct^{\beta'} + 2c y_1(t) t^{\hat{\beta}+\beta'}}{2} + \sqrt{\frac{(ct^{\beta'} + 2c y_1(t) t^{\hat{\beta}+\beta'})^2 + 4x_0^2}{4}} \leq ct^{\beta'} + 2c y_1(t) t^{\hat{\beta}+\beta'} + x_0 \quad (\text{A.6})$$

and plugging this into (A.5) we finally arrive at

$$y_1(t) \leq d(t, x_0) y_1(t) + f(t, x_0) + h(t) y_1(t)^2, \quad t \in I_1 = [0, \hat{t}_1],$$

where the functions have been defined in (A.2). Then, taking $a(t) = 2h(t)$ and $b(t, x_0) = 2f(t, x_0)$ there exists a $K_1 \geq 1$ such that for any $\hat{K} \geq K_1$ and $\hat{t}_1 = \hat{K}^{-1}$ we have

$$d(\hat{K}^{-1}, x_0) \leq \frac{1}{2}, \quad b(\hat{K}^{-1}, x_0) \leq \frac{(\hat{K})^{\hat{\beta}}}{2}, \quad 4a(\hat{K}^{-1})b(\hat{K}^{-1}, x_0) < 1.$$

Hence, we have the conditions [Lemma A.4](#) and as a consequence we claim that $y_1(\hat{t}_1) \leq (\hat{K})^{\hat{\beta}} = \hat{t}_1^{-\hat{\beta}}$. Let us fix such a \hat{K} such that in addition $x_0 \leq c\hat{K}^{1-\beta'}/(1-\beta')$. Then, from [\(A.6\)](#), simply using the general notation for constants c , we get

$$x_1(\hat{t}_1) \leq 3c\hat{t}_1^{\beta'} + x_0 \leq \frac{4c\hat{K}^{1-\beta'}}{1-\beta'} =: \hat{x}_1.$$

Now we can repeat the same arguments than above in each interval $I_i, i = 2, 3, \dots$, by doing the corresponding suitable changes. In order to do that we need to rewrite the estimates [\(5.5\)](#) and [\(5.7\)](#) in those intervals. In particular, in I_i we have to take as initial condition $u_n(\hat{t}_{i-1})$ and t can be estimated by the length of the interval I_i which is nothing but $(Ki)^{-1}$. Taking $x_i(t) := \max\{1, \|u_n\|_{C, \hat{t}_{i-1}, t}\}$ with $x_i(\hat{t}_{i-1}) \geq 1$ and $y_i(t) := \|u_n\|_{\hat{\beta}, -\delta, \hat{t}_{i-1}, t}$, for $t \in I_i$, for induction we assume

$$x_{i-1}(\hat{t}_{i-1}) \leq 3c \sum_{j=1}^{i-1} (Kj)^{-\beta'} + x_0 \leq \frac{4c(K(i-1))^{1-\beta'}}{1-\beta'} =: \hat{x}_{i-1}$$

and choose $K > \hat{K}$ such that for $i = 2, 3, \dots$

$$\begin{aligned} d((Ki)^{-1}, \hat{x}_{i-1}) &= c(Ki)^{-\beta'} + 4c^3(Ki)^{-1-2\beta'} + 2c^2\hat{x}_{i-1}(Ki)^{-\beta'-1} \\ &\leq cK^{-\beta'} + 4c^3K^{-1-2\beta'} + \frac{8c^3}{1-\beta'}K^{1-\beta'}K^{-\beta'-1} \leq o(K^{-\varepsilon}) \leq \frac{1}{2} \\ f((Ki)^{-1}, \hat{x}_{i-1}) &= \bar{c}\hat{x}_{i-1}(Ki)^{\hat{\beta}-\delta} + c\hat{x}_{i-1}^2(Ki)^{\hat{\beta}-1} + c^2\hat{x}_{i-1}(Ki)^{-1+\hat{\beta}-\beta'} + c^3(Ki)^{-1-2\beta'+\hat{\beta}} + c(Ki)^{-\beta'+\hat{\beta}} \\ &\leq Co(K^{-\varepsilon})(Ki)^{\hat{\beta}} \leq \frac{(Ki)^{\hat{\beta}}}{4} \end{aligned}$$

for a constant C and a sufficiently small $\varepsilon > 0$ independent of K and i . For example, for the critical term in the expression of f given for the quadratic term, we have that

$$c\hat{x}_{i-1}^2(Ki)^{\hat{\beta}-1} \leq \frac{16c^3}{(1-\beta')^2}K^{2-2\beta'-1+\hat{\beta}}i^{2-2\beta'-1+\hat{\beta}} \leq \frac{16c^3}{(1-\beta')^2}K^{1-2\beta'}(Ki)^{\hat{\beta}} \leq Co(K^{-\varepsilon})(Ki)^{\hat{\beta}}$$

where this last inequality is true since $\beta' \in (1/2, 1)$.

Again, for $a(t) = 2h(t)$, $b(t, \hat{x}_{i-1}) = 2f(t, \hat{x}_{i-1})$, choosing K sufficiently large such that

$$4a((Ki)^{-1})b((Ki)^{-1}, \hat{x}_{i-1}) \leq 64c^3(Ki)^{-1-2\beta'-\hat{\beta}}\frac{(Ki)^{\hat{\beta}}}{4} < 1$$

we obtain by [Lemmas A.3](#) and [A.4](#) that $y_i(\hat{t}_i) \leq (Ki)^{\hat{\beta}}$. If we denote $\hat{t}_i - \check{t}_i =: \Delta t_i = (Ki)^{-1}$ the previous inequality can be rewritten as $y_i(\hat{t}_i) \leq (\Delta t_i)^{-\hat{\beta}}$, and similar to [\(A.6\)](#)

$$\begin{aligned} x_i(\hat{t}_i) &\leq c\Delta t_i^{\beta'} + 2cy_i(\hat{t}_i)\Delta t_i^{\hat{\beta}+\beta'} + x_{i-1}(\hat{t}_{i-1}) \leq 3c\Delta t_i^{\beta'} + x_{i-1}(\hat{t}_{i-1}) \\ &\leq x_0 + 3c \sum_{j=1}^i (Kj)^{-\beta'} \leq x_0 + 3cK^{-\beta'} \int_0^i r^{-\beta'} dr \leq x_0 + \frac{3cK^{-\beta'}i^{1-\beta'}}{1-\beta'} \leq \frac{4c(Ki)^{1-\beta'}}{1-\beta'} =: \hat{x}_i \end{aligned}$$

and therefore we obtain that $x_i(\hat{t}_i) \leq \hat{x}_i$. \square

Finally we present the proof of [Lemma 5.5](#):

Proof. Following the steps of [Proposition 5.2](#) we prove that each $u_n \in C^{\hat{\beta}}([0, T]; V_{-\delta})$ with $1/2 < \hat{\beta} < \check{\beta}$. Then we apply [Lemmas A.3–A.5](#) obtaining that this sequence is uniformly bounded in $C^{\hat{\beta}}([0, T]; V_{-\delta}) \cap C([0, T]; V)$. \square

References

[1] E.B. Gledzer, System of hydrodynamic type admitting two quadratic integrals of motion, Dokl. Akad. Nauk SSSR 209 (1973) 1046–1048; Sov. Phys. Dokl. 18 (1973) 216–217 (Engl. Transl.).
 [2] K. Ohkitani, M. Yamada, Lyapunov spectrum of a chaotic model of three-dimensional turbulence, J. Phys. Soc. Japan 56 (1987) 4210–4213.
 [3] V.S. L'vov, E. Podivilov, A. Pomyalov, I. Procaccia, D. Vandembroucq, Improved shell model of turbulence, Phys. Rev. E 58 (1998) 1811–1822.
 [4] P. Constantin, B. Levant, E.S. Titi, Analytic study of shell models of turbulence, Physica D 219 (2) (2006) 120–141.
 [5] D. Barbatto, M. Barsanti, H. Bessaih, F. Flandoli, Some rigorous results on a stochastic GOY model, J. Stat. Phys. 125 (3) (2006) 677–716.
 [6] H. Bessaih, B. Ferrario, Invariant Gibbs measures of the energy for shell models of turbulence; the inviscid and viscous cases, Nonlinearity 25 (2012) 1075–1097.
 [7] H. Bessaih, B. Ferrario, Invariant measures of Gaussian type for 2D turbulence, J. Stat. Phys. 149 (2) (2012) 259–283.
 [8] L. Fang, P. Sundar, F.G. Viens, Two-dimensional stochastic Navier–Stokes equations with fractional Brownian noise, Random Oper. Stoch. Equ. 21 (2) (2013) 135–158.
 [9] L.C. Young, An integration of Hölder type, connected with Stieltjes integration, Acta Math. 67 (1936) 251–282.
 [10] M. Zähle, Integration with respect to fractal functions and stochastic calculus. I, Probab. Theory Related Fields 111 (3) (1998) 333–374.
 [11] S.G. Samko, A.A. Kilbas, O.I. Marichev, Fractional integrals and derivatives: Theory and applications, Gordon and Breach Science Publishers, Switzerland and Philadelphia, PA, USA, 1993.
 [12] H. Bessaih, M.J. Garrido-Atienza, B. Schmalfuss, Random attractors for stochastic shell equations driven by fractional Brownian motion, in preparation.
 [13] D. Nualart, A. Răşcanu, Differential equations driven by fractional Brownian motion, Collect. Math. 53 (1) (2002) 55–81.
 [14] M.J. Garrido-Atienza, B. Masłowski, B. Schmalfuss, Random attractors for stochastic equations driven by a fractional Brownian motion, Int. J. Bifurcation Chaos 20 (9) (2010) 1–22.

- [15] D. Nualart, P.A. Vuillermot, Variational solutions for partial differential equations driven by a fractional noise, *J. Funct. Anal.* 232 (2006) 390–454.
- [16] B. Maslowski, D. Nualart, Evolution equations driven by a fractional Brownian motion, *J. Funct. Anal.* 202 (1) (2003) 277–305.
- [17] M.J. Garrido-Atienza, K. Lu, B. Schmalzfuss, Random dynamical systems for stochastic partial differential equations driven by a fractional Brownian motion, *Discrete Contin. Dyn. Syst. Ser. B* 14 (2) (2010) 473–493.
- [18] M.A. Diop, M.J. Garrido-Atienza, Retarded stochastic evolution systems driven by fractional Brownian motion with Hurst-parameter $H > 1/2$, *Nonlinear Analysis TMA* 97 (2014) 15–29.
- [19] Y. Chen, H. Gao, M.J. Garrido-Atienza, B. Schmalzfuss, Pathwise solutions of SPDEs and random dynamical systems, *Discrete Contin. Dyn. Syst. Ser. A* 34 (1) (2014) 79–98.
- [20] T. Cass, Z. Qian, J. Tudor, Non-linear evolution equations driven by rough paths. *Stochastic analysis and applications to finance*, in: *Interdiscip. Math. Sci.*, vol. 13, World Sci. Publ., Hackensack, NJ, 2012, pp. 1–18.
- [21] H. Gao, M.J. Garrido-Atienza, B. Schmalzfuss, Random attractors for stochastic evolution equations driven by fractional Brownian motion, *SIAM J. Math. Anal.* 46 (4) (2014) 2281–2309.
- [22] P. Constantin, B. Levant, E.S. Titi, Regularity of inviscid shell models of turbulence, *Phys. Rev. E* (3) 75 (1) (2007) 016304. 10 pp.
- [23] G.R. Sell, Y. You, *Dynamics of evolutionary equations*, in: *Springer Applied Mathematical Series*, Springer-Verlag, Berlin, 2002.
- [24] A. Pazy, *Semigroups of linear operators and applications to partial differential equations*, in: *Springer Applied Mathematical Series*, Springer-Verlag, Berlin, 1983.
- [25] M.I. Vishik, A.V. Fursikov, *Mathematical Problems of Statistical Hydromechanics*, Kluwer Academic Publishers, 1988.
- [26] P. Friz, M. Hairer, *A Course on Rough Paths, With An Introduction to Regularity Structures*, Universitext. Springer, Cham, 2014.
- [27] P. Friz, N. Victoir, *Multidimensional stochastic processes as rough paths. Theory and applications*, in: *Cambridge Studies of Advanced Mathematics*, vol. 120, Cambridge University Press, 2010.
- [28] L.H. Duc, S. Siegmund, B. Schmalzfuss, A note of the generation of random dynamical systems for fractional stochastic delay differential equations, *Stoch. Dyn.* 15 (3) (2015) 1550018.
- [29] H. Kunita, *Stochastic Flows and Stochastic Differential Equations*, Cambridge University Press, 1990.
- [30] R. Temam, *Navier–Stokes equations. Theory and numerical analysis*, in: *Studies in Mathematics and its Applications*, vol. 2, North-Holland Publishing Co., 1979.
- [31] H. Sohr, *The Navier–Stokes equations. An elementary functional analytic approach*, in: *Birkhäuser Advances Texts*, Birkhäuser Verlag, Basel–Boston–Berlin, 2001.