Quaternary Science Reviews 129 (2015) 178–195

Contents lists available at ScienceDirect

Quaternary Science Reviews

journal homepage: www.elsevier.com/locate/quascirev

Tracking rainfall in the northern Mediterranean borderlands during sapropel deposition

Samuel Toucanne [a, *], Charlie Morelle Angue Minto'o [b], Christophe Fontanier [a], Maria-Angela Bassetti [b], Stephan J. Jorry [a], Gwenaël Jouet [a]

[a] IFREMER, Laboratoire Environnements Sédimentaires, BP70, 29280 Plouzané, France
[b] Université de Perpignan, Lab. IMAGES (BatU), 52 Av. P. Alduy, 66860 Perpignan, France

ARTICLE INFO

Article history:
Received 25 March 2015
Received in revised form
10 July 2015
Accepted 1 October 2015
Available online xxx

Keywords:
Mediterranean
Rainfall
Interglacials
Sapropel

ABSTRACT

The role of mid-latitude precipitation in the hydrological forcing leading to the deposition of sapropels in the Mediterranean Sea remains unclear. The new GDEC-4-2 borehole, East Corsica margin (northern Tyrrhenian Sea), provides the first precisely dated evidence for enhanced rainfall in the Western Mediterranean during warm intervals of interglacial periods over the last 547 kyr. Comparison of GDEC-4-2 proxy records with pollen sequences and speleothems from the central and eastern Mediterranean reveals that these pluvial events were regional in character and occurred probably in response to the intensification of the Mediterranean storm track along the northern Mediterranean borderlands in autumn/winter. Our dataset suggests that the timing of maxima of the Mediterranean autumn/winter storm track precipitation coincide with that of the North African summer monsoon and sapropel deposition. Besides highlighting a close coupling between mid- and low-latitude hydrological changes, our findings suggest that during warm intervals of interglacial periods the reduced sea-surface water salinities, together with the high flux of nutrient and organic matter, produced by the monsoonal Nile (and wadi-systems) floods, were maintained throughout the winter by the Mediterranean rainfall. This provides an important additional constraint on the hydrological perturbation causing sapropel formation.

© 2015 Elsevier Ltd. All rights reserved.

1. Introduction

Recent understanding of paleohydrological-paleoclimatic conditions in the Mediterranean is based on the intercomparison of marine, lacustrine and cave records. As a prominent example, the integrated study of sapropel deposits, pollen sequences and speleothems reveals that precession, through its role on both the latitudinal migration of the Intertropical Convergence Zone (ITCZ), the African monsoon variability and the runoff from North Africa into the Mediterranean, has been a fundamental pacer of Mediterranean climate (Lourens et al., 1992; Rohling et al., 2015; Tzedakis, 2007; Tzedakis et al., 2009). Indeed, the intensification of the African summer monsoon during precession-driven northern hemisphere insolation maxima led to extra freshwater input into the Eastern Mediterranean due to enhanced runoff of both the Nile River (Ducassou et al., 2009; Revel et al., 2010; Rossignol-Strick, 1983, 1985; Scrivner et al., 2004) and the Central Saharan watershed (e.g., Irharhar, Sahabi and Kufrah rivers; Coulthard et al., 2013; Drake et al., 2011; Larrasoana et al., 2003; Osborne et al., 2008; Rohling et al., 2002, Fig. 1), that in turn caused a disruption in the basin's hydrological cycle, reduced deep-water ventilation and the sapropel deposition (Abu-Zied et al., 2008; Casford et al., 2003; Rohling, 1994; Rohling and Hilgen, 1991; Rohling et al., 2015; Rossignol-Strick, 1985). The sapropel deposition and, by extension, the hydrological cycle in the Mediterranean Sea has long been considered as a low-latitude signal. However, it is well-known that the atmospheric humidity and precipitation amount also increased over the eastern Mediterranean regions during the sapropel deposition (Milner et al., 2012; Roberts et al., 2006; Rohling, 1994; Rohling and Hilgen, 1991; Rohling et al., 2015; Tzedakis, 2007), as seen in the 250 kyr-long speleothem record for the Soreq Cave, Israel (Bar-Matthews et al., 2003, 2000). Interestingly, these speleothems reveal that the isotopic

* Corresponding author.
E-mail address: stoucann@ifremer.fr (S. Toucanne).

http://dx.doi.org/10.1016/j.quascirev.2015.10.016
0277-3791/© 2015 Elsevier Ltd. All rights reserved.




Fig. 1. Location of the main sites mentioned in the text, including the GDEC-4-2 borehole (red circle). The blue and green circles (with associated coloured names) indicate the marine and continental (i.e., speleothems, pollen sequences) records, respectively. See the main text for references. The modern (approximate) position of the Azores High and of the ITCZ (red bands) is shown. The dashed red line over North Africa show the maximum northward displacement of the ITCZ over the last million years (e.g., Tuenter et al., 2003). Course of the Central Saharan rivers according to Coulthard et al. (2013). The orange arrow shows the Mediterranean storm track. (For interpretation of the references to colour in this figure caption, the reader is referred to the web version of this article.)

composition of the precipitation over the Eastern Mediterranean during sapropel deposition was distinct from African monsoonal composition and had a Mediterranean origin (Bar-Matthews et al., 2000). This result is supported for the last insolation maxima and sapropel S1 (~7–9 thousand years ago, ka) by a concomitant humid interval over the Aegean Sea (Kotthoff et al., 2008), in Turkey (Göktürk et al., 2011) and in the northern Red Sea (Arz et al., 2003) that originated from enhancement of rainfall from Mediterranean sources. Similar evidence from northeast Greece and the Tenagghi Philippon peatland were also recently presented for the penultimate interglacial interval [Marine Isotope Stage (MIS) 5] and sapropel S5 (~125 ka) (Milner et al., 2012; Tzedakis, 2007). These findings definitely challenge the idea for a northward extension of the African monsoon over the Mediterranean basin during insolation maxima. The enhancement of the hydrological activity during sapropel deposition is also recorded in the Western Mediterranean, up to 42–43°N. The isotopic composition of a stalagmite collected from Antro del Corchia Cave, Central Italy (Zanchetta et al., 2007), and the study of lacustrine and marine sequences from around the Italian Peninsula (Ariztegui et al., 2000; Magny et al., 2013) at the time of Sapropel S1 support this idea. Similar features are observed during sapropel deposition dating from MIS 5e (sapropel S5) and MIS 7a (sapropel S7, ~195 ka) during which significant decrease in salinity occurred in the Eastern Tyrrhenian Sea (Kallel et al., 2000). Results from a stalagmite collected in the Argentarola Cave, Tyrrhenian coast of Italy, also reveal the occurrence of a wet period during the penultimate glacial period (precisely MIS 6d, ~170–180 ka) that corresponds chronologically to the deposition of sapropel S6 (Bard et al., 2002). Taken together, these evidences highlight that freshwater runoff during sapropel events was not restricted to the Eastern Mediterranean Sea and to the output of both the Nile River and the Central Saharan watershed, but was rather widespread over the entire Mediterranean Sea due to increased rainfall (Bard et al., 2002; Kallel et al., 2000, 1997). Recent numerical modelling focussing on the time intervals coinciding with the deposition of sapropels also support this assumption (Bosmans et al., 2015; Kutzbach et al., 2014). However, relatively few studies are based upon long records that contain multiple precession cycles and intervals for sapropel deposition, especially in the Western Mediterranean. Here, we directly address this gap by studying a high-resolution ~550 kyr long sedimentary archives from the Northern Tyrrhenian Sea (Figs. 1–3), well-known as a highly-sensitive region for paleoclimate reconstructions (e.g., Allen et al., 1999; Bard et al., 2002; Brauer et al., 2007).




Fig. 2. Schematic dip section across the East Corsica margin, from the Golo River mouth to the Golo basin, illustrating the stacking of the last five sedimentary sequences bounded by major discontinuities on the shelf and attributed to 100 kyr-glacio-eustatic cyles. The stratigraphic scheme attests the presence of transgressive and highstand (i.e., interglacial) accumulations, imprinted by stacked continuous and regional seismic reflections representing thick interglacial sedimentary intervals (ca. 10–15 m). Note that the borehole GDEC-4-2 is located between the North Golo canyon and the South Golo canyon. This makes GDEC-4-2 well suited for continuously recording sediment input variability from the Golo River throughout the last climate cycles.

2. Material and methods

2.1. The GDEC-4-2 borehole, East Corsica margin

This study is based on borehole GDEC-4-2 (N42°31′23.2, E9°42′59.5), a 125.7 m continuous core (99.9% recovery) drilled at 492 m water depth in the Northern Tyrrhenian Sea (Fig. 1) by the R/V 'Bavenit' (FUGRO) during the 2009 GOLODRILL cruise. The GDEC-4-2 borehole was drilled on the upper continental slope of the Golo basin, East Corsica (Fig. 2), directly off the Golo River (89 km long, drainage basin of ca. 1214 km^2). The Golo is a short, mountainous river (maximum altitude of ca. 2700 m), and is an highly efficient sediment routing system (i.e., Golo source-to-sink system), constituting the main sediment source to the adjacent margin (e.g., Calvès et al., 2013; Forzoni et al., 2015; Sømme et al., 2011). The cored interval is composed only of hemipelagic sediments (mainly silty-clay, with some carbonate-rich intervals), and provides a chronostratigraphic framework for East Corsica margins where seismic and sequence stratigraphic organisation are well established (e.g., Calvès et al., 2013 and references therein). The interpretation of high-resolution seismic Sparker lines illustrates the stacking of the last five sedimentary sequences bounded by major discontinuities on the shelf (Fig. 2). The sedimentary sequences are attributed to 100 kyr-glacio-eustatic cyles, as demonstrated at several margins around the Western Mediterranean (Jouet et al., 2006; Rabineau et al., 2005; Ridente et al., 2009; Sierro et al., 2009). The regressive deposits represent the most significant element constituting upper slope sequences, but the stratigraphic scheme also attests to the presence of transgressive and highstand accumulations, imprinted by stacked continuous and regional seismic reflections representing thick interglacial sedimentary intervals (ca. 10–15 m). Sequence architecture on the upper slope confirms that borehole GDEC-4-2 well records the major sedimentary events that form strata on East Corsica continental margin, in relation to sea-level and climate changes. As no erosion occurred along the recovered interval, it represents a good balance between sediment supply and accommodation space. The well-preserved depositional sequences directly archive the variable influence of the continental supply, at a maximum distance of 15 km (during highstand conditions) offshore from the Golo river outlet (Fig. 2). This makes borehole GDEC-4-2 well suited for continuously recording sediment input variability from the Golo River and the island of Corsica throughout the last climate cycles.

2.2. Analytical methods

2.2.1. Sedimentology and sediment flux

Physical properties of the GDEC-4-2 borehole which include gamma-ray attenuation density were determined every 1 cm using a Geotek Multi Sensor Core Logger (MSCL). The borehole was then sampled continuously at 5–20 cm intervals (i.e., 218- to 870-year time resolution). Weight percent $CaCO_3$ were measured on 1412 samples (Figs. 3d and 4). Bulk sediments were acidified with 3 M hydrochloric acid and analyses were performed with an Aquitaine Technique Innovation automated calcimeter. Analytical precision is estimated to be ±2%. Grain-size analyses (Fig. 5d) were performed on 1002 samples using a Coulter LS200 laser microgranulometer with no chemical pre-treatment of the bulk sediment. Micro particle size standard (15, 35, 500 μm) were used as controls to verify the performance of the measurement system. The terrigenous inputs at site GDEC-4-2 were quantified by the calculation of the terrigenous flux or Mass Accumulation Rates (MAR_t, in g cm^{-2} kyr^{-1}; Fig. 5b) according to the following formula: MAR = LSR × DBD × (1 − carbonate content), with LSR: Linear Sedimentation Rate (cm kyr^{-1}) and DBD: Dry Bulk Density (g cm^{-3}), which has been calculated assuming a mean grain density of 2.65 g cm^{-3} and an interstitial water density of 1.024 g cm^{-3} (Cremer et al., 1992) as follows: DBD = 2.65 × (1.024 − D_{wet})/




Fig. 3. (A) Greenland synthetic $\delta^{18}O$ record (GLT-syn; Barker et al., 2011); (B) $\delta^{18}O$ of *G. bulloides* from the combined record of core MD01-2444 and MD01-2443, Iberian margin (Hodell et al., 2013); (C) Composite $\delta^{18}O$ of *G. bulloides*, *G. ruber* (white) and *N. pachyderma* (dextral) of the GDEC-4-2 borehole (green tones). The chronology for GDEC-4-2 was constructed by aligning this planktonic $\delta^{18}O$ record to the synthetic Greenland (GL_T-syn) record of Barker et al. (2011). The triangles at the bottom part of the figure indicate the tie points (see Table 1 for details). The yellow (continuous) and purple (dashed) lines correspond to the $\delta^{18}O$ of *G. bulloides* at site MD01-2472, East Corsica margin (Toucanne et al., 2012) and at site ODP 975, West Balearic Basin (Pierre et al., 1999), respectively. The $\delta^{18}O$ at site ODP 975 is offset by −1‰ to facilitate the comparison with the GDEC-4-2 and MD01-2472 records; (D) XRF Ca/Ti (log scale; blue line) and weight percent calcium carbonate (wgt. $\%CaCO_3$; red line) of the GDEC-4-2 borehole. The black line corresponds to the wgt. $\%CaCO_3$ (−5% for comparison with the GDEC-4-2 dataset) for core LC06, Balearic Abyssal Plain (Hoogakker et al., 2004). Note the opposite evolution of the wgt. $\%CaCO_3$ during interglacial warm intervals (i.e, MIS 5a, 5c, 5e) at sites GDEC-4-2 ($\%CaCO_3$ lows) and LC06 ($\%CaCO_3$ highs). The red arrows highlight the unexpected low levels for carbonate content (i.e., '$CaCO_3$ paradox', see the main text for details); (E) Composite $\delta^{18}O$ of *C. pachyderma*, *C. kullenbergi* and *C. wuellerstorfi* of the GDEC-4-2 borehole. Light grey vertical bands indicate interglacial conditions *s.l.*, while dark bands indicate interglacial warm intervals and the interglacial *s.s.* (i.e., MIS 1, MIS 5e, MIS 7c, 7e, MIS 9, MIS11 and MIS 13c). Terminations (T.) I to VI according to Barker et al. (2011). Scheme of marine isotope substages according to Railsback et al. (2015). (For interpretation of the references to colour in this figure caption, the reader is referred to the web version of this article.)

(1.024–2.65). Wet bulk densities (D_{wet}) were derived from gamma-ray attenuation density measurements.

2.2.2. X-ray fluorescence analysis

X-ray fluorescence (XRF) core scanning provides rapid high-resolution records of chemical composition on split sediment cores (Richter et al., 2006). The bulk intensity of major elements for borehole GDEC-4-2 was analysed using an Avaatech XRF core scanner at the Institut Français de Recherche pour l'Exploitation de la Mer (IFREMER), Brest (France). XRF data were collected every 1 cm along the entire length of the borehole, with a count time of 10 s, by setting the voltage to 10 kV (no filter) and the intensity to 600 μA. Only data for Calcium (Ca) and Titanium (Ti) elements are reported in this study (Figs. 3d, 4 and 5a). It is commonly admitted




Fig. 4. Weight percent calcium carbonate (weight %$CaCO_3$) versus XRF Ca/Ti (log scale) (n = 1412, r = 0.91/p < 0.01).

that elemental Ti is related to terrigenous-siliciclastic components (clays, heavy minerals), while Ca mainly reflects the carbonate content (calcite and aragonite) in the sediment (Richter et al., 2006). Results are presented in log ratios of element intensities (Weltje and Tjallingii, 2008).

2.2.3. Stable isotopes

Planktonic (920 samples) and benthic (774 samples) foraminiferal $^{18}O/^{16}O$ and $^{13}C/^{12}C$ ratios ($\delta^{18}O$ and $\delta^{13}C$ respectively, expressed in ‰ versus Vienna Pee-Dee Belemnite, VPDB) were measured at the Laboratoire des Sciences de l'Environnement et du Climat (LSCE), Gif-sur-Yvette (France), on Finnigan Δ+, and OPTIMA and Elementar Isoprime GV mass spectrometers. VPDB is defined with respect to the NBS-19 calcite standard ($\delta^{18}O = -2.20$‰ and $\delta^{13}C = +1.95$‰). The mean external reproducibility (1σ) of carbonate standards is ±0.05‰ for $\delta^{18}O$ and ±0.03‰ for $\delta^{13}C$. Measured NBS-18 $\delta^{18}O$ is −23.27 ± 0.10‰ VPDB and measured NBS-18 $\delta^{13}C$ is −5.01 ± 0.03‰ VPDB. The three mass spectrometers used for these measurements are calibrated together with respect to NBS-19, NBS-18 and other in-house carbonate standards. The obtained correction does not exceed 0.15‰. Measurements were performed on the epipelagic species *Globigerina bulloides* and *Globigerinoides ruber* (white) from the 250–315 μm size fraction, the mesopelagic *Neogloboquadrina pachyderma* (dextral) from the 200–250 μm size fraction, and the epifaunal *Cibicides wuellerstorfi*, *Cibicidoides pachyderma* and *Cibicidoides kullenbergi* found in the >150 μm size fraction (Fig. 3c, e and 6d). The samples (ca. 50 μg minimum) were cleaned in a methanol ultrasonic bath for a few seconds and roasted under vacuum at 380 °C for 45 min prior to analysis, in order to eliminate impurities (Duplessy, 1978). A correction factor of +1.14‰ for $\delta^{18}O$ was applied to the isotope results from *G. ruber* to account for the relatively constant offset (i.e., vital and habitat preferences) with regard to *N. pachyderma* and *G. bulloides*. This correction factor, calculated on 15 paired analyses of *N. pachyderma* and *G. ruber*, is in agreement with that used for the Upper Pleistocene on ODP 653 and 654, Central Tyrrhenian Sea (Vergnaud-Grazzini et al., 1990). Such a correction factor also allows the alignment of the composite GDEC-4-2 planktonic $\delta^{18}O$ record onto the $\delta^{18}O$ *G. bulloides* record at site MD01-2472 (Fig. 3c) (Toucanne et al., 2012), ca. 15 km north of site GDEC-4-2. For the same reasons, a correction factor of −0.08‰ for $\delta^{18}O$ and of +0.42‰ for $\delta^{13}C$ (calculated on 9 paired analyses of *Cibicidoides pachyderma* and *Cibicides wuellerstorfi*, and 21 paired analyses of *Cibicidoides kullenbergi* and *Cibicides wuellerstorfi*) was applied to the isotope results from *Cibicidoides pachyderma* and *Cibicidoides kullenbergi* with regard to *Cibicides wuellerstorfi*.

2.2.4. Benthic foraminifera assemblages

A total of 299 sediment samples were analysed for their benthic foraminiferal abundance (Fig. 6b, c and 7b). Samples were sieved through a screen with a 150-μm mesh, and thereafter the sieve residues were dried in an oven (50 °C). Foraminifera were sorted from dried sediments and stored in Plummer slides. When possible, at least 250 individuals per sediment interval were counted (Murray, 2006). If necessary, samples were divided into sub-fractions using an Otto splitter. Only 18 intervals presented total foraminiferal stocks comprised between 105 and 249 individuals. In order to reconstruct the paleoenvironmental conditions prevailing at the seafloor, the relative abundance of various ecological assemblages was calculated. First, the relative abundance of foraminiferal species thriving in neritic ecosystems (e.g., *Ammonia* spp., *Elphidium* spp., *Haynesina* spp., *Rosalina* spp.) (Goineau et al., 2012; Murray, 2006) was determined. Their contribution at our upper-slope site is likely related to downslope transfer by hydrosedimentary processes (e.g., nepheloid layers and turbidity currents from shelf). Therefore, they were removed from foraminiferal census data, and the relative abundance of other species (assumedly autochthonous on the upper slope) was recalculated. Each upper-slope taxon presents a modal bathymetrical distribution which is mainly controlled by the organic matter flux at the seafloor and the bottom water oxygenation (Gooday, 2003; Murray, 2006; Jorissen et al., 2007). Among those taxa, foraminiferal species which can thrive along well-ventilated slope supplied by high organic matter flux were regrouped in the 'eutrophic assemblage' (*Amphicoryna scalaris*, *Bolivina spathulata*, *Bolivina alata*, *Bulimina marginata*, *Bulimina costata*, *Chilostomella* spp., *Globobulimina* spp., *Hyalinea balthica*, *Trifarina angulosa*, *Valvulineria bradyana*, *Pseudoclavulina crustata*) (De Rijk et al., 2000; Fontanier et al., 2008b, 2002; Mojtahid et al., 2009). Note that a special attention was paid to the so-called 'deep infaunal group', which consists of *Chilostomella* spp. and *Globobulimina* spp. Both highly specialized taxa are able to live in well-ventilated as well as oxygen-depleted ecosystems where degraded organic detritus accumulates (Abu-Zied et al., 2008; Fontanier et al., 2014, 2005).

3. Chronology

The chronology for GDEC-4-2 was developed by aligning the planktonic $\delta^{18}O$ (influenced by local hydrography and global ice volume changes) and both the weight percent $CaCO_3$ and XRF-log(Ca/Ti) records (influenced by carbonate productivity and detrital sedimentation) to the NGRIP ice core isotopes from Greenland for the last 60 ka (GICC05 chronology; Rasmussen et al., 2006; Svensson et al., 2008) and to the synthetic Greenland (GL_T-syn) record of Barker et al. (2011) from 60 to ~550 ka (Table 1 and Fig. 3). The GL_T-syn record was constructed from the EPICA Dome C (East Antarctica) ice core, using the bipolar-seesaw model, and placed on the absolute 'Speleo-Age' (i.e., uranium-thorium based) timescale (Barker et al., 2011). Such an approach was successfully used to produce an accurate long-term chronology on the southwestern Iberian margin (Fig. 3b) (i.e., composite record from cores MD99-2344 and MD99-2343; Hodell et al., 2013; Margari et al., 2014). Previous works at this site have shown that oxygen isotope variability in planktonic foraminifera closely matches the ice core records of temperature over Greenland during the last glacial period (Shackleton et al., 2000), and that prominent lows in Ca/Ti correspond to cold stadials in both the Greenland ice core and the GL_T-syn records (Hodell et al., 2013). We applied the same




Fig. 5. (A) XRF Ca/Ti (log scale) of the GDEC-4-2 borehole (blue line) and of the combined record of core MD01-2444 and MD01-2443 (orange line), Iberian margin (Hodell et al., 2013). The red arrows highlight the %$CaCO_3$ lows identified in Fig. 3 (i.e., '$CaCO_3$ paradox', see the main text for details); (B) Terrigenous flux (i.e., mass accumulation rates) at site GDEC-4-2 (grey line). The blue line depicts terrigenous flux normalised to the changing distance between the Golo river mouth and the GDEC-4-2 borehole with regard to sea-level changes (see Fig. 5e). The red arrows highlight unexpected high terrigenous flux during interglacials *s.s.* (see the main text for details); (C) XRF Ti/Ca (log scale) of the GDEC-4-2 borehole; (D) Abundance of the silt fraction (10–63 μm) at site GDEC-4-2; (E) $U^{k'}_{37}$-SST (sea-surface temperature) of the combined record of core MD01-2444 and MD01-2443 (orange line), Iberian margin (Martrat et al., 2007). The blue shaded interval depicts the 95% probability interval for the probability maximum of the Red Sea relative sea-level (RSL) dataset (core KL09; Grant et al., 2012; Grant et al., 2014); (F) Composite $\delta^{18}O$ of *G. bulloides*, *G. ruber* (white) and *N. pachyderma* (dextral) of the GDEC-4-2 borehole (green tones). Light grey bands indicate interglacial conditions *s.l.*, while dark bands indicate interglacial warm intervals and the interglacial *s.s.* (i.e., MIS 1, MIS 5e, MIS 7c, 7e, MIS 9, MIS11 and MIS 13c). Terminations (T.) I to VI according to Barker et al. (2011). Scheme of marine isotope substages according to Railsback et al. (2015). (For interpretation of the references to colour in this figure caption, the reader is referred to the web version of this article.)

methodology for GDEC-4-2 because of the striking resemblance between the planktonic $\delta^{18}O$ and Ca/Ti records from the Corsican margin with those from the Iberian margin (Figs. 3 and 5a). This results from the close linkage of the Mediterranean climate oscillations with North Atlantic climate changes (Cacho et al., 1999; Martrat et al., 2004). The synchronisation of GDEC-4-2 with the synthetic Greenland record reveals that the 125.7 m long marine sedimentary archive encompasses the last 547 kyr with a mean




Fig. 6. (A) Composite $\delta^{18}O$ of *G. bulloides*, *G. ruber* (white) and *N. pachyderma* (dextral) of the GDEC-4-2 borehole (green tones); (B) Abundance of the benthic foraminifera eutrophic group; (C) Abundance of the deep-infaunal foraminifera (red line) and eccentricity of the Earth's orbit (dashed grey line; Laskar et al., 2004). The interglacials (*s.s.*) terrigenous flux at site GDEC-4-2 are also shown (blue diamonds). These fluxes can be compared since similar sea-level highstand conditions prevailed during these intervals; (D) Composite $\delta^{13}C$ of *C. pachyderma* ($\delta^{13}C_{Cp}$), *C. kullenbergi* ($\delta^{13}C_{Ck}$) and *C. wuellerstorfi* ($\delta^{13}C_{Cw}$) of the GDEC-4-2 borehole. Depletions of both $\delta^{13}C_{Cp}$ and $\delta^{13}C_{Ck}$ signals below 0.3‰, and of the $\delta^{13}C_{Cw}$ signal below 0.9‰ are likely related to episodes of enhanced organic-matter mineralization within the sediment. These events are represented by vertical blue bands; (E) June insolation for 65°N (black line) and precession (dashed grey line) (Laskar et al., 2004). Light grey bands indicate interglacial conditions *s.l.*, while dark bands indicate interglacial warm intervals and the interglacial *s.s.* (i.e., MIS 1, MIS 5e, MIS 7c, 7e, MIS 9, MIS11 and MIS 13c). Terminations (T.) I to VI according to Barker et al. (2011). Chronology for sapropel (S) layers (speleothem age-scale) according to Ziegler et al. (2010) until Sapropel S9 and Konijnendijk et al. (2014) thereafter. Scheme of marine isotope substages according to Railsback et al. (2015). (For interpretation of the references to colour in this figure caption, the reader is referred to the web version of this article.)

sedimentation rate of 23 cm kyr^{-1}. It corresponds to the highest resolution continuous marine sedimentary archive of the last five climatic cycles in the Mediterranean Sea obtained to date. The placement of the GDEC-4-2 sequence on the speleothem age-scale permits the correlation of the GDEC-4-2 sequences to key reference data coming from the Mediterranean (e.g., Bar-Matthews et al., 2003, 2000; Ziegler et al., 2010) and the North Atlantic (Hodell et al., 2013), which are also placed on the speleothem age-scale. Note that the chronology of the uppermost part of GDEC-4-2 is supported by radiocarbon ages performed on shells of planktonic foraminifera (mainly *Globigerina bulloides*) picked from the bulk sediment (Table 2). The ages were corrected for a marine reservoir effect of 400 years, except for those from the period 15–17 ka for which a correction of 800 years was applied (Siani et al., 2001). Radiocarbon ages were calibrated to calendar years using CALIB Rev 7.0.0 (Stuiver and Reimer, 1993) and the IntCal13 calibration curve (Reimer et al., 2013).

Table 1
Chronological framework for GDEC-4-2. The ages of isotope events from the North-GRIP ice core (0–60 ka interval, GICC05 chronology; Rasmussen et al., 2006; Svensson et al., 2008) and the synthetic Greenland (GL_T-syn, 60–550 ka interval) record of Barker et al. (2011) were used to calibrate isotope events in the core. GI is Greenland Interstadial; MIS is Marine Isotope Stage; T is Termination. Scheme of marine isotope substages according to Railsback et al. (2015). Radiocarbon ages (i.e., CALIB 7 Age) refer to Table 2. Ages marked by an asterik (chronological inversion) are not included in the age model.

Stratigraphic event	GDEC-4-2 – depth (cmbsf)	CALIB 7 age (ka BP)	NGRIP GICC05 age (ka BP)	GLT-syn age (ka BP)	Summary age (ka BP)
Radiocarbon dates	62.5	5.517*	_	_	5.517
Radiocarbon dates	118.5	5.322*	_	_	5.322
Radiocarbon dates	170.5	9.989	_	_	9.989
Radiocarbon dates	198.0	8.926*	_	_	8.926
YD/Holocene transition	230.0	_	11.70	_	11.7
Radiocarbon dates	277.8	13.117	_	_	13.117
Radiocarbon dates	347.0	13.954	_	_	13.954
Onset GI-1/T.I	360.5	_	14.70	_	14.7
Radiocarbon dates	399.5	15.522	_	_	15.522
Radiocarbon dates	589.5	18.160	_	_	18.160
Radiocarbon dates	744.5	21.304	_	_	21.304
GI-2 peak	860	_	23.4	_	23.38
Radiocarbon dates	855.5	23.935	_	_	23.935
Radiocarbon dates	981	26.767	_	_	26.767
Onset GI-3	1050	_	27.82	_	27.82
Onset GI-4	1110	_	28.92	_	28.92
Radiocarbon dates	1179.5	32.359	_	_	32.359
Radiocarbon dates	1274.5	25.545*	_	_	25.545
Onset GI-7	1300	_	35.52	_	35.52
Radiocarbon dates	1363.5	37.516	_	_	37.516
Onset GI-8	1430.5	_	38.26	_	38.26
Radiocarbon dates	1454.5	39.758	_	_	39.758
Onset GI-10	1498.5	_	41.5	_	41.50
Onset GI-11	1560	_	43.4	_	43.40
Onset GI-12	1675	_	46.9	_	46.90
Onset GI-14	1890	_	53.85	_	53.85
Onset GI-18	2340	_	_	64.1	64.7
Onset GI-19	2685	_	_	71.7	71.7
Onset GI-20	2760	_	_	75.7	75.7
Onset GI-21/MIS 5a	2960	_	_	83.7	83.7
Onset MIS5b	3090	_	_	90.7	90.7
Onset MIS5c	3345	_	_	109.8	109.8
Mid-MIS5d	3398			111.3	111.3
Onset MIS5d	3517			118	118
Mid-MIS5e (2)	3585			122.9	122.9
Mid-MIS5e (1)	3680			125.35	125.35
Onset MIS5e/T.II	3840	_	_	131	131
Mid-MIS6 (2)	4430	_	_	146.5	146.5
Mid-MIS6 (1)	4850	_	_	151	151
Onset MIS6d	5780	_	_	178.5	178.5
Onset MIS7a	6290	–	_	199	199
Onset MIS7c	6888	_	_	217	217
Onset MIS7e/T.III	7300	_	_	243	243
Mid-MIS8	7990	_	_	264.3	264.3
Onset MIS8	8380	_	_	280.5	280.5
Onset MIS9a	8500	_	_	291.4	291.4
Onset MIS9b	8690	_	_	298	298
Onset MIS9d	8890	_	_	321	321
Onset MIS9e/T.IV	9210	_	_	336	336
Mid-MIS-10	9490	_	_	351	351
Onset MIS10	10,350	_	_	396.5	396.5
Mid-MIS11c (2)	10,520	_	_	413	413
Mid-MIS11c (1)	10,780	_	_	423	423
Onset MIS11e/T.V	10,988	_	_	431	431
Mid-MIS12	11,550	_	_	460.5	460.5
Onset MIS13a	12,010	_	_	495.5	495.5
Onset MIS13b	12,190	_	_	520.5	520.5
Base GDEC-4-2 (MIS14)	12,510	_	_	542.8	542.8

4. Results and discussion

4.1. The GDEC-4-2 borehole: a high-resolution record of paleoenvironmental changes in the Western Mediterranean

4.1.1. Glacial-interglacial climate signature recorded in oxygen isotopes

The prominent features of the marine oxygen isotope records known for the last five climatic cycles (e.g., Hodell et al., 2013; Lisiecki and Raymo, 2005) are reproduced in oxygen isotope measurements from site GDEC-4-2 (Fig. 3c and e). The foraminiferal $\delta^{18}O$ show values ranging from −0.49‰ (−1.63‰ without correction for *G. ruber*) to 4.11‰ for planktonic species [*G. bulloides*, *G. ruber* (w) and *N. pachyderma* (d)], and from 0.75‰ to 4.18‰ for benthic species (*Cibicides wuellerstorfi*, *Cibicidoides pachyderma* and *Cibicidoides kullenbergi*). These two end-members characterise interglacial (MIS 1, 5, 7, 9, 11 and 13; light isotopes values) and glacial (MIS 2–4, 6, 8, 10 and 12; heavy values) climate conditions,

Table 2
Radiocarbon ages for GDEC-4-2. The age dates were corrected for a marine reservoir effect of 400 years, except for those from the period 15–17 ka (marked by an asterisk) for which a correction of 800 years was applied (cf. Siani et al., 2001). Radiocarbon ages were calibrated to calendar years using CALIB 7.0.0 and the IntCal13 calibration curve (Reimer et al., 2013).

Core number	Depth (cmbsf)	Material	Lab code	Corrected ^{14}C Age (yr BP)	Cal BP age Ranges (2σ)	Cal BP age Median probability	Data origin
GDEC-4-2	62.5	Bulk planktic	Poz-40548	4760 ± 35	5458–5588	5517	This study
GDEC-4-2	118.5	Bulk planktic	Poz-40549	4600 ± 35	5373–5461	5322	This study
GDEC-4-2	170.5	Bulk planktic	Poz-33942	8860 ± 50	9760–10,172	9989	This study
GDEC-4-2	198	Bulk planktic	Poz-40550	8050 ± 50	8746–9090	8926	This study
GDEC-4-2	277.8	Bulk planktic	Poz-40590	11,250 ± 70	12,995–13,272	13,117	This study
GDEC-4-2	347	Bulk planktic	Poz-40591	12,100 ± 60	13,773–14,118	13,954	This study
GDEC-4-2	399.5	Bulk planktic	Poz-40593	12,980 ± 70*	15,269–15,776	15,522*	This study
GDEC-4-2	589.5	Bulk planktic	Poz-40551	14,940 ± 80	17,933–18,373	18,160	This study
GDEC-4-2	744.5	Bulk planktic	Poz-40552	17,620 ± 90	20,986–21,613	21,304	This study
GDEC-4-2	855.5	Bulk planktic	Poz-40554	19,890 ± 120	23,605–24,253	23,935	This study
GDEC-4-2	981	Bulk planktic	Poz-40555	22,470 ± 240	26,179–27,298	26,767	This study
GDEC-4-2	1179.5	Bulk planktic	Poz-40594	27,980 ± 250	31,572–33,124	32,359	This study
GDEC-4-2	1274.5	Bulk planktic	Poz-40595	21,270 ± 350	24,600–26,167	25,545	This study
GDEC-4-2	1363.5	Bulk planktic	Poz-40597	33,300 ± 500	36,272–38,674	37,516	This study
GDEC-4-2	1454.5	Bulk planktic	Poz-40598	35,200 ± 600	38,535–41,072	39,758	This study

respectively. The glacial-interglacial transitions (i.e., terminations) show significant depletions of ca. 2.0–3.5‰ for planktonic oxygen isotopes and of ca. 1.5–2 to 3‰ for benthic signals. The most significant isotopic depletions (for both planktonic and benthic foraminifera) are recorded during the terminations T.I (i.e., transition from MIS 2 to 1, centred at ~15 ka), T.II (MIS 6–5 transition, ~130 ka) and T.IV (MIS 10–9 transition, ~337 ka). Terminations T.III (MIS 8–7 transition, ~243 ka), T.V (MIS 12–11 transition, ~433 ka) and T.VI (MIS 14–13 transition, ~528 ka) show moderate isotopic depletions in comparison, either due to 'heavy' isotope values for the interglacial periods (e.g., MIS 13c and MIS 7e) or 'light' values for the preceding glacial intervals (e.g., MIS 12 and the second part of MIS 8). Such isotopic variability characterizing terminations is observed in the high-resolution planktonic records from the Iberian margin (Hodell et al., 2013) and the Algero-Balearic Basin (ODP 975; Pierre et al., 1999) (Fig. 3b and c). This reveals the regional relevance of the 547-kyr long GDEC-4-2 oxygen isotopes record. Further comparison between the planktonic $\delta^{18}O$ at these sites highlights that during MIS5, 6 and 7, the GDEC-4-2 site records some pronounced isotopic fluctuations (compared to the ODP 975 planktonic $\delta^{18}O$ record especially; Fig. 3c), with the same order of magnitude of those recorded during terminations. Indeed, MIS 5 and 7 interglacials show high-amplitude oscillations in the planktonic $\delta^{18}O$ of ca. 1.5–2.7‰ (up to 2‰ for benthic isotopes) (Fig. 3c). Interestingly, a similar isotope fluctuation (ca. 2.5‰ and 1.5‰ for planktonic and benthic foraminifera, respectively) is recorded in the first half of glacial MIS 6 (i.e., MIS 6d), with planktonic $\delta^{18}O$ (down to 0.1‰ between 163 and 179 ka) reaching similar levels to those recorded during MIS 5e (ca. 0‰) and MIS 7a (0.3‰) and 7c (−0.2‰). These shifts strongly exceed the attendant sea-level changes related isotopes fluctuations [e.g., ca. 1.0–1.2‰ for planktonic oxygen isotopes (Rohling et al., 2014) and ca. 0.87‰–1.4‰ for benthic isotopes (Cacho et al., 2006; Waelbroeck et al., 2002) for a 110–130 m sea-level change]. This suggests that in addition to changes in global ice-sheet volume, the $\delta^{18}O$ records at site GDEC-4-2 reflect significant surface (and deep water) temperature and/or salinity changes. These changes in surface water hydrography off Corsica closely mirror the climate variability, including recurrent and widespread millennial cooling/warming events during interglacial intervals recognized both on the Iberian margin (Desprat et al., 2006; Hodell et al., 2013; Martrat et al., 2004; Roucoux et al., 2006; Sánchez Goñi et al., 1999) and in the Greenland synthetic record (Barker et al., 2011). This strongly supports the direct connection between climate changes at high-latitudes and over the Mediterranean Sea (e.g., Cacho et al., 1999; Martrat et al., 2007). In addition, the subtantial shifts observed in the planktonic $\delta^{18}O$ between MIS 7e and MIS 5 highlight the direct impact of the precession cyclicity at the GDEC-4-2 site, as previously shown in the central Adriatic (PRAD1-2 borehole; Piva et al., 2008). This astronomical forcing, showing prominent fluctuations during these intervals (Berger, 1978; Laskar et al., 2004) and widely observed in the Mediterranean records, is interpreted as a low-latitude signal (Hodell et al., 2013; Lourens et al., 1992; Sánchez Goñi et al., 2008; Tzedakis, 2007). Taken together, this confirms that the Corsica Trough and the northern Tyrrhenian Sea are climatically sensitive regions of the Mediterranean Sea (e.g., Hayes et al., 2005; Kuhlemann et al., 2008; Toucanne et al., 2012).

4.1.2. *Sediment supplies and the 'CaCO$_3$ paradox'*

Sedimentation at site GDEC-4-2 shows the alternation of silty-clay and carbonate-rich (i.e., $CaCO_3$ > 30%) silty-clay deposited by hemipelagic processes. The carbonate content on the upper continental slopes is generally controlled by the surface-water carbonate productivity and the dilution by terrigenous sediments (Cremer et al., 1992; Hoogakker et al., 2004). Downcore fluctuations of the weight percent $CaCO_3$ at site GDEC-4-2 range from 7 to 45% and follow the local (i.e., site GDEC-4-2) and regional (i.e., ODP 975 and combined record of core MD01-2444 and MD01-2443) $\delta^{18}O$ records, with generally higher (lower) weight percent $CaCO_3$ during interglacial *s.l.* (glacial) periods. This carbonate signal is well-correlated with the high-resolution XRF log (Ca/Ti) record ($r = 0.91/p < 0.01$; Fig. 4) that reflects varying proportions of biogenic (Ca) and detrital (Ti) sediment supply (Richter et al., 2006). Besides confirming that log (Ca/Ti) provides a reliable proxy for weight percent $CaCO_3$ (Hodell et al., 2013), our results reveal that glacial-interglacial $CaCO_3$ cycles in the Northern Tyrrhenian Sea (Fig. 3d) are similar to those found in the Atlantic Ocean (e.g., Balsam and McCoy, 1987; Helmke and Bauch, 2001; Ruddiman, 1971). In the Western Mediterranean, this variability on glacial-interglacial time scales is assumed to reflect variable dilution by clays (see Hoogakker et al., 2004 for a thorough discussion). During glacial intervals, cold and dry climatic conditions (i.e., low vegetation cover and enhanced soil erodibility) combined with a lowered sea level (i.e., closeness of the sediment source) promote enhanced fluvial input to the sea (Blum and Törnqvist, 2000; Bonneau et al., 2014; Hoogakker et al., 2004). Such a pattern has been described for the Golo River for the last glacial period (Calvès et al., 2013), and is

illustrated for the last 547 kyr by the calculation of the terrigenous flux at site GDEC-4-2 (Fig. 5b). The latter reveals generally higher riverine input during glacials (sea-level lowstands, ≤ -60 m relative to the present day) and lower input during interglacials *s.l.* (sea-level highstands, ≥ -60 m). This also confirmed by the close correlation between the terrigenous flux and the XRF log (Ti/Ca), a reliable proxy for river discharge (e.g., Arz et al., 1998; Toucanne et al., 2009) (Fig. 5c). Thus, the variability in the terrigenous flux explains the glacial-interglacial periodicity observed in the carbonate content at site GDEC-4-2. The reliable agreement between the variations in log (Ca/Ti) at site GDEC-4-2 and those recorded on the Iberian margin (Fig. 5a), where variations in weight percent $CaCO_3$ are ascribed to variable dilution by terrigenous input (Hodell et al., 2013; Thomson et al., 1999), strengthens our interpretation.

Surprisingly, some weight percent $CaCO_3$ and log (Ca/Ti) lows are observed during interglacials *s.s.* (i.e., MIS 5e, MIS 7c and 7e, MIS 9e, MIS 11, MIS 13c; Figs. 3d, 5a and 7c and 8d) with the exception of MIS 1. Both weight percent $CaCO_3$ and Ca/Ti ratios show only moderate values during MIS 1, precisely between 7.5 and 10 ka, and higher carbonate level are described thereafter. Weight percent $CaCO_3$ and log (Ca/Ti) lows observed during previous interglacials *s.s.* coincide with both the lightest values of the local planktonic $\delta^{18}O$, and sea-surface temperature peaks (ca. 19–23 °C) on the Iberian margin and in the Alboran Sea (Martrat et al., 2007, 2014) (Fig. 5e,f and 8e–g). They usually follow (with the exception of MIS 9d) an abrupt millennial Ca/Ti increase that occurs at the end of the termination (e.g., T.V and T.VI) or during the first millennia of the interglacial *s.s.* (e.g., MIS 5e, MIS 7e). These carbonate content lows, also observed during the successive warm intervals that follow the interglacials *s.s.* (e.g., MIS 5c, MIS 7c, MIS 13a), precede maxima in Ca/Ti values. The latter are observed either at the transition between the interglacials *s.s.* and the subsequent millennial cooling events (e.g., MIS 7e-7d transition) or during these cooling events (e.g., MIS 5d, MIS 9d). The carbonate content at site GDEC-4-2 during interglacials *s.s.* thus follows an M-shape pattern that is particularly well-expressed during MIS 5e, MIS7e and MIS11 (Figs. 3d, 5a and 7c). This feature is unexpected considering both the carbonate productivity and the amount of sediment delivery to the Western Mediterranean Sea during interglacial periods. Indeed, interglacial conditions favoured the biogenic carbonate productivity in the Western Mediterranean Sea (Hoogakker et al., 2004) (see Fig. 3d, core LC06, for MIS 5), and the input of river-derived clays to continental margins usually reached a minima at that time. This is primarily caused by the development of the vegetation which tends to increase the stability of soil, and highstand conditions of sea level which favoured the trapping of river sediment inputs on the inner continental shelves (Blum and Törnqvist, 2000). Thus, these conditions (i.e., high carbonate productivity and low dilution by clays) explain the high carbonate content observed in the Balearic Abyssal Plain (Hoogakker et al., 2004) (Fig. 3d) and on the Iberian margin (Hodell et al., 2013) (Fig. 5a) during interglacial conditions. Our data point out a '$CaCO_3$ paradox' on the East Corsica margin during these intervals. For illustrating this '$CaCO_3$ paradox' we compare Ca/Ti values at site GDEC-4-2 with those of the Iberian margin (Hodell et al., 2013) (Fig. 5a). Strong differences in the carbonate content are observed in these two sites at time of interglacial warm intervals and especially during interglacials *s.s.* (e.g., MIS 5a, 5e; MIS 7a, 7c, 7e; MIS 9a, 9c, 9e; MIS 11). This '$CaCO_3$ paradox' is solved through the quantification of the terrigenous flux (Fig. 5b). It reveals that (*i*) input of Golo-derived clays did not reach a minimum at the time of the interglacials *s.s.* but during the second half of the interglacial periods *s.l.* (e.g., MIS 5c, MIS 7d, MIS 9c-d), and (*ii*) except for MIS 1, interglacials *s.s.* sediment flux were equivalent to (or even exceeded; see MIS 7c and MIS 11) those observed during the preceding glacial interval. This indicates that the Golo River activity during interglacials *s.s.* was important enough to produce unexpected high terrigenous flux at site GDEC-4-2 despite the sea-level highstand conditions. The carbonate content lows observed during interglacials *s.s.* and more generally during warm intervals of interglacial periods thus result from a dilution effect by the Golo River terrigenous input. The latter is also confirmed by concomitant increases of the silt fraction at site GDEC-4-2 (Fig. 5d). The evolution of the silt fraction over the studied interval closely matches the Ti/Ca ratios, thus reinforcing their interpretation as proxies for river discharge.

4.1.3. Benthic foraminifera assemblages and carbon isotopes

Benthic foraminifera assemblages and carbon isotopes are used to test the assumption described above for enhanced Golo River discharges during warm intervals of interglacial periods. The relative abundance of the foraminiferal eutrophic group gives reliable insights regarding the Golo River input and the surface-water productivity exported to our bathyal site. Indeed, the relative contribution of the eutrophic species varies over the last 547 kyr in phase with the glacial-interglacial variability (Fig. 6b). Higher abundances of the eutrophic group (abundance of up to 60–80%) are generally found during glacials compared to interglacial *s.l.* periods. This variability likely results from the sea-level variability, with sea-level lowstand conditions enhancing deposition of terrestrial organic matter supplied by the Golo River on the upper slope. Moreover, the depth-dependant exported productivity may increase significantly when sea level is close to the shelf break, which induces a likely increase of eutrophic group abundance. This relationship between the foraminiferal faunas and eustatic changes is corroborated by the highest abundance of the foraminiferal eutrophic group recorded during MIS12 (Fig. 6b), that corresponds to the most extreme lowstand conditions recognized for the last climatic cycles (Grant et al., 2014; Waelbroeck et al., 2002). Minima in the abundance of the eutrophic group (ca. 20%) do not coincide strictly with the interglacials *s.s.* and the attendant sea-level highstand conditions, but match with the second part of the interglacials *s.l.*, following the same pattern of sediment flux as described above. In contrast, some peaks in the abundance of the eutrophic taxa (ca. 50–70%; i.e., equivalent or higher than that observed during glacials) are also recorded at the beginning of the interglacials (e.g., MIS1, MIS5e, MIS7c, MIS9e). It suggests that significant input of organic matter occurs at GDEC-4-2 site during these periods of sea-level highstand. Such an exportation of organic matter may be related to enhanced discharge of the Golo River and the increase of productivity exported to the seabed. This assumption is supported by the concomitant increase in the abundance of the deep infaunal group composed of *Chilostomella* spp. and *Globobulimina* spp, both species being known to proliferate in oxygenated ecosystems where organic detritus focuses (Fontanier et al., 2014, 2005, 2008a).

The benthic $\delta^{13}C$ at site GDEC-4-2 relies mainly on *Cibicidoides pachydermus* ($\delta^{13}C_{Cp}$) and *Cibicidoides kullenbergi* ($\delta^{13}C_{Ck}$) (Fig. 6d) which are shallow infaunal species (Eberwein and Mackensen, 2006; Fontanier et al., 2002, 2013). Their $\delta^{13}C$ signatures, primarily constrained by the $\delta^{13}C$ of bottom water (ca. 0.9‰ for the modern Levantine Intermediate Water -LIW- in the Western Mediterranean Sea; Pierre, 1999), shift to lower values in relation to organic detritus mineralization within the sediment (e.g., Eberwein and Mackensen, 2006; Fontanier et al., 2006a; Schmiedl et al., 2004). In other words, both the modern $\delta^{13}C_{Cp}$ and $\delta^{13}C_{Ck}$ signals echo the $\delta^{13}C_{DIC}$ of pore water, which are likely depleted during early diagenesis compared to the $\delta^{13}C$ of bottom water. For instance, offsets ranging between −0.3‰ and −0.6‰ were documented between the $\delta^{13}C_{DIC}$ of bottom water and $\delta^{13}C_{Cp}$ (Fontanier et al.,

2006b; Schmiedl et al., 2004). For *C. kullenbergi*, offsets between −0.3 and −0.5‰ were recorded by Griveaud (2007) and Licari and Mackensen (2005). We have arbitrarily inserted the higher above-mentioned thresholds of −0.6‰ related to the modern $\delta^{13}C_{DIC}$ of pore water in Fig. 6d. Depletions of both $\delta^{13}C_{Cp}$ and $\delta^{13}C_{Ck}$ signals above this limit (i.e., threshold of 0.3‰; and 0.9‰ for $\delta^{13}C_{Cw}$) are recorded throughout our record, especially during warm interglacial intervals and during the glacial MIS 6d. These events occur simultaneously with high abundances of the deep infaunal group (ca. 10–40%), supporting a higher rate of in-sediment organic matter mineralization in response to enhanced input of organic detritus during these periods. We assume that the synchronicity of benthic $\delta^{13}C$ depletions and peaks in the abundance of eutrophic and deep infaunal foraminifera, coupled with concomitant peaks in terrigenous input, supports the assumption for enhanced Golo River discharges during warm intervals of interglacial periods (interglacials *s.s.* especially) and during the glacial MIS 6d.

4.2. Correlation with Mediterranean records and paleoclimatic implications

4.2.1. Local to regional significance of the rainfall events recorded over Corsica

Our result suggesting enhanced Golo River discharges during interglacial warm intervals (interglacials *s.s.* especially) and the glacial MIS 6d implies a significant increase in precipitation amount over Corsica. This assumption is strongly supported through the correlation between the GDEC-4-2 proxy records for river discharge and paleoclimatic records from the Italian Peninsula and its surrounding seas. Indeed, increased flux of continental materials (i.e., sediment and organic matter) at site GDEC-4-2 correlate with prominent negative $\delta^{18}O$ excursions in both the Sulmona basin (Central Italy; Regattieri et al., 2015) and the Argentarola (Tyrrhenian coast of Italy; Bard et al., 2002), Antro del Corchia and Tana Che Urla (Central Italy; Drysdale et al., 2005; Regattieri et al., 2014; Zanchetta et al., 2007) cave speleothems (Fig. 7a), and with salinity decreases in the central Adriatic basin (Piva et al., 2008) and in the Eastern Tyrrhenian Sea (Kallel et al., 2000). This indicates that the high runoff recorded for the Golo River during warm intervals of interglacial periods (interglacials *s.s.* especially) and the glacial MIS 6d likely originate from enhanced regional rainfall. Such 'pluvial' conditions explain the ca. 1.5–2.7‰ depletions observed in the planktonic $\delta^{18}O$ during MIS 5 and MIS 7 (Figs. 3c, 5f and 6a). As a result, the GDEC-4-2 borehole can be seen as a valuable record for rainfall variability in the Western Mediterranean.

The comparison of our results with paleoclimatic records located further east suggests a wider significance for the GDEC-4-2 'runoff/rainfall' record presented here. The comparison of the GDEC-4-2 proxy records for river discharge with the 250-kyr composite isotopic record of speleothems from the Soreq and Peqiin caves (Israel) reveals that periods characterised by increased Corsican runoff since MIS 7 (i.e., MIS1, MIS 5a, 5c, 5e; MIS 6d; MIS 7a, 7c, 7e) coincide with high rainfall regimes (i.e., low $\delta^{18}O$) in the eastern Mediterranean region (Fig. 7a) (Ayalon et al., 2002; Bar-Matthews et al., 2003). The runoff increase recorded at site GDEC-4-2 during MIS 5e is also synchronous with a significant salinity decrease at site KS205, northwest Ionian Sea (Fig. 8e) (Rohling et al., 2002). In agreement with the conclusions of Bard et al. (2002) for MIS 6 and Kallel et al. (2000) for the 200–60 ka interval, this shows that both western and eastern Mediterranean basins experienced simultaneously wetter conditions over the last 250 ka, precisely during interglacial warm intervals (interglacials *s.s.* especially) and the glacial MIS 6d. The lack of pre-MIS 7 speleothem-based rainfall records in the Eastern Mediterranean precludes a robust inter-basin comparison for the basal part of the GDEC-4-2 borehole. Nevertheless, the above correlation for the last 250 ka as well as palynological evidence showing relatively high water availability during interglacials in Northern Greece (Fig. 7f) (Tenaghi Philippon; Tzedakis et al., 2006; Tzedakis et al., 2009) and in Northern Levant (Yammouneh sequence; Gasse et al., 2014) encourages us to identify the GDEC-4-2 events (high -silty-terrigenous input, high abundance of foraminiferal eutrophic and deep infaunal taxa, and benthic $\delta^{13}C$ depletions) over the 547-250-kyr period as pluvial episodes over the Mediterranean basin (Fig. 6). As a result, we assume that Mediterranean 'pluvial' phases also occurred during MIS 9 (MIS 9e especially), MIS 11 and MIS 13 (MIS 13a and 13b) (Figs. 3, 5 and 6). This highlights interglacial 'pluvial' periods as a distinctive feature of the Mediterranean climate (Bar-Matthews et al., 2003) and confirms that (minima in) precession (i.e., winter insolation minima and summer insolation maxima) has been a pacemaker for rainfall variability in the Mediterranean basin (e.g., Lourens et al., 1992; Sánchez Goñi et al., 2008; Tzedakis, 2007). The trend in the evolution of the GDEC-4-2 proxy records for river discharge (especially the terrigenous fluxes and the abundance of the deep infaunal foraminifera) also attests to the impact of eccentricity modulation of precession (and insolation) on the runoff/rainfall activity. Indeed, maxima of the 100-kyr eccentricity cycles correlate with periods of highest Golo River discharge (Figs. 6c and 7b). This astronomical forcing, previously identified as a pacemaker for annual rainfall (Sierro et al., 2000) and lake-level fluctuations in Neogene sequences of the Western Mediterranean (Abels et al., 2009; Valero et al., 2014; Van Vugt et al., 2001), likely explains the significant difference in amplitude of the Golo River discharges between MIS 1 (low-amplitude), MIS 5e (moderate to high) and MIS 7c (very-high) for example.

4.2.2. Source and timing of Mediterranean rainfall events recorded at the GDEC-4-2 site

The Mediterranean pluvial phases recorded at site GDEC-4-2 (as well as in the Israeli speleothems over the last 250 kyr; Bar-Matthews et al., 2003) are synchronous with increased river runoff from the Libyan-Egyptian sector of North Africa. This is nicely illustrated at high-resolution for MIS 5e through the comparison between the GDEC-4-2 proxy records for river discharge and the *G. ruber* $\delta^{18}O$ record of ODP 967 and ODP 971 (Fig. 8f,g) located close to the outflow of the Nile and the Libyan river systems, respectively (Rohling et al., 2002). For the last climatic cycles, the relationship between Mediterranean 'pluvial' phases and the runoff from North African rivers is supported by the correlation between the GDEC-4-2 dataset and the XRF Ba/Al ratio from ODP 968 (Fig. 7d) (Eastern Mediterranean; Ziegler et al., 2010) that reveals that the timing of 'pluvial' phases coincides with that of sapropel deposition. Indeed, the sapropel deposition, as a direct consequence of a profound disruption in the basin's hydrological cycle, is directly linked to increased river summer floods from the Central Saharan watersheds (e.g., Irharhar, Sahabi and Kufrah rivers; Fig. 1) and the Nile River (Larrasoana et al., 2003; Osborne et al., 2008; Rohling et al., 2002; Rohling and Hilgen, 1991; Rossignol-Strick, 1985; Rossignol-Strick et al., 1982; Scrivner et al., 2004). Increased summer floods from the Nile and Central North African rivers are linked to the intensification and northward extension of the African summer monsoon at time of minima in the precession index (i.e., during northern hemisphere summer insolation maxima), but the monsoon rains only penetrate to 25°N (Rohling et al., 2002; Rohling and Hilgen, 1991; Tuenter et al., 2003; Tzedakis, 2007). In addition, terrestrial pollen sequences from Italy, Greece, Turkey and the Levant reveals the expansion of Mediterranean sclerophyllous vegetation, indicative of increased summer aridity, in the northern Mediterranean borderlands during interglacial




Fig. 7. (A) $\delta^{18}O$ record of the Soreq (Central Israel, black line; Bar-Matthews et al., 2003), Peqiin (Northern Israel, grey line; Bar-Matthews et al., 2003), Antro del Corchia (Central Italy, orange line; Drysdale et al., 2005; Zanchetta et al., 2007), Tana Che Urla (Central Italy, purple line; Regattieri et al., 2014) and Argentarola (Tyrrhenian coast of Italy, green line; Bard et al., 2002) speleothems. The $\delta^{18}O$ record of the Sulmona basin (Central Italy, pink line; Regattieri et al., 2015) is also shown; (B) Abundance of the benthic foraminifera deep-infaunal group (red line) and eccentricity of the Earth's orbit (dashed grey line; Laskar et al., 2004). The interglacials (*s.s.*) terrigenous flux at site GDEC-4-2 are also shown (blue diamonds). These fluxes can be compared since similar sea-level highstand conditions prevailed during these intervals; (C) XRF Ca/Ti (log scale) of the GDEC-4-2 borehole; (D) XRF Ba/Al (log scale) from ODP 968 (Southern Cyprus; Ziegler et al., 2010). High Ba/Al ratios are very characteristic for sapropel layers. Chronology for sapropel layers (speleothem age-scale) according to Ziegler et al. (2010); (E) $\delta^{18}O$ record of the Sanbao-Hulu caves (Wang et al., 2008); (F) Tenaghi Philippon temperate tree pollen percentages (Tzedakis et al., 2006); (G) June insolation for 65°N (black line) and precession (dashed grey line) (Laskar et al., 2004). Seasonality at 65°N (green line) is shown through the difference JJA (June, July, August) − DJF (December, January, February). Vertical blue bands indicate the timing for sapropel deposition (Ziegler et al., 2010). Terminations (T.) I to III according to Barker et al. (2011). Scheme of marine isotope substages according to Railsback et al. (2015). (For interpretation of the references to colour in this figure caption, the reader is referred to the web version of this article.)




Fig. 8. (A) June insolation (blue line) and seasonality (JJA-DJF, purple line) for 45°N (Laskar et al., 2004); (B) Alboran Sea SST, ODP 976 and ODP 977 (Martrat et al., 2014); (C) Pollen percentage data for sclerophyllous Mediterranean at Tenaghi Philippon (Milner et al., 2012); (D) XRF Ca/Ti (log scale; continuous line) and abundance of the silt fraction (dashed line) at site GDEC-4-2; (E, F, G) $\delta^{18}O$ of *G. ruber* (white) in core KS205 (Ionian Sea), ODP 971 (Levantine Sea) and ODP 967 (Southern Cyprus), all plotted versus the ODP 971-equivalent depth scale (see Rohling et al., 2002 for details). Conversion on the speleothem age-scale is based on the alignment of the $\delta^{18}O$ of *G. ruber* in ODP 967 (i.e., enhanced Nile outflow at time of Sapropel S5; Rohling et al., 2002) on the Ba/Al ratio (Sapropel S5 layer, see Fig. 8h) of Ziegler et al. (2010) in ODP 968 (located ca. 30 km north); (H) XRF Ba/Al (log scale) from ODP 968 (Southern Cyprus; Ziegler et al., 2010); (I) Paleorainfall $\delta^{18}O$ values at Soreq Cave (Central Israel; Bar-Matthews et al., 2003); (J) Continental climatic episodes (Ducassou et al., 2009 and references therein). Chronology (speleothem age-scale) for Sapropel S5 according to Ziegler et al. (2010).

sapropel deposition (Milner et al., 2012; Tzedakis, 2007; Tzedakis et al., 2003) (see Fig. 8c for MIS 5e and the Sapropel S5 deposition). Taken together, these results suggest that the Mediterranean 'pluvial' events recorded at site GDEC-4-2 probably occurred in autumn/winter (i.e., during winter insolation minima) (Bosmans et al., 2015; Fletcher and Sánchez Goñi, 2008; Kotthoff et al., 2008; Rohling et al., 2015; Sánchez Goñi et al., 2008; Tzedakis, 2007). This is supported for the last interglacial by pollen-based climate reconstructions of temperature and precipitation seasonality from Lake Accesa (Italy), ca. 100 km east of site GDEC-4-2 (Peyron et al., 2011). These 'pluvial' events probably originated from the cyclogenesis mechanisms in the Mediterranean (Reale et al., 2001; Trigo et al., 2002, 1999). Nowadays, the northern Mediterranean borderlands (including, from west to east: site GDEC-4-2, the Corchia and Argentarola caves, the KS205 marine record, the terrestrial pollen sequences cited above and the Soreq and Peqiin caves) are under the influence of the autumn/winter (i.e., October–March) Mediterranean storm track, with the Gulf of Genoa and the Aegean Sea consisting of two of the most active cyclogenetic regions in the Mediterranean realm. The cyclogenesis over these regions can occur consecutively as the result of the same major synoptic system, usually of North Atlantic origin, crossing central Europe (Trigo et al., 2002, 1999). We showed previously that enhanced interglacial runoff of the Golo River, interpreted as enhanced rainfall over Corsica, coincides with the maximum rainfall amount in Israel (Bar-Matthews et al., 2003). By considering both the Mediterranean origin of the Levant precipitation (Bar-Matthews et al., 2003, 2000) and the pollen-based reconstructions of precipitation seasonality in the northern Mediterranean borderlands (Milner et al., 2012; Peyron et al., 2011; Tzedakis, 2007), we thus assume that the GDEC-4-2 sequence records the upstream activity of the autumn/winter Mediterranean storm track over the last 547 kyr. In other words, the GDEC-4-2 borehole is the first sequence containing multiple precession cycles that highlights enhanced autumn/winter rainfall over the northern Mediterranean borderlands during interglacial warm intervals, in particular during interglacials *s.s.*. Our results support recent modelling experiments showing increased winter precipitation in regions between 30°N and 45°N over the Mediterranean and Middle East during periods of maximum orbitally-forced insolation seasonality (Kutzbach et al., 2014). Our results also reconcile the speleothem-based snapshots for rainfall activity in the northern Mediterranean borderlands during interglacial warm intervals (Bar-Matthews et al., 2003, 2000; Drysdale et al., 2005; Göktürk et al., 2011; Zanchetta et al., 2007) with the incongruent evidence (i.e., increased -summer- aridity during interglacial *s.s.*) from long-term pollen sequences (see Tzedakis, 2007 for a thorough review).

4.2.3. *Rainfall events over the northern Mediterranean borderlands and the African monsoon*

The above lines of evidence highlight a direct relationship between the rainfall variability over the northern Mediterranean borderlands and the precession forcing. By considering the long-

known link between the precession forcing and the low-latitude climate (e.g., Hilgen, 1991; Rossignol-Strick, 1985; Trauth et al., 2009; Tzedakis et al., 2009), we can hypothesize strong teleconnections between the Mediterranean storm track (and more generally the mid-latitude atmospheric circulation pattern) and the monsoon activity. The coincidence in the timing of the increased Mediterranean (autumn/winter) storm track precipitation with that of the North African (summer) monsoon during warm intervals of interglacial periods, shown by the correlation of 'pluvial' events over Corsica and the increased runoff of the Nile River (Figs. 7 and 8), highlights a close link between mid- and low-latitude hydrological changes. Moreover, these climatic events coincide with increased rainfall over East Asia in response to enhanced (summer) monsoon (Fig. 7e) (Jo et al., 2014; Wang et al., 2008). The concomitant changes in the precipitation patterns over the northern Mediterranean borderlands, the North Africa continent and East Asia likely reflect shifts in the mean latitudinal position of the ITCZ, with an attendant climatic response at mid-latitudes.

This relationship between mid- and low-latitude hydrological changes is nicely depicted for Southern Europe and North Africa at centennial to millennial time-scale during MIS 5e, especially between ca. 121 and 129 ka (Fig. 8). During this period of enhanced summer monsoon over North Africa, a sustained (ca. 800 yr) relaxation of the ITCZ penetration leading to a monsoon disruption occurred at around 124–125 ka (after alignment on the speleothem age-scale; Rohling et al., 2002). At the same time, while enhanced winter rainfall dominates over the 122–127 kyr interval, a sustained decrease in riverine inputs occurred off Corsica (Fig. 8d). This short-lived event, accompanied by a ca. 2‰ increase of the *G. ruber* $\delta^{18}O$ in the Ionian Sea (Fig. 8e) (Rohling et al., 2002), highlights a reduction in the winter precipitation and, by extension, a decrease of the Atlantic–Mediterranean cyclogenesis activity. This close relationship between the ITCZ motion, the North Africa summer monsoon and the rainfall activity over the northern Mediterranean borderlands, evident for the whole interval (i.e., 121–129 ka; Fig. 8), points out a positive correlation between the intensity of the North African summer monsoon and the cyclogenesis activity in the Mediterranean Sea (i.e., the stronger the North Africa summer monsoon, the stronger the Mediterranean winter rainfall). This indicates that climate conditions over North Africa and the Mediterranean in summer could possibly force the cyclogenesis activity on the northern Mediterranean borderlands during the successive winter. At present, it is well-known that the northward shift of the ITCZ over North Africa during summer (with the associated reinforcement of the Hadley circulation and the eastwards expansion of the Azores subtropical high) induces hot and dry conditions in the western Mediterranean region (Baldi et al., 2004; Xoplaki et al., 2003). These climatic conditions could also originate from the Indian summer monsoon and its impact on subsidence over the Mediterranean Sea through the westward propagation of Rossby waves (Marzin and Braconnot, 2009; Rodwell and Hoskins, 1996, 2001). By considering the extreme northward position of the ITCZ and the strong seasonality pattern (i.e., summer insolation maxima versus winter insolation minima) during interglacial *s.s.*, the summer conditions over the Mediterranean adding to the subsequent rapid climate deterioration occurring in autumn/winter could have increased the air-sea temperature contrast and, correspondingly, the necessary heat and moisture fluxes required for the development and strengthening of cyclonic circulations over the northern Mediterranean borderlands (Kutzbach et al., 2014; Meijer and Tuenter, 2007; Trigo et al., 2002, 1999; Tuenter et al., 2003). This allowed the autumn/winter rainy westerlies to reach the Mediterranean area more frequently. This pattern could explain the intimate relationship observed between the North African summer monsoon and the Mediterranean autumn/winter rainfall during MIS 5e, while also giving insights into the mechanism for the transfer of energy between the tropics and higher latitudes (Rohling et al., 2002).

4.3. Mediterranean rainfall events and sapropel deposition

The origin of sapropel deposits, and precisely the source(s) for freshwater, nutrients and continental organic matter, has long been debated (e.g., Rohling et al., 2015). The meltwater runoff from the Fennoscandian ice sheet into the Eastern Mediterranean by way of the Black and Aegean seas was first proposed (Ryan, 1972; Williams et al., 1978). Later, Rossignol-Strick et al. (1982) and Rossignol-Strick (1985) identified heavy Nile River summer floods due to increased monsoonal summer precipitation over Ethiopia as the main forcing for sapropel deposition. The contribution from Macedonian, Greek and Turkish rivers, in addition to that of the Nile, was then pointed out (Cramp et al., 1988; Rossignol-Strick, 1987; Shaw and Evans, 1984; Wijmstra et al., 1990). Although these studies only focused on the Late Pleistocene and Sapropel S1, it has been widely accepted that precipitation over the northern borderlands of the Eastern Mediterranean also played a significant role for sapropel deposition (Rohling and Hilgen, 1991; Rossignol-Strick, 1987). This assumption was ultimately supported through the 250-kyr isotopic record of cave speleothems from the Levant that revealed that the periods for deposition of Sapropel S1 to S9 were characterized by enhanced rainfall of Mediterranean origin in the eastern Mediterranean land and sea regions (Bar-Matthews et al., 2003, 2000). At the other side of the Mediterranean realm, a few studies revealed that increased precipitation over the Western Mediterranean occurs simultaneously with the formation of Sapropel S1 (Ariztegui et al., 2000; Kotthoff et al., 2008; Magny et al., 2013; Zanchetta et al., 2007), Sapropel S4 (Regattieri et al., 2015), Sapropel S5 (Drysdale et al., 2005; Kallel et al., 2000), Sapropel S6 (Bard et al., 2002) and Sapropel S7 (Kallel et al., 2000), thus suggesting that increased rainfall during sapropel deposition was not restricted to the Eastern Mediterranean. Although compelling, each of these records were discontinuous in character. The correlation of the 547-kyr GDEC-4-2 sequence for Golo River runoff with the long speleothem records from the Levant (Bar-Matthews et al., 2003), as well as with the continental and marine (short) records scattered in the central and eastern Mediterranean regions (Ariztegui et al., 2000; Bard et al., 2002; Cramp et al., 1988; Emeis et al., 2000; Göktürk et al., 2011; Kallel et al., 2000; Kotthoff et al., 2008; Milner et al., 2012; Regattieri et al., 2015; Rossignol-Strick, 1987; Shaw and Evans, 1984; Wijmstra et al., 1990; Zanchetta et al., 2007) confirms that increased rainfall was widespread over the northern Mediterranean borderlands during sapropel deposition.

Our evidence for enhanced rainfall activity along the northern Mediterranean borderlands during warm intervals of interglacial periods provides an additional constraint on the role of the mid-latitude storm tracks on the forcing leading to sapropel deposition. First by increasing the flux of continental organic matter to the oceanic basin through river runoff, with possible subsequent positive effects on the primary productivity and the marine organic matter flux to the seafloor (e.g., Rohling and Hilgen, 1991). Second, by creating the necessary hydrological changes leading to disruption in the basin's hydrological cycle and reduced intermediate and deep water ventilation (Meijer and Tuenter, 2007). Indeed, it has been shown that the Nile input was not the only factor affecting the freshwater balance of the Eastern Mediterranean during sapropel events (Scrivner et al., 2004). Based on these considerations, we propose the following scenario for interglacials *s.s.*: precession-driven Northern Hemisphere insolation maxima, through the northward shift of the ITCZ, led to heavy summer rainfall

(June–September; Janowiak, 1988) over Northern Africa and to subsequent summer–autumnal (August–October; Conway and Hulme, 1993) Nile (and wadi-systems) floods into the Eastern Mediterranean (Coulthard et al., 2013; Rohling et al., 2002; Rossignol-Strick, 1985). At seasonal scale, warm and arid conditions prevailed in summer over southern Europe and the Mediterranean until the rapid arrival of autumn/winter conditions due to the extreme seasonality (i.e., precession minima). The Mediterranean storm track intensified and rainfall was enhanced over the northern Mediterranean borderlands until the next spring and the rapid recovery of hot dry conditions over the Mediterranean. Thus, high flux of freshwater, nutrient and continental organic matter entered the Mediterranean Sea from the Northern African margin and the northern Mediterranean borderlands in summer/autumn and autumn/winter, respectively. This indicates that on an annual scale, the rainfall over the northern Mediterranean Sea maintained the reduced sea-surface water salinities and the high-flux of nutrient and organic matter in the Mediterranean Sea initially caused by the Nile (and wadi-systems) floods, until the next spring. This pattern prevailed as the insolation/seasonality is high and deeply affects, through its duration, the basin's hydrological cycle up to the development of widespread bottom anoxic conditions. It is difficult to determine whether Nile floods or rainfall over the northern Mediterranean borderlands is the dominant forcing of changes in stratification and deep water formation (Bosmans et al., 2015). Meijer and Tuenter (2007) have previously shown that the effects of increased rainfall over the northern Mediterranean borderlands are possibly of equal or greater importance than that of increased Nile discharge. A prerequisite for such an impact is that rainfall had an extra-Mediterranean origin, since local rainfall (i.e., convective precipitation) cannot have affected the intermediate and deep water ventilation significantly (Rohling et al., 2015). Such an origin is questioned by recent numerical modelling (Bosmans et al., 2015) but there is support from the North Atlantic source of modern extreme-rainfall episodes over the northern Mediterranean borderlands (e.g., Celle-Jeanton et al., 2004; Duffourg and Ducrocq, 2013; Reale et al., 2001), from studies showing surface ocean dilution (Emeis et al., 2003; Kallel et al., 2000, 1997) and from changes in the sources of precipitation in the Western Mediterranean at time of sapropel formation (e.g., Bard et al., 2002; Drysdale et al., 2004; Zanchetta et al., 2007).

5. Conclusion

The 125 m-long GDEC-4-2 borehole provides a continuous high-resolution paleoclimatic record on the East Corsica margin, northern Tyrrhenian Sea, over the last 547 kyr. Sedimentological, geochemical and micropaleotological analysis reveal a close coupling between river runoff and climate changes over the studied period, and during warm intervals of interglacial periods especially. The main results of our study are the following:

1. Warm intervals of interglacial periods and interglacials *s.s.* are characterised by high terrigenous flux and benthic foraminifera indicative of high organic matter input. This evidence points to high sediment discharges by East Corsica rivers in response to high precipitation levels;
2. Comparison of GDEC-4-2 proxy records for river discharge with pollen sequences and speleothems from the central and eastern Mediterranean reveals that the interglacial wet conditions recorded off Corsica were regional in character. Considering the present-day Mediterranean climate pattern and palynological evidence, our dataset likely records the activity of the autumn/winter Mediterranean storm track along the northern Mediterranean borderlands and its intensification during interglacials *s.s.*;
3. If the correlation between the rainfall activity over the northern Mediterranean borderlands and the interglacial warm intervals confirms the precessional component of the Mediterranean climate, the GDEC-4-2 record is long enough to highlight the impact of eccentricity modulation of precession (and insolation) on the Mediterranean cyclogenesis;
4. Our dataset reveals a correlation between the timing of maxima of the Mediterranean autumn/winter storm track precipitation and that of the North African summer monsoon. Millennial-scale examination of the penultimate interglacial *s.s.* (MIS 5e) suggests that climate conditions over North Africa (and possibly India) and the Mediterranean in summer could force the cyclogenesis activity on the northern Mediterranean borderlands during the successive autumn/winter. This highlights a close coupling between low- and mid-latitude hydrological changes;
5. Finally, our findings suggest that during warm intervals of interglacial periods the reduced sea-surface water salinities, together with the high flux of nutrient and organic matter, produced by the monsoonal (i.e., summer/autumn) Nile (and wadi-systems) floods, were maintained throughout the winter by the Mediterranean rainfall. This identifies rainfall over the northern Mediterranean borderlands, in addition to the river runoff from the Libyan–Egyptian sector of North Africa, as a possible forcing on sapropel deposition.

Acknowledgements

The authors are very grateful to A. Roubi, M. Rovere, C. Marin, M. Montferrand, L. Blanchet (IFREMER) and J. Duprat for their technical support; E. Michel, F. Dewilde, G. Isguder (LSCE) for their expert analytical work about stable isotopes measurements; D. Hodell and P.C. Tzedakis that kindly provided data from the Iberian margin and Tenaghi Philippon, respectively; T. Caley, Y. Dixit, N. Freslon, T. Konijnendijk, M. Rogerson, M.F. Sánchez Goñi and M. Ziegler for invaluable discussions at various stages of this work; and three anonymous reviewers who provide helpful review of the manuscript. The authors also thank the crew and scientific team (G. Calvès, G. Dan-Unterseh, L. Droz, I. Jégou, G. Lericolais, E. Marchès, T. Marsset) of the GOLODRILL cruise (R/V 'Bavenit' – FUGRO) for the recovery of the GDEC-4-2 borehole. This study has been realised in the frame of the "GOLO PROGRAM", a research consortium between IFREMER, TOTAL, EXXONMOBIL and FUGRO.

References

Abels, H.A., Abdul Aziz, H., Calvo, J.P., Tuenter, E., 2009. Shallow lacustrine carbonate microfacies document orbitally paced lake-level history in the Miocene Teruel Basin (North-East Spain). Sedimentology 56 (2), 399–419.
Abu-Zied, R.H., Rohling, E.J., Jorissen, F.J., Fontanier, C., Casford, J.S.L., Cooke, S., 2008. Benthic foraminiferal response to changes in bottom-water oxygenation and organic carbon flux in the eastern Mediterranean during LGM to recent times. Mar. Micropaleontol. 67, 46–68.
Allen, J.R.M., et al., 1999. Rapid environmental changes in southern Europe during the last glacial period. Nature 400, 740–743.
Ariztegui, D., et al., 2000. Palaeoclimate and the formation of sapropel S1: inferences from late quaternary lacustrine and marine sequences in the central Mediterranean region. Palaeogeogr. Palaeoclimatol. Palaeoecol. 158 (3–4), 215–240.
Arz, H.W., Pätzold, J., Wefer, G., 1998. Correlated millennial-scale changes in surface hydrography and terrigenous sediment yield inferred from Last-Glacial marine deposits off northeastern Brazil. Quat. Res. 50 (2), 157–166.
Arz, H.W., Lamy, F., Pätzold, J., Müller, P.J., Prins, M.A., 2003. Mediterranean moisture source for an early-Holocene humid period in the Northern Red Sea. Science 300, 118–121.
Ayalon, A., Bar-Matthews, M., Kaufman, A., 2002. Climatic conditions during marine oxygen isotope stage 6 in the eastern Mediterranean region from the isotopic composition of speleothems of Soreq Cave, Israel. Geology 30 (4), 303–306.

Baldi, M., Meneguzzo, F., Dalu, G.A., Maracchi, G., Pasqui, M., Capecchi, V., Crisci, A., Piani, F., 2004. Guinea GULF SST and Mediterranean summer climate: analysis of the interannual variability. Bull. Am. Meteorol. Soc. 4167–4187.
Balsam, W.L., McCoy, F.W., 1987. Atlantic sediments: glacial/interglacial comparisons. Paleoceanography 2 (5), 531–542.
Bar-Matthews, M., Ayalon, A., Gilmour, M., Matthews, A., Hawkesworth, C.J., 2003. Sea–land oxygen isotopic relationships from planktonic foraminifera and speleothems in the Eastern Mediterranean region and their implication for paleorainfall during interglacial intervals. Geochim. Cosmochim. Acta 67 (17), 3181–3199.
Bar-Matthews, M., Ayalon, A., Kaufman, A., 2000. Timing and hydrological conditions of Sapropel events in the Eastern Mediterranean, as evident from speleothems, Soreq cave. Isr. Chem. Geol. 169 (1–2), 145–156.
Bard, E., Delaygue, G., Rostek, F., Antonioli, F., Silenzi, S., Schrag, D.P., 2002. Hydrological conditions over the western Mediterranean basin during the deposition of the cold Sapropel 6 (ca. 175 kyr BP). Earth Planet. Sci. Lett. 202, 481–494.
Barker, S., Knorr, G., Edwards, R.L., Parrenin, F., Putnam, A.E., Skinner, L.C., Wolff, E., Ziegler, M., 2011. 800,000 years of abrupt climate variability. Science 334 (6054), 347–351.
Berger, A., 1978. Long-term variations of daily insolation and quaternary climatic changes. J. Atmos. Sci. 35 (12), 2362–2367.
Blum, M.D., Törnqvist, T.E., 2000. Fluvial responses to climate and sea-level change: a review and look forward. Sedimentology 47, 2–48.
Bonneau, L., Jorry, S.J., Toucanne, S., Silva Jacinto, R., Emmanuel, L., 2014. Millennial-scale response of a Western Mediterranean river to late quaternary climate changes: a view from the deep sea. J. Geol. 122 (6), 687–703.
Bosmans, J.H.C., Drijfhout, S.S., Tuenter, E., Hilgen, F.J., Lourens, L.J., Rohling, E.J., 2015. Precession and obliquity forcing of the freshwater budget over the Mediterranean. Quat. Sci. Rev. 123, 16–30.
Brauer, A., Allen, J.R.M., Mingram, J., Dulski, P., Wulf, S., Huntley, B., 2007. Evidence for last interglacial chronology and environmental change from Southern Europe. Proc. Natl. Acad. Sci. 104 (2), 450–455.
Cacho, I., Grimalt, J.O., Pelejero, C., Canals, M.I., Sierro, F.J., Flores, J.A., Shackleton, N., 1999. Dansgaard-Oeschger and Heinrich events imprints in Alboran Sea paleotemperatures. Paleoceanography 14 (6), 698–705.
Cacho, I., Shackleton, N.J., Elderfield, H., Sierro, F.J., Grimalt, J.O., 2006. Glacial rapid variability in deep-water temperature and d^{18}O from the Western Mediterranean Sea. Quat. Sci. Rev. 25, 3294–3311.
Calvès, G., et al., 2013. Inferring denudation and climatic variations from the sediment record; an example of the last glacial cycle record of the Golo basin and watershed, East Corsica, western Mediterranean Sea. Basin Res. 25 (2), 197–218.
Casford, J.S.L., Rohling, E.J., Abu-Zied, R.H., Fontanier, C., Jorissen, F.J., Leng, M.J., Schmiedl, G., Thomson, J., 2003. A dynamic concept for eastern Mediterranean circulation and oxygenation during sapropel formation. Palaeogeogr. Palaeoclimatol. Palaeoecol. 190, 103–119.
Celle-Jeanton, H., Travi, Y., Blavoux, B., 2004. Isotopic typology of the precipitation in the Western Mediterranean region at three different time scales. Geophys. Res. Lett. 28, 1215–1218.
Conway, D., Hulme, M., 1993. Recent fluctuations in precipitation and runoff over the Nile sub-basins and their impact on main Nile discharge. Clim. change 25 (2), 127–151.
Coulthard, T.J., Ramirez, J.A., Barton, N., Rogerson, M., Brücher, T., 2013. Were rivers flowing across the Sahara during the last interglacial? Implications for human migration through Africa. PLoS One 8 (9). http://dx.doi.org/10.1371/journal.pone.0074834.
Cramp, A., Collins, M., West, R., 1988. Late pleistocene Holocene sedimentation in the NW Aegean Sea: a palaeo- climatic palaeoceanographic reconstruction. Palaeogeogr. Palaeoclimatol. Palaeoecol. 68, 61–77.
Cremer, M., Grousset, F., Faugeres, J.C., Duprat, J., Gonthier, E., 1992. Sediment flux patterns in the northeastern Atlantic: variability since the last interglacial. Mar. Geol. 104 (1–4), 31–53.
De Rijk, S., Jorissen, F.J., Rohling, E.J., Troelstra, S.R., 2000. Organic flux control on bathymetric zonation of Mediterranean benthic foraminifera. Mar. Micropaleontol. 40 (3), 151–166.
Desprat, S., Sanchez Goni, M.F., Turon, J.L., Duprat, J., Malaize, B., Peypouquet, J.P., 2006. Climatic variability of marine isotope stage 7: direct land-sea-ice correlation from a multiproxy analysis of a north-western Iberian margin deep-sea core. Quat. Sci. Rev. 25, 1010–1026.
Drake, N.A., Blench, R.M., Armitage, S.J., Bristow, C.S., White, K.H., 2011. Ancient watercourses and biogeography of the Sahara explain the peopling of the desert. Proc. Natl. Acad. Sci. 108 (2), 458–462.
Drysdale, R.N., Zanchetta, G., Hellstrom, J.C., Fallick, A.E., Zhao, J.X., 2005. Stalagmite evidence for the onset of the last interglacial in southern Europe at 129±1 ka. Geophys. Res. Lett. 32 (24) http://dx.doi.org/10.1029/2005GL024658.
Drysdale, R.N., Zanchetta, G., Hellstrom, J.C., Fallick, A.E., Zhao, J.X., Isola, I., Bruschi, G., 2004. Palaeoclimatic implications of the growth history and stable isotope (δ^{18}O and δ^{13}C) geochemistry of a middle to late pleistocene stalagmite from central-western Italy. Earth Planet. Sci. Lett. 227 (3–4), 215–229.
Ducassou, E., Migeon, S., Mulder, T., Murat, A., Capotondi, L., Bernasconi, S.M., Mascle, J., 2009. Evolution of the Nile deep-sea turbidite system during the late quaternary: influence of climate change on fan sedimentation. Sedimentology 56, 2061–2090.
Duffourg, F., Ducrocq, V., 2013. Assessment of the water supply to Mediterranean heavy precipitation: a method based on finely designed water budgets. Atmos. Sci. Lett. 14 (3), 133–138.
Duplessy, J.C., 1978. Isotopes studies. In: Gribbin, J. (Ed.), Climatic Change. Cambridge Univ. Press, London/New York, pp. 46–67.
Eberwein, A., Mackensen, A., 2006. Regional primary productivity differences off Morocco (NW-Africa) recorded by modern benthic foraminifera and their stable carbon isotopic composition. Deep Sea Res. Part I Oceanogr. Res. Pap. 53, 1379–1405.
Emeis, K.C., Struck, U., Schulz, H.-M., Rosenberg, R., Bernasconi, S., Erlenkeuser, H., Sakamoto, T., Martinez-Ruiz, F., 2000. Temperature and salinity variations of Mediterranean Sea surface waters over the last 16,000 years from records of planctonic stable isotopes and alkenone unsaturation ratios. Palaeogeogr. Palaeoclimatol. Palaeoecol. 158, 259–280.
Emeis, K.C., et al., 2003. Eastern Mediterranean surface water temperatures and δ^{18}O composition during deposition of sapropels in the late quaternary. Paleoceanography 18 (1), 1005. http://dx.doi.org/10.1029/2000PA000617.
Fletcher, W.J., Sánchez Goñi, M.F., 2008. Orbital- and sub-orbital-scale climate impacts on vegetation of the western Mediterranean basin over the last 48,000 yr. Quat. Res. 70, 451–464.
Fontanier, C., Biscara, L., Mamo, B., Delord, E., 2014. Deep-sea foraminifera in an area around the Cassidaigne Canyon (NW Mediterranean) affected by bauxite discharge. Mar. Biodivers. 1–12. http://dx.doi.org/10.1007/s12526-014-0281-9.
Fontanier, C., Jorissen, F.J., Licari, L., Alexandre, A., Anschutz, P., Carbonel, P., 2002. Live benthic foraminiferal faunas from the Bay of Biscay: faunal density, composition, and microhabitats. Deep Sea Res. Part I – Oceanogr. Res. Pap. 49 (4), 751–785.
Fontanier, C., Jorissen, F.J., Anschutz, P., Chaillou, G., 2006a. Seasonal variability of benthic foraminiferal faunas at 1000 m depth in the Bay of Biscay. J. Foraminifer. Res. 36, 61–76.
Fontanier, C., Jorissen, F.J., Chaillou, G., Anschutz, P., Grémare, A., Griveaud, C., 2005. Live foraminiferal faunas from a 2800 m deep lower canyon station from the Bay of Biscay: faunal response to focusing of refractory organic matter. Deep Sea Res. Part I Oceanogr. Res. Pap. 52, 1189–1227.
Fontanier, C., Mackensen, A., Jorissen, F.J., Anschutz, P., Licari, L., Griveaud, C., 2006b. Stable oxygen and carbon isotopes of live benthic foraminifera from the Bay of Biscay: microhabitat impact and seasonal variability. Mar. Micropaleontol. 58 (3), 159–183.
Fontanier, C., Jorissen, F.J., Geslin, E., Zaragosi, S., Duchemin, G., Laversin, M., Gaultier, M., 2008a. Live and dead foraminiferal faunas from the Saint-Tropez Canyon (Bay of Fréjus): observations based on in situ and incubated cores. J. Foraminifer. Res. 38, 137–156.
Fontanier, C., et al., 2008b. Live foraminifera from the open slope between Grand Rhône and Petit Rhône Canyons (Gulf of Lions, NW Mediterranean). Deep Sea Res. Part I – Oceanogr. Res. Papers 55 (11), 1532–1553.
Fontanier, C., et al., 2013. Live (stained) benthic foraminifera off Walvis Bay (Namibia): a deep-sea ecosystem under the influence of benthic nepheloid layer. J. Foraminifer. Res. 43, 50–66.
Forzoni, A., Storms, J.E.A., Reimann, T., Moreau, J., Jouet, G., 2015. Non-linear response of the Golo River system, Corsica, France, to late quaternary climatic and sea level variations. Quat. Sci. Rev. 121, 11–27.
Gasse, F., et al., 2014. Hydroclimatic changes in northern Levant over the past 400,000 years. Quat. Sci. Rev. 111, 1–8.
Goineau, A., et al., 2012. Temporal variability of live (stained) benthic foraminiferal faunas in a river-dominated shelf-faunal response to rapid changes of the river influence (Rhône prodelta, NW Mediterranean). Biogeosciences 9, 1367–1388.
Gooday, A.J., 2003. Benthic foraminifera (protista) as tools in deep-water palaeoceanography: environmental influences on faunal characteristics. Adv. Mar. Biol. 46, 1–90.
Göktürk, O.M., Fleitmann, D., Badertscher, S., Cheng, H., Edwards, R.L., Leuenberger, M., Fankhauser, A., Tüysüz, O., Kramers, J., 2011. Climate on the southern Black Sea coast during the Holocene: implications from the Sofular Cave record. Quat. Sci. Rev. 30 (19), 2433–2445.
Grant, K.M., Rohling, E.J., Bar Matthews, M., Ayalon, A., Medina Elizalde, M., Bronk Ramsey, C., Satow, C., Roberts, A.P., 2012. Rapid coupling between ice volume and polar temperature over the past 150,000 years. Nature 491, 744–747.
Grant, K.M., et al., 2014. Sea-level variability over five glacial cycles. Nat. Commun. 5, 5076.
Griveaud, C., 2007. Influence des conditions écologiques sur la composition isotopique (δ^{13}C, δ^{18}O) du test de foraminifères benthiques actuels (Unpublished PhD thesis). Université d'Angers, p. 326.
Hayes, A., Kucera, M., Kallel, N., Sbaffi, L., Rohling, E.J., 2005. Glacial Mediterranean Sea surface temperatures based on planktonic foraminiferal assemblages. Quat. Sci. Rev. 24, 999–1016.
Helmke, J.P., Bauch, H.A., 2001. Glacial-interglacial relationship between carbonate components and sediment reflectance in the North Atlantic. Geo-Marine Lett. 21 (1), 16–22.
Hilgen, F.J., 1991. Astronomical calibration of Gauss to Matuyama sapropels in the Mediterranean and implication for the geomagnetic polarity time scale. Earth Planet. Sci. Lett. 104, 226–244.
Hodell, D.A., Crowhurst, S., Skinner, L.C., Tzedakis, P.C., Margari, V., Channell, J.E.T., Kamenov, G., Maclachlan, S., Rothwell, R.G., 2013. Response of Iberian Margin sediments to orbital and suborbital forcing over the past 420 ka. Paleoceanography 28, 185–199. http://dx.doi.org/10.1002/paleo.20017.
Hoogakker, B.A.A., Rothwell, R.G., Rohling, E.J., Paterne, M., Stow, D.A.V., Herrle, J.O., Clayton, T., 2004. Variations in terrigenous dilution in western Mediterraean Sea pelagic sediments in response to climate change during the last glacial

cycle. Mar. Geol. 211, 21–43.
Janowiak, J.E., 1988. An investigation of interannual rainfall variability in Africa. J. Clim. 1 (3), 240–255.
Jo, K.N., Woo, K.S., Yi, S., Yang, D.Y., Lim, H.S., Wang, Y., Chen, G., Edwards, R.L., 2014. Mid-latitude interhemispheric hydrologic seesaw over the past 550,000 years. Nature 508, 378–382.
Jorissen, F.J., Fontanier, C., Thomas, E., 2007. Chapter seven paleoceanographical proxies based on deep-sea benthic foraminiferal assemblage characteristics. Dev. Mar. Geol. 1, 263–325.
Jouet, G., Berné, S., Rabineau, M., Bassetti, M.A., Bernier, P., Dennielou, B., Sierro, F.J., Flores, F.J., Taviani, M., 2006. Shoreface migrations at the shelf edge and sea-level changes around the Last Glacial Maximum (Gulf of Lions, NW Mediterranean). Mar. Geol. 234 (1–4), 21–42.
Kallel, N., Duplessy, J.C., Labeyrie, L., Fontugne, M., Paterne, M., Montacer, M., 2000. Mediterranean pluvial periods and sapropel formation over the last 200 000 years. Palaeogeogr. Palaeoclimatol. Palaeoecol. 157, 45–58.
Kallel, N., Paterne, M., Duplessy, J.C., Vergnaud-Grazzini, C., Pujol, C., Labeyrie, L., Arnold, M., Fontugne, M., Pierre, C., 1997. Enhanced rainfall in the mediterranean region during the last sapropel event. Oceanol. Acta 20 (5), 697–712.
Konijnendijk, T.Y.M., Ziegler, M., Lourens, L.J., 2014. Chronological constraints on pleistocene sapropel depositions from high-resolution geochemical records of ODP Sites 967 and 968. Newsletters Stratigr. 47 (3), 263–282.
Kotthoff, U., Pross, J., Müller, U.C., Peyron, O., Schmiedl, G., Schulz, H., Bordon, A., 2008. Climate dynamics in the borderlands of the Aegean Sea during formation of sapropel S1 deduced from a marine pollen record. Quat. Sci. Rev. 27 (7), 832–845.
Kuhlemann, J., Rohling, E., Krumrei, I., Kubik, P.W., Ivy-Ochs, S., Kucera, M., 2008. Regional synthesis of Mediterranean atmospheric circulation during the Last Glacial Maximum. Science 321, 1338–1340.
Kutzbach, J.E., Chen, G., Cheng, H., Edwards, R.L., Liu, Z., 2014. Potential role of winter rainfall in explaining increased moisture in the Mediterranean and Middle East during periods of maximum orbitally-forced insolation seasonality. Clim. Dyn. 42 (3–4), 1079–1095.
Larrasoana, J.C., Roberts, A.P., Rohling, E.J., Winklhofer, M., Wehausen, R., 2003. Three million years of monsoon variability over the northern Sahara. Clim. Dyn. 21, 689–698. http://dx.doi.org/10.1007/s00382-003-0355-z.
Laskar, J., Robutel, P., Joutel, F., Gastineau, M., Correia, A.C.M., Levrard, B., 2004. A long-term numerical solution for the insolation quantities of the Earth. Astronomy Astrophysics 428 (1), 261–285.
Licari, L.N., Mackensen, A., 2005. Benthic foraminifera off West Africa (11N to 321S): do live assemblages from the topmost sediment reliably record environmental variability? Mar. Micropaleontol. 55, 205–233.
Lisiecki, L.E., Raymo, M.E., 2005. A pliocene-pleistocene stack of 57 globally distributed benthic $\delta^{18}O$ records. Paleoceanography 20, PA1003. http://dx.doi.org/10.1029/2004PA001071.
Lourens, L.J., Hilgen, F.J., Gudjonsson, L., Zachariasse, W.J., 1992. Late pliocene to early pleistocene astronomically forced sea surface productivity and temperature variations in the Mediterranean. Mar. Micropaleontol. 19, 49–78.
Magny, M., et al., 2013. North–south palaeohydrological contrasts in the central Mediterranean during the Holocene: tentative synthesis and working hypotheses. Clim. Past 9, 2043–2071. http://dx.doi.org/10.5194/cp-9-2043-2013.
Margari, V., Skinner, L.C., Hodell, D.A., Martrat, B., Toucanne, S., Grimalt, J.O., Gibbard, P.L., Lunkka, J.P., Tzedakis, C., 2014. Land-ocean changes on orbital and millenial time scales and the penultimate glaciation. Geology 42 (3), 183–186.
Martrat, B., et al., 2004. Abrupt temperature changes in the Western Mediterranean over the past 250,000 years. Science 306 (5702), 1762–1765.
Martrat, B., Grimalt, J.O., Shackleton, N.J., De Abreu, L., Hutterli, M.A., Stocker, T.F., 2007. Four climate cycles of recurring deep and surface water destabilizations on the Iberian margin. Science 317 (5837), 502–507.
Martrat, B., Jimenez-Amat, P., Zahn, R., Grimalt, J.O., 2014. Similarities and dissimilarities between the last two deglaciations and interglaciations in the North Atlantic region. Quat. Sci. Rev. 99, 122–134.
Marzin, C., Braconnot, P., 2009. Variations of Indian and African monsoons induced by insolation changes at 6 and 9.5 kyr BP. Clim. Dyn. 33, 215–231.
Meijer, P.T., Tuenter, E., 2007. The effect of precession-induced changes in the Mediterranean freshwater budget on circulation at shallow and intermediate depth. J. Mar. Syst. 68, 349–365.
Milner, A.M., Collier, R.E.L., Roucoux, K.H., Müller, U.C., Pross, J., Kalaitzidis, S., Christanis, K., Tzedakis, P.C., 2012. Enhanced seasonality of precipitation in the Mediterranean during the early part of the last interglacial. Geology 40, 919–922.
Mojtahid, M., Jorissen, F., Lansard, B., Fontanier, C., Bombled, B., Rabouille, C., 2009. Spatial distribution of live benthic foraminifera in the Rhône prodelta: faunal response to a continental-marine organic matter gradient. Mar. Micropaleontol. 70 (3–4), 177–200.
Murray, J.W., 2006. Ecology and Application of Benthic Foraminifera. Cambridge University Press, Cambridge, UK, p. 426.
Osborne, A.H., Vance, D., Rohling, E.J., Barton, N., Rogerson, M., Fello, N., 2008. A humid corridor across the Sahara for the migration of early modern humans out of Africa 120,000 years ago. Proc. Natl. Acad. Sci. 105 (43), 16444–16447.
Peyron, O., Goring, S., Dormoy, I., Kotthoff, U., Pross, J., de Beaulieu, J.L., Drescher-Schneider, R., Magny, M., 2011. Holocene seasonality changes in the central Mediterranean region reconstructed from the pollen sequences of Lake Accesa (Italy) and Tenaghi Philippon (Greece). Holocene 21, 131–146.
Pierre, C., 1999. The oxygen and carbon isotope distribution in the Mediterranean water masses. Mar. Geol. 153 (1–4), 41–45.
Pierre, C., Belanger, P., Saliège, J.F., Urrutiaguer, M.J., Murat, A., 1999. Paleoceanography of the western Mediterranean during the pleistocene: oxygen and carbon isotope records at site 975. In: Zahn, R., Comas, M.C., Klaus, A. (Eds.), Proceedings of the Ocean Drilling Program, p. 481.
Piva, A., Asioli, A., Andersen, N., Grimalt, J.O., Schneider, R., Trincardi, F., 2008. Climatic cycles as expressed in sediments of the PROMESS1 borehole PRAD1-2, central Adriatic, for the last 370 kyr: 2. Paleoenvironmental evolution. Geochem. Geophys. Geosystems 9 (3), Q03R02. http://dx.doi.org/10.1029/2007GC001785.
Rabineau, M., Berné, S., Aslanian, D., Olivet, J.L., Joseph, P., Guillocheau, F., Bourillet, J.F., Le Drezen, E., Granjeon, D., 2005. Sedimentary sequences in the Gulf of Lion: a record of 100,000 years climatic cycles. Mar. Pet. Geol. 22 (6), 775–804.
Railsback, L.B., Gibbard, P.L., Head, M.J., Voarintsoa, N.R.G., Toucanne, S., 2015. An optimized scheme of lettered marine isotope substages for the last 1.0 million years, and the climatostratigraphic nature of isotope stages and substages. Quat. Sci. Rev. 111 (1), 94–106.
Rasmussen, S.O., et al., 2006. A new Greenland ice core chronology for the last glacial termination. J. Geophys. Res. 111, D06102. http://dx.doi.org/10.1029/2005JD006079.
Reale, O., Feudale, L., Turato, B., 2001. Evaporative moisture sources during a sequence of floods in the Mediterranean region. Geophys. Res. Lett. 28 (10), 2085–2088.
Regattieri, E., Giaccio, B., Zanchetta, G., Drysdale, R.N., Galli, P., Nomade, S., Peronace, E., Wulf, S., 2015. Hydrological variability over the Apennines during the early last glacial precession minimum, as revealed by a stable isotope record from Sulmona basin, central Italy. J. Quat. Sci. 30 (1), 19–31.
Regattieri, E., Zanchetta, G., Drysdale, R.N., Isola, I., Hellstrom, J.C., Roncioni, A., 2014. A continuous stable isotope record from the penultimate glacial maximum to the last interglacial (159–121 ka) from Tana Che Urla Cave (Apuan Alps, central Italy). Quat. Res. 82 (2), 450–461.
Reimer, P.J., et al., 2013. Intcal13 and marine13 radiocarbon age calibration curves 0-50,000 years cal BP. Radiocarbon 55 (4), 1869–1887.
Revel, M., et al., 2010. 100,000 years of African monsoon variability recorded in sediments of the Nile margin. Quat. Sci. Rev. 29 (11–12), 1342–1362.
Richter, T.O., Van Der Gaast, S., Koster, B., Vaars, A., Gieles, R., De Stigter, H.C., De Haas, H., Van Weering, T.C.E., 2006. The Avaatech XRF core scanner: technical description and applications to NE Atlantic sediments. In: Rothwell, R.G. (Ed.), New Techniques in Sediment Core Analysis. Geological Society, Special Publications, pp. 39–50.
Ridente, D., Trincardi, F., Piva, A., Asioli, A., 2009. The combined effect of sea level and supply during Milankovitch cyclicity: evidence from shallow-marine $\delta^{18}O$ records and sequence architecture (Adriatic margin). Geology 37, 1003–1006.
Roberts, D.H., Chiverrell, R.C., Innes, J.B., Horton, B.P., Brooks, A.J., Thomas, G.S.P., Turner, S., Gonzalez, S., 2006. Holocene sea levels, last glacial maximum glaciomarine environments and geophysical models in the northern Irish Sea Basin, UK. Mar. Geol. 231 (1–4), 113–128.
Rodwell, M.J., Hoskins, B.J., 1996. Monsoons and the dynamics of deserts. Q. J. R. Meteorol. Soc. 122, 1385–1404.
Rodwell, M.J., Hoskins, B.J., 2001. Subtropical anticyclones and summer monsoons. J. Clim. 14, 3192–3211.
Rohling, E.J., 1994. Review and new aspects concerning the formation of Mediterranean sapropels. Mar. Geol. 122, 1–28.
Rohling, E.J., et al., 2002. African monsoon variability during the previous interglacial maximum. Earth Planet. Sci. Lett. 202, 61–75.
Rohling, E.J., Foster, G.L., Grant, K.M., Marino, G., Roberts, A.P., Tamisiea, M.E., Williams, F., 2014. Sea-level and deep-sea-temperature variability over the past 5.3 million years. Nature 508, 477–482.
Rohling, E.J., Hilgen, F.J., 1991. The eastern Mediterranean climate at times of sapropel formation: a review. Geol. Mijnb. 70, 253–264.
Rohling, E.J., Marino, G., Grant, K.M., 2015. Mediterranean climate and oceanography, and the periodic development of anocix events (sapropels). Earth Sci. Rev. 143, 62–97.
Rossignol-Strick, M., 1983. African monsoons, an immediate climate response to orbital insolation. Nature 30, 446–449.
Rossignol-Strick, M., 1985. Mediterranean quaternary sapropels, an immediate response of the african monsoon to variation of insolation. Palaeogeogr. Palaeoclimatol. Palaeoecol. 49, 237–263.
Rossignol-Strick, M., 1987. Rainy periods and bottom water stagnation initiating brine accumulation metal concentrations: 1. the late quaternary. Paleoceanography 2, 333–360.
Rossignol-Strick, M., Nesteroff, W., Olive, P., Vergnaud-Grazzini, C., 1982. After the deluge, Mediterranean stagnation and sapropel formation. Nature 295, 105–110.
Roucoux, K.H., Tzedakis, P.C., de Abreu, L., Shackleton, N.J., 2006. Climate and vegetation changes 180,000 to 345,000 years ago recorded in a deep-sea core off Portugal. Earth Planet. Sci. Lett. 249 (3), 307–325.
Ruddiman, W.F., 1971. Pleistocene sedimentation in the equatorial Atlantic: stratigraphy and faunal paleoclimatology. Geol. Soc. Am. Bull. 82 (2), 283–302.
Ryan, W.B.F., 1972. Stratigraphy of late quaternary sediments in the eastern Mediterranean. In: Stanley, D.J. (Ed.), The Mediterranean Sea: a Natural Sedimentation Laboratory, pp. 149–169 (Stroudsburg, Pennsylvania, Dowden, Hutchinson and Ross).
Sánchez Goñi, M.F., Eynaud, F., Turon, J.L., Shackleton, N.J., 1999. High resolution

palynological record off the Iberian margin: direct land-sea correlation for the last interglacial complex. Earth Planet. Sci. Lett. 171 (1), 123–137.
Sánchez Goñi, M.F., Landais, A., Fletcher, W.J., Naughton, F., Desprat, S., Duprat, J., 2008. Contrasting impacts of Dansgaard-Oeschger events over a western European latitudinal transect modulated by orbital parameters. Quat. Sci. Rev. 27 (11–12), 1136–1151.
Schmiedl, G., Pfeilsticker, M., Hemleben, C., Mackensen, A., 2004. Environmental and biological effects on the stable isotope composition of recent deep-sea benthic foraminifera from the western Mediterranean Sea. Mar. Micropaleontol. 51 (1–2), 129–152.
Scrivner, A.E., Vance, D., Rohling, E.J., 2004. New neodymium isotope data quantify Nile involvement in Mediterranean anoxic eposides. Geology 32 (7), 565–568.
Shackleton, N.J., Hall, M.A., Vincent, E., 2000. Phase relationships between millennial scale events 64,000 to 24,000 years ago. Paleoceanography 15, 565–569.
Shaw, H.F., Evans, G., 1984. The nature, distribution and origin of a sapropelic layer in sediments of the Cilicia Basin, northeastern Mediterranean. Mar. Geol. 61, 1–12.
Siani, G., Paterne, M., Michel, E., Sulpizio, R., Sbrana, A., Arnold, M., Haddad, G., 2001. Mediterranean sea surface radiocarbon reservoir age changes since the last glacial maximum. Science 294, 1917–1920.
Sierro, F.J., Ledesma, S., Flores, F.J., Torrescusa, S., del Olmo, W.M., 2000. Sonic and gamma-ray astrochronology: cycle to cycle calibration of Atlantic climatic records to Mediterranean sapropels and astronomical oscillations. Geology 28 (8), 695–698.
Sierro, F.J., et al., 2009. Phase relationship between sea level and abrupt climate changes. Quat. Sci. Rev. 28 (25–26), 2867–2881.
Sømme, T.O., Piper, D.J., Deptuck, M.E., Helland-Hansen, W., 2011. Linking onshore-offshore sediment dispersal in the Golo source-to-sink system (Corsica, France) during the late quaternary. J. Sediment. Res. 81 (2), 118–137.
Stuiver, M., Reimer, P.J., 1993. Extended ^{14}C data-base and revised Calib 3.0 ^{14}C age calibration program. Radiocarbon 35, 215–230.
Svensson, A., et al., 2008. A 60,000 year Greenland stratigraphic ice core chronology. Clim. Past 4, 47–57.
Thomson, J., Nixon, S., Summerhayes, C.P., Schonfeld, J., Zahn, R., Grootes, P., 1999. Implications for sedimentation changes on the Iberian margin over the last two glacial/interglacial transitions from (^{230}Th excess) systematics. Earth Planet. Sci. Lett. 165 (3–4), 255–270.
Toucanne, S., et al., 2009. Timing of massive 'Fleuve Manche' discharges over the last 350 kyr: insights into the European Ice Sheet oscillations and the European drainage network from MIS 10 to 2. Quat. Sci. Rev. 28 (13–14), 1238–1256.
Toucanne, S., et al., 2012. A 130,000-year record of levantine intermediate water flow variability in the Corsica Trough, western Mediterranean Sea. Quat. Sci. Rev. 33, 55–73.
Trauth, M.H., Larrasoaña, J.C., Mudelsee, M., 2009. Trends, rhythms and events in plio-pleistocene African climate. Quat. Sci. Rev. 28, 399–411.
Trigo, I.F., Bigg, G.R., Davies, T.D., 2002. Climatology of cyclogenesis mechanisms in the Mediterranean. Mon. Weather Rev. 130 (3), 549–569.
Trigo, I.F., Davies, T.D., Bigg, G.R., 1999. Objective climatology in the Mediterranean region. J. Clim. 12, 1685–1696.
Tuenter, E., Weber, S.L., Hilgen, F.J., Lourens, L.J., 2003. The response of the African summer monsoon to remote and local forcing due to precession and obliquity. Glob. Planet. Change 36, 219–235.
Tzedakis, P.C., 2007. Seven ambiguities in the Mediterranean palaeoenvironmental narrative. Quat. Sci. Rev. 26, 2042–2066.
Tzedakis, P.C., Hooghiemstra, H., Pälike, H., 2006. The last 1.35 million years at Tenaghi Philippon: revised chronostratigraphy and long-term vegetation trends. Quat. Sci. Rev. 25 (23–24), 3416–3430.
Tzedakis, P.C., McManus, J.F., Hooghiemstra, H., Oppo, D.W., Wijmstra, T.A., 2003. Comparison of changes in vegetation in northeast Greece with records of climate variability on orbital and suborbital frequencies over the last 450 000 years. Earth Planet. Sci. Lett. 212 (1–2), 197–212.
Tzedakis, P.C., Pälike, H., Roucoux, K.H., De Abreu, L., 2009. Atmospheric methane, southern European vegetation and low-mid latitude links on orbital and millenial timescales. Earth Planet. Sci. Lett. 277, 307–317.
Valero, L., Garcès, M., Cabrera, L., Costa, E., Saez, A., 2014. 20 Myr of eccentricity paced lacustrine cycles in the Cenozoic Ebro Basin. Earth Planet. Sci. Lett. 408, 183–193.
Van Vugt, N., Langereis, C.G., Hilgen, F.J., 2001. Orbital forcing in pliocene-pleistocene Mediterranean lacustrine deposits: dominant expression of eccentricity versus precession. Palaeogeogr. Palaeoclimatol. Palaeoecol. 172, 193–205.
Vergnaud-Grazzini, C., Saliège, J.F., Urrutiager, M.J., Iannace, A., 1990. Oxygen and carbon isotope stratigraphy of ODP Hole 653A and site 654: the pliocene-pleistocene glacial history recorded in the Tyrrhenian Basin (West Mediterranean). In: Kastens, K.A., Mascle, J., et al. (Eds.), Proceedings of the Ocean Drilling Program, Scientific Results. Tyrrhenian Sea, vol. 107, pp. 361–386 (Ocean Drilling Program, College Station, TX).
Waelbroeck, C., Labeyrie, L., Michel, E., Duplessy, J.C., McManus, J.F., Lambeck, K., Balbon, E., Labracherie, M., 2002. Sea-level and deep water temperature changes derived from benthic foraminifera isotopic records. Quat. Sci. Rev. 21 (1–3), 295–305.
Wang, Y., et al., 2008. Millennial- and orbital-scale changes in the East Asian monsoon over the past 224,000 years. Nature 451 (7182), 1090–1093.
Weltje, G.J., Tjallingii, R., 2008. Calibration of XRF core scanners for quantitative geochemical logging of sediment cores: theory and application. Earth Planet. Sci. Lett. 274, 423–438.
Wijmstra, T.A., Young, R., Witte, H.J.L., 1990. An evaluation of the climatic conditions during the late quaternary in northern Greece by means of multivariate analysis of palynological data and comparison with recent phytosociological and climatic data. Geol. Mijnb. 69, 243–251.
Williams, D.F., Thunell, R.C., Kennett, J.P., 1978. Periodic freshwater flooding and stagnation of the eastern Mediterranean Sea during the late quaternary. Science 201, 252–254.
Xoplaki, E., Gonzalez-Rouco, F.J., Luterbacher, J., Wanner, H., 2003. Mediterranean summer air temperature variability and its connection to the large-scale atmospheric circulation and SSTs. Clim. Dyn. 20, 723–739.
Zanchetta, G., Drysdale, R.N., Hellstrom, J.C., Fallick, A.E., Isola, I., Gagan, M.K., Pareschi, M.T., 2007. Enhanced rainfall in the Western Mediterranean during deposition of sapropel S1: stalagmite evidence from Corchia cave (Central Italy). Quat. Sci. Rev. 26 (3–4), 279–286.
Ziegler, M., Tuenter, E., Lourens, L.J., 2010. The precession phase of the boreal summer monsoon as viewed from the eastern Mediterranean (ODP Site 968). Quat. Sci. Rev. 29, 1481–1490.