



# Warfarin treatment and risk of myocardial infarction — A cohort study of patients with atrial fibrillation treated in primary health care

Per Wändell <sup>a,b,\*</sup>, Axel C. Carlsson <sup>a,c</sup>, Martin J. Holzmänn <sup>d,e</sup>, Johan Ärnlöv <sup>c,f</sup>, Sven-Erik Johansson <sup>g</sup>, Jan Sundquist <sup>g</sup>, Kristina Sundquist <sup>g</sup>

<sup>a</sup> Division of Family Medicine, Department of Neurobiology, Care Science and Society, Karolinska Institutet, Huddinge, Sweden

<sup>b</sup> Academic Primary Healthcare Centre, Stockholm County Council, Huddinge, Sweden

<sup>c</sup> Department of Medical Sciences, Cardiovascular Epidemiology, Uppsala University, Uppsala, Sweden

<sup>d</sup> Department of Emergency Medicine, Karolinska University Hospital, Stockholm, Sweden

<sup>e</sup> Department of Internal Medicine, Karolinska Institutet, Stockholm, Sweden

<sup>f</sup> School of Health and Social Studies, Dalarna University, Falun, Sweden

<sup>g</sup> Centre for Primary Health Care Research, Lund University, Malmö, Sweden

## ARTICLE INFO

### Article history:

Received 12 May 2016

Accepted 8 July 2016

Available online 9 July 2016

### Keywords:

Atrial fibrillation  
Myocardial infarction  
Gender  
Follow-up  
Co-morbidity  
Anticoagulants  
Antiplatelets

## ABSTRACT

**Objective:** To study the risk of myocardial infarction (MI) in patients with atrial fibrillation (AF) treated in primary health care with warfarin or acetylsalicylic acid (ASA, aspirin).

**Methods:** The study population included subjects ( $n = 12,283$ ) 45 years or older diagnosed with AF who were treated in 75 primary care centres in Sweden between 2001 and 2007. MI was defined as a hospital stay for MI during 2001 through 2010 registered in the Swedish Patient Register. Associations between warfarin or ASA treatment and incident MI were explored using Cox regression analysis, by estimating hazard ratios (HRs) and 95% confidence intervals (95% CIs). Adjustment was made for age, socio-economic factors and cardio-vascular co-morbidity. **Results:** Persistent treatment ("per protocol" treatment) with warfarin alone was present among 32.4% of women and 37.4% of men, and with ASA alone among 30.0% of women and 28.1% of men. The fully adjusted HRs for MI, compared to those with no antithrombotic treatment, with warfarin treatment for women were 0.26 (95% CI 0.16–0.41) and for men 0.28 (95% CI 0.20–0.39); and the corresponding HRs for those treated with ASA were for women 0.57 (95% CI 0.37–0.87), and for men 0.44 (95% CI 0.31–0.63). The fully adjusted HR for MI when comparing patients with warfarin treatment to those with ASA treatment was for women 0.46 (95% CI 0.27–0.80), and for men 0.58 (95% CI 0.38–0.89).

**Conclusions:** Warfarin seems to prevent MI among AF patients in a primary healthcare setting, which emphasizes the importance of persistent anticoagulant treatment in those patients.

© 2016 Elsevier Ireland Ltd. All rights reserved.

## Key messages

### What is already known about this subject?

In patients with atrial fibrillation there is an uncertainty whether anticoagulants are superior to antiplatelets in preventing myocardial infarction.

### What does this study add?

We found anticoagulants to be superior to antiplatelets among both men and women with atrial fibrillation in preventing myocardial infarction.

### How might this impact on clinical practice?

Persistent anticoagulant treatment is important among patients with atrial fibrillation not only in preventing stroke but also in preventing myocardial infarction.

## 1. Introduction

Atrial fibrillation (AF) is considered as a major health problem [1], with an increasing trend of incidence and prevalence globally [2–4]. In Sweden, around 2% of the population are diagnosed with AF [5]. The most important complication of AF is ischemic stroke [6,7], estimated to be 5 times as common as in individuals without AF [8].

Myocardial infarction (MI) is a risk factor for AF, and incident AF is present in 6–21% among patients with an acute MI [9]. Considering the reverse causation, AF could be associated with an increased risk of MI, and a higher risk of MI among women with AF has indeed been

\* Corresponding author at: Division of Family Medicine, NVS Department, Karolinska Institutet, Alfred Nobels Allé 12, 141 83 Huddinge, Sweden.  
E-mail address: [per.wandell@ki.se](mailto:per.wandell@ki.se) (P. Wändell).

reported [10]. Interestingly, the mortality risk estimates were higher although not statistically significant in an American study comparing patients with AF and MI with patients with AF alone [10].

Among prescribed pharmacotherapies to AF patients, anticoagulant (predominantly warfarin) therapy has benefits over antiplatelet (mostly acetylsalicylic acid, i.e. ASA, aspirin) therapy [11]. This is because anticoagulant treatment is superior in preventing strokes [12]. Before the introduction of the non-vitamin K antagonist oral anticoagulants (NOACs), warfarin was the most commonly prescribed oral anticoagulant used to prevent stroke in patients with AF [13]. With regard to the risk of MI in patients with AF, a Cochrane report found a reduced, although not significantly reduced risk in patients with oral anticoagulant compared to antiplatelet therapy; the OR was 0.69, 95% CI 0.47 to 1.01, while the mortality risk was similar for oral anticoagulants and antiplatelets [14]. A review of stroke-preventive studies regarding the effect of warfarin compared to non-warfarin anticoagulants found a significant reduction in MI [15].

Many studies on AF patients are based on samples from hospitals although many patients with AF are cared for at their primary health care centres. Out of all patients recorded with an AF diagnosis in Stockholm County in Sweden, 64% had their AF diagnosis reported in the primary healthcare records [5]. Thus, it is important to study the risks and benefits of different therapies prescribed to patients with AF within this setting.

The objective of the present study was to explore the risk of MI associated with warfarin and ASA treatment compared with no antithrombotic treatment in women and men with AF in a large cohort treated in primary health care. Secondary aims were to explore the mortality risk in women and men with AF who experienced a MI with women and men who did not experience a MI, and to explore mortality among patients with MI in relation to treatment with warfarin or ASA.

## 2. Methods

### 2.1. Design

This study was performed using individual-level patient data from 75 Swedish primary health care centres (PHCCs). The majority of the centres were located in Stockholm County ( $n = 48$ ). Men and women visiting any of the participating PHCCs between 2001 and 2007 were included in the study. We used Extractor software ([http://www.sll.se/SLPOtemplates/SLPOPPage1\\_10400.aspx](http://www.sll.se/SLPOtemplates/SLPOPPage1_10400.aspx); accessed 19 September 2010) to collect individual files from the electronic patient records (EPR) at the PHCCs. Individual identification numbers were replaced by serial numbers to ensure anonymity. The EPR files from the PHCCs were linked to Swedish national registers [16]. The registers used were: The Total Population register (which contains data on, e.g., age and education); The Inpatient Register (hospital admissions); and the Cause of Death Register. These registers contain individual-level population data for all residents registered in Sweden. Thus, a new research database was created, containing individual clinical patient data from a total of 1,098,420 subjects registered at these 75 PHCCs, linked to national demographic and socio-economic data. A follow-up was performed using the Swedish Cause of Death Register, which has been shown to be almost complete, 99.8%, and lacking data only for a few emigrants from Sweden to other countries and thus lost to follow-up [17].

### 2.2. Study population and co-morbidities

The study included all patients with diagnosed AF, identified by the presence of the ICD-10 code (10th version of the WHO International Classification of Diseases) for atrial fibrillation (I48) in patients' medical records. The following related cardiovascular disorders/co-morbidities were used as covariates: hypertension (I10–I15), heart failure (CHF; I50 and I110), cerebrovascular diseases (CVD; I60–I69), and diabetes mellitus (E10–E14). Presence of coronary heart disease (CHD; I20–I25) was noted, but we did not have access to reliable data as regards earlier MI before first AF diagnosis. In total, 6646 men and 5637 women who were aged 45 years or older at the time of AF diagnosis and who visited any of the 75 participating PHCCs from 1 January 2001 until 31 December 2007, and had data on neighbourhood socio-economic status, were included in the study [18].

### 2.3. Outcome variable

Time to first MI after registration of AF diagnosis during the assessment period until 31 December 2010, was defined as having an ICD-10 code indicating an acute myocardial infarction (I21) in the Patient Register (hospital admissions) or The Cause of Death Register [19].

In addition, time to mortality from first AF diagnosis to death was registered as a secondary outcome.

### 2.4. Demographic and socio-economic variables

Sex: Men and women.

Age was categorized as follows: 45–54, 55–64, 65–74, 75–84 and  $\geq 85$  years. Individuals younger than 45 years were excluded (AF was rare in individuals below 45 years of age, who are not representative of AF patients in general).

The neighbourhood socio-economic status (SES) areas were categorized into three groups according to the neighbourhood index: more than one standard deviation (SD) below the mean (high SES or low deprivation level), more than one SD above the mean (low SES or high deprivation level), and within one SD of the mean (middle SES or deprivation level) [20]. The neighbourhood summary index was based on information about female and male residents aged 20 to 64 years because this age group represents those who are among the most socio-economically active in the population. The index was based on the following four variables: low educational status ( $< 10$  years of formal education); income from all sources, including interest and dividends, that is  $< 50\%$  of the median individual income; unemployment (excluding full-time students, those completing military service, and early retirees); and receipt of social welfare. We also registered change of neighbourhood SES during the study period, i.e. until 31 December 2007.

Educational attainment was categorized as  $\leq 9$  years (partial or complete compulsory schooling), 10–12 years (partial or complete secondary schooling) and  $> 12$  years (attendance at college and/or university).

Marital status was characterized as married, unmarried, divorced or widowed.

### 2.5. Antithrombotic treatment

Treatment with an antithrombotic drug was defined as a prescription noted in the electronic patient record in primary health care 2001 to 2007. The prescribed antithrombotic drugs were classified as "intention-to-treat" ("ITT") if ever present before the years of the first MI, or if present at any time among subjects not experiencing a MI. The prescribed warfarin was classified as "per-protocol" ("PP") if present the year before and the year of first MI, or present among subjects not experiencing a MI if present at least 50% of actual years after first recorded year of AF. Prescriptions of antithrombotic treatment were classified into anticoagulant treatment, i.e. of warfarin (B01AA03), and of antiplatelet agents (B01AC). Antiplatelet agents were classified into ASA (B01AC06, and ASA combined with dipyridamole, B01AC30), or clopidogrel (B01AC04), also including related drugs (ticlopidine, B01AC05), with only marginal prescription.

### 2.6. Statistical analysis

Baseline characteristics for all included men and women, as well as for those with a recorded MI, were presented as mean (SD) if continuous, and as frequencies if categorical.

We also made stratified analyses in subjects classified as not having a "per-protocol" prescription of antithrombotic drugs. Thus, we were able to estimate the risk of MI when not being on antithrombotic treatment.

We also estimated the incidence rates of MI per 100 person-years at risk for men and women. As a sensitivity analysis we assessed the incidence rate for MI for subjects with no antithrombotic treatment. The age-adjusted relative risk of MI for patients on "ITT" and "PP" warfarin or ASA treatment was analysed using Cox proportional hazard regression analysis, and presented as hazard ratios (HR) with 95% confidence intervals (CI). Adjustments were also made for socio-economic factors (educational level, marital status and neighbourhood SES), and also for co-morbidity (hypertension, CHF, CVS and diabetes). Models were checked for interactions by the Breslow method for ties. When interaction was present, interaction terms were used. Model specification was also tested.

In secondary analysis, Cox regression was used for estimating mortality risk in patients with MI, with patients without a MI as referents, with HRs and 95% CI, for men and women separately, with adjustments for age and sex, for socio-economic factors and for co-morbidity. Furthermore, Cox regression was used for estimating mortality risk in patients with MI in relation to antithrombotic treatment, with adjustment as stated above.

The study was approved by the regional ethics boards at Karolinska Institutet and Lund University.

## 3. Results

The characteristics of the men and women with AF treated in primary care without ( $n = 11,283$ ) or with a MI during follow-up ( $n = 1000$ ) are shown in Table 1. The mean follow-up time was 5.6 years (SD 2.5), and in total the analyses included 68,172 patient years.

Antithrombotic treatment by different combinations of drugs according to per-protocol analysis is shown in Table 2. Only a few patients had treatment with two drugs, the most common being warfarin and ASA (3.5% among women, and 4.6% among men).

Incidence rates of first MI by sex are shown in Table 3. Women showed a non-significantly lower risk of a MI than men, fully adjusted

**Table 1**

Data on subjects aged 45 + years (n = 12,283) with a diagnosis of atrial fibrillation and without (n = 12,121) or with (n = 162) myocardial infarction (MI) after AF diagnosis in primary care from 1 January 2001 to 31 December 2007.

	Women		p-Value	Men		p-Value
	Without MI n = 5165	With MI n = 472		Without MI n = 6118	With MI n = 528	
Age (years), mean (SD)	<b>76.9 (9.4)</b>	<b>79.8 (8.0)</b>	<b>&lt;0.001</b>	<b>71.8 (10.2)</b>	<b>75.6 (8.8)</b>	<b>&lt;0.001</b>
Age group (years)			<b>&lt;0.001</b>			<b>&lt;0.001</b>
45–54	<b>104 (2.0)</b>	<b>1 (0.2)</b>		<b>356 (5.8)</b>	<b>14 (2.7)</b>	
55–64	<b>498 (9.6)</b>	<b>23 (4.9)</b>		<b>1167 (19.1)</b>	<b>55 (10.4)</b>	
65–74	<b>1191 (23.1)</b>	<b>75 (15.9)</b>		<b>1910 (31.2)</b>	<b>132 (25.0)</b>	
75–84	<b>2288 (44.3)</b>	<b>246 (52.2)</b>		<b>2079 (34.0)</b>	<b>261 (49.4)</b>	
85 +	<b>1084 (21.0)</b>	<b>127 (26.9)</b>		<b>606 (9.9)</b>	<b>66 (12.5)</b>	
Neighbourhood SES			0.052			0.028
High	1802 (34.9)	146 (30.9)		2456 (40.1)	200 (37.9)	
Middle	2544 (49.3)	233 (49.4)		2799 (45.8)	231 (43.8)	
Low	819 (15.9)	93 (19.7)		863 (14.1)	97 (18.4)	
Marital status			<b>0.003</b>			<b>&lt;0.001</b>
Married	<b>1548 (30.1)</b>	<b>115 (24.5)</b>		<b>3652 (59.9)</b>	<b>298 (56.7)</b>	
Unmarried	<b>373 (7.3)</b>	<b>26 (5.5)</b>		<b>590 (9.7)</b>	<b>40 (7.6)</b>	
Divorced	<b>732 (14.2)</b>	<b>60 (12.8)</b>		<b>947 (15.5)</b>	<b>74 (14.1)</b>	
Widowed	<b>2489 (48.4)</b>	<b>268 (57.1)</b>		<b>909 (14.9)</b>	<b>114 (21.7)</b>	
Educational level			<b>&lt;0.001</b>			<b>&lt;0.001</b>
Compulsory school	<b>2363 (51.9)</b>	<b>236 (59.8)</b>		<b>2248 (38.8)</b>	<b>238 (48.2)</b>	
Secondary school	<b>1502 (33.0)</b>	<b>126 (31.9)</b>		<b>2194 (37.9)</b>	<b>173 (35.0)</b>	
College/university	<b>691 (15.2)</b>	<b>33 (8.3)</b>		<b>1354 (23.4)</b>	<b>83 (16.8)</b>	
AF-related disease						
Hypertension	2456 (49.3)	241 (51.1)	0.46	2558 (41.8)	241 (45.6)	0.087
CHD	<b>966 (18.7)</b>	<b>210 (44.5)</b>	<b>&lt;0.001</b>	<b>1083 (17.7)</b>	<b>255 (48.3)</b>	<b>&lt;0.001</b>
Heart failure	1017 (20.1)	116 (24.6)	0.020	<b>1022 (16.7)</b>	<b>133 (25.2)</b>	<b>&lt;0.001</b>
Valvular disease	244 (4.7)	33 (7.0)	0.029	269 (4.7)	25 (4.7)	0.72
Cardiomyopathy	29 (0.6)	1 (0.2)	0.32	55 (0.9)	5 (1.0)	0.91
CVS	621 (12.0)	48 (10.2)	0.23	693 (11.3)	49 (9.3)	0.15
First ischemic stroke	722 (14.0)	46 (9.8)	0.10	665 (10.9)	32 (6.1)	<b>0.001</b>
First intracranial haemorrhage	66 (1.3)	1 (0.2)	0.042	93 (1.5)	2 (0.4)	0.033
Diabetes mellitus	<b>973 (18.8)</b>	<b>120 (25.4)</b>	<b>0.001</b>	<b>1178 (19.3)</b>	<b>134 (25.4)</b>	<b>0.001</b>
Depression	565 (10.9)	62 (13.1)	0.15	377 (6.2)	35 (6.6)	0.67
Anxiety	286 (5.5)	27 (5.7)	0.87	166 (2.7)	17 (3.2)	0.50
Drugs						
Ever warfarin	<b>2504 (48.5)</b>	<b>187 (39.6)</b>	<b>&lt;0.001</b>	3444 (56.3)	272 (51.5)	0.034
Warfarin ITT	<b>2410 (46.7)</b>	<b>158 (33.5)</b>	<b>&lt;0.001</b>	<b>3320 (54.3)</b>	<b>241 (45.6)</b>	<b>&lt;0.001</b>
Warfarin PP	<b>1760 (34.1)</b>	<b>67 (14.2)</b>	<b>&lt;0.001</b>	<b>2389 (39.1)</b>	<b>97 (18.4)</b>	<b>&lt;0.001</b>
Ever ASA	<b>2961 (57.3)</b>	<b>330 (69.9)</b>	<b>&lt;0.001</b>	<b>3145 (51.4)</b>	<b>352 (66.7)</b>	<b>&lt;0.001</b>
ASA ITT	<b>2593 (50.2)</b>	<b>296 (62.7)</b>	<b>&lt;0.001</b>	<b>2761 (45.1)</b>	<b>299 (56.6)</b>	<b>&lt;0.001</b>
ASA PP	1568 (30.4)	125 (26.5)	0.079	1742 (28.5)	124 (23.5)	0.014
Ever clopidogrel	<b>156 (3.0)</b>	<b>40 (8.5)</b>	<b>&lt;0.001</b>	<b>148 (2.4)</b>	<b>59 (11.2)</b>	<b>&lt;0.001</b>
Clopidogrel ITT	108 (2.1)	8 (1.7)	0.56	107 (1.8)	10 (1.9)	0.81
Clopidogrel PP	52 (1.1)	4 (0.9)	1.0	46 (0.8)	6 (1.1)	0.34

Prescription of warfarin was classified as “intention-to-treat” (“ITT”) if ever present before the years of MI, or present among subjects without MI. Prescription of warfarin was classified as “per-protocol” (“PP”) if present the year before and the year of first MI, or present among subjects without MI if present during at least three years, at least 50% of actual years after first recorded year of AF, or during both 2006 and 2007. Significant differences (on the level  $p < 0.01$ ) are shown in bold.

HR 0.88 (95% CI 0.71–1.10). The fully adjusted HRs for MI, compared to those with no antithrombotic treatment, were with PP warfarin treatment for women 0.26 (95% CI 0.16–0.41), and for men 0.28 (95% CI

0.20–0.39); and the corresponding HRs were for those treated with PP ASA for women 0.57 (95% CI 0.37–0.87), and for men 0.44 (95% CI 0.31–0.63). In the comparison between PP warfarin and PP ASA treatment, the fully adjusted HR was for women 0.46 (95% CI 0.27–0.80), and for men 0.58 (95% CI 0.38–0.89). The number needed to treat (NNT) for PP warfarin treatment to prevent one MI was 58.7 per year in women and 62.4 per year in men. The NNT for treatment with PP ASA to prevent one MI was 140 per year in women and 178 per year in men.

Mortality risks for women and men with MI or without MI were estimated (Supplementary Table 1), with incidence rates per 100 person-years at risk for women with MI 9.584 (95% CI 8.481–10.830), and women without MI 5.758 (95% CI 5.492–6.037); and for men with MI 9.198 (95% CI 8.176–10.347), and men without MI 4.720 (95% CI 4.502–4.950). In fully adjusted Cox regression models, women with MI vs. those without MI had a higher risk, HR 1.38 (95% CI 1.18–1.61), as men with MI vs. those without MI, HR 1.52 (95% CI 1.33–1.74). Women had a lower mortality risk than men after MI, fully adjusted HR 0.79 (95% CI 0.64–0.97).

When estimating the mortality risk among patients with PP warfarin, PP ASA or PP any antithrombotic treatment was non-significant in

**Table 2**

Data on antithrombotic treatment (according to “PP”) in subjects aged 45 + years (n = 12,283) with a diagnosis of atrial fibrillation. Number of patients (percentage).

Treatment	Women n = 5637	Men n = 6646
No treatment	2281 (40.5)	2581 (38.8)
ASA	1480 (26.3)	1540 (23.2)
Clopidogrel	37 (0.7)	23 (0.4)
ASA + clopidogrel	12 (0.2)	16 (0.2)
Warfarin	1621 (28.8)	2167 (32.6)
Warfarin + ASA	199 (3.5)	306 (4.6)
Warfarin + clopidogrel	5 (0.1)	9 (0.1)
Warfarin + ASA + clopidogrel	2 (0.0)	4 (0.1)

Prescription was classified as “per-protocol” (“PP”) if present the year before and the year of MI, or present among subjects without MI if present during at least three years, at least 50% of actual years after first recorded year of AF, or during both 2006 and 2007.

**Table 3**  
Cox regression for risk of first myocardial infarction (MI) in women and men with atrial fibrillation treated with warfarin, using those without warfarin treatment as referents. Models shown by “per protocol” analysis (PP) for warfarin and for ASA treatment. Incidence rate per 100 person-years at risk.

	Events/at risk (n)	Incidence rate (95% CI)	Warfarin			ASA		
			Model 1	Model 2	Model 3	Model 1	Model 2	Model 3
MI								
Women	472/5637	1.530 (1.398–1.674)	0.26 (0.20–0.35)	0.27 (0.17–0.42)	0.26 (0.16–0.41)	0.54 (0.44–0.68)	0.61 (0.40–0.93)	0.57 (0.37–0.87)
Men	528/6646	1.415 (1.299–1.541)	0.29 (0.23–0.37)	0.30 (0.21–0.42)	0.28 (0.20–0.39)	0.48 (0.39–0.60)	0.47 (0.33–0.66)	0.44 (0.31–0.63)

Prescription of warfarin and ASA was classified as “per-protocol” (“PP”) if present the year before and the year of MI, or present among subjects without MI if present at least 50% of actual years after first recorded year of AF. For warfarin, patients on ASA or clopidogrel were excluded from analyses, and for ASA, patients on warfarin or clopidogrel were excluded.

Model 1 age-adjusted, and Model 2 as Model 1 but also adjusted for socio-economic factors (neighbourhood socio-economic status, educational level and marital status; for women also interaction terms between age and neighbourhood socio-economic status and marital status, respectively; for men also interaction terms between age and marital status), and Model 3 as Model 2 but also adjusted for cardiovascular co-morbidity (hypertension, cerebrovascular disease, CHF and diabetes; also including interaction terms between age and diabetes).

fully adjusted models (Supplementary Table 2), HRs 0.94 (95% CI 0.72–1.24), 0.88 (95% CI 0.69–1.14) and 0.89 (95% CI 0.73–1.09), respectively.

Risk of MI was also assessed in relation to scores on CHADS<sub>2</sub> and CHA<sub>2</sub>DS<sub>2</sub>-VAsC, with values for all patients and for those with no antithrombotic treatment according to “per protocol” analysis (Supplementary Table 3), and for all patients with no antithrombotic treatment (Supplementary Table 4). Incidence rates per 100 person-years at risk exceeded 2 for patients with no antithrombotic treatment, both in general and according to “per protocol” analysis, at CHADS<sub>2</sub> scores  $\geq 2$  for women, and  $\geq 3$  for men, and at CHA<sub>2</sub>DS<sub>2</sub>-VAsC scores  $\geq 4$  for women, and  $\geq 3$  for men.

#### 4. Discussion

The main finding was that warfarin treatment in patients with AF was associated with a lower risk of incident MI in comparison to patients with no antithrombotic treatment, as well as to patients with ASA treatment. The results remained significant after adjustments for socio-economic factors and relevant cardiovascular co-morbidities. The magnitude of the effect of warfarin treatment in preventing MI was remarkably high.

According to our findings it seems relevant to primarily use warfarin with regard to the risk of MI among AF patients, since the risk estimates compared to those without antiplatelet and antithrombotic treatment were higher than for warfarin. The use of warfarin is also in accord with current stroke prevention recommendations for patients with AF [11]. A similar suggestion was also proposed beyond the first year after an acute coronary syndrome in a recent systematic review [21].

Lower risk of MI was also found for patients treated with ASA, which has previously been shown to be effective in the prevention of MI [22,23], even if the effect seemed to lower among women than among men. Interestingly, some therapy recommendations suggest that both warfarin and ASA should be used simultaneously [24]. A large Danish study found no benefits of combining antiplatelet therapy with warfarin on recurrent coronary events or thromboembolism, but a significantly increased risk of bleeding complications [25]. It is also suggested that the risks associated with the combination of ASA with anticoagulation in patients with AF outweigh the benefit [26].

We have previously shown that warfarin is more often prescribed to women and men living in high socio-economic neighbourhoods [20]. Moreover, both individual-level and neighbourhood-level socio-economic status may affect the prescribing of warfarin as well as the mortality rate [27]. It is thus possible that the lower MI risk seen among those prescribed warfarin can be explained by factors associated with having warfarin prescribed, rather than the warfarin itself. Yet the association found between warfarin treatment and reduced risk of MI remained significant when we adjusted for several factors including education level, marital status and neighbourhood socio-economic status suggesting that the lower MI risk is explained by protective effects of warfarin itself. There may, however, be residual confounding

that we could not adjust for which may explain differences between those prescribed warfarin and those not prescribed warfarin [20,28].

As expected, the mortality was higher among AF patients with MI than AF patients without MI during follow-up, even if the increased mortality risk was not as high as could have been expected, i.e. the relative risk was around 1.4 for women and 1.5 for men. In general the MI mortality in Sweden is decreasing [29], as in other EU countries [30]. Surprisingly, we found no mortality-reducing effect from antithrombotic treatment in general after MI. The reason for the non-significant findings is puzzling, but we could speculate that they are explained by non-adherence, or treatment by specialists in open care, e.g. cardiologists (we only had prescription data from primary care).

The incidence rates of MI after exclusion of patients on effective warfarin treatment could be expected to reflect the true incidence of MI of this cohort of primary healthcare patients, with 2.4 events per 100 patient-years among women and 2.3 among men. The incidence rate overall was 1.5 events per 100 patient-years among women and 1.4 among men. In Sweden in 2013, the incidence rate of a first MI was 1.1% among women aged 80–84 years; and 1.3% among men aged 75–79, with the corresponding figure in Stockholm County being 1.1%, reflecting the mean ages in the cohort of women and men during the time period [31].

The risk of MI among AF patients was lower in women than in men, HR 0.89, but non-significant. The aforementioned US study found a significantly higher MI risk among women [10].

There are certain limitations of this study. This is an observational study, and prescription of warfarin may have been influenced by other factors than we recorded, i.e. confounding by indication may be one explanation for the high magnitude of the preventive effect of warfarin on first stroke [32]. An earlier Swedish study concluded that “warfarin-treated patients are highly selected and that decisions not to treat elderly, frail high-risk patients are at higher risk of having complicating co-morbidities and a poor prognosis” [12]. Our data were extracted from electronic patient records in primary health care, and data may have been incomplete, e.g. for listings of diagnoses. However, we could expect the diagnoses of cardiovascular diseases and diabetes to be more accurate and complete than many other diagnoses, and less than 2% of the total number of diagnoses was missing [33]. Besides, we used hospital data for the diagnosis of MI. We had no data available on the type of atrial fibrillation (paroxysmal, persistent, permanent). We had no data on electro-conversion of AF, nor had we information on procedures such as catheter ablation or Cox-Maze operations. Furthermore, data on the severity of the cardiovascular co-morbidities, e.g. NYHA classification of congestive heart failure, were not available. However, since the variables available in the present study were obtained from primary health care electronic patient records they are made by active clinicians. We had no data on time in therapeutic INR range (TIR). Our analyses of PP-treatment are attempts to try to reflect a regular treatment, while analyses of ITT or not reflect a more crude division. In the statistical analyses, it was not possible to find a balanced model when trying to use a propensity score analysis.

Despite the limitations, one of the key strengths of this study is the linkage of clinical data from individual patients to national demographic and socio-economic data with less than 1% missing data. The clinical data were also highly complete, and studies using only hospital patients may underestimate the burden of co-morbidities [5]. For example, most patients with hypertension (70%) and diabetes (55%) are exclusively diagnosed in primary healthcare [34]. The comprehensive nature of our data made it possible to analyse men and women from all educational backgrounds and marital statuses. Another strength is the sample size of the study, i.e. 6646 men and 5637 women, and 68,000 person-years at risk analysed.

In conclusion, our results show that warfarin prevents MI when used to treat AF patients in a primary healthcare setting, and emphasize the importance of persistent anticoagulant treatment in those patients.

## Disclosures

The authors have no conflict of interest to disclose.

## Acknowledgements

This work was supported by grants to Kristina Sundquist and Jan Sundquist from the Swedish Research Council (K2012-70X-15428-08-3) as well as ALF funding (88009) to Jan Sundquist and Kristina Sundquist from Region Skåne.

Research reported in this publication was also supported by the National Heart, Lung, and Blood Institute of the National Institutes of Health under Award Number R01HL116381 to Kristina Sundquist. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

## Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.ijcard.2016.07.119>.

## References

- [1] A.S. Go, E.M. Hylek, K.A. Phillips, Y. Chang, L.E. Henault, J.V. Selby, et al., Prevalence of diagnosed atrial fibrillation in adults: national implications for rhythm management and stroke prevention: the AnTicoagulation and Risk Factors in Atrial Fibrillation (ATRIA) study, *JAMA* 285 (2001) 2370–2375.
- [2] S.S. Chugh, R. Havmoeller, K. Narayanan, D. Singh, M. Rienstra, E.J. Benjamin, et al., Worldwide epidemiology of atrial fibrillation: a Global Burden of Disease 2010 Study, *Circulation* 129 (2014) 837–847.
- [3] F. Rahman, G.F. Kwan, E.J. Benjamin, Global epidemiology of atrial fibrillation, *Nat. Rev. Cardiol.* 11 (2014) 639–654.
- [4] J. Ball, M.J. Carrington, J.J. McMurray, S. Stewart, Atrial fibrillation: profile and burden of an evolving epidemic in the 21st century, *Int. J. Cardiol.* 167 (2013) 1807–1824.
- [5] T. Forslund, B. Wettermark, P. Wandell, M. von Euler, J. Hasselstrom, P. Hjerdahl, Risk scoring and thromboprophylactic treatment of patients with atrial fibrillation with and without access to primary healthcare data: experience from the Stockholm health care system, *Int. J. Cardiol.* 170 (2013) 208–214.
- [6] F.R. Hobbs, C.J. Taylor, G. Jan Geersing, F.H. Rutten, J.R. Brouwer, European Primary Care Cardiovascular Society (EPCCS) consensus guidance on stroke prevention in atrial fibrillation (SPAF) in primary care, *Eur. J. Prev. Cardiol.* (2015).
- [7] L.S. Wann, A.B. Curtis, K.A. Ellenbogen, N.A. Estes, M.D. Ezekowitz, W.M. Jackman, et al., Management of patients with atrial fibrillation (compilation of 2006 ACCF/AHA/ESC and 2011 ACCF/AHA/HRS recommendations): a report of the American College of Cardiology/American Heart Association Task Force on practice guidelines, *Circulation* 127 (2013) 1916–1926.
- [8] P.A. Wolf, R.D. Abbott, W.B. Kannel, Atrial fibrillation as an independent risk factor for stroke: the Framingham study, *Stroke* 22 (1991) 983–988.
- [9] J. Schmitt, G. Duray, B.J. Gersh, S.H. Hohnloser, Atrial fibrillation in acute myocardial infarction: a systematic review of the incidence, clinical features and prognostic implications, *Eur. Heart J.* 30 (2009) 1038–1045.
- [10] E.Z. Soliman, M.M. Safford, P. Muntner, Y. Khodneva, F.Z. Dawood, N.A. Zakai, et al., Atrial fibrillation and the risk of myocardial infarction, *JAMA Intern. Med.* 174 (2014) 107–114.
- [11] J.T. Zhang, K.P. Chen, S. Zhang, Efficacy and safety of oral anticoagulants versus aspirin for patients with atrial fibrillation: a meta-analysis, *Medicine (Baltimore)* 94 (2015), e409.
- [12] T. Forslund, B. Wettermark, P. Wandell, M. von Euler, J. Hasselstrom, P. Hjerdahl, Risks for stroke and bleeding with warfarin or aspirin treatment in patients with atrial fibrillation at different CHA(2)DS(2)-VASc scores: experience from the Stockholm region, *Eur. J. Clin. Pharmacol.* 70 (2014) 1477–1485.
- [13] A. Dogliotti, E. Paolasso, R.P. Giugliano, Current and new oral antithrombotics in non-valvular atrial fibrillation: a network meta-analysis of 79 808 patients, *Heart* 100 (2014) 396–405.
- [14] M.I. Aguilar, R. Hart, L.A. Pearce, Oral anticoagulants versus antiplatelet therapy for preventing stroke in patients with non-valvular atrial fibrillation and no history of stroke or transient ischemic attacks, *Cochrane Database Syst. Rev.* (2007) CD006186.
- [15] G.Y. Lip, D.A. Lane, Does warfarin for stroke thromboprophylaxis protect against MI in atrial fibrillation patients? *Am. J. Med.* 123 (2010) 785–789.
- [16] K. Sundquist, X. Li, Coronary heart disease risks in first- and second-generation immigrants in Sweden: a follow-up study, *J. Intern. Med.* 259 (2006) 418–427.
- [17] T. Almgren, L. Wilhelmsen, O. Samuelsson, A. Himmelmann, A. Rosengren, O.K. Andersson, Diabetes in treated hypertension is common and carries a high cardiovascular risk: results from a 28-year follow-up, *J. Hypertens.* 25 (2007) 1311–1317.
- [18] P. Wandell, A.C. Carlsson, K. Sundquist, S.E. Johansson, J. Sundquist, Effect of cardiovascular drug classes on all-cause mortality among atrial fibrillation patients treated in primary care in Sweden: a cohort study, *Eur. J. Clin. Pharmacol.* 69 (2013) 279–287.
- [19] N.D. Aberg, H.G. Kuhn, J. Nyberg, M. Waern, P. Friberg, J. Svensson, et al., Influence of cardiovascular fitness and muscle strength in early adulthood on long-term risk of stroke in Swedish men, *Stroke* 46 (2015) 1769–1776.
- [20] A.C. Carlsson, P. Wandell, D. Gasevic, J. Sundquist, K. Sundquist, Neighborhood deprivation and warfarin, aspirin and statin prescription – a cohort study of men and women treated for atrial fibrillation in Swedish primary care, *Int. J. Cardiol.* 187 (2015) 547–552.
- [21] A.Y. Chou, G.B. Mancini, Warfarin monotherapy in atrial fibrillation patients with stable coronary disease one year after myocardial infarction/stent: two birds with one stone? *Can. J. Cardiol.* 31 (2015) 1070–1072.
- [22] S. Halvorsen, F. Andreotti, J.M. ten Berg, M. Cattaneo, S. Coccheri, R. Marchioli, et al., Aspirin therapy in primary cardiovascular disease prevention: a position paper of the European Society of Cardiology working group on thrombosis, *J. Am. Coll. Cardiol.* 64 (2014) 319–327.
- [23] C.H. Hennekens, J.E. Dalen, Aspirin in the primary prevention of cardiovascular disease: current knowledge and future research needs, *Trends Cardiovasc. Med.* 24 (2014) 360–366.
- [24] S.C. Smith Jr., E.J. Benjamin, R.O. Bonow, L.T. Braun, M.A. Creager, B.A. Franklin, et al., AHA/ACC secondary prevention and risk reduction therapy for patients with coronary and other atherosclerotic vascular disease: 2011 update: a guideline from the American Heart Association and American College of Cardiology Foundation, *Circulation* 124 (2011) 2458–2473.
- [25] M. Lamberts, G.H. Gislason, G.Y. Lip, J.F. Lassen, J.B. Olesen, A.P. Mikkelsen, et al., Antiplatelet therapy for stable coronary artery disease in atrial fibrillation patients taking an oral anticoagulant: a nationwide cohort study, *Circulation* 129 (2014) 1577–1585.
- [26] G.C. Flaker, M. Gruber, S.J. Connolly, S. Goldman, S. Chaparro, A. Vahanian, et al., Risks and benefits of combining aspirin with anticoagulant therapy in patients with atrial fibrillation: an exploratory analysis of stroke prevention using an oral thrombin inhibitor in atrial fibrillation (SPORTIF) trials, *Am. Heart J.* 152 (2006) 967–973.
- [27] P. Wandell, A.C. Carlsson, D. Gasevic, J. Sundquist, K. Sundquist, Neighbourhood socio-economic status and all-cause mortality in adults with atrial fibrillation: a cohort study of patients treated in primary care in Sweden, *Int. J. Cardiol.* 202 (2016) 776–781.
- [28] A.C. Carlsson, P. Wandell, K. Sundquist, S.E. Johansson, J. Sundquist, Differences and time trends in drug treatment of atrial fibrillation in men and women and doctors' adherence to warfarin therapy recommendations: a Swedish study of prescribed drugs in primary care in 2002 and 2007, *Eur. J. Clin. Pharmacol.* 69 (2013) 245–253.
- [29] J. Berg, L. Björck, G. Lappas, M. O'Flaherty, S. Capewell, A. Rosengren, Continuing decrease in coronary heart disease mortality in Sweden, *BMC Cardiovasc. Disord.* 14 (2014) 9.
- [30] M. Nichols, N. Townsend, P. Scarborough, M. Rayner, Trends in age-specific coronary heart disease mortality in the European Union over three decades: 1980–2009, *Eur. Heart J.* 34 (2013) 3017–3027.
- [31] D. Falkstedt, J. Stejerschantz Forsberg, Myocardial infarction and stroke (Hjärtinfarkt och stroke), in: M. Backhans, J. Stejerschantz Forsberg, A. Lager (Eds.), Public Health Report 2015, (Folkhälsorapport 2015), Centrum för epidemiologi och samhällsmedicin, Stockholms läns landsting, Stockholm 2015, pp. 15–20 (in Swedish).
- [32] J.S. de Koning, N.S. Klazinga, P.J. Koudstaal, A. Prins, G.J. Borsboom, J.P. Mackenbach, The role of 'confounding by indication' in assessing the effect of quality of care on disease outcomes in general practice: results of a case-control study, *BMC Health Serv. Res.* 5 (2005) 10.
- [33] K. Sundquist, A. Chaikiat, V.R. Leon, S.E. Johansson, J. Sundquist, Country of birth, socioeconomic factors, and risk factor control in patients with type 2 diabetes: a Swedish study from 25 primary health-care centres, *Diabetes Metab. Res. Rev.* 27 (2011) 244–254.
- [34] A.C. Carlsson, P. Wandell, U. Osby, R. Zarrinkoub, B. Wettermark, G. Ljunggren, High prevalence of diagnosis of diabetes, depression, anxiety, hypertension, asthma and COPD in the total population of Stockholm, Sweden – a challenge for public health, *BMC Public Health* 13 (2013) 670.