



Contents lists available at ScienceDirect

## International Journal of Cardiology

journal homepage: [www.elsevier.com/locate/ijcard](http://www.elsevier.com/locate/ijcard)

Letter to the Editor

## Effect of acute aerobic exercise on serum BDNF levels in patients with Chagas heart disease

Henrique Silveira Costa <sup>a,\*</sup>, Márcia Maria Oliveira Lima <sup>b</sup>, Marconi Gomes Silva <sup>a</sup>, Maria Clara Noman Alencar <sup>a</sup>, Maria Carmo Pereira Nunes <sup>a</sup>, Elizabeth Ribeiro Silva Camargos <sup>c</sup>, Patrícia Massara Martinelli <sup>c</sup>, Manoel Otávio Costa Rocha <sup>a</sup>

<sup>a</sup> Postgraduate Course of Infectious Diseases and Tropical Medicine, Department of Internal Medicine, Medical School and Hospital das Clínicas of the Federal University of Minas Gerais (UFMG), Belo Horizonte, Brazil

<sup>b</sup> Healthy and Biological Sciences Faculty, Physical Therapy School, Federal University of the Jequitinhonha and Mucuri Valleys (UFVJM), Diamantina, Brazil

<sup>c</sup> Department of Morphology, Institute of Biological Sciences, Federal University of Minas Gerais (UFMG), Belo Horizonte, Brazil

## ARTICLE INFO

## Article history:

Received 17 January 2014

Received in revised form 9 April 2014

Accepted 12 April 2014

Available online xxxx

## Keywords:

Chagas heart disease

Brain derived neurotrophic factor

Cardiopulmonary exercise testing

Acute aerobic exercise

Chagas disease is an important endemic disease in Latin America and an emerging disease in non-endemic countries [1]. Chagas heart disease (CHD), the most important clinical manifestation of this infection, causes heart failure, complex arrhythmias, thromboembolism and sudden death [2]. A diffuse damage in the autonomic nervous system is a pathological feature commonly present in variable degrees in patients with CHD [3].

The neurotrophin Brain Derived Neurotrophic Factor (BDNF) plays important functions in the nervous system, such as neuronal survival and maintenance, synaptic plasticity and memory processing [4]. Moreover, BDNF has an important role in metabolic events [5], being susceptible to regulation by physical activity. Recently, it has been demonstrated that Chagas disease patients exhibit higher serum BDNF levels compared to healthy subjects [6]. Interestingly, in patients with Chagas cardiomyopathy underwent 12-week-aerobic training, the best functional capacity was associated to higher serum BDNF levels [7]. However, the acute effect of aerobic exercise on the BDNF levels

in CHD remains unknown, as well as its relationship to ventricular dysfunction and exercise intensity.

Patients were recruited from an Outpatient Reference Center for Chagas Disease in the state of Minas Gerais, Brazil. The study was approved by the Institutional Review Committee and the subjects gave informed consent.

Inclusion criteria were the diagnosis of Chagas disease determined by positive serology, clinical, electrocardiographic or echocardiographic findings compatible with CHD, age between 30 and 60 years and sedentary lifestyle according to the International Physical Activity Questionnaire (IPAQ). Exclusion criteria were the presence of systemic or heart disease by any other causes or co-morbidities, the use of cardiac pacemaker, blood transfusion within six months and the use of antidepressant medication.

All patients underwent clinical evaluation, echocardiography, the Cardiopulmonary Exercise Testing (CPET) and blood sampling for measurement of serum BDNF. Patients were classified into 2 groups: dilated cardiomyopathy characterized by the echocardiographic finding of a dilated left ventricle with impaired ventricular systolic function, and non-dilated heart disease defined as a normal left ventricular dimensions and function. The CPET, gold standard in the assessment of functional capacity, was performed on a treadmill ramp protocol and the intensity of exercise was determined as the percentage of maximum heart rate ( $HR_{max} = 220$  minus the patient age) during the test. For moderate exercise the percentage was set at 60–79% and for high intensity above 80% of the  $HR_{max}$ . Blood samples were taken at rest and immediately after the CPET. Serum BDNF levels were determined by ELISA (Enzyme Linked Immuno Sorbent Assay), according to R&D Systems protocol (Minneapolis, MN, USA).

G Power software, version 3.1.0, was used to determine the sample size as 28 individuals, considering an alpha error of 0.05 and a statistical power of 95%. Thirty CHD patients ( $47.85 \pm 8.71$  years, 20 males) were selected for this study. Parametric paired t-test and Pearson correlation test and non-parametric Mann–Whitney and Spearman Rank correlation test were performed for data analysis, with significance levels at 0.001% and 0.05%.

In the overall study population, there was a significant decrease in serum BDNF levels after acute exercise ( $p = 0.006$ ). The clinical

\* Corresponding author at: Rua dos Guajajaras, 1172/407, Centro, 30.180-100, Belo Horizonte, Minas Gerais, Brazil. Tel.: +55 31 32754205.

E-mail address: [henriquesilveira@yahoo.com.br](mailto:henriquesilveira@yahoo.com.br) (H.S. Costa).

**Table 1**

Clinical data from CHD patients stratified according to left ventricular systolic function.

Variable	Non-dilated group (N = 16)	Dilated group (N = 14)	P value
BMI (kg/m <sup>2</sup> )	26.95 ± 4.63	26.85 ± 4.55	0.802
LVEF (%)	64.57 ± 5.50	39.50 ± 9.06	<b>&lt;0.001</b>
VO <sub>2peak</sub> (mL/kg/min)	29.98 ± 6.23	21.28 ± 7.54	<b>0.001</b>
BDNF at rest (pg/mL)	15,083.10 ± 4541.22	13,203.54 ± 3527.27	0.845
BDNF after exercise (pg/mL)	12,804.17 ± 4605.47	10,870.73 ± 5387.26	0.769
Δ BDNF (pg/mL)	−2278.93 ± 4183.27	−2334.76 ± 6376.03	0.902

Data presented as mean and standard deviation (M ± SD). BMI = body mass index, LVEF = left ventricular ejection fraction; VO<sub>2peak</sub> = peak oxygen uptake; Δ BDNF = changes in BDNF after exercise compared to rest; pg/mL = picograms per milliliter. The values highlighted in bold are statistically significant (P < 0.05).

characteristics of the CHD patients stratified according to left ventricular systolic function are presented in Table 1.

There was no significant change in serum BDNF levels after exercise in the dilated group (p = 0.136). In contrast, the non-dilated group

showed a significant decrease in serum BDNF levels (p = 0.038) (Fig. 1A). Concerning the intensity of physical exercise, patients who underwent CPET at moderate intensity (n = 11; all patients in dilated group) exhibited no changes in serum BDNF levels (p = 0.477). However, the patients who underwent CPET at high intensity (n = 19, 16 in the non-dilated group and 3 in the dilated group) had a significant decrease in BDNF levels (p < 0.001) (Fig. 1B).

To the best of our knowledge, this is the first study to demonstrate the effect of acute aerobic exercise on serum BDNF levels in patients with CHD. The main findings are the decrease in serum BDNF after acute aerobic exercise and the exercise intensity-dependent BDNF response.

All patients in the non-dilated group underwent CPET at high intensity, probably because they had higher functional capacity and lower exercise intolerance than those with left ventricular dysfunction. We hypothesized that both, the dysautonomia associated to CHD and the high intensity exercise, could contribute to a stress condition leading to a decrease in the serum BDNF levels after treadmill exercise.


Aerobic exercise at high intensity increases serum glucocorticoids such as cortisol that can negatively affect hippocampal plasticity [8] and consequently reduces BDNF levels. A significant reduction in BDNF levels after maximal exercise performed immediately after warm-up activity was also found by Rojas Vega et al. [9] in patients with a spinal cord injury. This previous study suggests that maximal effort can reduce BDNF levels by decreasing their production or release, increasing their degradation or stimulating their uptake from the blood into the brain.

A single session of moderate aerobic exercise may not be the stimulus needed to cause changes in BDNF levels in patients with CHD, since these patients have a severe autonomic dysfunction. However, the regular practice of moderate aerobic exercise might affect BDNF expression. Lima et al. [7] showed a significant increase in BDNF levels in patients with Chagas cardiomyopathy who had increased functional capacity after 12 weeks (36 sessions) of moderate exercise.

Currently, it has been suggested that high-intensity aerobic exercise is more effective to increase peak oxygen uptake in heart failure patients [10]. However, we observed that acute exercise performed at this intensity can reduce serum BDNF levels in Chagas heart disease patients, which could adversely affect neural and metabolic properties of this neurotrophic factor. In conclusion, our results suggest caution in prescribing high-intensity aerobic exercise for Chagas heart disease patients, as well as the importance of monitoring these patients during exercise. To confirm this, further longitudinal studies are necessary in encompassing a greater number of chagasic patients and exercise training at different levels of intensity.

## References

- [1] Nunes MC, Dones W, Morillo CA, Encina JJ, Ribeiro AL. Chagas disease: an overview of clinical and epidemiological aspects. *J Am Coll Cardiol* 2013;62(9):767–76.
- [2] Rocha MO, Teixeira MM, Ribeiro AL. An update on the management of Chagas cardiomyopathy. *Expert Rev Anti Infect Ther* 2007;5(4):727–43.
- [3] Marin-Neto JA, Cunha-Neto E, Maciel BC, Simoes MV. Pathogenesis of chronic Chagas heart disease. *Circulation* 2007;115(9):1109–23.



**Fig. 1.** Changes in serum BDNF before (white box) and after (gray box) acute exercise. A) Non-dilated and dilated group; B) chagasic patients who underwent Cardiopulmonary Exercise Test (CPET) at moderate and high intensity. \*p < 0.05; \*\*p < 0.001.

- [4] Lanni C, Stanga S, Racchi M, Govoni S. The expanding universe of neurotrophic factors: therapeutic potential in aging and age-associated disorders. *Curr Pharm Des* 2010;16(6):698–717.
- [5] Gomez-Pinilla F, Vaynman S, Ying Z. Brain-derived neurotrophic factor functions as a metabotrophin to mediate the effects of exercise on cognition. *Eur J Neurosci* 2008;28(11):2278–87.
- [6] Martinelli PM, da Costa Rocha MO, Teixeira AL, do Carmo Pereira Nunes M, da Silva Camargos ER. Brain-derived neurotrophic factor is up regulated in chronic Chagas disease. *Int J Cardiol* 2011;149(2):277–8.
- [7] Lima MM, Nunes MC, Nascimento B, et al. Improvement of the functional capacity is associated with BDNF and autonomic modulation in Chagas disease. *Int J Cardiol* 2013;167(5):2363–6.
- [8] Shen X, Li A, Zhang Y, et al. The effect of different intensities of treadmill exercise on cognitive function deficit following a severe controlled cortical impact in rats. *Int J Mol Sci* 2013;14(11):21598–612.
- [9] Rojas Vega S, Abel T, Lindschulten R, Hollmann W, Bloch W, Struder HK. Impact of exercise on neuroplasticity-related proteins in spinal cord injured humans. *Neuroscience* 2008;153(4):1064–70.
- [10] Ismail H, McFarlane JR, Dieberg G, Smart NA. Exercise training program characteristics and magnitude of change in functional capacity of heart failure patients. *Int J Cardiol* 2014;171(1):62–5.