



Identification and characterization of estrogen receptor-related receptor alpha and gamma in human glioma and astrocytoma cells

Mukesh K. Gandhari^a, Chester R. Frazier^a, Julia S. Hartenstein^a, Jean-Francois Cloix^b, Michel Bernier^{a,*}, Irving W. Wainer^{a,*}

^a Laboratory of Clinical Investigation, Biomedical Research Center, National Institute on Aging, National Institutes of Health, 251 Bayview Boulevard, Suite 100, Baltimore, MD 21224, USA

^b Laboratoire de Neurobiologie, Université d'Orléans, BP 6759, 45067 Orléans Cedex 2, France

ARTICLE INFO

Article history:

Received 31 August 2009

Accepted 2 October 2009

Keywords:

DY131
Biochanin A
XCT790
4-Hydroxytamoxifen
AAB-TATA-Luc
3xERE-TATA-Luc
T98G
U87MG
A-172 cells
ERRγ
ERRα

ABSTRACT

The purpose of this study was to examine expression and function of estrogen receptor-related receptors (ERRs) in human glioma and astrocytoma cell lines. These estrogen receptor-negative cell lines expressed ERRα and ERRγ proteins to varying degree in a cell context dependent manner, with U87MG glioma cells expressing both orphan nuclear receptors. Cell proliferation assays were performed in the presence of ERR isoform-specific agonists and antagonists, and the calculated EC₅₀ and IC₅₀ values were consistent with previous reported values determined in other types of cancer cell lines. Induction of luciferase expression under the control of ERR isoform-specific promoters was also observed in these cells. These results indicate that ERRα and ERRγ are differentially expressed in these tumor cell lines and likely contribute to agonist-dependent ERR transcriptional activity.

Published by Elsevier Ireland Ltd.

1. Introduction

In humans, gliomas form the vast majority of malignant brain tumors and are among the most lethal (Wrensch et al., 2006). Gliomas are tumors of glial cells, and among them astrocytomas represent about half of the central nervous system cancers (Trojan et al., 2007). Glioblastomas, the highest grade of gliomas, are highly invasive and exhibit spontaneous resistance to chemotherapeutic drugs, which is partly due to a lack of specificity of the drugs and to the inability to achieve their efficacious targeting concentrations. As a result, the various therapeutic means have thus far been inefficient in significantly improving patient survival. So new ways of treatment are constantly being investigated (Prados et al., 2006; Stupp et al., 2006).

It has been previously reported that a subset of estrogen receptor (ER)-negative breast cancers can respond to tamoxifen (Goldenberg and Froese, 1982; Gelmann, 1996), whereas some ERα-positive breast cancers are tamoxifen resistant (Ingle et al.,

1991; Jaiyesimi et al., 1995). These and other findings led to the search for novel tamoxifen-binding nuclear receptors and resulted in the identification of the estrogen related receptors ERRα and ERRγ (Giguère et al., 1988; Heard et al., 2000). It was later found that the antiestrogen tamoxifen was a competitive inhibitor of ERRγ, while being inactive toward ERRα (Coward et al., 2001). A recent study has showed that tamoxifen can inhibit the proliferation of a series of malignant ER-negative glioma cells (Hui et al., 2004); however, no attempt was made to establish the presence of ERR in these gliomas.

ERRα, ERRβ, and ERRγ form a subfamily of orphan nuclear receptors that have significant amino acid homology with the estrogen receptors ERα and ERβ (Ariazi and Jordan, 2006). As with other nuclear receptors ERRs are organized into modular domains with a less characterized N-terminal domain, a highly conserved DNA binding domain (DBD), and a potential ligand binding domain (LBD). ERRs share high homology in their DBD, and are constitutively active without binding to natural estrogen (Giguère, 2002; Horard and Vanacker, 2003). Like ERs, all ERRs bind to the classic ERE motif (AGGTCANNNTGACCT), suggesting that ERRs may be involved in related ER-mediated signaling pathways through association with coregulatory proteins and regulation of target gene expression. ERRs also bind to ERE-related response elements

* Corresponding authors. Tel.: +1 410 558 8199; fax: +1 410 558 8381.

E-mail addresses: Bernierm@mail.nih.gov (M. Bernier), Wainerlr@grc.nia.nih.gov (I.W. Wainer).

with extended half-site core sequences (TNAAGGTCA; ERRE/SF-1RE; Razaque et al., 2004), indicating that ERRs may also have their own independent regulatory pathways or functions distinct from ERs. Therefore, it is likely that ERRs may regulate a broad spectrum of genes in target cells.

Previous studies have demonstrated that ERRs are constitutively active and that this activity is controlled by coactivator concentration rather than small ligands (Greschik et al., 2002; Kallen et al., 2004). Indeed, it has been shown that a ligand is not required for the ERR γ LBD to adopt a transcriptionally active conformation and interact with the PPAR γ coactivator-1 (PGC-1) (Hentschke et al., 2002). In addition, while ERRs share significant LBD homology with ERs, they have different responses to ER agonists and antagonists. For example, the ER agonist estradiol is inactive in ERR expressing cell lines (Giguère et al., 1988; Hong et al., 1999), and diethyl stilbestrol, a structurally unrelated ER agonist, exerts antagonistic properties on all three ERR subtypes (Tremblay et al., 2001). While 4-hydroxytamoxifen (4-OHT) does not affect ERR α , it acts instead as an ERR β and ERR γ antagonist (Coward et al., 2001).

The tissue-specific distribution, expression levels, and function of various ERR subtypes has been reported previously (Giguère, 2008; Bookout et al., 2006). However, the relative quantity and/or repertoire of ERR subtypes in glioma and astrocytoma tumor lines has not been addressed to date. The objectives of this study were to examine the expression of ERRs in a number of human ER α -negative glioma and astrocytoma cell lines. Moreover, the effect of known isoform-specific agonists and antagonists of ERR α and ERR γ on cellular proliferation and ERR reporter activity was investigated. These results could lead to new therapeutic approaches for the treatment of malignant gliomas and astrocytomas.

2. Materials and methods

2.1. Cell culture

All cell lines were purchased from American Type Culture Collection (Manassas, VA). T98G, U87MG, U138, U118MG, U373MG, and LN229 cells were cultured in phenol red-free minimal essential medium (α -MEM) (Invitrogen, Carlsbad, CA), while the A-172 and MCF-7 cells were maintained in phenol red-free DMEM (Invitrogen). Both media were supplemented with penicillin (25 U/ml), streptomycin (25 U/ml), L-glutamine (Invitrogen), pyruvic acid, non-essential amino acids, and 5% charcoal/dextran-treated fetal bovine serum (Hyclone, Logan, Utah). Cells were maintained at 37 °C in a humidified incubator with 5% CO₂.

2.2. Western blotting

Cells grown at approximately 70–80% confluence were lysed in lysis buffer (10 mM Tris-HCl, pH 7.4, 0.15 M NaCl, 5 mM EDTA, pH 8.0, 1% Triton X-100) supplemented with a protease inhibitor cocktail (Complete EDTA-free protease inhibitor mixture tablets; Roche Molecular Biochemical, Indianapolis, IN). Insoluble material was removed by centrifugation (12,000 \times g for 20 min at 4 °C) and the precleared lysates were aliquoted and stored at –70 °C until analysis. The protein concentration of the lysates was measured with the BCA protein assay kit (Pierce Biotechnology, Inc., Rockford, IL) using bovine serum albumin as standard, and samples were normalized for protein content. An equal amount of 3 \times sample buffer was added, and samples (70 μ g) were separated by 10% SDS-polyacrylamide gel electrophoresis and transferred to polyvinylidene difluoride membranes using iBLOT (Invitrogen). The membranes were blocked with 5% milk in Tris-buffered saline/0.1% Tween-20, and then probed with primary antibodies at optimal dilutions, followed by horseradish peroxidase-conjugated secondary antibodies. The ECL Western blotting detection system (GE Healthcare, Piscataway, NJ) was used for visualization. The primary antibodies used were: rabbit polyclonal antibodies against ERR α and ERR γ (Affinity BioReagents, Golden, CO) and mouse monoclonal anti- β -actin (Abcam, Cambridge, MA).

2.3. MTS assay

The non-radioactive CellTiter 96[®] Aqueous cell proliferation assay was performed according to the manufacturer's protocol (Promega, Madison, WI). Briefly, cells were seeded in 96-well plates at a density of 3500 cells/well and cultured in complete phenol red-free medium for 24 h. Test compounds dissolved in dimethyl sulfoxide (DMSO) and mixed with culture medium were added to the cells in dif-

ferent concentrations. The compounds were as followed: DY131 (Tocris Bioscience, Ellisville, MO) [25, 50, 75, 100, 125, 150 and 175 nM]; 4-OHT (Sigma-Aldrich, St. Louis, MO) [0.1, 0.25, 0.5, 0.75, 1 and 2 μ M]; 5,7-dihydroxy-4'-methoxyisoflavone (biochanin A) (Sigma-Aldrich) [25, 50, 75, 100, 125 and 250 nM]; and XCT790 (Sigma-Aldrich) [0.1, 0.5, 1, 2, 3 and 4 μ M]. Control cultures were treated with DMSO. The maximum concentration of DMSO added to the medium was 0.5% and was found not to be cytotoxic. After 48 h, 20 μ l of 0.5 mg/ml MTS solution was added to each well, and the cultures were further incubated for 1 h. The absorbance was measured at 490 nm with a microplate reader (Thermo Scientific, USA). Change in growth rate was calculated as follows: [$A_{490\text{nm}}$ of treated cells/ $A_{490\text{nm}}$ of control cells]. Three wells were used for each treatment, and the experiments were repeated three times. EC₅₀ for agonists DY131 and biochanin A, and IC₅₀ for antagonists 4-OHT and XCT790 were calculated using the Graphpad Prism software (La Jolla, CA).

2.4. Luciferase assays

For reporter assays, cells were approximately 70% confluent at the time of transfection in 6-well plates. For each dish, 0.2 μ g ERR reporter plasmid (AAB-TATA-Luc or 3X ERE-TATA-Luc; both kindly provided by Christina T. Teng, National Institute of Environmental Health Sciences, Research Triangle Park, NC) and 0.1 μ g pCMV β -gal vector (Clontech Laboratories, Mountain View, CA) were mixed and used for transient transfection using LipofectAMINE 2000. After overnight incubation, the medium was replaced with complete medium with or without test compounds. Twenty-four hours later, cells were harvested and aliquots of each lysate were assayed for luciferase and β -gal activities according to the manufacturer's instructions (Promega). The luciferase activity was normalized to that of β -gal. These experiments were performed in three separate experiments each in triplicate.

2.5. Statistical analyses

All the experiments were performed in triplicate and repeated three times. Unless otherwise indicated, the data represent the means \pm SE. Differences between mean values were compared statistically by Student's *t* test and ANOVA. *P* < 0.05 was considered statistically significant.

3. Results and discussion

3.1. Estrogen receptor-related receptor expression in glioblastoma and astrocytoma cell lines

A comparative analysis of the relative expression levels of ERRs was performed in seven human ER-negative glioblastoma and astrocytoma cell lines using Western immunoblotting (Fig. 1). The breast cancer cell line MCF-7 was used as positive control as it expresses ERR α and ERR γ (Cheung et al., 2005). The results demonstrated cell-type specific expression of ERR α and ERR γ proteins. High constitutive levels of ERR α expression were observed in A-172 and U87MG cells, and significantly lower abundance in T98G, U138MG, and LN229 cells (Fig. 1). In contrast, the levels of ERR γ were the highest in T98G and U87MG cells, but almost undetectable in the other cell lines. These findings demonstrate that these ER-negative glioma and astrocytoma cell lines express a unique repertoire of ERR proteins, which could be reflective of distinct cellular functions.




Fig. 1. Constitutive expression of cellular ERR α and ERR γ proteins. The human brain tumor cell lines T98G, A-172, U87MG, U138MG, U118MG, U373MG and LN229, and the MCF-7 breast cancer cells were lysed and processed for Western blot analysis. The relative units (Rel) show the average of triplicate experiments, where the densitometric quantitation of the signals corresponding to ERR α and ERR γ proteins were normalized to that of β -actin.




Fig. 2. Effect of ERR agonists and antagonists on glioma cell proliferation. A-172, T98G and U87MG cells were incubated with the indicated concentrations of the ERR γ agonist, DY131 (A), ERR γ antagonist, 4-OHT (B), ERR α agonist, biochanin A (C), or ERR α antagonist, XCT790 (D) for 48 h followed by another 60 min incubation in the presence of 0.5 mg/ml MTS. Reduction of MTS was determined colorimetrically. (E) U87MG cells were incubated either with vehicle, 1 μ M 4-OHT, 1.5 μ M XCT790 or the combination 4-OHT plus XCT790 for 48 h followed by MTS treatment. Values are means \pm SE from 3 independent experiments performed in triplicate. Statistical analysis by ANOVA, with ^a $P < 0.05$ and ^b $P < 0.01$ vs. vehicle-treated controls.

3.2. Role of ERRs in cell proliferation

The constitutive expression of ERR α and/or ERR γ in T98G, A-172 and U87MG cells make them suitable models for the study of the role of ERRs in astrocytoma and glioblastoma cell proliferation. The non-radioactive MTS cell proliferation assay can be substituted for [³H]thymidine incorporation. Consequently, compounds blocking cell proliferation will decrease the reduction of the soluble tetrazolium salt, MTS, to formazan product in intact cells.

The ERR γ agonist DY131, which has been shown to induce a ~3- to 4-fold increase in ERR γ transcriptional activity in CV-1 cells (Yu and Forman, 2005), and 4-OHT were used to characterize the role of endogenous ERR γ in cell proliferation. Treatment of T98G and U87MG cells with DY131 for 48 h increased cell proliferation in a dose-dependent manner, with a maximum at 0.175 μ M and EC₅₀ values of 0.054 ± 0.016 μ M and 0.062 ± 0.001 μ M in T98G and U87MG cells, respectively (Fig. 2A). Blocking the ERR γ activity with 4-OHT markedly reduced cell proliferation, with IC₅₀ values of 1.58 ± 0.21 μ M and 1.46 ± 0.16 μ M in T98G and U87MG cells, respectively ($P \leq 0.01$) (Fig. 2B). These values are consistent with the previously reported IC₅₀ value of 4.2 ± 2.0 μ M in MCF-7 human breast cancer cells (Davis et al., 2008). The A-172 cells did not respond to either DY131 or 4-OHT, consistent with their lack of ERR γ protein (see Fig. 1).

Three isoflavones (daidzein, biochanin A and genistein) and one flavone (6,3',4'-trihydroxyflavone) have been identified as specific ERR α agonists, capable at inducing ERR α transcriptional activity in HeLa cells (Suetsugi et al., 2003). Of these, biochanin A was the most potent phytoestrogen tested. In contrast, the compound XCT790 has been found to inhibit ERR α transcriptional activity in the human breast cancer derived MCF-7 cell line, while promoting proteosomal degradation of this orphan nuclear receptor (Lanvin et al., 2007). Treatment of A-172 and U87MG cells with biochanin A significantly increased cell proliferation after 48 h, with a maximum at 0.250 μ M and EC₅₀ values of 0.046 ± 0.001 μ M and 0.045 ± 0.002 μ M in A-172 and U87MG cells, respectively (Fig. 2C). These values are 10-fold lower than the previously reported EC₅₀ value of 0.46 μ M in MCF-7 cells (Joung et al., 2003). The ERR α antagonist XCT790 produced a significant inhibition of cell proliferation in A-172 and U87MG cells, with IC₅₀ values of 2.1 ± 0.19 μ M and 3.7 ± 0.4 μ M, respectively (Fig. 2D), in close agreement with the reported IC₅₀ value of ~0.40 μ M in various cell-based assays (Willy et al., 2004). The T98G cells did not respond to either biochanin A or XCT790, consistent with their lack of ERR α protein (see Fig. 1).

Because U87MG cells contain both ERR α and ERR γ , we evaluated the effect of the combination 4-OHT plus XCT790 on cell proliferation. When used at concentrations at or near IC₅₀, both compounds acted in an additive fashion with more than 67% inhi-

bition as compared to the 24% and 29% inhibition when 4-OHT and XCT790 were added alone (Fig. 2E).

3.3. *In vitro* transcriptional activity of ERRs

To confirm that the endogenously expressed ERRs are transcriptionally active, a luciferase reporter construct containing either the 3xERE-TATA-Luc or AAB-TATA-Luc (Zhang and Teng, 2007) was transfected into T98G, A-172 and U87MG cells along with a plasmid encoding β -galactosidase for normalization (Fig. 3A). The AAB-TATA-Luc promoter activity was greater in T98G and U87MG cells than in A-172 cells, in agreement with an earlier report showing that ERR γ is specific for AAB elements (Zhang and Teng, 2007). Conversely, 3xERE-TATA-Luc promoter activity level was significantly higher in A-172 and U87MG cells when compared to T98G cells,

consistent with a previous report showing that ERR α is specific for 3xERE element (Zhang and Teng, 2007).

We examined the ability of ERR γ agonist and antagonist to modulate AAB-TATA-Luc promoter activity in T98G and U87MG cells. The ligand DY131 (175 nM) stimulated the promoter activity level by ~2.5-fold, while the antagonist 4-OHT (2 μ M) blocked the constitutive ERR γ transcriptional activity (Fig. 3B). Moreover, the ability of ERR α agonist and antagonist to modulate the 3xERE-TATA-Luc promoter activity was examined in A-172 and U87MG cells. The promoter activity level was stimulated approximately 2-fold with biochanin A (250 nM), but markedly inhibited upon cell treatment with XCT790 (2 μ M) (Fig. 3C).

It has been recently reported that the transcriptional activity of ectopically expressed ERR γ construct was dependent on AP-1/c-Jun complex, indicating a non-nuclear action of ERR γ (Riggins et al., 2008). Of interest, we sequenced the two promoters used in our study (3xERE-TATA-Luc and AAB-TATA-Luc), and could not find the consensus AP-1 sequence (TGA^{C/G}TC^{C/G}A). However, several 'AP-1 like' sites were present especially in the AAB-TAT-Luc construct (Supplemental Fig. S1). Future experiments aimed at determining the role of these 'AP-1 like' sites in ERR transcriptional activity is warranted but these fall outside the scope of the current study.

Given that ERR α and ERR γ are differentially expressed both in normal and tumor tissues, it is tempting to speculate that some of the cellular processes involved in the control of gene expression may depend on the expression profile of ERR isoforms. On the other hand, protein dimerization modulates the transcriptional activities of ERRs. Although active as a monomer, ERR γ homodimer exhibits enhanced transcriptional activity, whereas its heterodimerization with ERR α results in impaired activities of both ERR α and ERR γ (Huppunen and Aarnisalo, 2004). Therefore, formation of ERR γ -ERR α heterodimers in U87MG cells may well direct some gene expression program distinct from that of ERR γ homodimers or ERR α homodimers in T98G and A-172 cells, respectively.

4. Conclusion and implications

There have been a number of clinical studies exploring the use of tamoxifen in the treatment of gliomas, both as monotherapy or in combination with DNA alkylating agents or radiotherapy (Parney and Chang, 2003; Robins et al., 2006). The results have been varied and, on the whole, disappointing as the use of high dose tamoxifen appears to have limited efficacy in a majority of the patients. However, there continues to be interest in the use of tamoxifen and other selective estrogen receptor modulators (SERMs) in the treatment of gliomas, even though these tumors are estrogen receptor-negative. A recent study by Hui et al. (2004) demonstrated that both tamoxifen and CC-8490, a benzopyrone with SERM activity, displayed antiglioma activity *in vitro* and *in vivo*. The ability of tamoxifen to inhibit glioma cell proliferation and induce apoptosis has been associated with inhibition of protein kinase C (PKC) activity (Baltuch et al., 1995; Mastronardi et al., 1998) and the data from the study of Hui et al. suggests that the antiglioma activity of SERMs can be enhanced through the inhibition of the NF- κ B pathway (Hui et al., 2004).

It is of interest that the majority of the *in vitro* data reported by Hui et al. was obtained using the U87MG cell line and the *in vivo* xenograft model was also based upon implanted U87MG cells (Hui et al., 2004). The data from this study indicates that the U87MG cells express high levels of both ERR α and ERR γ (Fig. 1) and that these proteins are responsive to isoform-specific inhibitors (Figs. 2 and 3). This suggests that the SERM related antiglioma activity observed by Hui et al. (2004) might be due in part to the antagonist activity of these compounds at the level of ERR γ . In addition, the data from this study demonstrate that ERR α and ERR γ were differentially




Fig. 3. Constitutive and ligand-induced ERR promoter activity in human brain tumor cell lines. (A) A-172, T98G and U87MG cells were transiently transfected with either the 3xERE-TATA-Luc or AAB-TATA-Luc plasmid together with β -galactosidase vector. Twenty-four hours later, the cells were incubated in medium supplemented with 5% charcoal/dextran-treated serum for 24 h. (B) T98G and U87MG cells were transfected with the AAB-TATA-Luc plasmid followed by a 24-h incubation either with DMSO, 175 nM DY131, or 2 μ M 4-OHT. (C) A-172 and U87MG cells were transfected with the 3xERE-TATA-Luc plasmid, followed by a 24-h incubation either with vehicle (DMSO), 250 nM biochanin A, or 2 μ M XCT790. Results are expressed as means \pm SD of a single experiment performed in triplicate dishes. Results are representative of three separate experiments with similar results.

expressed at the protein levels in the three glioma and astrocytoma cell lines examined, and that the A172 and T98G cell lines contained only ERR α or ERR γ protein, respectively. Our results suggest that the variable response to tamoxifen in the clinical treatment of glioblastomas may reflect the variable expression of ERR α or ERR γ and that better outcomes may be obtained through the use of tamoxifen and an ERR α antagonist. The expression of ERR α and ERR γ in tumor biopsy samples is currently in progress and the results will be reported elsewhere.

Acknowledgments

The work was supported entirely by the Intramural Research Program of the NIH, National Institute on Aging.

Appendix A. Supplementary data

Supplementary data associated with this article can be found, in the online version, at doi:10.1016/j.mce.2009.10.001.

References

- Ariazi, E.A., Jordan, V.C., 2006. Estrogen-related receptors as emerging targets in cancer and metabolic disorders. *Curr. Top. Med. Chem.* 6, 203–215.
- Baltuch, G.H., Dooley, N.P., Villemure, J.G., Yong, V.W., 1995. Protein kinase C and growth regulation of malignant gliomas. *Can. J. Neurol. Sci.* 22, 264–271.
- Bookout, A.L., Jeong, Y., Downes, M., Yu, R.T., Evans, R.M., Mangelsdorf, D.J., 2006. Anatomical profiling of nuclear receptor expression reveals a hierarchical transcriptional network. *Cell* 126, 789–799.
- Cheung, C.P., Yu, S., Wong, K.B., Chan, L.W., Lai, F.M., Wang, X., Suetsugi, M., Chen, S., Chan, F.L., 2005. Expression and functional study of estrogen receptor-related receptors in human prostatic cells and tissues. *J. Clin. Endocrinol. Metab.* 90, 1830–1844.
- Coward, P., Lee, D., Hull, M.V., Lehmann, J.M., 2001. 4-Hydroxytamoxifen binds to and deactivates the estrogen-related receptor gamma. *Proc. Natl. Acad. Sci. U.S.A.* 98, 8880–8884.
- Davis, D.D., Díaz-Cruz, E.S., Landini, S., Kim, Y.W., Brueggemeier, R.W., 2008. Evaluation of synthetic isoflavones on cell proliferation, estrogen receptor binding affinity, and apoptosis in human breast cancer cells. *J. Steroid Biochem. Mol. Biol.* 108, 23–31.
- Gelmann, E.P., 1996. Tamoxifen induction of apoptosis in estrogen receptor-negative cancers: new tricks for an old dog? *J. Natl. Cancer Inst.* 88, 224–226.
- Giguère, V., Yang, N., Segui, P., Evans, R.M., 1988. Identification of a new class of steroid hormone receptors. *Nature* 331, 91–94.
- Giguère, V., 2002. To ERR in the estrogen pathway. *Trends Endocrinol. Metab.* 13, 220–225.
- Giguère, V., 2008. Transcriptional control of energy homeostasis by the estrogen-related receptors. *Endocr. Rev.* 29, 677–696.
- Goldenberg, G.J., Froese, E.K., 1982. Drug and hormone sensitivity of estrogen receptor-positive and -negative human breast cancer cells in vitro. *Cancer Res.* 42, 5147–5151.
- Greschik, H., Wurtz, J.M., Sanglier, S., Bourguet, W., van Dorsselaer, A., Moras, D., Renaud, J.P., 2002. Structural and functional evidence for ligand-independent transcriptional activation by the estrogen-related receptor 3. *Mol. Cell* 9, 303–313.
- Heard, D.J., Norby, P.L., Holloway, J., Vissing, H., 2000. Human ERR γ , a third member of the estrogen receptor-related receptor (ERR) subfamily of orphan nuclear receptors: tissue-specific isoforms are expressed during development and in the adult. *Mol. Endocrinol.* 14, 382–392.
- Hentschke, M., Süsens, U., Borgmeyer, U., 2002. PGC-1 and PERC, coactivators of the estrogen receptor-related receptor γ . *Biochem. Biophys. Res. Commun.* 299, 872–879.
- Hong, H., Yang, L., Stallcup, M.R., 1999. Hormone-independent transcriptional activation and coactivator binding by novel orphan nuclear receptor ERR3. *J. Biol. Chem.* 274, 22618–22626.
- Horard, B., Vanacker, J.M., 2003. Estrogen receptor-related receptors: orphan receptors desperately seeking a ligand. *J. Mol. Endocrinol.* 31, 349–357.
- Hui, A.M., Zhang, W., Chen, W., Xi, D., Purow, B., Friedman, G.C., Fine, H.A., 2004. Agents with selective estrogen receptor (ER) modulator activity induce apoptosis in vitro and in vivo in ER-negative glioma cells. *Cancer Res.* 64, 9115–9123.
- Hupponen, J., Aarnisalo, P., 2004. Dimerization modulates the activity of the orphan nuclear receptor ERR γ . *Biochem. Biophys. Res. Commun.* 314, 964–970.
- Ingle, J.N., Twito, D.I., Schaid, D.J., Cullinan, S.A., Krook, J.E., Mailliard, J.A., Tschetter, L.K., Long, H.J., Gerstner, J.G., Windschitl, H.E., et al., 1991. Combination hormonal therapy with tamoxifen plus fluoxymesterone versus tamoxifen alone in postmenopausal women with metastatic breast cancer. An updated analysis. *Cancer* 67, 886–891.
- Jaiyesimi, I.A., Buzdar, A.U., Decker, D.A., Hortobagyi, G.N., 1995. Use of tamoxifen for breast cancer: twenty-eight years later. *J. Clin. Oncol.* 13, 513–529.
- Joung, K.E., Kim, Y.W., Sheen, Y.Y., 2003. Assessment of the estrogenicity of isoflavonoids, using MCF-7-ERE-Luc cells. *Arch. Pharm. Res.* 26, 756–762.
- Kallen, J., Schlaeppli, J.M., Bitsch, F., Filipuzzi, I., Schilb, A., Riou, V., Graham, A., Strauss, A., Geiser, M., Fournier, B., 2004. Evidence for ligand-independent transcriptional activation of the human estrogen-related receptor α (ERR α): crystal structure of ERR α ligand binding domain in complex with peroxisome proliferator-activated receptor coactivator-1 α . *J. Biol. Chem.* 279, 49330–49337.
- Lanvin, O., Bianco, S., Kersual, N., Chalbos, D., Vanacker, J.M., 2007. Potentiation of ICI182,780 (Fulvestrant)-induced estrogen receptor- α degradation by the estrogen receptor-related receptor- α inverse agonist XCT1790. *J. Biol. Chem.* 282, 28328–28334.
- Mastronardi, L., Farah, J.O., Puzzilli, F., Ruggeri, A., 1998. Tamoxifen modulation of carboplatin cytotoxicity in human U-138 glioma cell line. *Clin. Neurol. Neurosurg.* 100, 89–93.
- Parney, I.F., Chang, S.M., 2003. Current chemotherapy for glioblastoma. *Cancer J.* 9, 149–156.
- Prados, M.D., Lamborn, K., Yung, W.K., Jaeckle, K., Robins, H.I., Mehta, M., Fine, H.A., Wen, P.Y., Cloughesy, T., Chang, S., Nicholas, M.K., Schiff, D., Greenberg, H., Junck, L., Fink, K., Hess, K., Kuhn, J., 2006. North American Brain Tumor Consortium A phase 2 trial of irinotecan (CPT-11) in patients with recurrent malignant glioma: a North American Brain Tumor Consortium study. *Neuro-oncol.* 8, 189–193.
- Razzaque, M.A., Masuda, N., Maeda, Y., Endo, Y., Tsukamoto, T., Osumi, T., 2004. Estrogen receptor-related receptor- γ has an exceptionally broad specificity of DNA sequence recognition. *Gene* 340, 275–282.
- Riggins, R.B., Lan, J.P., Zhu, Y., Klimach, U., Zwart, A., Cavalli, L.R., Haddad, B.R., Chen, L., Gong, T., Xuan, J., Ethier, S.P., Clarke, R., 2008. ERR γ mediates tamoxifen resistance in novel models of invasive lobular breast cancer. *Cancer Res.* 68, 8908–8917.
- Robins, H.I., Won, M., Seiferheld, W.F., Schultz, C.J., Choucair, A.K., Brachman, D.G., Demas, W.F., Metha, M.P., 2006. Phase 2 trial of radiation plus high-dose tamoxifen for glioblastoma multiforme: RTOG protocol BR-0021. *Neuro-oncol.* 8, 47–52.
- Stupp, R., Hegi, M.E., van den Bent, M.J., Mason, W.P., Weller, M., Mirmanoff, R.O., Cairncross, J.G., European Organization for Research and Treatment of Cancer Brain Tumor and Radiotherapy Groups; National Cancer Institute of Canada Clinical Trials Group 2006. Changing paradigms—an update on the multidisciplinary management of malignant glioma. *Oncologist* 11, 165–180.
- Suetsugi, M., Su, L., Karlsberg, K., Yuan, Y.C., Chen, S., 2003. Flavone and isoflavone phytoestrogens are agonists of estrogen-related receptors. *Mol. Cancer Res.* 1, 981–991.
- Tremblay, G.B., Kunath, T., Bergeron, D., Lapointe, L., Champigny, C., Bader, J.A., Rossant, J., Giguère, V., 2001. Diethylstilbestrol regulates trophoblast stem cell differentiation as a ligand of orphan nuclear receptor ERR β . *Genes Dev.* 15, 833–838.
- Trojan, J., Cloix, J.F., Ardourel, M.Y., Chatel, M., Anthony, D.D., 2007. Insulin-like growth factor type I biology and targeting in malignant gliomas. *Neuroscience* 145, 795–811.
- Willy, P.J., Murray, I.R., Qian, J., Busch, B.B., Stevens Jr., W.C., Martin, R., Mohan, R., Zhou, S., Ordentlich, P., Wei, P., Sapp, D.W., Horlick, R.A., Heyman, R.A., Schulman, I.G., 2004. Regulation of PPAR γ coactivator 1 α (PGC-1 α) signaling by an estrogen-related receptor α (ERR α) ligand. *Proc. Natl. Acad. Sci. U.S.A.* 101, 8912–8917.
- Wrench, M., Rice, T., Miike, R., McMillan, A., Lamborn, K.R., Aldape, K., Prados, M.D., 2006. Diagnostic, treatment, and demographic factors influencing survival in a population-based study of adult glioma patients in the San Francisco Bay Area. *Neuro-oncol.* 8, 12–26.
- Yu, D.D., Forman, B.M., 2005. Identification of an agonist ligand for estrogen-related receptors ERR β / γ . *Bioorg. Med. Chem. Lett.* 15, 1311–1313.
- Zhang, Z., Teng, C.T., 2007. Interplay between estrogen-related receptor α (ERR α) and γ (ERR γ) on the regulation of ERR α gene expression. *Mol. Cell. Endocrinol.* 264, 128–141.