

Is depression associated with increased oxidative stress? A systematic review and meta-analysis

Catherine N. Black^{a,d,*}, Mariska Bot^{a,d}, Peter G. Scheffer^b,
Pim Cuijpers^{c,d}, Brenda W.J.H. Penninx^{a,d}

^a Department of Psychiatry, VU University Medical Center, Amsterdam, The Netherlands

^b Department of Clinical Chemistry, Metabolic Laboratory, VU University Medical Center, Amsterdam, The Netherlands

^c Department of Clinical Psychology, VU University Amsterdam, The Netherlands

^d EMGO Institute for Health and Care Research, Amsterdam, The Netherlands

Received 25 July 2014; received in revised form 23 September 2014; accepted 24 September 2014

KEYWORDS

Depression;
Major depressive
disorder;
Bipolar disorder;
Oxidative stress;
8-Hydroxy-2'-
deoxyguanosine
(8-OHdG);
F2-isoprostanes

Summary

Background: It has been suggested that depressed persons have increased oxidative stress and decreased anti-oxidant defences. 8-Hydroxy-2'-deoxyguanosine (8-OHdG) and F2-isoprostanes, measures of oxidative DNA and lipid damage respectively, are among the most reliable oxidative stress markers, but studies on their association with depression show conflicting results. This meta-analysis quantifies the association between depression and these markers and explores factors that may explain inconsistencies in the results.

Methods: A systematic literature search was conducted in PubMed, EMBASE and PsycINFO. Studies assessing the association of 8-OHdG or F2-isoprostanes with elevated depressive symptoms, major depressive disorder (MDD) or bipolar disorder (BD) were pooled in two random-effect models.

Results: The pooled effect size (Hedges' g) for the association of depression with oxidative stress was 0.31 ($p=0.01$, $I^2=75\%$) for 8-OHdG (10 studies, 1308 subjects) and 0.48 ($p=0.001$, $I^2=73\%$) for F2-isoprostanes (8 studies, 2471 subjects), indicating that both markers are increased in depression. There was no indication of publication bias for either marker. The F2-isoprostane results did not differ by type of depression, biological specimen, laboratory method or quality, however subgroup analyses in the 8-OHdG studies showed significantly stronger associations in plasma/serum vs. urine samples ($p<0.01$), in measurements performed with immuno-assay vs. chromatography–mass spectrometry ($p<0.01$) and weaker associations in high quality studies vs. low ($p=0.02$).

* Corresponding author at: Department of Psychiatry, VU University Medical Center, Postbus 74077, 1070 BB Amsterdam, The Netherlands. Tel.: +31 20 788 4597.

E-mail address: c.black@ggzingeest.nl (C.N. Black).

Conclusion: This meta-analysis finds that oxidative stress, as measured by 8-OHdG and F2-isoprostanes, is increased in depression. Larger-scale studies are needed to extend the evidence on oxidative stress in depression, and examine the potential impact of treatment.

© 2014 Elsevier Ltd. All rights reserved.

1. Introduction

Depression is a leading cause of morbidity worldwide (Vos et al., 2012). Depression is highly prevalent (Kessler et al., 2003) and has a profound impact on functioning and quality of life (Bijl and Ravelli, 2000) as well as on somatic health. Sufferers are at higher risk of diseases that are usually associated with increasing age such as cardiovascular disease (Nicholson et al., 2006), obesity (Luppino et al., 2010), diabetes (Mezuk et al., 2008), cancer (Chida et al., 2008), cognitive impairment (Barnes et al., 2006) and have a higher all-cause mortality rate (Cuijpers et al., 2014). It is hypothesized that increased metabolic stress and accelerated cellular ageing may be underlying pathways that contribute to this poorer physical health in individuals with depression (Wolkowitz et al., 2011b). A fast growing body of evidence suggests the involvement of a specific component of metabolic stress, oxidative stress, in the pathophysiology of depression (Maes et al., 2011).

Oxidative stress refers to the biologically damaging effects of free radicals (Valko et al., 2007). The production of free radicals, or reactive oxygen species (ROS), is a normal process in aerobic metabolism and ROS perform a number of physiological roles in cellular signalling and in the defence against pathogens. However, when present in excess, ROS cause damage to lipids, proteins and DNA, and can ultimately result in cell death. Oxidative stress is a well-recognized mechanism in ageing and disease. It has been shown to play a role in the pathophysiology of – among others – cardiovascular disease, diabetes mellitus, cancer and Alzheimer's disease (Valko et al., 2007). Additionally, there is evidence suggesting that oxidative stress may be increased in a number of psychiatric disorders, including depression (Pandya et al., 2013).

A recent meta-analysis pooling data from studies with different oxidative stress markers suggests oxidative stress is increased and antioxidant defences are decreased in depression (Palta et al., 2014). In line with these findings, increased nitric oxide (NO) and lipid peroxidation, as measured by thiobarbituric acid reactive substance (TBARS) assay, have also been found in patients with bipolar disorder, however these patients did not differ from controls in anti-oxidant enzymes levels (Andreazza et al., 2008). Overall, these studies suggest that oxidative stress is increased in major depressive disorder and bipolar disorder.

There is a wide range of oxidative stress biomarkers and laboratory techniques available, each of which has its own strengths and limitations (Dalle-Donne et al., 2006). To date there is no consensus on the most appropriate biomarkers of oxidative stress in general and the validity of many of those in use is to still be established. ROS have a short half-life, making measurement difficult. Levels of antioxidants,

vitamins or anti-oxidant enzymes are informative, but reflect only one side of redox homeostasis, leaving the question unanswered whether decreased levels are actually also indicative of increased oxidative damage. Studies show quite consistently that lipid peroxidation reflected by malondialdehyde (MDA) measured with the TBARS assay is increased in depression (Palta et al., 2014) and in bipolar disorder (Andreazza et al., 2008). However this commonly used method also has well recognized limitations: MDA is not a specific product of lipid peroxidation, and the TBARS assay itself can generate MDA, causing overestimation of levels. MDA therefore cannot be considered an optimal representation of oxidative stress in vivo (Meagher and FitzGerald, 2000; Dalle-Donne et al., 2006).

The current study focusses on two important measures of oxidative damage that have already been widely studied in somatic disease and are the subject of an increasing number of recent publications on depression: 8-hydroxy-2'-deoxyguanosine (8-OHdG) and F2-isoprostanes. The majority of the currently available literature on these markers in depression was not included, or not yet available for inclusion, in the previous meta-analyses on this subject. 8-OHdG and F2-isoprostanes reflect oxidative damage to DNA and lipids respectively. 8-OHdG is an oxidized derivative of deoxyguanosine and it is both the most abundant and most investigated DNA lesion. It has recognized mutagenic properties and has been linked to – among others – the development of cancer (Valavanidis et al., 2009). F2-isoprostanes, oxidized derivatives of arachidonic acid, have come to be considered the preferred approach to assess oxidative stress in vivo and lipid peroxidation in particular (Niki, 2014).

Several studies have found elevated levels of F2-isoprostanes (Dimopoulos et al., 2008; Yager et al., 2010; Chung et al., 2013) and 8-OHdG (Irie et al., 2005; Forlenza and Miller, 2006) in patients with depression, but these findings have not been consistent (Yi et al., 2012; Rawdin et al., 2013). Earlier studies did not systematically explore to what extent the (conflicting) findings are due to e.g. the laboratory methods, biological specimens used for oxidative stress, or the extent to which studies took potential confounders such as health and lifestyle factors into account. The present study extends the current evidence-base by systematically meta-analysing the literature on two robust markers of oxidative stress, 8-OHdG and F2-isoprostanes, and their association with depression (major depressive disorder, bipolar disorder and elevated depressive symptoms). In addition, by conducting subgroup analyses based on type of depression, biological specimen, laboratory method used to measure oxidative stress, correction for confounders and the quality of studies, this study aims to identify factors that contribute to the inconsistent findings of individual studies.

2. Methods

2.1. Literature search and study selection

Systematic searches of the literature were conducted in the databases PubMed, EMBASE and PsycINFO up to January 8th 2014 with search terms covering major depressive disorder, bipolar disorder and depressive symptoms combined with 8-OHdG and F2-isoprostanes respectively, taking into account a wide range of synonyms used for these markers. A full list of search terms is reported in [Appendix A](#). No limitations in the search strategy were set. The search results were reviewed by two independent reviewers (CNB and MB) by screening title and abstract, followed by a full text review. Disagreements were settled by discussion.

Studies were eligible for inclusion if they:

- (1) contained a measurement of 8-OHdG or F2-isoprostanes in any body fluid or tissue in live human adult subjects;
- (2) defined major depressive disorder (MDD) or bipolar disorder (BD) according to DSM-IV or ICD-10 criteria or assessed depressive symptoms using a validated instrument;
- (3) reported (or were able to provide) sufficient information to calculate an effect size for the difference between levels of the oxidative stress markers in control subjects and subjects with depression.

An assessment of the references of the included studies as well as a search of their citations in the PubMed database was performed to identify any additional studies.

2.2. Data extraction and risk of bias assessment

Two authors (CNB and MB) independently extracted the study characteristics (including among others biological specimen, age and sex distribution) and results from each study using a predesigned collection form (see [Appendix B](#)). The results extracted included means, standard deviations of oxidative stress markers in patient and control groups (or alternative results sufficient to calculate an effect size) and an assessment of correction for potential confounders (age, sex, ethnicity, socio-economic status, smoking, alcohol use, body mass index (BMI), physical activity, somatic disease and antidepressant or mood stabilizer use). Authors of studies that did not contain sufficient information (means, standard deviations or standard errors and number of subjects) to calculate an effect size were contacted with a request for additional data. Authors of studies who only reported results on depressive symptoms as continuous variables were requested to perform additional analyses using a dichotomized classification of a depressed and control group based on the appropriate cut-off for the instrument used to assess depressive symptoms. Authors of studies whose original articles did not contain adjusted results were contacted with a request to perform additional analyses adjusting for as many of the following confounders if available: age, sex, socio-economic status or education, ethnicity, smoking, alcohol use, BMI; physical activity, presence of somatic disease and antidepressant or mood stabilizer use. If multiple adjusted analyses had been conducted, the results

corrected for the largest number of potential confounders were included in the meta-analysis.

The quality of the studies was determined by assessing the risk of bias by two independent reviewers (CNB and MB) in three domains: selection bias, information bias and confounding. Disagreements were settled by discussion. Risk of bias assessment in observational studies requires consideration of the risks specific to the subject of study. For this purpose we adapted a tool by [Hayden et al. \(2006\)](#) (see [Appendix B](#)). Low risk of bias was defined as a score of 4.5 or higher (on a 5 point scale).

2.3. Meta-analysis

Analyses were performed with comprehensive meta-analysis (CMA) software version 2.2.064. Effect sizes (Hedges' g) were calculated and pooled using a random effects model, as considerable heterogeneity was expected. A p -value < 0.05 was considered significant.

Possible publication bias was tested by inspecting the funnel plot, by the statistical significance of the Egger's test of the intercept ([Sterne and Egger, 2001](#)) and [Duval and Tweedie's \(2000\)](#) trim and fill procedure.

To examine heterogeneity, the I^2 -statistic was calculated. Values of 25%, 50% and 75% indicate low, moderate and high heterogeneity, respectively ([Higgins et al., 2003](#)). The 95% confidence intervals around I^2 ([Ioannidis et al., 2007](#)) were calculated using the non-central chi-squared-based approach within the heterogi module for STATA version 11.0 for Mac.

When a minimum of three samples per subgroup was available, analyses were performed categorized by: type of depression (MDD, BD, depressive symptoms above the questionnaire cut-off level), laboratory method for measuring oxidative stress (chromatography [coupled to either mass-spectrometry or electrochemical detection] vs. immunoassay), biological specimen used (urine, blood[product] or other), low or high risk of bias score, and adjusted versus unadjusted results. Studies that accounted for at least age, sex and one life-style variable (either by matching, restriction, adjustment or testing for baseline group differences) were defined as corrected for confounding.

3. Results

3.1. Search results, study characteristics and risk of bias scoring

3.1.1. 8-OHdG

After removal of duplicates, 79 records were assessed based on title and abstract. The full-text of 39 of these records was retrieved for further assessment. Of these studies 6 did not report sufficient information in the original articles to calculate an effect size and none was provided by authors on request ([Irie et al., 2001, 2002, 2003](#); [Maes et al., 2009](#); [Iida et al., 2011](#); [Ceylan et al., 2013](#)). In 2 studies both 8-OHdG and depression were measured but the association between the two was not the focus of study and therefore not analyzed ([Wu et al., 2009](#); [Cepnija et al., 2011](#)), nor was this data provided on request. In total 10 studies met




Figure 1 Flowchart of the selection of studies on 8-OHdG or F2-isoprostanes in depression (major depressive disorder, bipolar disorder and depressive symptoms).

all criteria for inclusion; 4 studies on depressive symptoms (Kupper et al., 2009; Wei et al., 2009a,b; Yi et al., 2012), 2 studies on MDD (Irie et al., 2005; Forlenza and Miller, 2006), 3 studies on BD (Ceylan et al., 2012; Soeiro-de-Souza et al., 2013; Huzayyin et al., 2014) and 1 study with both MDD and BD patients (Jorgensen et al., 2013) (Fig. 1). Two of the included studies (Yi et al., 2012; Jorgensen et al., 2013) provided additional data on request.

The 10 studies (Irie et al., 2005; Forlenza and Miller, 2006; Kupper et al., 2009; Wei et al., 2009a,b; Yi et al., 2012; Ceylan et al., 2012; Soeiro-de-Souza et al., 2013; Jorgensen et al., 2013; Huzayyin et al., 2014) included a total of 579 subjects with depression (332 with depressive symptoms [scoring above the cut-off of the instrument used], 141 with MDD, 106 with BD) and 729 controls. All included studies were published between 2005 and 2014. The studies include samples from the general population ($N=2$), psychiatric in- and outpatient clinics ($N=4$), hospital oncology ($N=2$) and heart failure departments ($N=1$) and 1 unreported source

(Table 1). Risk of bias was scored as low in 3 studies and high in 7 studies (Appendix C).

3.1.2. F2-isoprostanes

After removal of duplicates, 221 records were assessed based on title and abstract. The full-text of 30 of these records was retrieved for further assessment. Of these studies 1 did not report sufficient information in the original article to calculate an effect size and none was provided by authors on request (Freund-Levi et al., 2011). In one study both F2-isoprostanes and depression were measured but the association between the two was not the focus of study and therefore not analyzed nor provided on request (Janicki-Deverts et al., 2009). The samples of Rawdin et al. (2013) and Wolkowitz et al. (2011a) overlap, leaving 8 original samples eligible for inclusion, 3 on depressive symptoms (Chung et al., 2009; Segal et al., 2012; Milaneschi et al., 2013) and 5 on MDD (Fig. 1). Six of the included studies provided additional data upon request (Chung et al., 2009, 2013; Yager et al., 2010; Wolkowitz et al., 2011a; Segal et al., 2012; Milaneschi et al., 2013).

The 8 studies (Dimopoulos et al., 2008; Chung et al., 2009, 2013; Yager et al., 2010; Wolkowitz et al., 2011a; Segal et al., 2012; Pomara et al., 2012; Milaneschi et al., 2013) include a total of 293 subjects with depression (144 subjects with depressive symptoms, 149 with MDD) and 2178 controls. All included studies were published between 2008 and 2013. The studies include samples of adults from the general population ($N=2$), elderly adults from the general population ($N=3$), psychiatric outpatients ($N=1$), systemic lupus erythematosus (SLE) patients ($N=1$) and fibromyalgia patients ($N=1$) (Table 2). Risk of bias was scored as low in 3 studies and high in 5 studies (Appendix C).

3.2. Meta-analyses

3.2.1. 8-OHdG

The overall effect size (Hedges' g) including all 10 studies on 8-OHdG in the meta-analysis was 0.31 (95% CI 0.06, 0.56), $I^2 = 75\%$ (95% CI 58–86%) indicating 8-OHdG is significantly increased in depression (Fig. 2a and Table 3). Egger's test for publication bias was not significant ($p=0.69$). The effect size adjusted for publication bias by Duval and Tweedie's trim and fill procedure (two trimmed studies) increased marginally to 0.37 (95% CI 0.13, 0.61) indicating no or little effects of publication bias (see Appendix D for funnel plot).

Subgroup analyses in the 8-OHdG studies showed significantly larger effect sizes for studies that were conducted in plasma or serum vs. those in urine ($p<0.001$) and studies that were performed with immuno-assays vs. those with chromatography ($p=0.006$). Studies with a low risk of bias showed a significantly lower effect size compared to studies with a higher risk of bias ($p=0.02$). Subgroup analyses by type of depression (depressive symptoms or BD vs. MDD) and results corrected and uncorrected for confounders revealed no significant differences (Table 3).

3.2.2. F2-isoprostanes

The overall effect size (Hedges' g) including all 8 studies on F2-isoprostanes in the meta-analysis was 0.48 (95%

Table 1 Study characteristics of the included studies on the association of 8-OHdG with major depressive disorder, bipolar disorder and depressive symptoms.

8-OHdG	Study design	Sample description	Depressive disorder	Patients N	Controls N	Diagnostic method	Biological specimen	8-OHdG analysis	Confounders ^d				Overall correction confounding	Effect size Hedges' g (95% CI)
									Socio-demographics	Life-style	Somatic disease	AD/mod stabilizers		
Forlenza and Miller (2006)	CC	General population	MDD	62	85	DSM-IV (DISH)	Serum	ELISA	+	+	+	+	+	0.47 (0.14, 0.80)
Irie et al. (2005)	CC	Psychiatric outpatients	MDD	30	60	DSM-IV	Leukocytes	HPLC/ED	—	—	—	—	—	0.76 (0.31, 1.21)
Jorgensen et al. (2013)	CC	Psychiatric department	MDD ^a	26	27	DSM-IV (MINI)	Urine	UPLC/MS	±	±	—	+	+	−0.07 (−0.81, 0.67)
	CC	Psychiatric department	MDD ^b	23	27	ICD-10 psychiatrist	Urine	UPLC/MS	±	±	+	—	+	−0.10 (−0.85, 0.65)
	CC	Psychiatric department	BD ^c	6	27	ICD-10 psychiatrist	Urine	UPLC/MS	±	±	+	—	+	−0.15 (−1.13, 0.82)
Ceylan et al. (2012)	CC	Not reported	BD	36	14	DSM-IV (SCID)	Blood nos	GC/MS	—	—	—	—	—	−0.23 (−0.83, 0.38)
Soeiro-de-Souza et al. (2013)	C	Psychiatric outpatients	BD	50	50	DSM-IV (SCID)	Plasma	ELISA	±	—	—	+	—	1.15 (0.74, 1.57)
Huzayyin et al. (2014)	CC	Specialist psychiatric clinic	BD	14	16	DSM-IV (SADS-L)	Lymphoblasts	ELISA	±	—	—	—	—	0.17 (−0.53, 0.87)
Kupper et al. (2009)	C	Heart failure outpatients	DS	38	72	BDI ≥ 10	Serum	ELISA	—	—	—	—	—	0.14 (−0.25, 0.53)
Wei et al. (2009a)	CC	Oncology department	DS	52	30	HAM-D ≥ 20	Serum	ELISA	±	±	+	—	+	0.81 (0.35, 1.28)
Wei et al. (2009b)	CC	Oncology department	DS	63	43	HAM-D ≥ 20	Serum	ELISA	+	—	+	—	+	0.64 (0.25, 1.04)
Yi et al. (2012)														
Males	C	Municipal workers	DS	105	196	CES-D ≥ 16	Urine	HPLC/ED	±	+	—	+	+	−0.01 (−0.25–0.23)
Females	C	Municipal workers	DS	74	136	CES-D ≥ 16	Urine	HPLC/ED	±	+	+	+	+	−0.09 (−0.37, 0.19)

BD, bipolar disorder; BDI, Beck Depression Inventory; C, cohort; CC, case–control; CES-D, Centre for Epidemiological Studies Depression Scale; DISH, Depression Interview and structured Hamilton Interview; DS, depressive symptoms; DSM-IV, Diagnostic and Statistical Manual of Mental Disorders IV; ELISA, enzyme-linked immunosorbent assay GC/MS, gas chromatography/mass spectrometry HAMD, Hamilton Depression Rating Scale; HPLC/ED, high-performance liquid chromatography/electrochemical detector; ICD-10, International Statistical Classification of Diseases and Related Health Problems; MDD, major depressive disorder; MINI, Mini International Neuropsychiatric Interview; NOS, not otherwise specified; SADS-L, Schedule for Affective Disorders and Schizophrenia, lifetime version; SCID, structured clinical interview for DSM-IV; UPL/MS, ultra performance liquid chromatography/mass spectrometry.

^a Subsample of patients in Jorgensen et al. (2013) described in the article as M-DEP (moderately depressed) patients.

^b Subsample of patients in Jorgensen et al. (2013) described in the article as S-DEP (severely depressed patients unipolar only).

^c Subsample of patients in Jorgensen et al. (2013) described in the article as S-DEP (severely depressed, bipolar only).

^d Socio-demographics include age, sex, socio-economic status (income, education or other), ethnicity. Lifestyle: smoking, alcohol, BMI, physical activity. Somatic disease: presence of (chronic) disease (cardiovascular, infectious, auto-immune or malignancy) that may influence oxidative stress levels. Antidepressant/mood stabilizers: current use. +, confounders is accounted for; —, confounder is not accounted for; ±, some, but not all of the confounders in the category have been accounted for. Studies are defined as corrected (+) if the have accounted for age, sex and at least on life-style variable.

Table 2 Study characteristics of the included studies on the association between F2-isoprostanes and major depressive disorder and depressive symptoms.

F2-isops	Study design	Sample description	Depressive disorder	Patients N	Controls N	Diagnostic method	Biological specimen	F2-isoprostane analysis	Confounders ^a				Overall correction confounding	Effect size Hedges' g (95% CI)
									Socio demographics	Life-style	Somatic disease	AD\mood stabilizers		
Chung et al. (2013)	CC	General population	MDD	18	36	DSM-IV (SCID)	Urine	GC/MS	±	±	+	+	+	1.12 (0.53, 1.72)
Dimopoulos et al. (2008)	CC	General population >60 years	MDD	33	33	DSM-IV psychiatrist	Plasma	ELISA	±	±	+	+	+	1.11 (0.60, 1.63)
Pomara et al. (2012)	CC	General population >60 years	MDD	28	19	DSM-IV (SCID)	CFS	ELISA	±	±	—	—	+	0.88 (0.28, 1.48)
Yager et al. (2010)	CC	General population	MDD	57	74	DSM-IV (DISH)	Serum	ELISA	+	+	+	+	+	0.45 (0.10, 0.80)
Wolkowitz et al. (2011a)	CC	Psychiatric outpatients	MDD	13	14	DSM-IV (SCID)	Plasma	GC/MS	+	+	+	+	+	−0.16 (−0.89, 0.58)
Chung et al. (2009)	CC	Fibromyalgia patients	DS	28	20	CES-D ≥ 16	Urine	GC/MS	—	—	+	—	—	0.37 (−0.20, 0.94)
Milaneschi et al. (2013)														
Males	C	General population	DS	31	996	GDS ≥ 5	Urine	RIA	+	+	+	—	+	0.30 (−0.06, 0.66)
Females	C	70–79 years	DS	52	896	and/or AD use	Urine	RIA	+	+	+	—	+	−0.02 (−0.30, 0.26)
Segal et al. (2012)	CC	SLE patients	DS	33	90	CES-D ≥ 16	Plasma	GC/MS	±	±	—	—	+	0.38 (−0.02, 0.78)

AD, antidepressant; C, cohort; CC, case–control; CES-D, Centre for Epidemiological Studies Depression Scale; CFS, cerebrospinal fluid; DISH, Depression Interview and structured Hamilton Interview; DS, depressive symptoms; DSM-IV, Diagnostic and Statistical Manual of Mental Disorders IV; ELISA, enzyme-linked immunosorbent assay; GC/MS, gas chromatography/mass spectrometry; GDS, geriatric depression scale; HAMD, Hamilton Depression Rating Scale; ICD-10, International Statistical Classification of Diseases and Related Health Problems; MDD, major depressive disorder; RIA, radio immuno assay; SCID, structured clinical interview for DSM-IV; SLE, systemic lupus erythematosus.

^a Socio-demographics include age, sex, socio-economic status (income, education or other), ethnicity. Lifestyle: smoking, alcohol, BMI, physical activity. Somatic disease: presence of (chronic) disease (cardiovascular, infectious, auto-immune or malignancy) that may influence oxidative stress levels. Antidepressant/mood stabilizers: current use. +, confounders is accounted for; —, confounder is not accounted for; ±, some, but not all of the confounders in the category have been accounted for. Studies are defined as corrected (+) if the have accounted for age, sex and at least on life style variable.

Table 3 Results of meta-analyses on the association of 8-OHdG and F2-isoprostanes with depressive symptoms, major depressive disorder and bipolar disorder.

8-OHdG		N studies	Hedges' g	95% CI	p	I ²	95% CI
All studies (depressive symptoms, MDD, BD)		10	0.31	0.06, 0.56	0.01	75%	58–86%
Egger's test for publication bias ^a		10			0.69		
Subgroup analyses		N samples	Hedges' g	95% CI	p ^f	I ²	95% CI
Depression ^b	Depressive symptoms	4	0.27	−0.06, 0.60	0.67	78%	48–91%
	MDD	3	0.37	0.01, 0.74	Reference	48%	0–83%
	BD	4	0.28	−0.48, 1.05	0.83	83%	55–93%
Biological specimen ^c	Urine	2	−0.05	−0.22, 0.12		0%	0–79%
	Plasma/serum	6	0.52	0.18, 0.87	<0.01	75%	43–89%
Laboratory method	Chromatography	4	0.05	−0.20, 0.31		50%	0–79%
	Immunoassay	6	0.58	−0.28, 0.88	<0.01	66%	19–86%
Correction for confounders	Unadjusted	5	0.43	−0.07, 0.92		80%	53–92%
	Adjusted	5	0.23	−0.04, 0.51	0.50	69%	36–85%
Risk of bias score ^d	"Low" risk of bias	3	0.06	−0.15, 0.27		35%	0–74%
	"High" risk of bias	7	0.53	0.19, 0.87	0.02	72%	40–87%
F2-isoprostanes ^e		N studies	Hedges' g	95% CI	p	I ²	95% CI
All studies (depressive symptoms, MDD)		8	0.48	0.19, 0.77	0.001	73%	47–86%
Egger's test for publication bias ^a		8			0.13		
Subgroup analyses		N samples	Hedges' g	95% CI	p ^f	I ²	95% CI
Depression	Depressive symptoms	3	0.24	−0.05, 0.53		56%	0–86%
	MDD	5	0.70	0.28, 1.12	0.08	66%	13–87%
Biological specimen ^c	Urine	3	0.41	−0.06, 0.89		80%	48–92%
	Plasma/serum	4	0.49	0.08, 0.89	0.82	67%	3–89%
Laboratory method	Immunoassay	4	0.51	0.10, 0.92		81%	57–92%
	Chromatography	4	0.45	0.00, 0.90	0.84	61%	0–87%
Correction for confounders		NA					
Risk of bias score ^d	"Low" risk of bias	3	0.51	−0.10, 1.11		74%	40–89%
	"High" risk of bias	5	0.46	0.11, 0.81	0.90	77%	24–93%

BD, bipolar disorder; DS, depressive symptoms; MD, major depressive disorder; NA, not applicable; no or insufficient studies to perform the subgroup analysis.

^a See [Appendix D](#) for funnel plots.

^b [Jorgensen et al. \(2013\)](#) is counted in two categories because it describes stratified analyses of both MDD and BD patients.

^c studies using specimens from others sources were excluded from this analysis ([Irie et al., 2005](#); [Huzayyin et al., 2014](#); [Pomara et al., 2012](#)).

^d See [Appendix C](#) for risk of bias scores.

^e Results of sensitivity analyses, excluding [Pomara et al. \(2012\)](#) with specimens from CSF (cerebrospinal fluid): all studies: Hedges' g 0.44, p 0.004; Egger's test: p 0.22. MDD: Hedges' g 0.66, p between subgroups 0.17; immunoassay: Hedges' g 0.44, p between subgroups 0.98, "High" risk of bias: Hedges' g 0.39, p between subgroups 0.75.

^f p value between subgroups.




Figure 2 (a) Meta-analysis of 8-OHdG in depression (major depressive disorder [MDD], bipolar disorder [BD] and depressive symptoms [DS]) with effect sizes Hedges' g and 95% confidence intervals for the comparison of 8-OHdG levels with controls. (b). Meta-analysis of F2-isoprostanes in depression (major depressive disorder [MDD] and depressive symptoms [DS]) with effect sizes Hedges' g and 95% confidence intervals for the comparison of F2-isoprostane levels with controls.

CI 0.19, 0.77), $I^2 = 73\%$ (95% CI 47–86%), indicating that F2-isoprostanes are significantly increased in depression (Fig. 2b and Table 3). Egger's test was not significant ($p = 0.13$) and the effect size adjusted for publication bias by Duval and Tweedie's trim and fill procedure (no trimmed studies) was unchanged, indicating no significant effects of publication bias (see Appendix D for funnel plot).

A sensitivity analysis was performed excluding one study (Pomara et al., 2012) as this study measured oxidative stress centrally (in cerebrospinal fluid) as opposed to peripheral measurements (blood and urine) used in all others. Exclusion of this study did not affect the overall results (Hedges' g 0.44, $p = 0.004$; Table 3). Subgroup analyses in F2-isoprostanes showed a trend level difference between studies assessing depressive symptoms vs. MDD with the MDD studies showing a larger effect size ($p = 0.08$). Subgroup analyses by biological specimen used, laboratory method for oxidative stress measurement and risk of bias score revealed no significant differences (Table 3). No subgroup analyses were performed based on correction for confounders as only one study was defined as uncorrected.

4. Discussion

This meta-analysis found that both oxidative stress markers, 8-OHdG and F2-isoprostanes, are increased in subjects with depression (major depressive disorder, bipolar disorder and depressive symptoms) compared to controls, with effect sizes in the small to moderate range (Cohen, 1988). In addition, subgroup analyses of the 8-OHdG studies revealed that some of the variation in the results may be explained by the type of the biological specimen and/or the laboratory method for oxidative stress measurement, and that studies with lower risk of bias reported significantly smaller effect sizes. Findings for F2-isoprostanes however, did not differ when analyzed by type of depression, biological specimen, laboratory method or quality.

The results indicate that depression is associated with increased oxidative damage to DNA and lipids (reflected by 8-OHdG and F2-isoprostanes respectively). These findings are in line with previous meta-analytic studies that reported decreased anti-oxidants and anti-oxidant enzymes in unipolar depression, and increased oxidative stress in both uni- and bipolar depression (Andreazza et al., 2008; Palta et al., 2014). Previously DNA damage in bipolar disorder has been reported measured by increased levels of DNA strand breakage with the comet assay (Andreazza et al., 2007). Oxidative damage to proteins determined by the protein carbonyl assay has not yet been as widely studied, but increased levels have been reported in MDD and BD (Kapczynski et al., 2011; Magalhaes et al., 2012), however not consistently (Andreazza et al., 2009; Gubert et al., 2013). The oxidative stress markers used in this meta-analysis are among the most robust markers of oxidative stress currently available and have recognized roles in the pathophysiology various somatic diseases such as cardiovascular disease, cancer and diabetes (Dalle-Donne et al., 2006; Valavanidis et al., 2009). 8-OHdG is a biologically important mutagenic DNA lesion, while F2-isoprostanes are known to be increased in atherosclerotic lesions and may also be biologically active in the pathogenesis of atherosclerosis (Ho et al., 2013). It should be noted that although both markers reflect oxidative damage they are not necessarily associated with each other. One study included in our meta-analysis (Yager et al., 2010) found no correlation between the two, suggesting that 8-OHdG and F2-isoprostanes reflect specific aspects of oxidative imbalance.

Although the main findings of this study are unlikely to be greatly influenced by publication bias, considerable heterogeneity between studies was found for both 8-OHdG and F2-isoprostanes. This heterogeneity may be explained by several factors. In the studies that examined 8-OHdG, patients in three bipolar disorder studies (Ceylan et al., 2012; Soeiro-de-Souza et al., 2013; Huzayyin et al., 2014) were not all currently depressed, but some were in a manic or euthymic state. This may account for the fact that two of these studies' results have the highest and lowest effect sizes included in the meta-analysis. In addition, the subgroup analyses in 8-OHdG studies demonstrated significant differences between studies based on risk of bias, with the studies with lower risk of bias finding smaller effects. There was also a significant difference in effect size in the subgroup analyses by biological specimen (higher in

plasma/serum than in urine) and by laboratory method used to measure oxidative stress (higher in immuno-assays than chromatography). With only one exception however, all the measurements using chromatography were done in urine samples, whereas all the immuno-assays for the measurement of 8-OHdG were done in plasma or serum. Therefore, it cannot be determined whether this difference is based on the biological specimen or laboratory technique. 8-OHdG levels determined by immuno-assays are higher than those measured by chromatography, with the latter considered the gold standard. The correlation between the two methods is generally high (Yoshida et al., 2002) and so may not necessarily affect the strength of the association with depression. It has been reported that urinary 8-OHdG levels are more stable than those in plasma or serum, and therefore possibly more reliable (Matsumoto et al., 2008). Although the subgroup analyses may help to explain the heterogeneity observed, the findings should be interpreted with considerable caution, as the number of studies in the subgroups is small.

This meta-analysis confirms oxidative stress markers are increased in subjects with depression in cross-sectional studies, but the underlying mechanisms explaining this link need to be examined further. Many behavioural factors that are related to increased exposure to ROS (smoking, alcohol use, overweight, physical activity) (Maritim et al., 2003; Dalle-Donne et al., 2006; Valko et al., 2007) are also associated with depression (Glassman et al., 1990; Abu-Omar et al., 2004; Sullivan et al., 2005; Luppino et al., 2010). It cannot be ruled out that the association is partially driven by these or other lifestyle confounders. The majority of the included studies took some, but few took all of these potential confounders into account. The results of studies that did correct for confounders did not differ significantly from studies that did not, strengthening the observation that an association between depression and oxidative stress is present independent of these life-style factors. The observed increased levels of oxidative stress in depression might be understood within the concepts of allostasis and allostatic load. The former refers to the physiological adaptation to physical, psychological, social and environmental stressors. The latter refers to the physical "wear and tear" induced by prolonged exposure to the stress response (McEwen and Wingfield, 2003). Depression has been found to be associated with increased "wear and tear" or accelerated cellular ageing reflected by decreased telomere length in depressed patients (Verhoeven et al., 2013). Dysregulations in the major stress systems (hypothalamic-pituitary-adrenal axis activity, autonomic nervous system function and inflammatory functions) have been demonstrated in depression (Penninx et al., 2013) and these could be contributing to increased oxidative stress. Oxidative stress is closely related to the inflammatory pathway in particular. Pro-inflammatory cytokines are produced in reaction to oxidative stress and oxidative stress in turn amplifies the inflammatory response. High cortisol levels have been associated with increased levels of oxidative damage (Joergensen et al., 2011). The damage caused by this allostatic load experienced during mood episodes is hypothesized to render an individual more vulnerable to developing a following episode and at higher risk to develop somatic disease (Maes et al., 2011; Grande et al., 2012). The brain is particularly vulnerable to

oxidative damage due to its high oxygen consumption and low anti-oxidants defences. There is evidence from post-mortem studies suggesting that in depression oxidative stress is increased (Wang et al., 2009; Che et al., 2010; Michel et al., 2012) and anti-oxidants are decreased (Gawryluk et al., 2011) in the brain.

This meta-analysis' strength lies in the focus on robust oxidative stress markers and in the comprehensive search that was conducted to identify all studies on the association of 8-OHdG and F2-isoprostanes with depression. Through additional data requests the number of studies eligible for inclusion was increased, and where possible adjusted data was obtained, mitigating the effects of possible publication bias and increasing the reliability of the overall results.

There are also a number of important limitations. As is apparent from the number of included studies and subjects the evidence on 8-OHdG and F2-isoprostanes in depression is limited. The studies that were excluded because they only reported on depressive symptoms as a continuous measure (Irie et al., 2001, 2003; Iida et al., 2011) all found a positive association. Therefore it is most likely that they would not have altered the overall result. Furthermore, the covariates used in the adjusted analyses varied widely. Therefore it was not possible to assess the effects of adjustment on the results satisfactorily. Information on diet and anti-oxidant supplement use was not available for most studies and therefore it cannot be ruled out this factor may have had an effect. In addition, the use of antidepressants or mood stabilizers could also be sources of confounding that were often not investigated. There is some evidence to suggest that antidepressants, lithium and other mood stabilizers protect against oxidative stress (Behr et al., 2012; Khairova et al., 2012; de Sousa et al., 2014). Studies addressing their effects on F2-isoprostanes however found an increase (Chung et al., 2013) or no effect on oxidative stress levels (Rawdin et al., 2013).

In conclusion, the finding that both 8-OHdG and F2-isoprostanes are increased in depression strongly suggests that depression is accompanied by increased oxidative damage. This finding supports the hypothesis that increased metabolic stress is present in depression, which could potentially contribute to the higher risk of somatic morbidity and mortality in sufferers. There is a need for future larger scale studies on oxidative stress in depression, in which the role of treatment effects should be addressed.

Role of the funding sources

Catherine N. Black and Brenda W.J.H. Penninx are supported through an NWO-VICI grant (number 91811602). The grant provider had no involvement in the study design; in the data collection, analysis or interpretation of data; in the writing of the report; or in the decision to submit the article for publication.

Conflict of interest

All authors confirm they have no financial or other conflicts of interest to disclose.

Acknowledgements

We would like to express our thanks and acknowledge the contribution of all the following authors who provided us with additional data for the meta-analysis. Cecilia P. Chung, MD, MPH, Assistant Professor of Medicine, Division of Rheumatology, Vanderbilt University Medical Center. Anders Jørgensen, MD, PhD, Psychiatric Centre Copenhagen, University Hospital of Copenhagen. Yuri Milaneschi PhD, Department of Psychiatry and EMGO Institute for Health and Care Research, VU University Medical Center/GGZ inGeest, Amsterdam, The Netherlands & Longitudinal Studies Section, Clinical Research Branch, National Institute on Aging, Baltimore, Maryland, United States of America. Gregory Miller, PhD Professor, Departments of Psychology and Medical Social Sciences Faculty Fellow, Cells to Society: The Center on Social Disparities and Health at The Institute for Policy Research, Northwestern University. Barbara M. Segal, MD, University of Minnesota, Minneapolis, MN, USA. C. Michael Stein, MBChB, Professor of Medicine, Professor of Pharmacology, Vanderbilt University Medical Center. Owen M. Wolkowitz, MD, Professor of Psychiatry, University of California San Francisco (UCSF) School of Medicine. Siyan Yi, MD, MHSc, PhD, Research Fellow, Asia Health Policy, The Walter H. Shorenstein Asia-Pacific Research Center, Freeman Spogli Institute for International Studies Stanford University. Tetsuya Mizoue, Department of Epidemiology and Prevention Clinical Research Center, National Center for Global Health and Medicine.

Appendices A, B, C, D. Supplementary data

Supplementary data associated with this article can be found, in the online version, at <http://dx.doi.org/10.1016/j.psyneuen.2014.09.025>.

References

- Abu-Omar, K., Rutten, A., Lehtinen, V., 2004. Mental health and physical activity in the European Union. *Soz. Präventivmed.* 49, 301–309.
- Andreazza, A.C., Kauer-Sant'anna, M., Frey, B.N., Bond, D.J., Kapczinski, F., Young, L.T., Yatham, L.N., 2008. Oxidative stress markers in bipolar disorder: a meta-analysis. *J. Affect. Disord.* 111, 135–144.
- Andreazza, A.C., Frey, B.N., Erdtmann, B., Salvador, M., Rombaldi, F., Santin, A., Goncalves, C.A., Kapczinski, F., 2007. DNA damage in bipolar disorder. *Psychiatry Res.* 153, 27–32.
- Andreazza, A.C., Kapczinski, F., Kauer-Sant'anna, M., Walz, J.C., Bond, D.J., Goncalves, C.A., Young, L.T., Yatham, L.N., 2009. 3-Nitrotyrosine and glutathione antioxidant system in patients in the early and late stages of bipolar disorder. *J. Psychiatry Neurosci.* 34, 263–271.
- Barnes, D.E., Alexopoulos, G.S., Lopez, O.L., Williamson, J.D., Yaffe, K., 2006. Depressive symptoms, vascular disease, and mild cognitive impairment: findings from the Cardiovascular Health Study. *Arch. Gen. Psychiatry* 63, 273–279.
- Behr, G.A., Moreira, J.C.F., Frey, B.N., 2012. Preclinical and clinical evidence of antioxidant effects of antidepressant agents: implications for the pathophysiology of major depressive disorder. *Oxid. Med. Cell Longev.* 2012, 609421.
- Bijl, R.V., Ravelli, A., 2000. Psychiatric morbidity, service use, and need for care in the general population: results of The Netherlands Mental Health Survey and Incidence Study. *Am. J. Public Health* 90, 602–607.
- Cepnija, M., Derek, L., Unic, A., Blazev, M., Fistic, M., Kozaric-Kovacic, D., Franic, M., Romic, Z., 2011. Oxidative stress markers in patients with post-traumatic stress disorder. *Coll. Antropol.* 35, 1155–1160.
- Ceylan, D., Tuna, G., Kirkali, G., Dizdaronullu, M., Tunca, Z., Yesilyurt, S., Ozerdem, A., 2012. DNA damage in bipolar disorder: a quantitative assessment by using gas chromatography–isotope dilution mass spectrometry (GC–MS) null. *Bipolar Disord.* 14 (Suppl. 1), 66.
- Ceylan, D., Tuna, G., Kirkali, G., Tunca, Z., Dizdaronullu, M., Can, G., Arat, E., Ozerdem, A., 2013. Increased oxidative base damage to DNA in bipolar disorder: a cross-sectional case control study. *Bipolar Disord.* 15 (1), 55.
- Che, Y., Wang, J.F., Shao, L., Young, T., 2010. Oxidative damage to RNA but not DNA in the hippocampus of patients with major mental illness. *J. Psychiatry Neurosci.* 35, 296–302.
- Chida, Y., Hamer, M., Wardle, J., Steptoe, A., 2008. Do stress-related psychosocial factors contribute to cancer incidence and survival? *Nat. Clin. Pract. Oncol.* 5, 466–475.
- Chung, C.P., Schmidt, D., Stein, C.M., Morrow, J.D., Salomon, R.M., 2013. Increased oxidative stress in patients with depression and its relationship to treatment. *Psychiatry Res.* 206, 213–216.
- Chung, C.P., Titova, D., Oeser, A., Randels, M., Avalos, I., Milne, G.L., Morrow, J.D., Stein, C.M., 2009. Oxidative stress in fibromyalgia and its relationship to symptoms. *Clin. Rheumatol.* 28, 435–438.
- Cohen, J., 1988. *Statistical Power Analysis for the Behavioral Sciences*. Lawrence Erlbaum, Hillsdale, NJ.
- Cuijpers, P., Vogelzangs, N., Twisk, J., Kleiboer, A., Li, J., Penninx, B.W., 2014. Comprehensive meta-analysis of excess mortality in depression in the general community versus patients with specific illnesses. *Am. J. Psychiatry* 171, 453–462.
- Dalle-Donne, I., Rossi, R., Colombo, R., Giustarini, D., Milzani, A., 2006. Biomarkers of oxidative damage in human disease. *Clin. Chem.* 52, 601–623.
- de Sousa, R.T., Zarate, C.A.J., Zanetti, M.V., Costa, A.C., Talib, L.L., Gattaz, W.F., Haddo-Vieira, R., 2014. Oxidative stress in early stage bipolar disorder and the association with response to lithium. *J. Psychiatr. Res.* 50, 36–41.
- Dimopoulos, N., Piperi, C., Psarra, V., Lea, R.W., Kalofoutis, A., 2008. Increased plasma levels of 8-iso-PGF₂α and IL-6 in an elderly population with depression. *Psychiatry Res.* 161, 59–66.
- Duval, S., Tweedie, R., 2000. Trim and fill: a simple funnel-plot-based method of testing and adjusting for publication bias in meta-analysis. *Biometrics* 56, 455–463.
- Forlenza, M.J., Miller, G.E., 2006. Increased serum levels of 8-hydroxy-2'-deoxyguanosine in clinical depression. *Psychosom. Med.* 68, 1–7.
- Freund-Levi, Y., Cederholm, T., Basu, S., 2011. Oxidative stress and cox-mediated inflammation in patients with Alzheimer's disease and effects of N-3 fatty acid supplementation the omegaAD study. *Clin. Nutr.* 6 (Suppl. 1), 9.
- Gawryluk, J.W., Wang, J.F., Andreazza, A.C., Shao, L., Young, L.T., 2011. Decreased levels of glutathione, the major brain antioxidant, in post-mortem prefrontal cortex from patients with psychiatric disorders. *Int. J. Neuropsychopharmacol.* 14, 123–130.
- Glassman, A.H., Helzer, J.E., Covey, L.S., Cottler, L.B., Stetner, F., Tipp, J.E., Johnson, J., 1990. Smoking, smoking cessation, and major depression. *J. Am. Med. Assoc.* 264, 1546–1549.
- Grande, I., Magalhaes, P.V., Kunz, M., Vieta, E., Kapczinski, F., 2012. Mediators of allostasis and systemic toxicity in bipolar disorder. *Physiol. Behav.* 106, 46–50.
- Gubert, C., Stertz, L., Pfaffenseller, B., Panizzutti, B.S., Rezin, G.T., Massuda, R., Streck, E.L., Gama, C.S., Kapczinski, F., Kunz,

- M., 2013. Mitochondrial activity and oxidative stress markers in peripheral blood mononuclear cells of patients with bipolar disorder, schizophrenia, and healthy subjects. *J. Psychiatr. Res.* 47, 1396–1402.
- Hayden, J.A., Cote, P., Bombardier, C., 2006. Evaluation of the quality of prognosis studies in systematic reviews. *Ann. Intern. Med.* 144, 427–437.
- Higgins, J.P.T., Thompson, S.G., Deeks, J.J., Altman, D.G., 2003. Measuring inconsistency in meta-analyses. *Br. Med. J.* 327, 557–560.
- Ho, E., Karimi Galougahi, K., Liu, C., Bhindi, R., Figtree, G., 2013. Biological markers of oxidative stress: applications to cardiovascular research and practice. *Redox Biol.* 1, 483–491.
- Huzayyin, A.A., Andreadza, A.C., Turecki, G., Cruceanu, C., Rouleau, G.A., Alda, M., Young, L.T., 2014. Decreased global methylation in patients with bipolar disorder who respond to lithium. *Int. J. Neuropsychopharmacol.* 17, 561–569.
- Iida, T., Chikamura, C., Inoue, K., Ito, Y., Ishikawa, H., Teradaira, R., Ono, Y., 2011. Association of STAI and SDS scores with 8-hydroxydeoxyguanosine and serotonin levels in young women with depressive symptoms. *J. Neuropsychiatry Clin. Neurosci.* 23, E10.
- Ioannidis, J.P.A., Patsopoulos, N.A., Evangelou, E., 2007. Uncertainty in heterogeneity estimates in meta-analyses. *Br. Med. J.* 335, 914–916.
- Irie, M., Asami, S., Nagata, S., Ikeda, M., Miyata, M., Kasai, H., 2001. Psychosocial factors as a potential trigger of oxidative DNA damage in human leukocytes. *Jpn. J. Cancer Res.* 92, 367–376.
- Irie, M., Asami, S., Ikeda, M., Kasai, H., 2003. Depressive state relates to female oxidative DNA damage via neutrophil activation. *Biochem. Biophys. Res. Commun.* 311, 1014–1018.
- Irie, M., Asami, S., Nagata, S., Miyata, M., Kasai, H., 2002. Psychological mediation of a type of oxidative DNA damage, 8-hydroxydeoxyguanosine, in peripheral blood leukocytes of non-smoking and non-drinking workers. *Psychother. Psychosom.* 71, 90–96.
- Irie, M., Miyata, M., Kasai, H., 2005. Depression and possible cancer risk due to oxidative DNA damage. *J. Psychiatr. Res.* 39, 553–560.
- Janicki-Deverts, D., Cohen, S., Matthews, K.A., Gross, M.D., Jacobs, D.R.J., 2009. Socioeconomic status, antioxidant micronutrients, and correlates of oxidative damage: the Coronary Artery Risk Development in Young Adults (CARDIA) study. *Psychosom. Med.* 71, 541–548.
- Joergensen, A., Broedbaek, K., Weimann, A., Semba, R.D., Ferrucci, L., Joergensen, M.B., Poulsen, H.E., 2011. Association between urinary excretion of cortisol and markers of oxidatively damaged DNA and RNA in humans. *PLoS ONE* 6, e20795, <http://dx.doi.org/10.1371/journal.pone.0020795>.
- Jorgensen, A., Krogh, J., Miskowiak, K., Bolwig, T.G., Kessing, L.V., Fink-Jensen, A., Nordentoft, M., Henriksen, T., Weimann, A., Poulsen, H.E., Jorgensen, M.B., 2013. Systemic oxidatively generated DNA/RNA damage in clinical depression: associations to symptom severity and response to electroconvulsive therapy. *J. Affect. Disord.* 149, 355–362.
- Kapczinski, F., Dal-Pizzol, F., Teixeira, A.L., Magalhaes, P.V.S., Kauer-Sant'anna, M., Klamt, F., Moreira, J.C., de Bittencourt Pasquali, M.A., Fries, G.R., Quevedo, J., Gama, C.S., Post, R., 2011. Peripheral biomarkers and illness activity in bipolar disorder. *J. Psychiatr. Res.* 45, 156–161.
- Kessler, R.C., Berglund, P., Demler, O., Jin, R., Koretz, D., Merikangas, K.R., Rush, A.J., Walters, E.E., Wang, P.S., 2003. The epidemiology of major depressive disorder: results from the National Comorbidity Survey Replication (NCS-R). *J. Am. Med. Assoc.* 289, 3095–3105.
- Khairouva, R., Pawar, R., Salvatore, G., Jurueña, M.F., de Sousa, R.T., Soeiro-de-Souza, M.G., Salvador, M., Zarate, C.A., Gattaz, W.F., Hado-Vieira, R., 2012. Effects of lithium on oxidative stress parameters in healthy subjects. *Mol. Med. Rep.* 5, 680–682.
- Kupper, N., Gidron, Y., Winter, J., Denollet, J., 2009. Association between type D personality, depression, and oxidative stress in patients with chronic heart failure. *Psychosom. Med.* 71, 973–980.
- Luppino, F.S., de Wit, L.M., Bouvy, P.F., Stijnen, T., Cuijpers, P., Penninx, B.W.J.H., Zitman, F.G., 2010. Overweight, obesity, and depression: a systematic review and meta-analysis of longitudinal studies. *Arch. Gen. Psychiatry* 67, 220–229.
- Maes, M., Galecki, P., Chang, Y.S., Berk, M., 2011. A review on the oxidative and nitrosative stress (O&NS) pathways in major depression and their possible contribution to the (neuro)degenerative processes in that illness. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 35, 676–692.
- Maes, M., Mihaylova, I., Kubera, M., Uytterhoeven, M., Vrydags, N., Bosmans, E., 2009. Increased 8-hydroxy-deoxyguanosine, a marker of oxidative damage to DNA, in major depression and myalgic encephalomyelitis/chronic fatigue syndrome. *Neuroendocrinol. Lett.* 30, 715–722.
- Magalhaes, P.V.S., Jansen, K., Pinheiro, R.T., Colpo, G.D., da Motta, L.L., Klamt, F., da Silva, R.A., Kapczinski, F., 2012. Peripheral oxidative damage in early-stage mood disorders: a nested population-based case-control study. *Int. J. Neuropsychopharmacol.* 15, 1043–1050.
- Maritim, A.C., Sanders, R.A., Watkins, J.B., 2003. Diabetes, oxidative stress, and antioxidants: a review. *J. Biochem. Mol. Toxicol.* 17, 24–38.
- Matsumoto, Y., Ogawa, Y., Yoshida, R., Shimamori, A., Kasai, H., Ohta, H., 2008. The stability of the oxidative stress marker, urinary 8-hydroxy-2'-deoxyguanosine (8-OHdG), when stored at room temperature. *J. Occup. Health* 50, 366–372.
- McEwen, B.S., Wingfield, J.C., 2003. The concept of allostasis in biology and biomedicine. *Horm. Behav.* 43, 2–15.
- Meagher, E.A., FitzGerald, G.A., 2000. Indices of lipid peroxidation in vivo: strengths and limitations. *Free Rad. Biol. Med.* 28, 1745–1750.
- Mezuk, B., Eaton, W.W., Albrecht, S., Golden, S.H., 2008. Depression and type 2 diabetes over the lifespan: a meta-analysis. *Diabetes Care* 31, 2383–2390.
- Michel, T.M., Pulschen, D., Thome, J., 2012. The role of oxidative stress in depressive disorders. *Curr. Pharm. Des.* 18, 5890–5899.
- Milaneschi, Y., Cesari, M., Simonsick, E.M., Vogelzangs, N., Kanaya, A.M., Yaffe, K., Patrignani, P., Metti, A., Kritchevsky, S.B., Pahor, M., Ferrucci, L., Penninx, B.W.J.H., 2013. Lipid peroxidation and depressed mood in community-dwelling older men and women. *PLoS ONE* 8, e65406.
- Nicholson, A., Kuper, H., Hemingway, H., 2006. Depression as an aetiological and prognostic factor in coronary heart disease: a meta-analysis of 6362 events among 146 538 participants in 54 observational studies. *Eur. Heart J.* 27, 2763–2774.
- Niki, E., 2014. Biomarkers of lipid peroxidation in clinical material. *Biochim. Biophys. Acta* 1840, 809–817.
- Palta, P., Samuel, L.J., Miller, E.R., Szanton, S.L., 2014. Depression and oxidative stress: results from a meta-analysis of observational studies. *Psychosom. Med.* 76, 12–19.
- Pandya, C.D., Howell, K.R., Pillai, A., 2013. Antioxidants as potential therapeutics for neuropsychiatric disorders. *Prog. Neuropsychopharmacol. Biol. Psychiatry* 46, 214–223.
- Penninx, B.W.J.H., Milaneschi, Y., Lamers, F., Vogelzangs, N., 2013. Understanding the somatic consequences of depression: biological mechanisms and the role of depression symptom profile. *BMC Med.* 11, 129.
- Pomara, N., Bruno, D., Sarreal, A.S., Hernando, R.T., Nierenberg, J., Petkova, E., Sidtis, J.J., Wisniewski, T.M., Mehta, P.D., Pratico, D., Zetterberg, H., Blennow, K., 2012. Lower CSF amyloid beta peptides and higher F2-isoprostanes in cognitively

- intact elderly individuals with major depressive disorder. *Am. J. Psychiatry* 169, 523–530.
- Rawdin, B.J., Mellon, S.H., Dhabhar, F.S., Epel, E.S., Puterman, E., Su, Y., Burke, H.M., Reus, V.I., Rosser, R., Hamilton, S.P., Nelson, J.C., Wolkowitz, O.M., 2013. *Dysregulated relationship of inflammation and oxidative stress in major depression. Brain Behav. Immun.* 31, 143–152.
- Segal, B.M., Thomas, W., Zhu, X., Diebes, A., McElvain, G., Baechler, E., Gross, M., 2012. *Oxidative stress and fatigue in systemic lupus erythematosus. Lupus* 21, 984–992.
- Soeiro-de-Souza, M.G., Andreazza, A.C., Carvalho, A.F., Hado-Vieira, R., Young, L.T., Moreno, R.A., 2013. *Number of manic episodes is associated with elevated DNA oxidation in bipolar I disorder. Int. J. Neuropsychopharmacol.* 16, 1505–1512.
- Sterne, J.A., Egger, M., 2001. *Funnel plots for detecting bias in meta-analysis: guidelines on choice of axis. J. Clin. Epidemiol.* 54, 1046–1055.
- Sullivan, L.E., Fiellin, D.A., O'Connor, P.G., 2005. *The prevalence and impact of alcohol problems in major depression: a systematic review. Am. J. Med.* 118, 330–341.
- Valavanidis, A., Vlachogianni, T., Fiotakis, C., 2009. *8-Hydroxy-2'-deoxyguanosine (8-OHdG): a critical biomarker of oxidative stress and carcinogenesis. J. Environ. Sci. Health C: Environ. Carcinog. Ecotoxicol. Rev.* 27, 120–139.
- Valko, M., Leibfritz, D., Moncol, J., Cronin, M.T.D., Mazur, M., Telser, J., 2007. *Free radicals and antioxidants in normal physiological functions and human disease. Int. J. Biochem. Cell Biol.* 39, 44–84.
- Verhoeven, J., Révész, D., Epel, E., Lin, J., Wolkowitz, O., Penninx, B., 2013. *Major depressive disorder and accelerated cellular aging: results from a large psychiatric cohort study. Mol. Psychiatry*, <http://dx.doi.org/10.1038/mp.2013.151>.
- Vos, T., Flaxman, A.D., Naghavi, M., Lozano, R., Michaud, C., Ezzati, M., Shibuya, K., Salomon, J.A., Abdalla, S., Aboyans, V., Abraham, J., Ackerman, I., Aggarwal, R., Ahn, S.Y., Ali, M.K., Alvarado, M., Anderson, H.R., Anderson, L.M., Andrews, K.G., Atkinson, C., Baddour, L.M., Bahalim, A.N., Barker-Collo, S., Barrero, L.H., Bartels, D.H., Basanez, M.G., Baxter, A., Bell, M.L., Benjamin, E.J., Bennett, D., Bernabe, E., Bhalla, K., Bhandari, B., Bikbov, B., Bin Abdulhak, A., Birbeck, G., Black, J.A., Blencowe, H., Blore, J.D., Blyth, F., Bolliger, I., Bonaventure, A., Boufous, S., Bourne, B., Boussinesq, M., Braithwaite, T., Brayne, C., Bridgett, L., Brooker, S., Brooks, P., Brugha, T.S., Bryan-Hancock, C., Bucello, C., Buchbinder, R., Buckle, G., Budke, C.M., Burch, M., Burney, P., Burstein, R., Calabria, B., Campbell, B., Canter, C.E., Carabin, H., Carapetis, J., Carmona, L., Cella, C., Charlson, F., Chen, H., Cheng, A.T.-A., Chou, D., Chugh, S.S., Coffeng, L.E., Colan, S.D., Colquhoun, S., Colson, K.E., Condon, J., Connor, M.D., Cooper, L.T., Corriere, M., Cortinovis, M., de Vaccaro, K.C., Couser, W., Cowie, B.C., Criqui, M.H., Cross, M., Dabhadkar, K.C., Dahiya, M., Dahodwala, N., msere-Derry, J., Danaei, G., Davis, A., De Leo, D., Degenhardt, L., Dellavalle, R., Delossantos, A., Denenberg, J., Derrett, S., Des Jarlais, D.C., Dharmaratne, S.D., Dherani, M., az-Torne, C., Dolk, H., Dorsey, E.R., Driscoll, T., Duber, H., Ebel, B., Edmond, K., Elbaz, A., Ali, S.E., Erskine, H., Erwin, P.J., Espindola, P., Ewoigbokhan, S.E., Farzadfar, F., Feigin, V., Felson, D.T., Ferrari, A., Ferri, C.P., Fevre, E.M., Finucane, M.M., Flaxman, S., Flood, L., Foreman, K., Forouzanfar, M.H., Fowkes, F.G., Franklin, R., Fransen, M., Freeman, M.K., Gabbe, B.J., Gabriel, S.E., Gakidou, E., Ganatra, H.A., Garcia, B., Gaspari, F., Gillum, R.F., Gmel, G., Gosselin, R., Grainger, R., Groeger, J., Guillemin, F., Gunnell,
- D., Gupta, R., Haagsma, J., Hagan, H., Halasa, Y.A., Hall, W., Haring, D., Haro, J.M., Harrison, J.E., Havmoeller, R., Hay, R.J., Higashi, H., Hill, C., Hoen, B., Hoffman, H., Hotez, P.J., Hoy, D., Huang, J.J., Ibeanusi, S.E., Jacobsen, K.H., James, S.L., Jarvis, D., Jasrasaria, R., Jayaraman, S., Johns, N., Jonas, J.B., Karthikeyan, G., Kassebaum, N., Kawakami, N., Keren, A., Khoo, J.P., King, C.H., Knowlton, L.M., Kobusingye, O., Koranteng, A., Krishnamurthi, R., Lalloo, R., Laslett, L.L., Lathlean, T., Leasher, J.L., Lee, Y.Y., Leigh, J., Lim, S.S., Limb, E., Lin, J.K., Lipnick, M., Lipshultz, S.E., Liu, W., Loane, M., Ohno, S.L., Lyons, R., Ma, J., Mabweijano, J., MacIntyre, M.F., Malekzadeh, R., Mallinger, L., Manivannan, S., Marcenes, W., March, L., Margolis, D.J., Marks, G.B., Marks, R., Matsumori, A., Matzopoulos, R., Mayosi, B.M., McAnulty, J.H., McDermott, M.M., McGill, N., McGrath, J., Medina-Mora, M.E., Meltzer, M., Mensah, G.A., Merriman, T.R., Meyer, A.C., Miglioli, V., Miller, M., Miller, T.R., Mitchell, P.B., Mocumbi, A.O., Moffitt, T.E., Mokdad, A.A., Monasta, L., Monticco, M., Moradi-Lakeh, M., Moran, A., Morawska, L., Mori, R., Murdoch, M.E., Mwaniki, M.K., Naidoo, K., Nair, M.N., Naldi, L., Narayan, K.M.V., Nelson, P.K., Nelson, R.G., Nevitt, M.C., Newton, C.R., Nolte, S., Norman, P., Norman, R., O'Donnell, M., O'Hanlon, S., Olives, C., Omer, S.B., Ortblad, K., Osborne, R., Ozgediz, D., Page, A., Pahari, B., Pandian, J.D., Rivero, A.P., Patten, S.B., Pearce, N., Padilla, R.P., Perez-Ruiz, F., 2012. *Years lived with disability (YLDs) for 1160 sequelae of 289 diseases and injuries 1990–2010: a systematic analysis for the Global Burden of Disease Study 2010. Lancet* 380, 2163–2196.
- Wang, J.F., Shao, L., Sun, X., Young, L.T., 2009. *Increased oxidative stress in the anterior cingulate cortex of subjects with bipolar disorder and schizophrenia. Bipolar Disord.* 11, 523–529.
- Wei, Y.C., Zhou, F.L., He, D.L., Bai, J.R., Ding, H., Wang, X.Y., Nan, K.J., 2009a. *Oxidative stress in depressive patients with gastric adenocarcinoma. Int. J. Neuropsychopharmacol.* 12, 1089–1096.
- Wei, Y.C., Zhou, F.L., He, D.L., Bai, J.R., Hui, L.y., Wang, X.Y., Nan, K.J., 2009b. *The level of oxidative stress and the expression of genes involved in DNA-damage signaling pathways in depressive patients with colorectal carcinoma. J. Psychosom. Res.* 66, 259–266.
- Wolkowitz, O.M., Mellon, S.H., Epel, E.S., Lin, J., Dhabhar, F.S., Su, Y., Reus, V.I., Rosser, R., Burke, H.M., Kupferman, E., Compagnone, M., Nelson, J.C., Blackburn, E.H., 2011a. *Leukocyte telomere length in major depression: correlations with chronicity, inflammation and oxidative stress – preliminary findings. PLoS ONE* 6, e17837.
- Wolkowitz, O.M., Reus, V.I., Mellon, S.H., 2011b. *Of sound mind and body: depression, disease, and accelerated aging. Dialogues Clin. Neurosci.* 13, 25–39.
- Wu, I.C., Shiesh, S.C., Kuo, P.H., Lin, X.Z., 2009. *High oxidative stress is correlated with frailty in elderly Chinese. J. Am. Geriatr. Soc.* 57, 1666–1671.
- Yager, S., Fortenza, M.J., Miller, G.E., 2010. *Depression and oxidative damage to lipids. Psychoneuroendocrinology* 35, 1356–1362.
- Yi, S., Nanri, A., Matsushita, Y., Kasai, H., Kawai, K., Mizoue, T., 2012. *Depressive symptoms and oxidative DNA damage in Japanese municipal employees. Psychiatry Res.* 200, 318–322.
- Yoshida, R., Ogawa, Y., Kasai, H., 2002. *Urinary 8-oxo-7,8-dihydro-2'-deoxyguanosine values measured by an ELISA correlated well with measurements by high-performance liquid chromatography with electrochemical detection. Cancer Epidemiol. Biomarkers Prev.* 11, 1076–1081.