



Invariance properties of coHochschild homology

Kathryn Hess^a, Brooke Shipley^{b,*}

^a EPFL SV BMI UPHESS, Station 8, CH-1015 Lausanne, Switzerland

^b Department of Mathematics, Statistics, and Computer Science, University of Illinois at Chicago, 508 SEO m/c 249, 851 S. Morgan Street, Chicago, IL, 60607-7045, USA



ARTICLE INFO

Article history:

Received 17 June 2019

Received in revised form 4 June 2020

Available online 15 July 2020

Communicated by A. Solotar

MSC:

Primary: 19D55; 16T15; secondary:

16E40; 55U35; 55P43

Keywords:

Topological Hochschild homology

Coalgebras

ABSTRACT

The notion of Hochschild homology of a dg algebra admits a natural dualization, the coHochschild homology of a dg coalgebra, introduced in [23] as a tool to study free loop spaces. In this article we prove “agreement” for coHochschild homology, i.e., that the coHochschild homology of a dg coalgebra C is isomorphic to the Hochschild homology of the dg category of appropriately compact C -comodules, from which Morita invariance of coHochschild homology follows. Generalizing the dg case, we define the topological coHochschild homology (coTHH) of coalgebra spectra, of which suspension spectra are the canonical examples, and show that coTHH of the suspension spectrum of a space X is equivalent to the suspension spectrum of the free loop space on X , as long as X is a nice enough space (for example, simply connected.) Based on this result and on a Quillen equivalence established in [24], we prove that “agreement” holds for coTHH as well.

© 2020 Elsevier B.V. All rights reserved.

Contents

1. Introduction	2
2. CoHochschild homology for chain coalgebras	3
2.1. The dg cobar construction and its extensions	4
2.2. Properties of the dg coHochschild construction	8
3. Topological coHochschild homology of spectra	17
3.1. The general theory	17
3.2. coTHH of suspension spectra	19
3.3. Cobar, Bar, and loop spaces	21
Appendix A. Total complexes of cosimplicial suspension spectra	23
References	26

* Corresponding author.

E-mail addresses: kathryn.hess@epfl.ch (K. Hess), shipleyb@uic.edu (B. Shipley).

1. Introduction

For any commutative ring \mathbb{k} , the classical definition of Hochschild homology of \mathbb{k} -algebras [33] admits a straightforward extension to differential graded (dg) \mathbb{k} -algebras. In [37] McCarthy extended the definition of Hochschild homology in another direction, to \mathbb{k} -exact categories, seen as \mathbb{k} -algebras with many objects. As Keller showed in [29], there is a common refinement of these two extended definitions to dg categories, seen as dg algebras with many objects. This invariant of dg categories satisfies many useful properties, including “agreement” (the Hochschild homology of a dg algebra is isomorphic to that of the dg category of compact modules) [29, 2.4] and Morita invariance (a functor in the homotopy category of dg categories that induces an isomorphism between the subcategories of compact objects also induces an isomorphism on Hochschild homology) [47, 4.4].

The notion of Hochschild homology of a differential graded (dg) algebra admits a natural dualization, the *coHochschild homology* of a dg coalgebra, which was introduced by Hess, Parent, and Scott in [23], generalizing the non-differential notion of [13]. They showed in particular that the coHochschild homology of the chain coalgebra on a simply connected space X is isomorphic to the homology of the free loop space on X and that the coHochschild homology of a connected dg coalgebra C is isomorphic to the Hochschild homology of ΩC , the cobar construction on C .

In this article we establish further properties of coHochschild homology, analogous to the invariance properties of Hochschild homology recalled above. We first prove a sort of categorification of the relation between coHochschild homology of a connected dg coalgebra C and the Hochschild homology of ΩC , showing that there is a dg Quillen equivalence between the categories of C -comodules and of ΩC -modules (Proposition 2.8). We can then establish an “agreement”-type result, stating that the coHochschild homology of a dg coalgebra C is isomorphic to the Hochschild homology of the dg category spanned by certain compact C -comodules (Proposition 2.12). Thanks to this agreement result, we can show as well that coHochschild homology is a Morita invariant (Proposition 2.23), using the notion of Morita equivalence of dg coalgebras formulated in [4], which extends that of Takeuchi [46] and which we recall here. Proving these results required us to provide criteria under which a dg Quillen equivalence of dg model categories induces a quasi-equivalence of dg subcategories (Lemma 2.13); this technical result, which we were unable to find in the literature, may also be useful in other contexts.

The natural analogue of Hochschild homology for spectra, called *topological Hochschild homology (THH)*, has proven to be an important and useful invariant of ring spectra, particularly because of its connection to K-theory via the Dennis trace. Blumberg and Mandell proved moreover that THH satisfies both “agreement,” in the sense that THH of a ring spectrum is equivalent to THH of the spectral category of appropriately compact R -modules, and Morita invariance [5].

We define here an analogue of coHochschild homology for spectra, which we call *topological coHochschild homology (coTHH)*. We show that coTHH is homotopy invariant, as well as independent of the particular model category of spectra in which one works. We prove moreover that coTHH of the suspension spectrum $\Sigma_{\pm}^{\infty} X$ of a connected Kan complex X is equivalent to $\Sigma_{\pm}^{\infty} \mathcal{L}X$, the suspension spectrum of the free loop space on X , whenever X is *EMSS-good*, i.e., whenever $\pi_1 X$ acts nilpotently on the integral homology of the based loop space on X (Theorem 3.7).

This equivalence was already known for simply connected spaces X , by work of Kuhn [31] and Malkiewicz [34], though they did not use the term coTHH. The extension of the equivalence to EMSS-good spaces is based on new results concerning total complexes of cosimplicial suspension spectra, such as the fact that $\overline{\text{Tot}}(\Sigma^{\infty} Y^{\bullet}) \simeq \Sigma^{\infty} \overline{\text{Tot}} Y^{\bullet}$ whenever the homology spectral sequence for a cosimplicial space Y^{\bullet} with coefficients in \mathbb{Z} strongly converges (Corollary A.3). We also show that if X is an EMSS-good space, then the Anderson spectral sequence for homology with coefficients in \mathbb{Z} for the cosimplicial space $\text{Map}(S_{\bullet}^1, X)$ strongly converges to $H_*(\mathcal{L}X; \mathbb{Z})$ (Proposition A.4).

In [7], Bökstedt and Waldhausen proved that $\mathrm{THH}(\Sigma_+^\infty \Omega X) \simeq \Sigma_+^\infty \mathcal{L}X$ for simply connected X . It follows thus from Theorem 3.7 that if X is simply connected, then $\mathrm{THH}(\Sigma_+^\infty \Omega X) \simeq \mathrm{coTHH}(\Sigma_+^\infty X)$, analogous to the result for dg coalgebras established in [23]. Combining this result with the spectral Quillen equivalence between categories of $\Sigma_+^\infty \Omega X$ -modules and of $\Sigma_+^\infty X$ -comodules established in [24] and with THH-agreement [5], we obtain coTHH-agreement for simply connected Kan complexes X : $\mathrm{coTHH}(\Sigma_+^\infty X)$ is equivalent to THH of the spectral category of appropriately compact $\Sigma_+^\infty \Omega X$ -modules (Corollary 3.11).

We do not consider Morita invariance for coalgebra spectra in this article, as the duality requirement of the framework in [4] is too strict to allow for interesting spectral examples. We expect that a meaningful formulation should be possible in the ∞ -category context.

In parallel with writing this article, the second author collaborated with Bohmann, Gerhardt, Høgenhaven, and Ziegenhagen on developing computational tools for coHochschild homology, in particular an analogue of the Bökstedt spectral sequence for topological Hochschild homology constructed by Angeltveit and Rognes [2]. For C a coalgebra spectrum, the E_2 -page of this spectral sequence is the associated graded of the classical coHochschild homology of the homology of C with coefficients in a field \mathbb{k} , and the spectral sequence abuts to the \mathbb{k} -homology of $\mathrm{coTHH}(C)$. If C is connected and cocommutative, then this is a spectral sequence of coalgebras. In [6] the authors also proved a Hochschild-Kostant-Rosenberg-style theorem for coHochschild homology of cofree cocommutative differential graded coalgebras.

In future work we will construct and study an analogue of the Dennis trace map, with source the K-theory of a dg or spectral coalgebra C and with target its (topological) coHochschild homology.

Acknowledgments

Work on this article began while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California, during Spring 2014, partially supported by the National Science Foundation under Grant No. 0932078000. The second author was also supported by NSF grants DMS-1104396, DMS-1406468, and DMS-1811278 during this work. We would also like to thank the University of Illinois at Chicago, the EPFL, and the University of Chicago for their hospitality during research visits enabling us to complete the research presented in this article. The authors would also like to thank The Isaac Newton Institute for Mathematical Sciences, Cambridge, for support and hospitality during the Fall 2018 program “Homotopy harnessing higher structures” where this paper was finished. This work was supported by EPSRC grant No. EP/K032208/1.

2. CoHochschild homology for chain coalgebras

In this section we recall from [23] the coHochschild complex of a chain coalgebra over a field \mathbb{k} , which generalizes the definitions in [13] and in [28] and dualizes the usual definition of the Hochschild complex of a chain algebra. We establish important properties of this construction analogous to those known to hold for Hochschild homology: “agreement” (in the sense of [37]) and Morita invariance.

Notation 2.1. Throughout this section we work over a field \mathbb{k} and write \otimes to denote the tensor product over \mathbb{k} and $|v|$ to denote the degree of a homogeneous element v of a graded vector space.

- We denote the category of (unbounded) graded chain complexes over \mathbb{k} by $\mathrm{Ch}_{\mathbb{k}}$, the category of augmented, nonnegatively graded chain algebras (dg algebras) over \mathbb{k} by $\mathrm{Alg}_{\mathbb{k}}$, and the category of coaugmented, connected (and hence nonnegatively graded) chain coalgebras (dg coalgebras) by $\mathrm{Coalg}_{\mathbb{k}}$. All of these categories are naturally dg categories, i.e., enriched over $\mathrm{Ch}_{\mathbb{k}}$, with $\mathrm{Alg}_{\mathbb{k}}$ and $\mathrm{Coalg}_{\mathbb{k}}$ inheriting their enrichments from that of $\mathrm{Ch}_{\mathbb{k}}$.

- We apply the Koszul sign convention for commuting elements of a graded vector space or for commuting a morphism of graded vector spaces past an element of the source module. For example, if V and W are graded algebras and $v \otimes w, v' \otimes w' \in V \otimes W$, then

$$(v \otimes w) \cdot (v' \otimes w') = (-1)^{|w| \cdot |v'|} vv' \otimes ww'.$$

Furthermore, if $f : V \rightarrow V'$ and $g : W \rightarrow W'$ are morphisms of graded vector spaces, then for all $v \otimes w \in V \otimes W$,

$$(f \otimes g)(v \otimes w) = (-1)^{|g| \cdot |v|} f(v) \otimes g(w).$$

All signs in the formulas below follow from the Koszul rule. It is a matter of straightforward calculation in each case to show that differentials square to zero.

- The *desuspension* endofunctor s^{-1} on the category of graded vector spaces is defined on objects $V = \bigoplus_{i \in \mathbb{Z}} V_i$ by $(s^{-1}V)_i \cong V_{i+1}$. Given a homogeneous element v in V , we write $s^{-1}v$ for the corresponding element of $s^{-1}V$.
- Given chain complexes (V, d) and (W, d) , the notation $f : (V, d) \xrightarrow{\cong} (W, d)$ indicates that f induces an isomorphism in homology. In this case we refer to f as a *quasi-isomorphism*.
- [47, Section 2.3] A *quasi-equivalence* of dg categories is a dg functor $F : \mathcal{C} \rightarrow \mathcal{D}$ such that $F_{X, X'} : \text{hom}_{\mathcal{C}}(X, X') \rightarrow \text{hom}_{\mathcal{D}}(F(X), F(X'))$ is a quasi-isomorphism for all $X, X' \in \text{Ob } \mathcal{C}$ (i.e., F is *quasi-fully faithful*) and such that the induced functor on the *homology categories*, $H_0F : H_0\mathcal{C} \rightarrow H_0\mathcal{D}$, is essentially surjective, i.e., F is *quasi-essentially surjective*. The objects of the homology category $H_0\mathcal{C}$, which is a dg category in which the hom-objects have zero differential, are the same as those of \mathcal{C} , while hom-objects are given by the 0th-homology of the hom-objects of \mathcal{C} .
- Let T denote the endofunctor on the category of graded vector spaces given by

$$TV = \bigoplus_{n \geq 0} V^{\otimes n},$$

where $V^{\otimes 0} = \mathbb{k}$. An element of the summand $V^{\otimes n}$ of TV is denoted $v_1 | \cdots | v_n$, where $v_i \in V$ for all i .

- The coaugmentation coideal of any C in $\text{Coalg}_{\mathbb{k}}$ is denoted \overline{C} .
- We consistently apply the Einstein summation convention, according to which an expression involving a term with the same letter as a subscript and a superscript denotes a sum over that index, e.g., $c_i \otimes c^i$ denotes a sum of elementary tensors over the index i .

2.1. The dg cobar construction and its extensions

Let Ω denote the *cobar construction* functor from $\text{Coalg}_{\mathbb{k}}$ to $\text{Alg}_{\mathbb{k}}$, defined by

$$\Omega C = (T(s^{-1}\overline{C}), d_{\Omega})$$

where, if d denotes the differential on C , then

$$\begin{aligned} d_{\Omega}(s^{-1}c_1 | \cdots | s^{-1}c_n) &= \sum_{1 \leq j \leq n} \pm s^{-1}c_1 | \cdots | s^{-1}(dc_j) | \cdots | s^{-1}c_n \\ &\quad + \sum_{1 \leq j \leq n} \pm s^{-1}c_1 | \cdots | s^{-1}c_{ji} | s^{-1}c_j^i | \cdots | s^{-1}c_n, \end{aligned}$$

with signs determined by the Koszul rule, where the reduced comultiplication applied to c_j is $c_{ji} \otimes c_j^i$. A straightforward computation shows that ΩC is isomorphic to the totalization of the cosimplicial cobar construction if C is 1-connected (i.e., C is connected and $C_1 = 0$).

The graded vector space underlying ΩC is naturally a free associative algebra, with multiplication given by concatenation. The differential d_Ω is a derivation with respect to this concatenation product, so that ΩC is itself a chain algebra. Any chain algebra map $\alpha : \Omega C \rightarrow A$ is determined by its restriction to the algebra generators $s^{-1}\overline{C}$.

The following two extensions of the cobar construction play an important role below. Let $\text{Mix}_{C,\Omega C}$ and $\text{Mix}_{\Omega C,C}$ denote the categories of left C -comodules in the category of right ΩC -modules and of right C -comodules in the category of left ΩC -modules, respectively. We call the objects of these categories *mixed modules*. There are functors

$$\mathcal{P}_L : \text{Coalg}_{\mathbb{k}} \rightarrow \text{Mix}_{C,\Omega C} \quad \text{and} \quad \mathcal{P}_R : \text{Coalg}_{\mathbb{k}} \rightarrow \text{Mix}_{\Omega C,C},$$

which we call the *left and right based path constructions* on C (where left and right refer to the side of the C -coaction) and which are defined as follows.

$$\mathcal{P}_L C = (C \otimes T(s^{-1}\overline{C}), d_{\mathcal{P}_L}) \quad \text{and} \quad \mathcal{P}_R C = (T(s^{-1}\overline{C}) \otimes C, d_{\mathcal{P}_R}),$$

where

$$d_{\mathcal{P}_L}(e \otimes s^{-1}c_1 | \cdots | s^{-1}c_n) = de \otimes s^{-1}c_1 | \cdots | s^{-1}c_n \pm e \otimes d_\Omega(s^{-1}c_1 | \cdots | s^{-1}c_n) \pm e_j \otimes s^{-1}e^j | s^{-1}c_1 | \cdots | s^{-1}c_n$$

$$d_{\mathcal{P}_R}(s^{-1}c_1 | \cdots | s^{-1}c_n \otimes e) = d_\Omega(s^{-1}c_1 | \cdots | s^{-1}c_n) \otimes e \pm s^{-1}c_1 | \cdots | s^{-1}c_n \otimes de \pm s^{-1}c_1 | \cdots | s^{-1}c_n | s^{-1}e_j \otimes e^j,$$

where $\Delta(e) = e_j \otimes e^j$, and applying s^{-1} to an element of degree 0 gives 0. For every C in $\text{Coalg}_{\mathbb{k}}$, there are twisted extensions of chain complexes

$$\Omega C \xrightarrow{\eta \otimes 1} \mathcal{P}_L C \xrightarrow{1 \otimes \varepsilon} C \qquad \Omega C \xrightarrow{1 \otimes \eta} \mathcal{P}_R C \xrightarrow{\varepsilon \otimes 1} C,$$

which are dg analogues of the based pathspace fibration, where $\eta : \mathbb{k} \rightarrow C$ is the coaugmentation and $\varepsilon : \Omega C \rightarrow \mathbb{k}$ the obvious augmentation.

As proved in [38, Proposition 10.6.3], both $\mathcal{P}_L C$ and $\mathcal{P}_R C$ are homotopy equivalent to the trivial mixed module \mathbb{k} , for all C in $\text{Coalg}_{\mathbb{k}}$, via a chain homotopy defined in the case of $\mathcal{P}_R C$ by

$$h_R : \mathcal{P}_R C \rightarrow \mathcal{P}_R C : w \otimes e \mapsto \begin{cases} 0 & : |e| > 0 \text{ or } w = 1 \\ s^{-1}c_1 | \cdots | s^{-1}c_{n-1} \otimes c_n & : |e| = 0 \text{ and } w = s^{-1}c_1 | \cdots | s^{-1}c_n \end{cases}$$

and analogously in the case of $\mathcal{P}_L C$. Observe that, when restricted to the sub ΩC -module of elements in positive degree, h_R is a homotopy of left ΩC -modules, while h_L is a homotopy of right ΩC -modules.

The proposition below generalizes this contractibility result.

Proposition 2.2. *There are strong deformation retracts*

1. $\Omega C \begin{matrix} \xrightarrow{\sigma} \\ \xleftarrow{\pi} \end{matrix} \mathcal{P}_R C \square_C \mathcal{P}_L C$ in the category of left ΩC -modules, and
2. $C \begin{matrix} \xrightarrow{\iota} \\ \xleftarrow{\rho} \end{matrix} \mathcal{P}_L C \otimes_{\Omega C} \mathcal{P}_R C$ in the category of left C -comodules.

Proof. Note that the graded vector space underlying $\mathcal{P}_R C \square_C \mathcal{P}_L C$ is isomorphic to $T(s^{-1}\overline{C}) \otimes C \otimes T(s^{-1}\overline{C})$, while that underlying $\mathcal{P}_L C \otimes_{\Omega C} \mathcal{P}_R C$ is isomorphic to $C \otimes T(s^{-1}\overline{C}) \otimes C$.

In the ΩC -module case, we define left ΩC -module maps

$$\pi : \mathcal{P}_R C \square_C \mathcal{P}_L C \rightarrow \Omega C : v \otimes c \otimes w \mapsto \begin{cases} \varepsilon(c) \cdot vw & : |c| = 0 \\ 0 & : |c| \neq 0, \end{cases}$$

where $\varepsilon : C \rightarrow \mathbb{k}$ denotes the counit, and

$$\sigma : \Omega C \rightarrow \mathcal{P}_R C \square_C \mathcal{P}_L C : w \mapsto w \otimes 1 \otimes 1.$$

While it is obvious that $\pi\sigma$ is the identity, showing that $\sigma\pi$ is homotopic to the identity requires a new chain homotopy $h : \mathcal{P}_R C \square_C \mathcal{P}_L C \rightarrow \mathcal{P}_R C \square_C \mathcal{P}_L C$ defined by

$$h(1 \otimes c \otimes s^{-1}c_1 | \dots | s^{-1}c_n) = \begin{cases} \sum_{1 \leq i \leq n} \pm s^{-1}c_1 | \dots | s^{-1}c_{i-1} \otimes c_i \otimes s^{-1}c_{i+1} | \dots | s^{-1}c_n & : |c| = 0 \\ 0 & : |c| \neq 0, \end{cases}$$

then extended to a map of left $T(s^{-1}\overline{C})$ -modules. A straightforward computation shows that $Dh + hD = \text{Id} - \sigma\pi$ as desired, where D denotes the differential on $\mathcal{P}_R C \square_C \mathcal{P}_L C$.

Let Δ denote the comultiplication on C . In the C -comodule case, we define left C -comodule maps by

$$\iota : C \rightarrow \mathcal{P}_L C \otimes_{\Omega C} \mathcal{P}_R C : c \mapsto c_i \otimes 1 \otimes c^i,$$

where $\Delta(c) = c_i \otimes c^i$, and

$$\rho : \mathcal{P}_L C \otimes_{\Omega C} \mathcal{P}_R C \rightarrow C : c \otimes w \otimes c' \mapsto \begin{cases} c & : |w| = |c'| = 0 \\ 0 & : \text{else.} \end{cases}$$

It is obvious that $\rho\iota$ is equal to the identity and that $\iota\rho$ is chain homotopic to the identity as left C -comodules, via the chain homotopy $\text{Id}_{\mathcal{P}_L C \otimes_{\Omega C} \mathcal{P}_R C} h_R$, which itself respects the left C -coaction as well. \square

Our interest in the left and right based path constructions stems from the following proposition.

Proposition 2.3. *The pair of functors*

$$\text{Comod}_C \begin{array}{c} \xrightarrow{-\square_C \mathcal{P}_L C} \\ \xleftarrow{-\otimes_{\Omega C} \mathcal{P}_R C} \end{array} \text{Mod}_{\Omega C}$$

forms a dg-adjunction.

Proof. It is well known that the dg-enrichments of Comod_C and $\text{Mod}_{\Omega C}$ can be constructed as equalizers in $\text{Ch}_{\mathbb{k}}$, as follows. For right C -comodules N and N' with C -coactions ρ and ρ' ,

$$\underline{\text{Comod}}_C(N, N') = \lim \left(\underline{\text{Ch}}_{\mathbb{k}}(N, N') \begin{array}{c} \xrightarrow{\rho'_*} \\ \xrightarrow{\rho^* \circ (-\otimes C)} \end{array} \underline{\text{Ch}}_{\mathbb{k}}(N, N' \otimes C) \right),$$

where the underline denotes the hom-chain complex, as opposed to the hom-set. For right ΩC -modules M and M' with ΩC -actions α and α' ,

$$\text{Mod}_{\Omega C}(M, M') = \lim \left(\text{Ch}_{\mathbb{k}}(M, M') \begin{array}{c} \xrightarrow{\alpha'_* \circ (-\otimes \Omega C)} \\ \xrightarrow{\alpha^*} \end{array} \text{Ch}_{\mathbb{k}}(M \otimes \Omega C, M') \right).$$

It is easy to see from these constructions that both of the functors in the statement of the proposition are dg-enriched, essentially because on the underlying graded vector spaces, both functors are given by tensoring with a fixed object.

Since $\pi : \mathcal{P}_R C \square_C \mathcal{P}_L C \rightarrow \Omega C$ is actually a map of ΩC -bimodules, while $\iota : C \rightarrow \mathcal{P}_L C \otimes_{\Omega C} \mathcal{P}_R C$ is a map of C -bicomodules, there are dg-natural transformations

$$\text{Id} \cong -\square_C C \xrightarrow{-\square_C \iota} -\square_C (\mathcal{P}_L C \otimes_{\Omega C} \mathcal{P}_R C)$$

and

$$-\otimes_{\Omega C} (\mathcal{P}_R C \square_C \mathcal{P}_L C) \xrightarrow{-\otimes_{\Omega C} \pi} -\otimes_{\Omega C} \Omega C \cong \text{Id},$$

which provide the unit and counit of the adjunction. It is an easy exercise to verify the triangle inequalities. \square

The extension of the cobar construction that is the focus of this article is the *coHochschild complex* functor

$$\widehat{\mathcal{H}} : \text{Coalg}_{\mathbb{k}} \rightarrow \text{Ch}_{\mathbb{k}},$$

defined as follows [23]. Let C be a connected, coaugmented chain coalgebra with comultiplication $\Delta(c) = c_i \otimes c^i$. We then let

$$\widehat{\mathcal{H}}(C) = (C \otimes T(s^{-1}\overline{C}), d_{\widehat{\mathcal{H}}})$$

where

$$\begin{aligned} d_{\widehat{\mathcal{H}}}(e \otimes s^{-1}c_1 | \cdots | s^{-1}c_n) &= de \otimes s^{-1}c_1 | \cdots | s^{-1}c_n \pm e \otimes d_{\Omega}(s^{-1}c_1 | \cdots | s^{-1}c_n) \\ &\quad \pm e_j \otimes s^{-1}e^j | s^{-1}c_1 | \cdots | s^{-1}c_n \\ &\quad \pm e^i \otimes s^{-1}c_1 | \cdots | s^{-1}c_n | s^{-1}e_i, \end{aligned}$$

where $\Delta(e) = e_j \otimes e^j$, and applying s^{-1} to an element of degree 0 gives 0. The signs follow from the Koszul rule, as usual. As in the case of the cobar construction, it is not hard to show that $\widehat{\mathcal{H}}(C)$ is isomorphic to the totalization of a certain cosimplicial construction when C is 1-connected; see the analogue for spectra in Section 3.

For every C in $\text{Coalg}_{\mathbb{k}}$, there is a twisted extension of chain complexes

$$\Omega C \xrightarrow{\eta \otimes 1} \widehat{\mathcal{H}}(C) \xrightarrow{1 \otimes \varepsilon} C, \tag{2.1}$$

which is the dg analogue of the free loop fibration, where, as above, $\eta : \mathbb{k} \rightarrow C$ is the coaugmentation and $\varepsilon : \Omega C \rightarrow \mathbb{k}$ the obvious augmentation.

Remark 2.4. There is a natural and straightforward extension of the coHochschild complex of a chain coalgebra to a *cocyclic complex*, analogous to the extension of the Hochschild complex of a chain algebra to the cyclic complex. Moreover the construction of the coHochschild complex of a coalgebra C can be generalized to allow for coefficients in any C -bicomodule [23, Section 1.3].

2.2. *Properties of the dg coHochschild construction*

The result below provides a first indication of the close link between the Hochschild and coHochschild constructions.

Proposition 2.5. [21, Corollary 2.22] *Let $\mathcal{H} : \mathbf{Alg}_{\mathbb{k}} \rightarrow \mathbf{Ch}_{\mathbb{k}}$ denote the usual Hochschild construction. For any C in $\mathbf{Coalg}_{\mathbb{k}}$, there is a natural quasi-isomorphism*

$$\widehat{\mathcal{H}}(C) \xrightarrow{\cong} \mathcal{H}(\Omega C).$$

Remark 2.6. Another (and easier) way to obtain an algebra from a coalgebra C is to take its linear dual degreewise, denoted C^\vee . A straightforward computation shows that for any C in $\mathbf{Coalg}_{\mathbb{k}}$, the linear dual of the coHochschild complex of C is isomorphic to the Hochschild (cochain) complex of C^\vee , i.e.,

$$(\widehat{\mathcal{H}}(C))^\vee \cong \mathcal{H}(C^\vee).$$

We show below that Proposition 2.5 can be categorified, i.e., lifted to model categories of C -comodules and ΩC -modules. We then use this categorification to establish “agreement” and Morita invariance for coHochschild homology.

Convention 2.7. Henceforth, we fix the following model structures.

- Endow $\mathbf{Ch}_{\mathbb{k}}$ with the model structure for which cofibrations are degreewise injections, fibrations are degreewise surjections, and weak equivalences are quasi-isomorphisms.
- For any C in $\mathbf{Coalg}_{\mathbb{k}}$, the category $\mathbf{Mod}_{\Omega C}$ of right ΩC -modules is equipped with the model structure right-induced from $\mathbf{Ch}_{\mathbb{k}}$ by the forgetful functor

$$U : \mathbf{Mod}_{\Omega C} \rightarrow \mathbf{Ch}_{\mathbb{k}},$$

which exists by [41, 4.1]. The fibrations in $\mathbf{Mod}_{\Omega C}$ are exactly those module maps that are degreewise surjective, whence every object is fibrant. Every cofibration in $\mathbf{Mod}_{\Omega C}$ is a retract of a sequential colimit of module maps given by pushouts along morphisms of the form (injection) $\otimes\Omega C$.

- For any C in $\mathbf{Coalg}_{\mathbb{k}}$, the category \mathbf{Comod}_C of right C -comodules is equipped with the model structure left-induced from $\mathbf{Ch}_{\mathbb{k}}$ by the forgetful functor

$$U : \mathbf{Comod}_C \rightarrow \mathbf{Ch}_{\mathbb{k}},$$

which exists by [22, 6.3.7]; see also [19]. The cofibrations in \mathbf{Comod}_C are exactly those comodule maps that are degreewise injective, whence every object is cofibrant. Every retract of a sequential limit of comodule maps given by pullbacks along morphisms of the form (surjection) $\otimes C$ is a fibration in \mathbf{Comod}_C .

2.2.1. *Categorifying Proposition 2.5*

Proposition 2.8. *For any C in $\mathbf{Coalg}_{\mathbb{k}}$, the enriched adjunction*

$$\mathbf{Comod}_C \begin{array}{c} \xrightarrow{-\square_C \mathcal{P}_L C} \\ \xleftarrow{-\otimes_{\Omega C} \mathcal{P}_R C} \end{array} \mathbf{Mod}_{\Omega C}$$

is a Quillen equivalence.

Proof. To simplify notation, we write

$$L_C = -\square_C \mathcal{P}_L C \quad \text{and} \quad R_C = -\otimes_{\Omega C} \mathcal{P}_R C.$$

Observe that $R_C(M) \cong (M \otimes C, D_R)$ and $L_C(N) \cong (N \otimes \Omega C, D_L)$, for every M in $\text{Mod}_{\Omega C}$ and every N in Comod_C , where the differentials of these complexes are specified by

$$D_R(x \otimes c) = dx \otimes c \pm x \otimes dc \pm (x \cdot s^{-1}c_i) \otimes c^i$$

and

$$D_L(y \otimes w) = dy \otimes w \pm y \otimes d_{\Omega} w \pm y_j \otimes (s^{-1}c^j \cdot w).$$

Here, $\rho(y) = y \otimes + y_j \otimes c^j$ and $\Delta(c) = c_i \otimes c^i$, where ρ is the C -coaction on N and Δ the comultiplication on C , and the signs are determined by the Koszul rule.

We show first that $L_C \dashv R_C$ is a Quillen adjunction. If $j : N \rightarrow N'$ is a cofibration in Comod_C , i.e., a degreewise injective morphism of C -comodules, then there is a decomposition $N' = N \oplus V$ as graded vector spaces, since we are working over a field. It follows $L_C(N')$ can be built inductively as an ΩC -module from $L_C(N)$, as we explain below.

Suppose that $dv \in N$ and $\rho(v) - v \otimes 1 \in N \otimes C$ for all $v \in V$, and let B be a basis of V . There is a pushout diagram in $\text{Mod}_{\Omega C}$

$$\begin{array}{ccc} \coprod_{x \in B} S^{|x|-1} \otimes \Omega C & \xrightarrow{\iota} & L_C(N) \\ \downarrow & & \downarrow L_C(j) \\ \coprod_{x \in B} D^{|x|} \otimes \Omega C & \longrightarrow & L_C(N \oplus V) \end{array}$$

where S^m is the chain complex with only one basis element, which is in degree m , D^{m+1} has two basis elements, in degrees m and $m + 1$, with a differential linking the latter to the former, and ι maps the generator of $S^{|x|-1}$ to dx for every $x \in B$. It follows that $L_C(j)$ is a cofibration, in this special case.

In the general case, we use that any comodule is the filtered colimit of its finite-dimensional subcomodules [20, Lemma 1.1]. We can structure this filtered colimit more precisely as follows. For any $n \geq 1$, let $\{N'(i, n) \mid i \in \mathcal{J}_n\}$ denote the set of subcomodules of N' such that $N'(i, n)/N$ is of dimension n for all $i \in \mathcal{J}_n$, and set $N'(n) = \Sigma_{i \in \mathcal{J}_n} N'(i, n)$. The argument above shows that the injection $N \rightarrow N'(1)$ is a pushout along a morphism of the form (injection) $\otimes \Omega C$, and, more generally, that the inclusion $N'(n) \rightarrow N'(n + 1)$ is a pushout along a morphism of the form (injection) $\otimes \Omega C$ for all n , whence $j : N \rightarrow N' = \text{colim}_n N'(n)$ is a cofibration.

On the other hand, we can show by a spectral sequence argument that the functor L_C preserves all weak equivalences and therefore preserves trivial cofibrations. Any N in Comod_C admits a natural ‘‘primitive’’ filtration

$$F_0 N \subseteq F_1 N \subseteq F_2 N \subseteq \dots \subseteq N \tag{2.2}$$

as a C -comodule, i.e., $F_0 N = \ker(N \xrightarrow{\bar{\rho}} N \otimes C)$ and

$$F_m N = \ker(N \xrightarrow{\bar{\rho}^{(n)}} N \otimes C^{\otimes n})$$

for all $m \geq 1$, where $\bar{\rho} = \rho - N \otimes \eta$, and $\bar{\rho}^{(n)} = (\bar{\rho} \otimes C^{\otimes n-1})\bar{\rho}^{(n-1)}$. Note that this filtration is always exhaustive, since C is connected, and $(\rho \otimes C)\rho = (N \otimes \Delta)\rho$.

Consider the exhaustive filtration of $L_C(N)$ as an ΩC -module induced by applying L_C to the primitive filtration (2.2) of N :

$$(F_0N \otimes \Omega C, d \otimes 1 + 1 \otimes d_\Omega) \subseteq (F_1N \otimes \Omega C, D_L) \subseteq (F_2(N) \otimes \Omega C, D_L) \subseteq \dots \subseteq (N \otimes \Omega C, D_L).$$

The E_2 -term of the spectral sequence associated to this filtration, which converges to $H_*L_C(N)$, is isomorphic as a graded vector space to $H_*(N) \otimes H_*(\Omega C)$, from which it follows that a quasi-isomorphism of C -comodules induces an isomorphism on the E_2 -terms of the associated spectral sequences and thus on the E_∞ -terms as well.

Since $R_C(M)$ is cofibrant in \mathbf{Comod}_C and $L_C(N)$ is fibrant in $\mathbf{Mod}_{\Omega C}$ for every M in $\mathbf{Mod}_{\Omega C}$ and every N in \mathbf{Comod}_C , it follows from [26, Proposition 1.3.13(b)] that $L_C \dashv R_C$ is a Quillen equivalence if the unit $N \rightarrow R_C L_C N$ and counit $L_C R_C M \rightarrow M$ of the $L_C \dashv R_C$ adjunction are weak equivalences for every N (since they are all cofibrant) and every M (since they are all fibrant). To conclude, it suffices therefore to observe that for every M , there is a sequence of isomorphisms and weak equivalences in $\mathbf{Mod}_{\Omega C}$,

$$L_C R_C M = M \otimes_{\Omega C} \mathcal{P}_R C \square_C \mathcal{P}_L C \simeq M \otimes_{\Omega C} \Omega C \cong M,$$

where the weak equivalence is a consequence of Proposition 2.2(1), and that for every N , there is a sequence of isomorphisms and weak equivalence in \mathbf{Comod}_C ,

$$R_C L_C N = N \square_C \mathcal{P}_L C \otimes_{\Omega C} \mathcal{P}_R C \simeq N \square_C C \cong N,$$

where the weak equivalence follows from Proposition 2.2(2). \square

Remark 2.9. In Chapter 2 of his thesis [32], Lefèvre-Hasegawa defined a model structure on the category of *cocomplete* comodules over a *cocomplete*, coaugmented dg coalgebra C , of which the proposition above would seem to be a special case. Here, “cocomplete” means that the respective primitive filtration is exhaustive, which is not immediate if C is not connected. It seems, however, that Lefèvre-Hasegawa did not check that the category of cocomplete comodules is closed under limits, which we suspect is actually not true.

The proposition above also could be viewed as a special case [14, Proposition 3.15], which establishes a general Quillen equivalence between certain categories of coalgebras over a cooperad and algebras over an operad as mediated by a twisting morphism. One needs to show that the weak equivalences of [14] are the same as those in our model structure on \mathbf{Comod}_C , which follows from Proposition 2.2. We think there is merit in providing an explicit, independent proof in this special case, especially as it makes evident the “geometric” nature of the proof (using based path spaces).

Example 2.10. Proposition 2.8 implies that for every reduced simplicial set K ,

$$\mathbf{Comod}_{C_*(K)} \begin{array}{c} \xrightarrow{-\square_{C_*(K)} \mathcal{P}_L C_*(K)} \\ \perp \\ \xleftarrow{-\otimes_{\Omega C_*(K)} \mathcal{P}_R C_*(K)} \end{array} \mathbf{Mod}_{\Omega C_*(K)}$$

is a Quillen equivalence, where $C_*(K)$ denotes the normalized chain coalgebra of K with coefficients in \mathbb{k} . Moreover, if K is actually 1-reduced, there is a natural quasi-isomorphism of chain algebras

$$\alpha_K: \Omega C_*(K) \xrightarrow{\cong} C_*(\mathbb{G}K)$$

[45], where \mathbb{G} denotes the Kan loop group functor, which induces a Quillen equivalence

$$\text{Mod}_{\Omega C_*(K)} \begin{array}{c} \xrightarrow{\alpha_!} \\ \xleftarrow{\alpha^*} \end{array} \text{Mod}_{C_*(\mathbb{G}K)} .$$

It follows that if K is 1-reduced, there is a Quillen equivalence

$$\text{Comod}_{C_*(K)} \begin{array}{c} \xrightarrow{\quad} \\ \xleftarrow{\quad} \end{array} \text{Mod}_{C_*(\mathbb{G}K)} .$$

2.2.2. Agreement

The Quillen equivalence of Proposition 2.8 enables us to establish “agreement” for coHochschild homology, analogous to “agreement” for Hochschild homology, which we recall now. In [37], McCarthy extended the notion of Hochschild homology in a natural way to exact categories, seen as “rings with many objects,” and established “agreement” in this context: the Hochschild homology of the exact category of finitely generated projective modules over a ring R is isomorphic to the Hochschild homology of R itself.

Keller generalized the definition of Hochschild homology to dg categories (seen as dg algebras with many objects) in [29], [30] and showed that agreement still held in this more general setting: for any dg \mathbb{k} -algebra A , the Hochschild homology of A is isomorphic to that of the full dg subcategory dgfree_A of Mod_A [29, Theorem 2.4]. The objects of dgfree_A are finitely generated quasi-free A -modules, i.e., A -modules such that the underlying nondifferential graded module is free and finitely generated over the nondifferential graded algebra underlying A . Note that objects in dgfree_A are cofibrant in our chosen model structure on Mod_A .

For any C in $\text{Coalg}_{\mathbb{k}}$, the notion of agreement for coHochschild homology is expressed in terms of the full dg subcategory dgcofree_C of Comod_C , the objects of which are the fibrant C -comodules N such that there exists a quasi-isomorphism of ΩC -modules $L_C(N) \xrightarrow{\cong} M$, where M is an object of $\text{dgfree}_{\Omega C}$. Observe that a morphism $L_C(N) \rightarrow M$ is a quasi-isomorphism if and only if its transpose $N \rightarrow R_C(M)$ is a quasi-isomorphism, since $L_C \dashv R_C$ is a Quillen equivalence, all objects in Comod_C or in the image of R_C are cofibrant, and all objects in $\text{Mod}_{\Omega C}$ or in the image of L_C are fibrant.

Remark 2.11. The notation for the dg subcategory dgcofree_C is a bit abusive, since not all of its objects are actually quasi-cofree, i.e., such that the underlying nondifferential graded comodule is cofree over the nondifferential graded coalgebra underlying C . On the other hand, any quasi-cofree C -comodule that is “finitely cogenerated” over C is an object of dgcofree_C . More precisely, any quasi-cofree C -comodule is the limit of a tower of comodule maps given by pullbacks along morphisms of the form (surjection) $\otimes C$ and is therefore fibrant. Moreover, if N is “finitely cogenerated” by V , then its image under L_C is finitely generated by V .

Proposition 2.12. *Agreement holds for coHochschild homology of coalgebras, i.e., for every C in $\text{Coalg}_{\mathbb{k}}$,*

$$H_*(\widehat{\mathcal{H}}(C)) \cong H_*(\mathcal{H}(\text{dgcofree}_C)).$$

The key to the proof of agreement for coalgebras, as well as to establishing Morita invariance at the end of this section, is the following lemma, providing conditions under which dg Quillen equivalences induce quasi-equivalences of dg categories. We were unable to find this result in the literature, though we suspect it is well known.

Lemma 2.13. *Let $M \begin{array}{c} \xrightarrow{F} \\ \perp \\ \xleftarrow{G} \end{array} N$ be an enriched Quillen equivalence of dg model categories. Let M'*

and N' be full dg subcategories of M and N , respectively, where all objects of M' and of N' are fibrant and cofibrant.

- (1) Suppose that F restricts and corestricts to a functor $F' : \mathbf{M}' \rightarrow \mathbf{N}'$. If for every $Y \in \text{Ob } \mathbf{N}'$ there exists a weak equivalence $p_Y : \widehat{G(Y)} \xrightarrow{\simeq} G(Y)$ where $\widehat{G(Y)} \in \text{Ob } \mathbf{M}'$, then F' is a quasi-equivalence.
- (2) Suppose that G restricts and corestricts to a functor $G' : \mathbf{N}' \rightarrow \mathbf{M}'$. If for every $X \in \text{Ob } \mathbf{M}'$ there exists a weak equivalence $j_X : F(X) \xrightarrow{\simeq} \widehat{F(X)}$ where $\widehat{F(X)} \in \text{Ob } \mathbf{N}'$, then G' is a quasi-equivalence.

Proof. We prove (1) and leave the dual proof of (2) to the reader. By [26, Proposition 1.3.13], since $F \dashv G$ is a Quillen equivalence, the unit morphism $\eta_{X'} : X' \rightarrow G'F'(X')$ is a weak equivalence for every $X' \in \text{Ob } \mathbf{M}'$, since X' is cofibrant and $F'(X')$ is fibrant. It follows that

$$F'_{X', Y'} : \text{hom}_{\mathbf{M}'}(X', Y') \rightarrow \text{hom}_{\mathbf{N}'}(F'(X'), F'(Y'))$$

is a quasi-isomorphism for all $X', Y' \in \text{Ob } \mathbf{M}'$, since it factors as

$$\text{hom}_{\mathbf{M}'}(X', Y') \xrightarrow{\simeq} \text{hom}_{\mathbf{M}'}(X', G'F'(Y')) \cong \text{hom}_{\mathbf{N}'}(F'(X'), F'(Y')).$$

The first map above is a quasi-isomorphism because X' is cofibrant, and $\eta_{X'}$ is a weak equivalence between fibrant objects. We conclude that F' is quasi-fully faithful.

Let $Z' \in \text{Ob } \mathbf{N}'$. Since we can choose $p_{Z'}$ as a cofibrant replacement of $G'(Z')$, the composite

$$F'(\widehat{G(Z')}) \xrightarrow{F'(p_{Z'})} F'G'(Z') \xrightarrow{\varepsilon_{Z'}} Z'$$

is a model for the derived counit of the adjunction and therefore a weak equivalence. Both its source and target are objects in \mathbf{N}' and therefore fibrant (and cofibrant), whence $\text{hom}_{\mathbf{N}'}(W', \varepsilon_{Z'}F'(p_{Z'}))$ is a quasi-isomorphism for all $W' \in \text{Ob } \mathbf{N}'$, as all objects in \mathbf{N}' are cofibrant. By Exercise 6 in [47, Section 2.3], it follows that the homology class of $\varepsilon_{Z'}F'(p_{Z'})$ is an isomorphism in the homology category of \mathbf{N}' and thus that F' is quasi-essentially surjective. \square

Proof of Proposition 2.12. Observe that

$$\widehat{\mathcal{H}}(C) \simeq \mathcal{H}(\Omega C) \simeq \mathcal{H}(\text{dgfree}_{\Omega C}) \simeq \mathcal{H}(\text{dgcofree}_C),$$

where the first equivalence is given by Proposition 2.5 and the second by [29]. The third follows from Proposition 2.8 and Lemma 2.13(2). To see that all of the conditions of Lemma 2.13(2) are satisfied, note first that all objects of $\text{dgfree}_{\Omega C}$ and dgcofree_C are both fibrant and cofibrant. Moreover, the functor R_C restricts and corestricts to a dg functor from $\text{dgfree}_{\Omega C}$ to dgcofree_C , since the counit $L_C R_C \rightarrow \text{Id}$ of the Quillen equivalence $L_C \dashv R_C$ is a quasi-isomorphism on objects that are cofibrant and fibrant. Finally, dgcofree_C is defined precisely so that the remaining condition holds as well.

Because every quasi-equivalence of dg categories is a Morita equivalence [47, Section 4.4], and Hochschild homology of dg categories is an invariant of Morita equivalence [47, Section 5.2], we can conclude. \square

2.2.3. Morita invariance

Thanks to the “agreement” result established above, we can now show that coHochschild homology satisfies a property dual to the Morita invariance of Hochschild homology. The study of equivalences between categories of comodules over coalgebras over a field, commonly referred to as *Morita-Takeuchi theory*, was initiated by Takeuchi [46] and further elaborated and generalized by Farinati and Solotar [18] and Brzezinski and Wisbauer [11], among others. In [4], Berglund and Hess formulated a homotopical version of this theory, in terms of the following notion.

Definition 2.14. Let C, D be objects in $\text{Coalg}_{\mathbb{k}}$. A *braiding from C to D* is a pair (X, T) where X is in $\text{Ch}_{\mathbb{k}}$, and T is a morphism of chain complexes

$$T: C \otimes X \rightarrow X \otimes D$$

satisfying the following axioms.

(Pentagon axiom)

The diagram

$$\begin{array}{ccc}
 C \otimes X & \xrightarrow{T} & X \otimes D \\
 \Delta_C \otimes 1 \downarrow & & \downarrow 1 \otimes \Delta_D \\
 C \otimes C \otimes X & & X \otimes D \otimes D \\
 \searrow 1 \otimes T & & \nearrow T \otimes 1 \\
 & C \otimes X \otimes D &
 \end{array} \tag{2.3}$$

commutes.

(Counit axiom)

The diagram

$$\begin{array}{ccc}
 C \otimes X & \xrightarrow{T} & X \otimes D \\
 \downarrow \epsilon_C \otimes 1 & & \downarrow 1 \otimes \epsilon_D \\
 \mathbb{k} \otimes X & \xrightarrow{\cong} X & \xleftarrow{\cong} X \otimes \mathbb{k}
 \end{array} \tag{2.4}$$

commutes.

We write $(X, T): C \rightarrow D$ to indicate that (X, T) is a braiding from C to D .

Example 2.15 (Change of coalgebras). A morphism $f: C \rightarrow D$ in $\text{Coalg}_{\mathbb{k}}$ gives rise to a braiding $(\mathbb{k}, f): C \rightarrow D$ and thus to an adjunction

$$\text{Comod}_C \begin{array}{c} \xrightarrow{f_*} \\ \xleftarrow{f^*} \end{array} \text{Comod}_D, \tag{2.5}$$

which we call the *coextension/corestriction-of-scalars adjunction* or *change-of-corings adjunction* associated to f . The D -component of the counit of the $f_* \dashv f^*$ adjunction is f itself and that for every C -comodule (M, δ) ,

$$f_*(M, \delta) = (M, (1 \otimes f)\delta).$$

Since Comod_C is bicomplete for all coalgebras C (as $\text{Ch}_{\mathbb{k}}$ is locally presentable, and Comod_C is a category of coalgebras for the comonad $-\otimes C$), it follows from [4, Proposition 3.17] that every braiding $(X, T): C \rightarrow D$ gives rise to a $\text{Ch}_{\mathbb{k}}$ -adjunction

$$\text{Comod}_C \begin{array}{c} \xrightarrow{T_*} \\ \xleftarrow{T^*} \end{array} \text{Comod}_D, \quad T_* \dashv T^*,$$

such that the diagram

$$\begin{array}{ccc}
 \text{Comod}_C & \xrightarrow{T_*} & \text{Comod}_D \\
 \downarrow U & & \downarrow U \\
 \text{Ch}_{\mathbb{k}} & \xrightarrow{-\otimes X} & \text{Ch}_{\mathbb{k}}
 \end{array}$$

commutes, i.e., the endofunctor on $\text{Ch}_{\mathbb{k}}$ is just given by tensoring with X . Moreover, since we are working over a field and thus tensoring with any chain complex preserves both weak equivalences and equalizers, Proposition 3.31 in [4] implies that if X is dualizable with dual X^\vee , then $T^* = -\square_D(X^\vee \otimes C)$, where $-\square_D-$ denotes the cotensor product over D .

The following lemma, relating braidings to the adjunction $L_C \dashv R_C$ studied above, plays an important role at the end of this section.

Lemma 2.16. *For every braiding $(X, T): C \rightarrow D$, the dg adjunction*

$$\begin{array}{ccc}
 \text{Mod}_{\Omega C} & \xrightleftharpoons[\text{hom}_{\Omega D}(X \otimes \Omega D, -)]{-\otimes_{\Omega C}(X \otimes \Omega D)} & \text{Mod}_{\Omega D} \\
 & \perp & \\
 & &
 \end{array}$$

is a dg Quillen pair, satisfying the natural isomorphisms

$$(-\otimes_{\Omega C}(X \otimes \Omega D)) \circ L_C \cong L_D \circ T_*$$

and

$$R_C \circ \text{hom}_{\Omega D}(X \otimes \Omega D, -) \cong T^* \circ R_D.$$

Proof. Given our choice of model structures on module categories, it is easy to check that the adjunction above is indeed a Quillen pair. It suffices to establish the first isomorphism, since the second is then an immediate consequence. The computation is straightforward, given that the left ΩC -action on $X \otimes \Omega D$ induced by the braiding

$$T: C \otimes X \rightarrow X \otimes D : c \otimes x \mapsto x_i \otimes d^i$$

is specified by $s^{-1}c \cdot (x \otimes w) = x_i \otimes s^{-1}d^i \cdot w$ for all $c \in \overline{C}$, $x \in X$, and $w \in \Omega D$. \square

The coalgebraic analogue of Morita equivalence is defined as follows.

Definition 2.17. Let C, D be in $\text{Coalg}_{\mathbb{k}}$. If there is a braiding (X, T) from C to D such that $T_* \dashv T^*$ is a Quillen equivalence, then C and D are (homotopically) Morita-Takeuchi equivalent.

As a special case of [4, Theorem 4.16], we can describe Morita-Takeuchi-equivalent pairs of chain coalgebras in terms of the following notions, recalled from [4].

Definition 2.18. Let X be a dualizable chain complex and C a dg coalgebra. The canonical coalgebra associated to X and C is the dg coalgebra $X_*(C)$ with underlying chain complex

$$X_*(C) = X^\vee \otimes C \otimes X,$$

and comultiplication given by the composite

$$\begin{array}{ccc}
 X^\vee \otimes C \otimes X & \xrightarrow{1 \otimes \Delta \otimes 1} & X^\vee \otimes C \otimes C \otimes X \\
 & & \downarrow 1 \otimes 1 \otimes u \otimes 1 \otimes 1 \\
 & & (X^\vee \otimes C \otimes X) \otimes (X^\vee \otimes C \otimes X),
 \end{array}$$

where $u : \mathbb{k} \rightarrow X \otimes X^\vee$ is the coevaluation map. The *canonical braiding* $(X, T_C^{\text{univ}}) : C \rightarrow X_*(C)$ is defined by

$$T_C^{\text{univ}} = u \otimes 1 : C \otimes X \rightarrow X \otimes (X^\vee \otimes C \otimes X).$$

The *canonical adjunction associated to X and C* is the adjunction governed by the universal braided bi-module (X, T_C^{univ}) ,

$$\text{Comod}_C \begin{array}{c} \xrightarrow{(T_C^{\text{univ}})_*} \\ \perp \\ \xleftarrow{(T_C^{\text{univ}})^*} \end{array} \text{Comod}_{X_*(C)}. \tag{2.6}$$

We say that X *satisfies effective homotopic descent* if this adjunction is a Quillen equivalence.

Remark 2.19. The canonical braiding determined by a dualizable chain complex X and a dg coalgebra C is universal, in the sense that any braiding $(X, T) : C \rightarrow C'$ factors as $(X, T_C^{\text{univ}}) : C \rightarrow X_*(C)$ followed by the change-of-coalgebras braiding $(\mathbb{k}, g_T) : X_*(C) \rightarrow C'$, where g_T is given by the composite

$$X_*(C) \xrightarrow{X^\vee \otimes T} X^\vee \otimes X \otimes C' \xrightarrow{ev \otimes C'} \mathbb{k} \otimes C' \cong C'.$$

Definition 2.20. A morphism $g : C \rightarrow C'$ of dg coalgebras is *copure* if the counit $g_*g^*(M) \rightarrow M$ of the $g_* \dashv g^*$ adjunction is a weak equivalence for all fibrant C' -comodules M .

Remark 2.21. Since \mathbb{k} is fibrant (seen as a chain complex concentrated in degree 0) in $\text{Ch}_{\mathbb{k}}$, every coalgebra is fibrant as a comodule over itself. Thus, if $g : C \rightarrow C'$ is copure, then $g_*g^*(C') \rightarrow C'$ is a weak equivalence. Since $g_*g^*(C') \cong C$, seen as a C' -comodule via g , it follows that a copure coalgebra map is, in particular, a weak equivalence.

The next result is a special case of the second part of [4, Theorem 4.16].

Theorem 2.22. [4, Theorem 4.16] *Let C, C' be in $\text{Coalg}_{\mathbb{k}}$. If C and C' are Morita-Takeuchi equivalent via a braiding (X, T) such that X is dualizable, then X satisfies effective homotopic descent with respect to C , and $g_T : X_*(C) \rightarrow C'$ is a copure weak equivalence of corings.*

We can deduce the promised invariance of coHochschild homology from this description of Morita-Takeuchi equivalent coalgebras.

Proposition 2.23. *Let C, C' be in $\text{Coalg}_{\mathbb{k}}$. If C and C' are Morita-Takeuchi equivalent via a braiding (X, T) such that the total dimension of X is finite, then $\widehat{\mathcal{H}}(C) \simeq \widehat{\mathcal{H}}(C')$.*

Note that if X has finite total dimension, then it is certainly dualizable.

Proof. Since X satisfies effective homotopic descent with respect to C , the canonical adjunction

$$\text{Comod}_C \begin{array}{c} \xrightarrow{(T_C^{\text{univ}})_*} \\ \perp \\ \xleftarrow{(T_C^{\text{univ}})^*} \end{array} \text{Comod}_{X_*(C)},$$

is a dg Quillen equivalence.

It follows that

$$\widehat{\mathcal{H}}(C) \simeq \mathcal{H}(\text{dgcofree}_C) \simeq \mathcal{H}(\text{dgcofree}_{X_*(C)}) \simeq \widehat{\mathcal{H}}(X_*(C)) \simeq \widehat{\mathcal{H}}(C'),$$

where the first and third weak equivalences follow from Agreement (Proposition 2.12), and the last equivalence from the fact that g_T is copure and therefore a weak equivalence.

Lemma 2.13(2) and Lemma 2.16 suffice to establish the second equivalence, as we now show. To simplify notation, taking $D = X_*(C)$ in Lemma 2.16, let $F \dashv G$ denote the adjunction

$$(- \otimes_{\Omega C}(X \otimes \Omega X_*(C))) \dashv \text{hom}_{\Omega X_*(C)}(X \otimes \Omega X_*(C), -),$$

and let $T = T_C^{\text{univ}}$.

Since T^* is right Quillen, it preserves fibrant objects. Moreover, if N is a $X_*(C)$ -comodule such that there exists an $\Omega X_*(C)$ -module M and a quasi-isomorphism $j : N \xrightarrow{\simeq} R_{X_*(C)}M$, then

$$T^*(j) : T^*(N) \xrightarrow{\simeq} T^*(R_{X_*(C)}M)$$

is also a quasi-isomorphism, since all modules are fibrant, and T^* is a right Quillen functor. By Lemma 2.16

$$T^*(R_{X_*(C)}M) \cong R_C \circ G(M),$$

so there is a quasi-isomorphism

$$T^*(N) \xrightarrow{\simeq} R_C \circ G(M)$$

or, equivalently, a quasi-isomorphism

$$L_C T^*(N) \xrightarrow{\simeq} G(M).$$

Moreover, because

$$G(M) = \text{hom}_{\Omega X_*(C)}(X \otimes \Omega X_*(C), M) \cong \text{hom}(X, M) \cong X^\vee \otimes M,$$

if M is actually an object of $\text{dgfree}_{\Omega X_*(C)}$, then $G(M)$ is an object of $\text{dgfree}_{\Omega C}$. We conclude that T^* restricts and corestricts to a functor

$$T^* : \text{dgcofree}_{X_*(C)} \rightarrow \text{dgcofree}_C.$$

By Lemma 2.13(2), to verify that T^* is actually a quasi-equivalence, it remains to check that for every N in dgcofree_C , there is an $N' \in \text{dgcofree}_{X_*(C)}$ and a quasi-isomorphism $T_*(N) \xrightarrow{\simeq} N'$. If N is an object of dgcofree_C , then there is an object M in $\text{dgfree}_{\Omega C}$ and a quasi-isomorphism $j : L_C(N) \xrightarrow{\simeq} M$. Since F is left Quillen, and both $L_C(N)$ and M are cofibrant ΩC -modules, it follows that $F(j) : F(L_C(N)) \rightarrow F(M)$ is also

a quasi-isomorphism. By Lemma 2.16, $F(L_C(N)) \cong L_{X_*(C)}(T_*(N))$, whence there is a quasi-isomorphism $L_{X_*(C)}(T_*(N)) \xrightarrow{\cong} F(M)$, where $F(M)$ is an object of $\mathbf{dgfree}_{\Omega X_*(C)}$. Indeed, if M is quasi-free on V of finite total degree, then $F(M)$ is quasi-free on $V \otimes X$, which is also free of finite total degree. If $T_*(N)$ is actually fibrant, then it is itself an object of $\mathbf{dgcofree}_{X_*(C)}$, and we can set $N' = T_*(N)$. If not, then for any fibrant replacement of $T_*(N)$ in $\mathbf{Comod}_{X_*(C)}$ will be an object of $\mathbf{dgcofree}_{X_*(C)}$ and can play the role of N' . \square

3. Topological coHochschild homology of spectra

We now consider a spectral version of the constructions and results in section 2. Here we work in any monoidal model category of spectra. We show that our results are model invariant in Proposition 3.3 below.

3.1. The general theory

Let k be a commutative ring spectrum, C a k -coalgebra with comultiplication $\Delta : C \rightarrow C \wedge_k C$, and M a C -bicomodule with right coaction $\rho : M \rightarrow M \wedge_k C$ and left coaction $\lambda : M \rightarrow C \wedge_k M$. Henceforth we write \wedge for \wedge_k and $C^{\wedge n}$ for the n -fold smash product of C over k .

Definition 3.1. The *coHochschild complex* $\widehat{\mathcal{H}}(M, C)$ is the cosimplicial spectrum with

$$\widehat{\mathcal{H}}(M, C)^n = M \wedge C^{\wedge n}$$

and coface operators

$$d^i = \begin{cases} \rho \wedge \text{Id}_C^{\wedge n} & i = 0 \\ \text{Id}_M \wedge \text{Id}_C^{\wedge i-1} \wedge \Delta \wedge \text{Id}_C^{\wedge n-i} & 1 \leq i \leq n \\ \tau \circ (\lambda \wedge \text{Id}_C^{\wedge n}) & i = n + 1 \end{cases}$$

where $\tau : C \wedge M \wedge C^{\wedge n} \rightarrow M \wedge C^{\wedge n+1}$ cycles the first entry to the last entry. The codegeneracies involve the counit of C .

Note that one can take $M = C$ with $\lambda = \rho = \Delta$. In this case $\widehat{\mathcal{H}}(C, C) = \widehat{\mathcal{H}}(C)$ is the *cyclic cobar complex*.

Next we define the homotopy invariant notion of topological coHochschild homology. We use $\overline{\text{Tot}}X^\bullet$ to denote the totalization of a Reedy fibrant replacement of the cosimplicial spectrum X^\bullet . By [25, 19.8.7], this is a model of the homotopy inverse limit. *Topological coHochschild homology* is defined as the derived totalization of the coHochschild complex,

$$\text{coTHH}(M, C) = \overline{\text{Tot}}\widehat{\mathcal{H}}(M, C).$$

We abbreviate $\text{coTHH}(C, C)$ as $\text{coTHH}(C)$.

The next statement shows that coTHH is homotopy invariant.

Lemma 3.2. *Let \mathcal{C} be a monoidal model category of k -module spectra. If $f : C \rightarrow C'$ is a map of coalgebra spectra in \mathcal{C} such that in the underlying category of k -module spectra f is a weak equivalence, and C, C' are cofibrant, then the induced map $\text{coTHH}(C) \rightarrow \text{coTHH}(C')$ is a weak equivalence.*

Proof. Since C and C' are cofibrant, and \mathcal{C} is a monoidal model category, f induces a levelwise weak equivalence $\widehat{\mathcal{H}}(C) \rightarrow \widehat{\mathcal{H}}(C')$. Since homotopy inverse limits preserve levelwise weak equivalences, the statement follows. \square

In addition, coTHH is model independent.

Proposition 3.3. *Topological coHochschild homology is independent of the model of spectra used.*

Proof. This follows from Lemma 3.4 below, since any two monoidal model categories of spectra are connected by Quillen equivalences via a strong monoidal left adjoint. A universal approach to constructing these monoidal Quillen equivalences is described in [44, 4.7]; explicit constructions are given in [36, 0.1,0.2], [40, 5.1], and [35, 1.1,1.8]. This is also summarized in a large diagram in [41, 7.1]. \square

Lemma 3.4. *Let $L : \mathcal{C} \rightarrow \mathcal{D}$ be the left adjoint of a strong monoidal Quillen equivalence between two monoidal model categories of k -module spectra with \overline{L} the associated derived functor. Let C be a coalgebra spectrum that is cofibrant as an underlying k -module spectrum. Then $\text{coTHH}(LC)$ is weakly equivalent to $\overline{L}\text{coTHH}(C)$.*

Proof. Let R be the right adjoint to L . By [25, 15.4.1], levelwise prolongation, denoted L^\bullet, R^\bullet , induces a Quillen equivalence between the associated Reedy model categories of cosimplicial spectra. Let $\overline{L^\bullet}$ and $\overline{R^\bullet}$ denote the derived functors.

Since $LC \wedge LC \cong L(C \wedge C)$, it follows that LC is a coalgebra spectrum in \mathcal{D} and that $L^\bullet \widehat{\mathcal{H}}(C) \cong \widehat{\mathcal{H}}(LC)$. Since C is cofibrant, $\widehat{\mathcal{H}}(C)$ is levelwise cofibrant and so is its cofibrant replacement in the Reedy model structure. Since L preserves weak equivalences between cofibrant objects, it follows that $\overline{L^\bullet} \widehat{\mathcal{H}}(C)$ is weakly equivalent to $L^\bullet \widehat{\mathcal{H}}(C)$ and hence also to $\widehat{\mathcal{H}}(LC)$.

Applying $\overline{R^\bullet}$ to both sides of this equivalence, we have that $\overline{R^\bullet L^\bullet} \widehat{\mathcal{H}}(C)$ is weakly equivalent to $\overline{R^\bullet} \widehat{\mathcal{H}}(LC)$. Since $\overline{L^\bullet}$ and $\overline{R^\bullet}$ form an equivalence of homotopy categories, $\overline{R^\bullet L^\bullet}$ is naturally weakly equivalent to the identity and therefore

$$\widehat{\mathcal{H}}(C) \simeq \overline{R^\bullet L^\bullet} \widehat{\mathcal{H}}(C) \simeq \overline{R^\bullet} \widehat{\mathcal{H}}(LC). \tag{3.1}$$

Let $c^\bullet X$ denote the constant cosimplicial object on X . Since $L^\bullet(c^\bullet X) \cong c^\bullet(LX)$, the right adjoints also commute, i.e., $\lim R^\bullet X^\bullet \cong R \lim X^\bullet$, and so the associated derived functors also commute. In particular, $\overline{R}\text{coTHH}(LC)$ is weakly equivalent to the homotopy inverse limit of $\overline{R^\bullet} \widehat{\mathcal{H}}(LC)$. It follows from (3.1) that $\text{coTHH}(C) \simeq \overline{R}\text{coTHH}(LC)$. Since \overline{L} and \overline{R} form an equivalence of homotopy categories, this is equivalent to the statement in the lemma. \square

It turns out that coalgebras in spaces with respect to the Cartesian product or in pointed spaces with respect to the smash product are of a very restricted nature. The only possible co-unital coalgebra structure on a space is given by the diagonal $\Delta : X \rightarrow X \times X$. Similarly, for a pointed space, the only possible co-unital coalgebra structure exists on a pointed space of the form X_+ and is induced by the diagonal $\Delta_+ : X_+ \rightarrow X_+ \wedge X_+$.

It follows that strictly counital coalgebra spectra are also very restricted. Consider a symmetric spectrum Z . Since the zeroth level of $Z \wedge Z$ is the smash product of two copies of level zero of Z , the zeroth level of a co-unital coalgebra symmetric spectrum must have a disjoint base point, which we denote $(Z_0)_+$. In fact, even more structure is forced in any of the symmetric monoidal categories of spectra. Let \mathbf{Sp} refer to the \mathbb{S} -modules of [17] or any diagram category of spectra, including symmetric spectra (over simplicial sets or topological spaces, see [27,36]), orthogonal spectra (see [36,35]), Γ -spaces (see [42,9]), and \mathcal{W} -spaces (see [1]).

Proposition 3.5. [39] *In \mathbf{Sp} , co-unital coalgebras over the sphere spectrum are cocommutative. In fact, if C is a co-unital coalgebra over the sphere spectrum, then $\Sigma_{\mp}^\infty C_0 \rightarrow C$ is surjective.*

In an earlier version of this paper, we proved the special case of this proposition for symmetric spectra over simplicial sets. Because of Proposition 3.5, we focus on suspension spectra in the next section.

3.2. coTHH of suspension spectra

The main statement of this section is the geometric identification of coTHH of a suspension spectrum as the suspension spectrum of the free loop space, see Theorem 3.7. The proof of this statement is delayed to the following section. This main statement leads to a connection between coTHH and THH and another analogue of “agreement” in the sense of [37]. Throughout this section by *spaces* we mean simplicial sets.

Definition 3.6. For X a Kan complex, consider any model of the loop-path space fibration, $\Omega X \rightarrow PX \rightarrow X$. We say that a Kan complex X is an *EMSS-good* space if X is connected and $\pi_1 X$ acts nilpotently on $H_i(\Omega X; \mathbb{Z})$ for all i .

This term refers to the fact that the Eilenberg-Moore spectral sequence for the loop-path space fibration converges strongly by [15] for any EMSS-good space X . Note that if X is simply connected, then X is certainly EMSS-good.

Theorem 3.7. *If X is an EMSS-good space, then the topological coHochschild homology of its suspension spectrum is equivalent to the suspension spectrum of the free loop space:*

$$\text{coTHH}(\Sigma_+^\infty X) \simeq \Sigma_+^\infty \mathcal{L}X.$$

See also [31] and [34, 2.22] for earlier proofs of this statement for simply connected spaces. The proof of this theorem is given in Section 3.3 and relies on the proofs of [8, 4.1, 8.4] and generalizations discussed in Appendix A. It is likely that this can be further generalized to non-connected spaces X , see for example the proofs of [43, 3.1, 3.2].

For X simply connected, $\text{THH}(\Sigma_+^\infty \Omega X) \simeq \Sigma_+^\infty \mathcal{L}X$ by [7], implying the following corollary.

Corollary 3.8. *Let X be a simply connected Kan complex. There is a weak equivalence between the topological coHochschild homology of the suspension spectrum of X and the topological Hochschild homology of the suspension spectrum of the based loops on X :*

$$\text{coTHH}(\Sigma_+^\infty X) \simeq \text{THH}(\Sigma_+^\infty \Omega X).$$

As in the differential graded context, there is also a categorified version of this result. In [24, 5.4], somewhat more generally reformulated below in Proposition 3.9, we show that there is a Quillen equivalence between the categories of module spectra over $\Sigma_+^\infty \Omega X$ and of comodule spectra over $\Sigma_+^\infty X$. Recall from [24, 5.2] that the category of comodules over $\Sigma_+^\infty X$ admits a model structure, denoted there by $(\text{Comod}_{\Sigma_+^\infty X})_{\pi_*^s}^{\text{st}}$ because it is the stabilization of the category of X_+ -comodules with respect to π_*^s -equivalences. Since this is the only model structure we consider for this category in this paper, we denote it simply by $\text{Comod}_{\Sigma_+^\infty X}$. Weak equivalences in this structure induce stable equivalences on the underlying spectra by [24, 5.2 (1)].

The first part of the following result is a simplified version of the statement in [24, 5.4], setting $\mathcal{E}_* = \pi_*^s$. Note that, as above, choosing a base point for X determines a coaugmentation map from the sphere spectrum $\mathbb{S} \rightarrow \Sigma_+^\infty X$, which in turn determines a $\Sigma_+^\infty X$ -comodule structure on \mathbb{S} .

Proposition 3.9. [24, 5.4] *For X a connected space, there is a Quillen equivalence*

$$\text{Mod}_{\Sigma_+^\infty \Omega X} \begin{array}{c} \xrightarrow{L} \\ \xleftarrow{R} \end{array} \text{Comod}_{\Sigma_+^\infty X}$$

such that $L(\Sigma_+^\infty \Omega X)$ is weakly equivalent to the sphere spectrum as a comodule, and $R(\Sigma_+^\infty X)$ is weakly equivalent to the sphere spectrum as a module.

Proof. The statement of [24, 5.4] is formulated for the model of the loop space on X given by the Kan loop group, $\mathbb{G}X$, for X a reduced simplicial set.

Here instead we work with $\Sigma_+^\infty \Omega X$, where ΩX denotes any model of the loop space of the fibrant replacement of X (i.e., a Kan complex). Since $\Sigma_+^\infty \Omega X$ and $\Sigma_+^\infty \mathbb{G}X$ have the same homotopy type, their categories of modules are Quillen equivalent. Moreover any connected simplicial set is weakly equivalent to a reduced simplicial set, and replacing X by a weakly equivalent space induces Quillen equivalences on the respective categories of comodule spectra (see [24, 5.3]).

The original result is also formulated with respect to a chosen generalized homology theory \mathcal{E}_* , which we fix here to be stable homotopy, $\mathcal{E}_* = \pi_*^s$. As in [24, 5.14], since levelwise π_*^s -equivalences are stable equivalences, one can show that the weak equivalences in this model structure on $\text{Mod}_{\Sigma_+^\infty \Omega X}$ are the stable equivalences on the underlying spectra, i.e., this is the usual model structure on $\text{Mod}_{\Sigma_+^\infty \Omega X}$. (The proof of [24, 5.14] treats the special case where X is a point, but works verbatim for any X .)

Concerning the second part of the theorem, the left adjoint in [24, 5.4], $-\wedge_{\Sigma_+^\infty \mathbb{G}X} \Sigma_+^\infty \mathbb{P}X$, takes $\Sigma_+^\infty \mathbb{G}X$ to $\Sigma_+^\infty \mathbb{P}X$, which is weakly equivalent to \mathbb{S} since $\mathbb{P}X$ is contractible. Hence, $L(\Sigma_+^\infty \mathbb{G}X) \simeq \mathbb{S}$. On the other hand, the functor from comodules to modules is the stabilization of the composite of three functors given in [24, 4.14]. By [24, 3.11], since X_+ is the cofree X_+ -comodule on S^0 , the first of these functors takes X_+ to a retractive space $\text{Ret}_X(S^0)$ over X with total space $S^0 \times X$. The next functor is an equivalence of categories that takes $\text{Ret}_X(S^0)$ to $\text{Ret}_{\mathbb{P}X}(S^0)$ with a trivial $\mathbb{G}X$ -action, which is sent by the third functor to S^0 , the trivial, pointed $\mathbb{G}X$ -module. Upon stabilization, this computation implies that the Quillen equivalence on the spectral level sends the comodule $\Sigma_+^\infty X$ to the module \mathbb{S} , i.e., $R(\Sigma_+^\infty X) \simeq \mathbb{S}$. \square

As in Proposition 2.12 in the differential graded context, it follows from Corollary 3.8 and Proposition 3.9 that topological coHochschild homology for suspension spectra satisfies “agreement.” Here though, instead of considering finitely generated free modules, we consider the modules that are finitely built from the free module spectrum. Recall that a subcategory of a triangulated category is called *thick* if it is closed under equivalences, triangles, and retracts. Here we also use the same terminology to refer to the underlying subcategory of the model category corresponding to the thick subcategory of the derived category. For example, for R a ring spectrum, we consider $\text{Thick}_R(R)$, the underlying spectral category associated to the thick subcategory generated by R . In the literature, these modules are variously called “perfect,” “compact,” or “finitely built from R .”

Since $L(\Sigma_+^\infty \Omega X) \simeq \mathbb{S}$, and Quillen equivalences preserve thick subcategories, [5, 5.3, 5.9] implies the following.

Lemma 3.10. *The Quillen equivalence in Proposition 3.9 induces a weak equivalence*

$$\text{THH}(\text{Thick}_{\Sigma_+^\infty \Omega X}(\Sigma_+^\infty \Omega X)) \simeq \text{THH}(\text{Thick}_{\Sigma_+^\infty X}(\mathbb{S})).$$

It is a consequence of [5, 5.12] that for any ring spectrum R ,

$$\text{THH}(R) \simeq \text{THH}(\text{Thick}_R(R)).$$

The next corollary follows immediately from this equivalence for $R = \Sigma_+^\infty \Omega X$, together with Corollary 3.8 and Lemma 3.10.

Corollary 3.11. *Agreement holds for topological coHochschild homology of coalgebra spectra that are suspension spectra. That is, for any simply-connected Kan complex X ,*

$$\text{coTHH}(\Sigma_+^\infty X) \simeq \text{THH}(\text{Thick}_{\Sigma_+^\infty X}(\mathbb{S})).$$

Remark 3.12. Note that $\text{Thick}_{\Sigma_+^\infty X}(\mathbb{S})$ is the subcategory of compact objects in the category of comodules over $\Sigma_+^\infty X$. This follows from the Quillen equivalence in Proposition 3.9, since $\text{Thick}_{\Sigma_+^\infty \Omega X}(\Sigma_+^\infty \Omega X)$ is the subcategory of compact object for modules over $\Sigma_+^\infty \Omega X$. Since $R(\Sigma_+^\infty X)$ is weakly equivalent to \mathbb{S} , where R the right adjoint in Proposition 3.9, it follows that $\Sigma_+^\infty X$ is a compact comodule over itself if and only if \mathbb{S} is a compact module over $\Sigma_+^\infty \Omega X$. In [16, 5.6(2)], working over $H\mathbb{F}_p$ instead of \mathbb{S} , it is shown that there are examples where $H\mathbb{F}_p$ is not compact as a module over $H\mathbb{F}_p \wedge \Sigma_+^\infty \Omega X$, e.g., when $X = \mathbb{C}P^\infty$.

3.3. Cobar, Bar, and loop spaces

In this section we consider the Cobar and Bar constructions on a suspension spectrum and prove Theorem 3.7 from the last section about coTHH of a suspension spectrum. The proofs in this section rely on results about the convergence of spectral sequences for cosimplicial spaces that are established in Appendix A.

Let C be a k -coalgebra spectrum, N a left C -comodule with coaction $\lambda : N \rightarrow C \wedge N$, and M a right C -comodule with coaction $\rho : M \rightarrow M \wedge C$.

Definition 3.13. The *cobar complex* $\Omega^\bullet(M, C, N)$ is the cosimplicial spectrum with

$$\Omega(M, C, N)^n = M \wedge C^{\wedge n} \wedge N$$

with coface operators

$$d^i = \begin{cases} \rho \wedge \text{Id}_C^{\wedge n} \wedge \text{Id}_N & i = 0 \\ \text{Id}_M \wedge \text{Id}_C^{\wedge i-1} \wedge \Delta \wedge \text{Id}_C^{\wedge n-i} & 1 \leq i \leq n \\ \text{Id}_M \wedge \text{Id}_C^{\wedge n} \wedge \lambda & i = n + 1 \end{cases}$$

The codegeneracies involve the counit of C .

If C is a coaugmented k -coalgebra with coaugmentation $\eta : k \rightarrow C$, i.e., η is a homomorphism of coalgebras such that $\epsilon\eta = \text{Id}_C$, then η endows k with the structure of a C -bicomodule. In this case $\Omega(k, C, k) = \Omega^\bullet(C)$ is the *cobar complex* of C . Its derived totalization is the *cobar construction on C* :

$$\text{Cobar}(C) = \overline{\text{Tot}}\Omega^\bullet(C).$$

The following cosimplicial resolution of the mapping space plays an important role in the statements below.

Definition 3.14. Let W and Z be pointed simplicial sets with Z a Kan complex, and let $\text{Map}_*(W_\bullet, Z)$ be the cosimplicial space with $\text{Map}_*(W, Z)^n$ equal to a product of copies of Z indexed by the non-base point n -simplices in W , with cofaces and codegeneracies induced by those of W . The pointed mapping space Z^W agrees with the totalization of this cosimplicial space.

If $W = S^1 = \Delta[1]/\partial\Delta[1]$, then $\text{Map}_*(S^1_\bullet, Z)^n = Z^{\times n}$ for all n , and the totalization is ΩZ , a simplicial model for the based loop space on $|Z|$.

For X a pointed space, there is a canonical map $S^0 \rightarrow X$ that gives rise to a coaugmentation $\mathbb{S} \rightarrow \Sigma_+^\infty X$. Thus we can consider the cobar construction on $\Sigma_+^\infty X$.

Proposition 3.15. *If X is pointed and an EMSS-good space, then the cobar construction on the suspension spectrum of X is weakly equivalent to the suspension spectrum of the pointed loops on X :*

$$\text{Cobar}(\Sigma_+^\infty X) \simeq \Sigma_+^\infty \Omega X.$$

Proof. If we add a disjoint base point, then $\text{Map}_*(S^1_\bullet, X)_+$ has cosimplicial level n given by $(X^{\times n})_+ \cong (X_+)^{\wedge n}$. Applying the suspension spectrum functor, we see that $\Sigma^\infty \text{Map}_*(S^1_\bullet, X)_+$ agrees with the cobar complex $\Omega^\bullet(\Sigma^{\infty}_+ X)$.

By [15], if X is EMSS-good, the Eilenberg-Moore spectral sequence converges for ordinary homology with integral coefficients. This Eilenberg-Moore spectral sequence is the homology spectral sequence for the cosimplicial space $\text{Map}_*(S^1_\bullet, X)$. By Corollary A.3 the strong convergence of this spectral sequence implies that the total complex commutes with the suspension spectrum functor, i.e.,

$$\overline{\text{Tot}}\Sigma^\infty \text{Map}_*(S^1_\bullet, X) \simeq \Sigma^\infty \overline{\text{Tot}} \text{Map}_*(S^1_\bullet, X).$$

By Proposition A.6, we can add disjoint base points to this equivalence, obtaining that

$$\overline{\text{Tot}}\Sigma^\infty \text{Map}_*(S^1_\bullet, X)_+ \simeq \Sigma^\infty \overline{\text{Tot}} \text{Map}_*(S^1_\bullet, X)_+.$$

Since $\overline{\text{Tot}} \text{Map}_*(S^1_\bullet, X)_+ \simeq \Omega X_+$, we can conclude. \square

A dual to Proposition 3.15, with a considerably simpler proof, holds as well. The statement of the dual is formulated in terms of the *Kan classifying space functor*, $\overline{W}: \mathbf{sGp} \rightarrow \mathbf{sSet}_0$, from simplicial groups to reduced simplicial sets. A detailed definition of this functor can be found in [12], where it is also shown that \overline{W} factors as the composite $\text{codiag} \circ N$, where $N: \mathbf{sGp} \rightarrow \mathbf{ssSet}$ is the levelwise nerve functor from simplicial groups to bisimplicial sets, and $\text{codiag}: \mathbf{ssSet} \rightarrow \mathbf{sSet}$ is the *Artin-Mazur codiagonalization functor* [3]. The functor codiag is often called *Artin-Mazur totalization* and denoted Tot , which we avoid, due to the risk of confusion with the other notion of totalization that we employ in this article. As we do not make any computations based on the explicit and somewhat involved formula for codiag , we do not recall it here.

We also consider the *bar construction functor*, denoted Bar , which associates to any associative ring spectrum R a spectrum $\text{Bar}R = |B_\bullet R|$, where $| - |$ denotes geometric realization, and $\text{Bar}_\bullet R$ is the simplicial spectrum with $\text{Bar}_n R = R^{\wedge n}$, face maps built from the multiplication map of R , and degeneracies from its unit map.

Proposition 3.16. *For any simplicial group G , the bar construction on the ring spectrum $\Sigma^{\infty}_+ G$ is naturally weak equivalent to the suspension spectrum of the bar construction of G , i.e.,*

$$\text{Bar}(\Sigma^{\infty}_+ G) \simeq \Sigma^{\infty}_+ \overline{W}G.$$

Proof. Cegarra and Remedios proved in [12] that the obvious natural transformation from the diagonalization functor $\text{diag}: \mathbf{ssSet} \rightarrow \mathbf{sSet}$ to codiag is in fact a natural weak equivalence. It follows that for any simplicial group G there is a sequence of natural weak equivalences and isomorphisms

$$\Sigma^{\infty}_+ \overline{W}G \simeq \Sigma^{\infty}_+ \text{diag} NG \cong |\Sigma^{\infty}_+ NG| \cong |\text{Bar}_\bullet \Sigma^{\infty}_+ G| = \text{Bar}(\Sigma^{\infty}_+ G),$$

where straightforward computations suffice to establish the two isomorphisms. \square

The next lemma is the first step in the proof of Theorem 3.7. Note that we consider unpointed mapping spaces here.

Lemma 3.17. *For any space X , there is an isomorphism of cosimplicial spectra*

$$\text{coTHH}^\bullet(\Sigma^{\infty}_+ X) \cong \Sigma^{\infty}_+ \text{Map}(S^1_\bullet, X).$$

Proof. Since these are both cosimplicial suspension spectra, it is enough to establish the isomorphism on the 0th space level. The 0th space of $\text{coTHH}(\Sigma_+^\infty X)$ has n th cosimplicial level $(X_+)^{\wedge(n+1)} \cong (X^{\times(n+1)})_+$, which agrees with $\text{Map}(S_\bullet^1, X)_+^n$. In both cases, the coface maps are induced by diagonals on the appropriate factor (with one extra twist for d^{n+1}), while the codegeneracy maps are projections onto the appropriate factors. \square

Proof of Theorem 3.7. Proposition A.6 implies that it is sufficient to prove the statement with disjoint base points removed, so it suffices to show that

$$\overline{\text{Tot}}\Sigma^\infty \text{Map}(S_\bullet^1, X) \simeq \Sigma^\infty \overline{\text{Tot}} \text{Map}(S_\bullet^1, X).$$

By Corollary A.3, it is enough to know that the Anderson spectral sequence for homology with coefficients in \mathbb{Z} for the cosimplicial space $\text{Map}(S_\bullet^1, X)$ strongly converges. By Proposition A.4 this holds for X an EMSS-good space, as required in the hypotheses here. \square

Appendix A. Total complexes of cosimplicial suspension spectra

In this section we prove several useful results concerning cosimplicial spectra, their associated spectral sequences, and commuting certain homotopy limits and colimits. The most general statement, Proposition A.1, gives conditions in terms of convergence of the associated spectral sequence for commuting the derived total complex (a homotopy limit) with smashing with a spectrum (a homotopy colimit). In this paper, we need only the suspension spectrum case, stated in Corollary A.3. The convergence conditions in the hypothesis here are verified in Proposition A.4 for the Anderson spectral sequence for the cosimplicial space $\text{Map}(S_\bullet^1, X)$. These statements are then used in the proofs of Theorem 3.7 and Proposition 3.15 above. Proposition A.6 shows that a variation of Corollary A.3 holds even after adding base points.

Proposition A.1. *If the spectral sequence associated to the cosimplicial space Y^\bullet for the generalized homology theory D_* converges strongly, then*

$$\overline{\text{Tot}}(D \wedge Y^\bullet) \simeq D \wedge \overline{\text{Tot}}Y^\bullet.$$

Bousfield, in [8], shows that the following conditions imply strong convergence for such spectral sequences.

Proposition A.2. [8, 3.1] *Let R be a ring such that $R \subset \mathbb{Q}$ or $R = \mathbb{Z}/p$ for p a prime. If Y^\bullet is a cosimplicial space such that the associated homology spectral sequence with coefficients in R strongly converges to $H_*(\overline{\text{Tot}}Y^\bullet; R)$, then for each connective spectrum D with R -nilpotent coefficient groups $\pi_i D$, the spectral sequence associated to Y^\bullet for the generalized homology theory D_* converges strongly to $D_*(\overline{\text{Tot}}Y^\bullet)$.*

Since abelian groups are \mathbb{Z} -nilpotent, the following corollary of Propositions A.1 and A.2 holds.

Corollary A.3. *If the integral spectral sequence for the cosimplicial space Y^\bullet strongly converges, then*

$$\overline{\text{Tot}}(\Sigma^\infty Y^\bullet) \simeq \Sigma^\infty \overline{\text{Tot}}Y^\bullet.$$

Proof of Proposition A.1. Recall from [10, X.6.1] that there is a homotopy spectral sequence for any cosimplicial space Y^\bullet that converges to the homotopy of $\overline{\text{Tot}}Y^\bullet$ under mild conditions. This spectral sequence arises from the tower of fibrations given by $\{\text{Tot}^s(Y^\bullet)\}$ and has E_2 -term given by $\pi^s \pi_t Y^\bullet$.

Rector’s spectral sequence for computing the D_* -homology of a cosimplicial space is considered in [8, 2.4], where it is constructed as the homotopy spectral sequence for the cosimplicial spectrum given by $D \wedge Y^\bullet$.

The E_2 -term is therefore given by $\pi^s \pi_t(D \wedge Y^\bullet) \cong \pi^s D_t(Y^\bullet)$, and it abuts to $\pi_* \overline{\text{Tot}}(D \wedge Y^\bullet)$. By [8, 2.5], strong convergence for this spectral sequence implies that $D_*(\overline{\text{Tot}}Y^\bullet)$ is isomorphic to $\pi_* \overline{\text{Tot}}(D \wedge Y^\bullet)$. Strong convergence for the homology spectral sequence for D_* thus implies the statement in the proposition. \square

Our next goal is to prove the following strong convergence result.

Proposition A.4. *For X an EMSS-good space, the Anderson spectral sequence for homology with coefficients in \mathbb{Z} for the cosimplicial space $\text{Map}(S^1_\bullet, X)$ strongly converges to $H_*(\mathcal{L}X; \mathbb{Z})$.*

This strengthens the convergence results in [8, 4.2] that require X to be simply connected. We expect that Proposition A.5 should enable similar generalizations for other mapping spaces.

To prove Proposition A.4, we need the following definitions and result. A cosimplicial space is *R-strongly convergent* if the associated homology spectral sequence with coefficients in R strongly converges. If $R = \mathbb{Z}$ we often leave off the R . It is *R-pro-convergent* if the homology spectral sequence with coefficients in R converges to the associated tower of partial total spaces; see [8, 8.4] for details. In each application in this paper, the associated tower of partial total spaces is eventually constant, so pro-convergence is equivalent to strong convergence in cases relevant to us. The next result, from [8, 8.4] and generalized to non-contractible Y^\bullet in [43, 3.2], is formulated in terms of R -pro-convergent cosimplicial spaces.

Consider a pull-back square of cosimplicial spaces

$$\begin{array}{ccc} M^\bullet & \longrightarrow & Y^\bullet \\ \downarrow & & \downarrow f \\ X^\bullet & \longrightarrow & B^\bullet \end{array}$$

There are associated pull-back squares for each cosimplicial level n

$$\begin{array}{ccc} M^n & \longrightarrow & Y^n \\ \downarrow & & \downarrow \\ X^n & \longrightarrow & B^n \end{array}$$

and for each partial total space Tot_s

$$\begin{array}{ccc} \text{Tot}_s M & \longrightarrow & \text{Tot}_s Y \\ \downarrow & & \downarrow \\ \text{Tot}_s X & \longrightarrow & \text{Tot}_s B \end{array}$$

Proposition A.5. [8, 8.4], [43, 3.2] *Consider a pull-back square of cosimplicial spaces as above, with f a fibration and $X^\bullet, Y^\bullet,$ and B^\bullet fibrant. If $X^\bullet, Y^\bullet,$ and B^\bullet are R pro-convergent, and the Eilenberg-Moore spectral sequences for the pull-back squares above for each cosimplicial level n and each total level s strongly converge, then M^\bullet is R -pro-convergent.*

Proof of Proposition A.4. Since S^1 is the following pushout in simplicial sets

$$\begin{array}{ccc} * & \longrightarrow & \Delta[1] \\ \downarrow & & \downarrow \\ S^0 & \longrightarrow & S^1 \end{array},$$

the cosimplicial space $\text{Map}(S_{\bullet}^1, X)$ is a pull-back:

$$\begin{array}{ccc} \text{Map}(S_{\bullet}^1, X) & \longrightarrow & \text{Map}(\Delta[1]_{\bullet}, X) \\ \downarrow & & \downarrow \\ \text{Map}(S_{\bullet}^0, X) & \longrightarrow & \text{Map}(*_{\bullet}, X) \end{array} .$$

We use Proposition A.5 to establish strong convergence of the homology spectral sequence with coefficients in \mathbb{Z} for $\text{Map}(S_{\bullet}^1, X)$ by verifying the hypotheses listed there.

Since the inclusion $* \rightarrow \Delta[1]$ is a cofibration of simplicial sets, and X is fibrant, the righthand vertical map above is a fibration of cosimplicial spaces. Also because X is fibrant, the four corners are fibrant cosimplicial spaces.

The cosimplicial spaces in the two bottom corners are constant, with each level given by X and X^2 respectively. Hence the associated homology spectral sequences are strongly convergent. The cosimplicial space for the top right corner is equivalent to the cosimplicial space given by Rector’s geometric cobar construction for the pullback of the identity maps:

$$\begin{array}{ccc} & & X \\ & & \downarrow \\ X & \longrightarrow & X \end{array} .$$

By [8, 4.1], since the fibers here are trivial, the associated homology spectral sequence converges as long as X is connected.

Next we consider the Eilenberg-Moore spectral sequences associated to the pullbacks in each level. In level n the pullback is given by

$$\begin{array}{ccc} & & X^{n+2} \\ & & \downarrow \pi_{0, n+2} \\ X & \xrightarrow{\Delta} & X^2 \end{array} .$$

Again, we apply [8, 4.1]. Here the vertical map is projection onto the first and last factors, so the action of $\pi_1(X^2)$ on the homology of the fiber is trivial. Thus, the associated spectral sequence strongly converges.

Finally, we consider the Eilenberg-Moore spectral sequences associated to the pullbacks of partial total spaces. Tot_0 agrees with cosimplicial level zero, so it is covered above for $n = 0$. By [10, X.3.3], $\text{Tot}_s \text{Map}(Z_{\bullet}, Y) \cong \text{Map}(Z_{\bullet}^{[s]}, Y)$ where $Z_{\bullet}^{[s]}$ is the s -skeleton of Z . It follows that $\text{Tot} \cong \text{Tot}_s$ for all s in the bottom two corners. Also, since $\Delta[1]$ is one dimensional, $\text{Tot}_s \text{Map}(\Delta[1]_{\bullet}, X) \cong \text{Map}(\Delta[1]_{\bullet}, X)$ for $s \geq 1$. So we have the same pullback of partial total spaces for each $s \geq 1$,

$$\begin{array}{ccc} & & \text{Map}(\Delta[1], X) \\ & & \downarrow \\ X & \xrightarrow{\Delta} & X^2 \end{array} .$$

Choose a point in X^2 in the image of the diagonal map Δ , so that the fiber over that point is the pointed loop space ΩX . Since X is EMSS-good, the action of the fundamental group $\pi_1(X)$ on the homology $H_*(\Omega X)$ for the path loop space fibration is nilpotent. It follows that for the vertical fibration above, $\pi_1(X^2)$ also acts nilpotently on $H_*(\Omega X)$. Thus, by [8, 4.1], the associated spectral sequence strongly converges. \square

The following proposition considers the effect on the weak equivalence of Corollary A.3 of adding disjoint base points.

Proposition A.6. *If Y^\bullet is a cosimplicial space such that $\overline{\text{Tot}}\Sigma^\infty Y^\bullet$ is weakly equivalent to $\Sigma^\infty \overline{\text{Tot}}Y^\bullet$, then $\overline{\text{Tot}}\Sigma_+^\infty Y^\bullet$ is weakly equivalent to $\Sigma_+^\infty(\overline{\text{Tot}}Y^\bullet)$.*

Proof. To be definite, we work here in the underlying model category of symmetric spectra of simplicial sets [27], where the suspension spectrum $\Sigma^\infty X$ is always cofibrant. Thus we do not need to derive the coproducts below, but we do need to consider the derived product, denoted $\overline{\times}$.

Recall that in the homotopy category of spectra, the coproduct is equivalent to the product, i.e.,

$$W \vee Z \simeq W \overline{\times} Z$$

for all spectra W and Z .

Let $\mathbb{S} \cong \Sigma^\infty S^0$ denote the sphere spectrum. If $f\Sigma^\infty X$ is a fibrant replacement of $\Sigma^\infty X$ for some space X , then

$$\Sigma_+^\infty X \cong \Sigma^\infty X \vee \mathbb{S} \simeq \Sigma^\infty X \overline{\times} \mathbb{S} \simeq f\Sigma^\infty X \times f\mathbb{S}.$$

Let $f\Sigma_+^\infty Y^\bullet$ denote the fibrant replacement of $\Sigma_+^\infty Y^\bullet$ in the Reedy model category of cosimplicial spectra. By the argument above, $f\Sigma_+^\infty Y^\bullet$ is levelwise weakly equivalent to $f\Sigma^\infty(Y^\bullet) \times c^\bullet f\mathbb{S}$, where c^\bullet denotes the constant cosimplicial spectrum functor. Since totalization commutes with products,

$$\overline{\text{Tot}}\Sigma_+^\infty Y^\bullet \simeq \overline{\text{Tot}}\Sigma^\infty(Y^\bullet) \times \overline{\text{Tot}}c^\bullet f\mathbb{S} \simeq \overline{\text{Tot}}\Sigma^\infty(Y^\bullet) \times f\mathbb{S}.$$

The hypothesis of the proposition, together with the weak equivalence between products and coproducts, implies that

$$\overline{\text{Tot}}\Sigma^\infty(Y^\bullet) \times f\mathbb{S} \simeq \Sigma^\infty \overline{\text{Tot}}Y^\bullet \vee f\mathbb{S}.$$

Since $\mathbb{S} \rightarrow f\mathbb{S}$ is a trivial cofibration, this last term is weakly equivalent to $\Sigma_+^\infty(\overline{\text{Tot}}Y^\bullet)$, as desired. \square

References

- [1] D.W. Anderson, Convergent functors and spectra, in: Localization in Group Theory and Homotopy Theory, and Related Topics, Sympos., Battelle Seattle Res. Center, Seattle, Wash., 1974, in: Lecture Notes in Math., vol. 418, Springer, Berlin, 1974, pp. 1–5.
- [2] Vigeik Angeltveit, John Rognes, Hopf algebra structure on topological Hochschild homology, Algebraic Geom. Topol. 5 (2005) 1223–1290.
- [3] M. Artin, B. Mazur, On the van Kampen theorem, Topology 5 (1966) 179–189.
- [4] Alexander Berglund, Kathryn Hess, Homotopical Morita theory for corings, Isr. J. Math. 227 (1) (2018) 239–287.
- [5] Andrew J. Blumberg, Michael A. Mandell, Localization theorems in topological Hochschild homology and topological cyclic homology, Geom. Topol. 16 (2) (2012) 1053–1120.
- [6] Anna Marie Bohmann, Teena Gerhardt, Amalie Högenhaven, Brooke Shipley, Stephanie Ziegenhagen, Computational tools for topological coHochschild homology, Topol. Appl. 235 (2018) 185–213.
- [7] Marcel Bökstedt, Friedhelm Waldhausen, The map $BSG \rightarrow A(*) \rightarrow QS^0$, in: Algebraic Topology and Algebraic K-Theory, Princeton, N.J., 1983, in: Ann. of Math. Stud., vol. 113, Princeton Univ. Press, Princeton, NJ, 1987, pp. 418–431.
- [8] A.K. Bousfield, On the homology spectral sequence of a cosimplicial space, Am. J. Math. 109 (2) (1987) 361–394.
- [9] A.K. Bousfield, E.M. Friedlander, Homotopy theory of Γ -spaces, spectra, and bisimplicial sets, in: Geometric Applications of Homotopy Theory II, Proc. Conf., Evanston, Ill., 1977, in: Lecture Notes in Math., vol. 658, Springer, Berlin, 1978, pp. 80–130.
- [10] A.K. Bousfield, D.M. Kan, Homotopy Limits, Completions and Localizations, Lecture Notes in Mathematics, vol. 304, Springer-Verlag, Berlin-New York, 1972.
- [11] Tomasz Brzezinski, Robert Wisbauer, Corings and Comodules, London Mathematical Society Lecture Note Series, vol. 309, Cambridge University Press, Cambridge, 2003.

- [12] A.M. Cegarra, Josué Remedios, The relationship between the diagonal and the bar constructions on a bisimplicial set, *Topol. Appl.* 153 (1) (2005) 21–51.
- [13] Yukio Doi, Homological coalgebra, *J. Math. Soc. Jpn.* 33 (1) (1981) 31–50.
- [14] Gabriel C. Drummond-Cole, Joseph Hirsh, Model structures for coalgebras, *Proc. Am. Math. Soc.* 144 (4) (2016) 1467–1481.
- [15] W.G. Dwyer, Strong convergence of the Eilenberg-Moore spectral sequence, *Topology* 13 (1974) 255–265.
- [16] W.G. Dwyer, J.P.C. Greenlees, S. Iyengar, Duality in algebra and topology, *Adv. Math.* 200 (2) (2006) 357–402.
- [17] Anthony Elmendorf, Igor Kriz, Michael A. Mandell, Peter May, Rings, Modules, and Algebras in Stable Homotopy Theory, *Mathematical Surveys and Monographs*, vol. 47, American Mathematical Society, Providence, RI, 1997, With an appendix by M. Cole.
- [18] Marco A. Farinati, Andrea Solotar, Cyclic cohomology of coalgebras, coderivations and de Rham cohomology, in: *Hopf Algebras and Quantum Groups*, Brussels, 1998, in: *Lecture Notes in Pure and Appl. Math.*, vol. 209, Dekker, New York, 2000, pp. 105–129.
- [19] Richard Garner, Magdalena Kędziorek, Emily Riehl, Lifting accessible model structures, *J. Topol.* 13 (1) (2020) 59–76.
- [20] Ezra Getzler, Paul Goerss, A model category structure for differential graded coalgebras, unpublished.
- [21] Kathryn Hess, The Hochschild complex of a twisting cochain, *J. Algebra* 451 (2016) 302–356.
- [22] Kathryn Hess, Magdalena Kędziorek, Emily Riehl, Brooke Shipley, A necessary and sufficient condition for induced model structures, *J. Topol.* 10 (2) (2017) 324–369.
- [23] Kathryn Hess, Paul-Eugène Parent, Jonathan Scott, CoHochschild homology of chain coalgebras, *J. Pure Appl. Algebra* 213 (4) (2009) 536–556.
- [24] Kathryn Hess, Brooke Shipley, Waldhausen K -theory of spaces via comodules, *Adv. Math.* 290 (2016) 1079–1137.
- [25] Philip S. Hirschhorn, *Model Categories and Their Localizations*, *Mathematical Surveys and Monographs*, vol. 99, American Mathematical Society, Providence, RI, 2003.
- [26] Mark Hovey, *Model Categories*, *Mathematical Surveys and Monographs*, vol. 63, American Mathematical Society, Providence, RI, 1999.
- [27] Mark Hovey, Brooke Shipley, Jeff Smith, Symmetric spectra, *J. Am. Math. Soc.* 13 (1) (2000) 149–208.
- [28] ElHassan Idrissi, L’isomorphisme de Jones-McCleary et celui de Goodwillie sont des isomorphismes d’algèbres, *C. R. Acad. Sci. Paris Sér. I Math.* 331 (7) (2000) 507–510.
- [29] Bernhard Keller, On the cyclic homology of exact categories, *J. Pure Appl. Algebra* 136 (1) (1999) 1–56.
- [30] Bernhard Keller, On differential graded categories, in: *International Congress of Mathematicians*, vol. II, Eur. Math. Soc., Zürich, 2006, pp. 151–190.
- [31] Nicholas J. Kuhn, The McCord model for the tensor product of a space and a commutative ring spectrum, in: *Categorical Decomposition Techniques in Algebraic Topology*, Isle of Skye, 2001, in: *Progr. Math.*, vol. 215, Birkhäuser, Basel, 2004, pp. 213–236.
- [32] Kenji Lefèvre-Hasegawa, *Sur les A_∞ -catégories*, PhD thesis, Université Paris 7-Denis Diderot, 2003.
- [33] Jean-Louis Loday, *Cyclic Homology*, *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*, vol. 301, Springer-Verlag, Berlin, 1992, Appendix E by María O. Ronco.
- [34] Cary Malkiewich, Cyclotomic structure in the topological Hochschild homology of DX , *Algebraic Geom. Topol.* 17 (4) (2017) 2307–2356.
- [35] M.A. Mandell, J.P. May, Equivariant orthogonal spectra and S -modules, *Mem. Am. Math. Soc.* 159 (755) (2002), x+108.
- [36] M.A. Mandell, J.P. May, S. Schwede, B. Shipley, Model categories of diagram spectra, *Proc. Lond. Math. Soc.* (3) 82 (2) (2001) 441–512.
- [37] Randy McCarthy, The cyclic homology of an exact category, *J. Pure Appl. Algebra* 93 (3) (1994) 251–296.
- [38] Joseph Neisendorfer, *Algebraic Methods in Unstable Homotopy Theory*, *New Mathematical Monographs*, vol. 12, Cambridge University Press, Cambridge, 2010.
- [39] Maximilien Péroux, Brooke Shipley, Coalgebras in symmetric monoidal categories of spectra, *Homol. Homotopy Appl.* 21 (1) (2019) 1–18.
- [40] Stefan Schwede, S -modules and symmetric spectra, *Math. Ann.* 319 (3) (2001) 517–532.
- [41] Stefan Schwede, Brooke Shipley, Equivalences of monoidal model categories, *Algebraic Geom. Topol.* 3 (2003) 287–334.
- [42] Graeme Segal, Categories and cohomology theories, *Topology* 13 (1974) 293–312.
- [43] Brooke Shipley, Convergence of the homology spectral sequence of a cosimplicial space, *Am. J. Math.* 118 (1) (1996) 179–207.
- [44] Brooke Shipley, Monoidal uniqueness of stable homotopy theory, *Adv. Math.* 160 (2) (2001) 217–240.
- [45] R.H. Szczarba, The homology of twisted Cartesian products, *Trans. Am. Math. Soc.* 100 (1961) 197–216.
- [46] Mitsuhiro Takeuchi, Morita theorems for categories of comodules, *J. Fac. Sci., Univ. Tokyo, Sect. 1A, Math.* 24 (3) (1977) 629–644.
- [47] Bertrand Toën, Lectures on dg-categories, in: *Topics in Algebraic and Topological K-Theory*, in: *Lecture Notes in Math.*, vol. 2008, Springer, Berlin, 2011, pp. 243–302.