

## Accepted Manuscript

Title: Recommendation to reconsider examining cannabis subtypes together due to opposing effects on brain, cognition and behavior

Authors: Kristine Rømer Thomsen, Mette Buhl Callesen, Sarah W. Feldstein Ewing



PII: S0149-7634(17)30229-4  
DOI: <http://dx.doi.org/doi:10.1016/j.neubiorev.2017.05.025>  
Reference: NBR 2866

To appear in:

Received date: 18-3-2017  
Revised date: 15-5-2017  
Accepted date: 29-5-2017

Please cite this article as: Thomsen, Kristine Rømer, Callesen, Mette Buhl, Ewing, Sarah W. Feldstein, Recommendation to reconsider examining cannabis subtypes together due to opposing effects on brain, cognition and behavior. *Neuroscience and Biobehavioral Reviews* <http://dx.doi.org/10.1016/j.neubiorev.2017.05.025>

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

**Recommendation to reconsider examining cannabis subtypes together  
due to opposing effects on brain, cognition and behavior**

Dr. Kristine Rømer Thomsen<sup>1,\*</sup>, Dr. Mette Buhl Callesen<sup>1</sup>, and Dr. Sarah W. Feldstein Ewing<sup>2</sup>

<sup>1</sup> Centre for Alcohol and Drug Research, Department of Psychology and Behavioural Sciences, Aarhus University, Bartholins Allé 10, 8000 Aarhus C, Denmark.

<sup>2</sup> Department of Child and Adolescent Psychiatry, Oregon Health and Science University, Mail Code: DC7, 3181 SW Sam Jackson Park Rd, Portland, OR 97239-3098, USA.

\*Corresponding author: Kristine Rømer Thomsen, E-mail: krt.crf@psy.au.dk, Tel: +45 87165447

**Abstract**

Cannabis use represents a major public health issue throughout the globe. Yet, we still lack the most fundamental knowledge on long-term effects of cannabis on neural, cognitive, and behavioral function. Part of this stems from how cannabis has been measured historically. To this end, most empirical examinations of cannabis have consolidated all types of cannabis collectively. However, this approach obscures differences in how cannabinoids operate. In this commentary, we address the contrasting properties of tetrahydrocannabinol (THC) and cannabidiol (CBD) and their opposing effects on cognitive function. In addition, we address the increase in cannabis potency throughout the past two decades and how that impacts generalizability of early data to evaluations of contemporary public health. We underscore the urgent need for future research to disaggregate examination of THC from CBD, along with the importance of measuring cannabis potency to more effectively unravel its influence on cognitive function and other health issues.

**Keywords:** Cannabis; tetrahydrocannabinol; cannabidiol; potency; cognition; memory; learning.

This paper is a commentary on the review “A systematic review of the effect of cannabidiol on cognitive function: Relevance to schizophrenia” that evaluated preclinical and clinical literature on the effects of cannabidiol (CBD) on cognitive domains relevant to schizophrenia, and points to the ability of CBD to improve cognition across multiple disorders (Osborne et al., 2017). We seek to use the opportunity of this commentary to extend the implications of these findings to the field of addiction, where the convention in empirical examination is to consolidate all types of cannabis use collectively, despite considerable variations in nature of tetrahydrocannabinol (THC) and CBD. The timeliness of this is further supported by the increase in cannabis potency over the past two decades (Dujourdy and Besacier, 2017), which challenges generalizability of data gathered more than 20 years ago on contemporary estimates of the effects of cannabis on cognitive function.

For most quantitative brain and behavioral studies of cannabis, the standard empirical approach within addiction neuroscience and treatment is to collapse all types of cannabis use in collective examinations (National Academies of Sciences, 2017). Thus, individuals who use a wide range of products with varying levels of THC/CBD and across a vast range of potency are assessed together, and outcomes are reported without disaggregating effects for these different types of active ingredients, and/or levels of potency. Importantly, at this time, it is these *collective* outcomes that are guiding scientific debate and public policy decision-making throughout the globe regarding how cannabis affects brain, cognition, and behavior.

However, this approach obscures real-world differences in how cannabinoids operate. In actuality, cannabis use includes a range of subtypes that vary considerably in potency. This has become particularly evident with the introduction of high potency cannabis types to the market such as sinsemilla or “skunk” that contain high levels of THC (20%) and low levels of CBD (<0.5%)

(Osborne et al., 2017). Cannabis contains more than 80 different cannabinoid compounds, and growing evidence suggests that two of the main cannabinoids, THC and CBD (Figure 1), display opposing neural, cognitive, and behavioral effects, as reflected in recent comprehensive systematic reviews of CBD's antipsychotic properties (Iseger and Bossong, 2015) and ability to restore cognitive function (Osborne et al., 2017). While THC is a CB<sup>1</sup> and CB<sup>2</sup> receptor partial agonist (Pertwee, 2008), CBD is a negative allosteric modulator of the cannabinoid CB<sup>1</sup> receptor (Laprairie et al., 2015). Further, THC induces psychotic symptoms and anxiety in healthy volunteers (D'Souza et al., 2004; Morrison et al., 2009), whereas CBD demonstrates antipsychotic and anxiolytic effects (Leweke et al., 2012; Zuardi et al., 2009; Zuardi et al., 2006; Zuardi et al., 1995). Evidence from clinical studies indicates that THC impairs learning and memory (including working memory) in healthy volunteers and cannabis users (D'Souza et al., 2004; Morgan et al., 2012; Morrison et al., 2009), whereas CBD enhances learning and memory and inhibits THC-elicited decline in learning and memory (including working memory) (Das et al., 2013; Englund et al., 2013; Morgan et al., 2012). Moreover, a recent study of cannabis users implicates THC in impaired facial emotional recognition, while CBD improves facial emotional recognition and attenuates THC-induced impairment (Hindocha et al., 2015). These effects are mirrored in functional imaging studies, which reveal opposing acute effects of THC and CBD in areas pivotal to queried cognitive processes including emotional processing (amygdala); salience processing (striatum, hippocampus, prefrontal cortex); and processing of auditory and visual information (auditory and visual cortex) (Bhattacharyya et al., 2010; Borgwardt et al., 2008; Fusar-Poli et al., 2009; Winton-Brown et al., 2011). It is relevant to note however, that research investigating the impact of CBD on human functioning is still sparse and suffers from limitations. For instance, Morgan et al. (2012) base their conclusions on hair samples from users with varying levels of THC and CBD, and Das et al. (2013) only assessed CBD and not its effects on THC. In general, double-blind placebo controlled

assessments are needed to improve our knowledge on the impact of THC and CBD, as well as cannabis with and without CBD, on cognition and behavior. Nevertheless, the available studies still reflect critical and relevant cognitive differences that should be further explored in future studies.

At the same time, possible long-term detrimental effects of cannabis on cognitive function continue to be at the center of scientific debate (National Academies of Sciences, 2017). Recent prospective studies (Jackson et al., 2016; Mokrysz et al., 2016) and systematic reviews (Broyd et al., 2016; Curran et al., 2016; National Academies of Sciences, 2017) show limited to mixed findings on the impact of cannabis in cognitive processing, that contrast with highly-cited findings from Meier and colleagues (2012) linking heavy cannabis use with IQ decline. Importantly, none of these empirical studies disentangled the role of THC from CBD, and/or examined the role of potency (Feldstein Ewing et al., 2017). These types of examinations have yet to be carefully scrutinized in prospective designs to investigate possible long-term effects of cannabis on cognitive functioning. Similarly, an increased focus on disaggregating the role of THC from CBD, and/or the role of potency in studies of acute effects of cannabis, also promises more robust and consistent findings (National Academies of Sciences, 2017).

When examined on a subtype level, cannabis has historically been assigned to two predominant types: herbal cannabis and cannabis resin. These categories contain even more subtypes, which at a minimum, vary considerably in potency (e.g., cannabis plant; shatter). Further, the most comprehensive study available (Dujourdy and Besacier, 2017), which examined cannabis samples from five French forensic police laboratories over 25 years, reflects that during the past two decades, there has been a gradual increase in potency in both subtypes, documented through increases in THC levels as well as calculation of THC/CBD ratios. In herbal cannabis, mean THC

content rose from 2% in 1995 to 13% in 2016; and in the past four years, the increase in potency in cannabis resin has been dramatic (10% mean THC content in 2009 to 23% in mid-2016) (Dujourdy and Besacier, 2017). This escalation in cannabis potency is comparable to findings in other European countries (Niesink et al., 2015; Zamengo et al., 2015) and the US (National Academies of Sciences, 2017).

The dramatic increase in potency also presents scientific and public health challenges in the generalizability and extension of early cannabis data on current evaluations of the implication of cannabis use on brain, cognition, and behavior. For example, the dramatic increase in potency in cannabis resin the past four years poses specific challenges in the generalizability of cannabis resin data gathered prior to 2012. We propose that these nuances are, in fact, critical health issues that must be carefully examined and addressed in order to truly begin to coordinate and interpret early data on public health from data collected today. At a minimum, these data underscore the urgent need to disaggregate examination of THC from CBD, along with the importance of measuring potency in future studies of cannabis and its impact on cognitive function and other health related issues across the globe.

**Funding:**

This work was funded by a block grant for the Danish Ministry for Social Affairs and the Interior (KRT, MBC) and by 3R01AA023658-02S1 from the National Institutes of Health, USA (SWFE). The funding sources had no role in the writing of the manuscript or the decision to submit it for publication.

**Contributors:**

KRT conceptualized the commentary and wrote the first draft of the commentary. KRT and SWFE performed the literary searches. All authors contributed to and have approved the final manuscript.

### **Conflict of interest:**

The authors declare no conflict of interest.

### **References:**

- Bhattacharyya, S., Morrison, P.D., Fusar-Poli, P., Martin-Santos, R., Borgwardt, S., Winton-Brown, T., Nosarti, C., CM, O.C., Seal, M., Allen, P., Mehta, M.A., Stone, J.M., Tunstall, N., Giampietro, V., Kapur, S., Murray, R.M., Zuardi, A.W., Crippa, J.A., Atakan, Z., McGuire, P.K., 2010. Opposite effects of delta-9-tetrahydrocannabinol and cannabidiol on human brain function and psychopathology. *Neuropsychopharmacology* 35, 764-774.
- Borgwardt, S.J., Allen, P., Bhattacharyya, S., Fusar-Poli, P., Crippa, J.A., Seal, M.L., Fraccaro, V., Atakan, Z., Martin-Santos, R., O'Carroll, C., Rubia, K., McGuire, P.K., 2008. Neural basis of Delta-9-tetrahydrocannabinol and cannabidiol: effects during response inhibition. *Biol Psychiatry* 64, 966-973.
- Broyd, S.J., van Hell, H.H., Beale, C., Yucel, M., Solowij, N., 2016. Acute and Chronic Effects of Cannabinoids on Human Cognition-A Systematic Review. *Biol Psychiatry* 79, 557-567.
- Curran, H.V., Freeman, T.P., Mokrysz, C., Lewis, D.A., Morgan, C.J., Parsons, L.H., 2016. Keep off the grass? Cannabis, cognition and addiction. *Nat Rev Neurosci* 17, 293-306.
- D'Souza, D.C., Perry, E., MacDougall, L., Ammerman, Y., Cooper, T., Wu, Y.T., Braley, G., Gueorguieva, R., Krystal, J.H., 2004. The psychotomimetic effects of intravenous delta-9-tetrahydrocannabinol in healthy individuals: implications for psychosis. *Neuropsychopharmacology* 29, 1558-1572.

- Das, R.K., Kamboj, S.K., Ramadas, M., Yogan, K., Gupta, V., Redman, E., Curran, H.V., Morgan, C.J., 2013. Cannabidiol enhances consolidation of explicit fear extinction in humans. *Psychopharmacology (Berl)* 226, 781-792.
- Dujourdy, L., Besacier, F., 2017. A study of cannabis potency in France over a 25 years period (1992–2016). *Forensic Sci Int* 272, 72-80.
- Englund, A., Morrison, P.D., Nottage, J., Hague, D., Kane, F., Bonaccorso, S., Stone, J.M., Reichenberg, A., Brenneisen, R., Holt, D., Feilding, A., Walker, L., Murray, R.M., Kapur, S., 2013. Cannabidiol inhibits THC-elicited paranoid symptoms and hippocampal-dependent memory impairment. *J Psychopharmacol* 27, 19-27.
- Feldstein Ewing, S.W., Lovejoy, T.I., Choo, E.K., 2017. How Has Legal Recreational Cannabis Affected Adolescents in Your State? A Window of Opportunity. *Am J Public Health* 107, 246-247.
- Fusar-Poli, P., Crippa, J.A., Bhattacharyya, S., Borgwardt, S.J., Allen, P., Martin-Santos, R., Seal, M., Surguladze, S.A., O'Carroll, C., Atakan, Z., Zuardi, A.W., McGuire, P.K., 2009. Distinct effects of  $\Delta^9$ -tetrahydrocannabinol and cannabidiol on neural activation during emotional processing. *Arch Gen Psychiatry* 66, 95-105.
- Hindocha, C., Freeman, T.P., Schafer, G., Gardener, C., Das, R.K., Morgan, C.J., Curran, H.V., 2015. Acute effects of delta-9-tetrahydrocannabinol, cannabidiol and their combination on facial emotion recognition: a randomised, double-blind, placebo-controlled study in cannabis users. *Eur Neuropsychopharmacol* 25, 325-334.
- Iseger, T.A., Bossong, M.G., 2015. A systematic review of the antipsychotic properties of cannabidiol in humans. *Schizophr Res* 162, 153-161.
- Jackson, N.J., Isen, J.D., Khoddam, R., Irons, D., Tuvblad, C., Iacono, W.G., McGue, M., Raine, A., Baker, L.A., 2016. Impact of adolescent marijuana use on intelligence: Results from two longitudinal twin studies. *Proc Natl Acad Sci U S A* 113, E500-508.



- Laprairie, R.B., Bagher, A.M., Kelly, M.E., Denovan-Wright, E.M., 2015. Cannabidiol is a negative allosteric modulator of the cannabinoid CB1 receptor. *Br J Pharmacol* 172, 4790-4805.
- Leweke, F.M., Piomelli, D., Pahlisch, F., Muhl, D., Gerth, C.W., Hoyer, C., Klosterkötter, J., Hellmich, M., Koethe, D., 2012. Cannabidiol enhances anandamide signaling and alleviates psychotic symptoms of schizophrenia. *Transl Psychiatry* 2, e94.
- Meier, M.H., Caspi, A., Ambler, A., Harrington, H., Houts, R., Keefe, R.S., McDonald, K., Ward, A., Poulton, R., Moffitt, T.E., 2012. Persistent cannabis users show neuropsychological decline from childhood to midlife. *Proc Natl Acad Sci U S A* 109, E2657-2664.
- Mokrysz, C., Landy, R., Gage, S.H., Munafo, M.R., Roiser, J.P., Curran, H.V., 2016. Are IQ and educational outcomes in teenagers related to their cannabis use? A prospective cohort study. *J Psychopharmacol* 30, 159-168.
- Morgan, C.J., Gardener, C., Schafer, G., Swan, S., Demarchi, C., Freeman, T.P., Warrington, P., Rupasinghe, I., Ramoutar, A., Tan, N., Wingham, G., Lewis, S., Curran, H.V., 2012. Sub-chronic impact of cannabinoids in street cannabis on cognition, psychotic-like symptoms and psychological well-being. *Psychol Med* 42, 391-400.
- Morrison, P.D., Zois, V., McKeown, D.A., Lee, T.D., Holt, D.W., Powell, J.F., Kapur, S., Murray, R.M., 2009. The acute effects of synthetic intravenous Delta9-tetrahydrocannabinol on psychosis, mood and cognitive functioning. *Psychol Med* 39, 1607-1616.
- National Academies of Sciences, E., and Medicine, 2017. The health effects of cannabis and cannabinoids: The current state of evidence and recommendations for research, Washington, DC.
- Niesink, R.J., Rigter, S., Koeter, M.W., Brunt, T.M., 2015. Potency trends of Delta9-tetrahydrocannabinol, cannabidiol and cannabinol in cannabis in the Netherlands: 2005-15. *Addiction* 110, 1941-1950.

- Osborne, A.L., Solowij, N., Weston-Green, K., 2017. A systematic review of the effect of cannabidiol on cognitive function: Relevance to schizophrenia. *Neurosci Biobehav Rev* 72, 310-324.
- Pertwee, R.G., 2008. The diverse CB1 and CB2 receptor pharmacology of three plant cannabinoids: delta9-tetrahydrocannabinol, cannabidiol and delta9-tetrahydrocannabivarin. *Br J Pharmacol* 153, 199-215.
- Winton-Brown, T.T., Allen, P., Bhattacharyya, S., Borgwardt, S.J., Fusar-Poli, P., Crippa, J.A., Seal, M.L., Martin-Santos, R., Ffytche, D., Zuardi, A.W., Atakan, Z., McGuire, P.K., 2011. Modulation of auditory and visual processing by delta-9-tetrahydrocannabinol and cannabidiol: an FMRI study. *Neuropsychopharmacology* 36, 1340-1348.
- Zamengo, L., Frison, G., Bettin, C., Sciarrone, R., 2015. Cannabis potency in the Venice area (Italy): update 2013. *Drug Test Anal* 7, 255-258.
- Zuardi, A.W., Crippa, J.A., Hallak, J.E., Pinto, J.P., Chagas, M.H., Rodrigues, G.G., Dursun, S.M., Tumas, V., 2009. Cannabidiol for the treatment of psychosis in Parkinson's disease. *J Psychopharmacol* 23, 979-983.
- Zuardi, A.W., Hallak, J.E., Dursun, S.M., Morais, S.L., Sanches, R.F., Musty, R.E., Crippa, J.A., 2006. Cannabidiol monotherapy for treatment-resistant schizophrenia. *J Psychopharmacol* 20, 683-686.
- Zuardi, A.W., Morais, S.L., Guimaraes, F.S., Mechoulam, R., 1995. Antipsychotic effect of cannabidiol. *J Clin Psychiatry* 56, 485-486.

**Figure 1. Chemical structure of tetrahydrocannabinol (A) and cannabidiol (B).**

Printed with permission from Hayakawa K, Mishima K, Fujiwara M. - Pharmaceuticals. 2010

Jul;3(7):2197–212. <http://dx.doi.org/10.3390/ph3072197>, CC BY 3.0,

<https://commons.wikimedia.org/w/index.php?curid=40172369>

