



ELSEVIER

Contents lists available at ScienceDirect

Journal of Number Theory

www.elsevier.com/locate/jnt



Integral cohomology of certain Picard modular surfaces

Dan Yasaki

*Department of Mathematics and Statistics, The University of North Carolina at Greensboro,
Greensboro, NC 27412, USA*

ARTICLE INFO

Article history:

Received 7 September 2007
 Received in revised form 8 July 2013
 Accepted 8 July 2013
 Available online 12 September 2013
 Communicated by David Goss

MSC:

primary 11F75

Keywords:

Picard modular group
 Locally symmetric space
 Cohomology of arithmetic subgroups
 Spine

ABSTRACT

Let $\bar{\Gamma}$ be the Picard modular group of an imaginary quadratic number field k and let \mathcal{D} be the associated symmetric space. Let $\Gamma \subset \bar{\Gamma}$ be a congruence subgroup. We describe a method to compute the integral cohomology of the locally symmetric space $\Gamma \backslash \mathcal{D}$. The method is implemented for the cases $k = \mathbb{Q}(i)$ and $k = \mathbb{Q}(\sqrt{-3})$, and the cohomology is computed for various Γ .

© 2013 Elsevier Inc. All rights reserved.

1. Introduction

Let $\bar{\Gamma} = \mathrm{SU}(2, 1; \mathcal{O}_k)$ be the Picard modular group of an imaginary quadratic number field k and let \mathcal{D} be the associated symmetric space. Let $\Gamma \subset \bar{\Gamma}$ be a congruence subgroup. Although \mathcal{D} is 4-dimensional, the virtual cohomological dimension of $\bar{\Gamma}$ is 3. Hence the cohomology $H^i(\Gamma \backslash \mathcal{D})$ vanishes for $i > 3$.

If Γ is torsion-free, then $\Gamma \backslash \mathcal{D}$ is an Eilenberg–MacLane space for Γ . It follows that the group cohomology of Γ with trivial complex coefficients is isomorphic to the complex cohomology of the locally symmetric space,

E-mail address: d_yasaki@uncg.edu.

URL: http://www.uncg.edu/~d_yasaki/.

$$H^*(\Gamma; \mathbb{C}) \simeq H^*(\Gamma \backslash \mathcal{D}; \mathbb{C}). \quad (1)$$

In fact, (1) remains true while using complex coefficients when Γ has torsion, but is not true in general for integral coefficients.

A deep result of Franke [7] describes a relationship between cohomology and automorphic forms. The cohomology groups $H^*(\Gamma; \mathbb{C})$, or more generally $H^*(\Gamma; M)$ for any complex finite dimensional rational representation of $\mathrm{SU}(2, 1)$, decompose into cuspidal cohomology and Eisenstein cohomology. The cuspidal cohomology of Γ can be represented by cuspidal automorphic forms. It is possible to compute the space of cuspidal Picard modular forms with the Hecke action using the Jacquet–Langlands correspondence [5]. However, these methods will not compute the torsion classes in the integral cohomology of the group or the locally symmetric space. In this paper, we use topological methods to compute the torsion classes in the integral cohomology of the locally symmetric space $\Gamma \backslash \mathcal{D}$.

There is a 3-dimensional cell complex W , known as a spine, that can be used to compute the integral cohomology of the locally symmetric space $\Gamma \backslash \mathcal{D}$. The existence of such a W for general \mathbb{Q} -rank 1 groups is known [9], but there are few explicit examples for non-linear symmetric spaces. We outline a method of computing W for $\mathrm{SU}(2, 1; \mathcal{O}_k)$, where k is an imaginary quadratic number field with class number 1. The structure of the spine is computed in [10] for the Gaussian case $k = \mathbb{Q}(i)$ and is computed here for the Eisenstein case $k = \mathbb{Q}(\sqrt{-3})$. Falbel and Parker [6] do similar computations for the Eisenstein–Picard modular group, but with a different purpose. They exhibit fundamental domain for the action of $\bar{\Gamma}$ on \mathcal{D} . Using the structure and combinatorics of the fundamental domain, they deduce a presentation for $\bar{\Gamma}$.

The outline of the paper is as follows. We first recall the Picard modular group and associated symmetric space in Section 2. A method of computing the spine W is given in Section 3. Section 4 outlines the method of [2] and its implications in the context of our cell complex. The cell complex is computed in [10] for the Picard modular group over the Gaussian integers. We compute the cell complex and give the stabilizers for Picard modular group over the Eisenstein integers in Section 5. Finally, Section 6 gives the cohomology computation results.

2. Preliminaries

Let k be an imaginary quadratic field with discriminant D and ring of integers \mathcal{O} . Thus $k = \mathbb{Q}(\sqrt{D})$ and D is either square-free and $D \equiv 1 \pmod{4}$ or $D = 4D'$, where D' is square-free and $D' \equiv 2 \pmod{4}$ or $D' \equiv 3 \pmod{4}$. Then \mathcal{O} is generated by 1 and $\omega = (D + \sqrt{D})/2$. For $d = 1, 2, 3, 7, 11, 19, 43, 67, 163$, $\mathbb{Q}(\sqrt{-d})$ has class number $h(k) = 1$ and \mathcal{O} is a principal ideal domain. Fix an imaginary quadratic field k with class number 1.

2.1. The unitary group

Let V be a 3-dimensional k -vector space with an integral structure given by an \mathcal{O} -lattice $L \subset V$. Let $\mathcal{Q}: V \times V \rightarrow k$ be a non-degenerate Hermitian form on V which is \mathcal{O} -valued on L and whose signature is $(2, 1)$ on $V_{\mathbb{R}}$. Then $\Gamma = \mathrm{SU}(\mathcal{Q}, V)$ is a semisimple algebraic group defined over \mathbb{Q} whose group of real points is isomorphic to $\mathrm{SU}(2, 1)$. The Picard Modular Group of k is

$$\bar{\Gamma} = \mathbf{G}(\mathbb{Z}) = \{ \gamma \in \mathbf{G}(\mathbb{Q}) \mid \gamma L = L \}.$$

For this paper, fix an embedding of k in \mathbb{C} and a k -basis of V with respect to which \mathcal{Q} is represented by

$$\mathcal{Q}(u, v) = u^* C v, \quad \text{where } C = \begin{pmatrix} 0 & 0 & \sqrt{D}^{-1} \\ 0 & 1 & 0 \\ -\sqrt{D}^{-1} & 0 & 0 \end{pmatrix}.$$

In particular,

$$G = \mathbf{G}(\mathbb{R}) = \{ g \in \mathrm{SL}(3, \mathbb{C}) \mid g^* C g = C \}.$$

2.2. Symmetric space

Let θ denote the Cartan involution given by inverse conjugate transpose and let K be the fixed points under θ . Let $\mathcal{D} = G/K$ be the associated Riemannian symmetric space of non-compact type. The symmetric space $\mathcal{D} = G/K$ has many useful realizations.

Using horospherical coordinates (y, β, r) we can view the symmetric space as $\mathbb{R}_{>0} \times \mathbb{C} \times \mathbb{R}$ as follows. Let $P_0 \subseteq G$ be the rational parabolic subgroup of upper triangular matrices

$$P_0 = \left\{ \begin{pmatrix} y\zeta & \beta\zeta^{-2} & \frac{\zeta}{y} \left(r - \frac{|\beta|^2}{2\sqrt{D}} \right) \\ 0 & \zeta^{-2} & -\frac{\bar{\beta}\zeta}{\sqrt{D}y} \\ 0 & 0 & \zeta/y \end{pmatrix} \mid \zeta, \beta \in \mathbb{C}, |\zeta| = 1, r \in \mathbb{R}, y \in \mathbb{R}_{>0} \right\}.$$

P_0 acts transitively on \mathcal{D} , and every point $X \in \mathcal{D}$ can be written as pK for some $p \in P_0$. When p is written as above, the point $X = pK$ is independent of ζ , and so we will denote such a point $X = (y, \beta, r)$.

A line l in V is said to be *negative* if

$$\mathcal{Q}(v, v) < 0, \quad \text{where } v \text{ is any vector on } l.$$

Let \mathcal{L} denote the set negative lines in V . The group G acts transitively on \mathcal{L} , and the stabilizer of a negative line is a maximal compact subgroup of G . Thus we can identify \mathcal{D} with \mathcal{L} . A negative line is the span of a vector with non-zero third component, and

so each line can be identified with a vector $(z, u, 1)^t$. This identification gives the Siegel domain realization of \mathcal{D} ,

$$\mathcal{D} = \left\{ (z, u) \in \mathbb{C}^2 \mid |u|^2 < \frac{2 \operatorname{Im}(z)}{\sqrt{|D|}} \right\}.$$

In coordinates, one computes that

$$z = r + iy^2 - \frac{|\beta|^2}{2\sqrt{D}} \quad \text{and} \quad u = \frac{-\bar{\beta}}{\sqrt{D}}. \tag{2}$$

2.3. The cusps

The cusps correspond to rational isotropic lines in V . These lines are in 1–1 correspondence with the rational parabolic subgroups. In particular, every rational parabolic subgroup stabilizes a rational isotropic line. We associate to each rational parabolic subgroup P an isotropic vector $v_P = (n, p, q)^t \in \mathcal{O}^3$ from the line stabilized by P such that the ideal $(n, p, q) = \mathcal{O}$. Note that this vector is well-defined up to multiplication by \mathcal{O}^* . Thus for the rest of the paper, we identify vectors in \mathcal{O}^3 that differ by \mathcal{O}^* . The isotropic condition $\mathcal{Q}(v_P, v_P) = 0$ implies that $v_P = (n, p, q)^t$ satisfies

$$|p|^2 = \frac{1}{\sqrt{D}}(n\bar{q} - \bar{n}q). \tag{3}$$

Let \mathcal{P} denote the set of proper rational parabolic subgroups of G . These parabolic subgroups are conjugate to P_0 via elements of $\mathbf{G}(\mathbb{Q})$. There is a natural action of $\bar{\Gamma}$ on \mathcal{P} given by conjugation and denoted

$${}^g P := gPg^{-1}.$$

The quotient $\bar{\Gamma} \backslash \mathcal{P}$ is a finite set whose cardinality is the class number of $\bar{\Gamma}$. Zink has shown that $h(\bar{\Gamma}) = h(k)$ [11]. In particular, since $h(k) = 1$, every rational parabolic subgroup is conjugate to P_0 via an element of $\bar{\Gamma}$.

2.4. The spine

There is a $\bar{\Gamma}$ -invariant decomposition of \mathcal{D} into codimension 0 sets using exhaustion functions. The union \mathcal{D}_0 of the boundaries of these sets forms a contractible $\bar{\Gamma}$ -equivariant retraction of \mathcal{D} that is known as a spine. This construction is described for the general \mathbb{Q} -rank 1 case in [9] and is analogous to Ash’s well-rounded retraction [1] for linear symmetric spaces.

There is a collection of subsets of rational parabolic subgroups \mathcal{S} called strongly admissible subsets that give a decomposition of the spine

$$\mathcal{D}_0 = \coprod_{\substack{\mathcal{I} \in \mathcal{S} \\ |\mathcal{I}| > 1}} \mathcal{D}'(\mathcal{I}).$$

The group $\bar{\Gamma}$ preserves this decomposition. In particular,

$$\gamma \cdot \mathcal{D}'(\mathcal{I}) = \mathcal{D}'(\gamma \mathcal{I})$$

for every $\gamma \in \bar{\Gamma}$.

The decomposition of \mathcal{D}_0 may be refined to a regular cell complex W that has the property that the stabilizer of each cell fixes the cell pointwise. Thus, the cohomology of $\bar{\Gamma} \backslash \mathcal{D}$ can be described by finite combinatorial data.

2.5. Congruence subgroups

Let $N \in \mathcal{O}$ and let $\bar{k} = \mathcal{O}/N\mathcal{O}$. Let $\bar{V} = \bar{k}^3$ and $\mathbb{P}^2 = \mathbb{P}(\bar{V})$. By \mathbb{P}^2 , we mean the set of vectors $(x_1, x_2, x_3) \in V$ that are primitive in the sense that the ideal $(x_1, x_2, x_3) = \mathcal{O}$. When N is prime, this is the usual projective space over the residue field \bar{k} . The equivalence class of the vector (x_1, x_2, x_3) will be denoted $[x_1 : x_2 : x_3]$. We view these triples as row vectors, and so $\bar{\Gamma}$ acts on \bar{V} and \mathbb{P}^2 on the right.

Let $\Gamma_1(N) \subset \bar{\Gamma}$ denote the stabilizer of the vector $(0, 0, 1) \in V$, and let $\Gamma_0(N) \subset \bar{\Gamma}$ denote the stabilizer of the point $[0 : 0 : 1] \in \mathbb{P}$. Let $\Gamma(N)$ denote the kernel of the map $\mu: \bar{\Gamma} \rightarrow \text{SL}_3(\bar{k})$.

3. Computing the spine

In any given example, one must understand the exhaustion functions to compute the spine. The functions are parametrized by rational parabolic subgroups can be thought of as height functions with respect to the various cusps. The family of exhaustion functions are $\bar{\Gamma}$ -invariant in the sense that

$$f_{\gamma P}(X) = f_P(\gamma^{-1} \cdot X) \quad \text{for } \gamma \in \bar{\Gamma}. \tag{4}$$

These exhaustion functions for $k = \mathbb{Q}(i)$ are described in detail in [10]. Here, we give the exhaustion functions in coordinates for an imaginary quadratic field $k = \mathbb{Q}(\sqrt{D})$ with class number one.

3.1. The exhaustion functions

Let $X = (z, u) \in \mathcal{D}$ and let P be a rational parabolic subgroup of G with associated isotropic vector $v_P = (n, p, q)^t$. Then the exhaustion function f_P can be written as

$$f_0(X) := f_{P_0}(X) = \sqrt{-\frac{\sqrt{|D|}}{2} \mathcal{Q}(X, X)} = y, \tag{5}$$

$$f_P(X) = \frac{f_0(X)}{|\mathcal{Q}(\sqrt{D}X, v_P)|} = \frac{y}{|q\bar{z} + \sqrt{D}p\bar{u} - n|}. \tag{6}$$

These exhaustion functions are used to define the decomposition of \mathcal{D}_0 into sets $\mathcal{D}'(\mathcal{I})$ for $\mathcal{I} \subset \mathcal{P}$. Let $\mathcal{D}(\mathcal{I}) \subset \mathcal{D}$ be the set of $X \in \mathcal{D}$ such that $f_P(X) \geq f_Q(X)$ for every $P \in \mathcal{I}$ and $Q \in \mathcal{P} \setminus \mathcal{I}$ and $f_P(X) = f_{P'}(X)$ for every $P, P' \in \mathcal{I}$. Define $\mathcal{D}'(\mathcal{I}) \subset \mathcal{D}(\mathcal{I})$ to be the subset where the inequality is strict, $f_P(X) > f_Q(X)$ for every $P \in \mathcal{I}$ and $Q \in \mathcal{P} \setminus \mathcal{I}$. In other words, $\mathcal{D}'(\mathcal{I})$ consists of the points that are higher with respect to $P \in \mathcal{I}$ than any other cusps. The subset \mathcal{I} is *admissible* if $\mathcal{D}(\mathcal{I}) \neq \emptyset$ and *strongly admissible* if $\mathcal{D}'(\mathcal{I}) \neq \emptyset$.

3.2. First contacts

Definition 3.1. For a finite subset $\mathcal{I} \subset \mathcal{P}$, the *first contact* for \mathcal{I} is the subset of

$$\{X \in \mathcal{D} \mid f_P(X) = f_Q(X) \text{ for every } P, Q \in \mathcal{P}\}$$

where the exhaustion functions f_P attain their maximum.

Proposition 3.2. Let $P \neq P_0$ be a rational parabolic subgroup and let $v_P = (n, p, q)^t$ be the associated isotropic vector. Then the first contact for $\{P_0, P\}$ is the point $X = (z, u)$, where

$$z = \frac{n}{q} + \frac{i}{|q|} \quad \text{and} \quad u = \frac{p}{q}.$$

In particular, for the first contact X we have

$$f_0(X)^2 = \frac{1}{|q|} = \frac{1}{\sqrt{|D|} |\mathcal{Q}(v_0, v_P)|}.$$

More generally, the first contact \hat{X} for rational parabolic subgroups $\{Q, R\}$ satisfies

$$f_Q(\hat{X})^2 = \frac{1}{\sqrt{|D|} |\mathcal{Q}(v_Q, v_R)|}.$$

Proof. To find the first contact for $\{P_0, P\}$, we need to find the point $X = (z, u)$ such that $f_0(X)$ is maximal on the set

$$|\bar{z} + \sqrt{D}p\bar{u}/q - n/q| = 1/|q|.$$

Using (2) to express z in its real and imaginary parts,

$$|r - iy^2 - \sqrt{D}|u|^2/2 + \sqrt{D}\bar{u}p/q - n/q| = 1/|q|,$$

we see that we can pick r such that the real part of the expression is 0. Thus it suffices to consider points where

$$|y^2 + \sqrt{|D|}|u|^2/2 - \text{Im}(\sqrt{D}\bar{u}p/q) + \text{Im}(n/q)| = 1/|q|.$$

Since v_P is isotropic, the left side of the equation is

$$|y^2 + \sqrt{|D|}|p|^2/(2|q|^2) + \sqrt{|D|}|u|^2/2 - \text{Im}(\sqrt{D}\bar{u}p/q)| = 1/|q|,$$

and hence the first contact point occurs at the minimum of

$$\sqrt{|D|}|u|^2/2 - \text{Im}(\sqrt{D}\bar{u}p/q) = \sqrt{|D|}(|u|^2/2 - \text{Re}(\bar{u}p/q)).$$

One can compute that this implies that the first contact occurs when $u = p/q$. Plugging back in, we get that $y = 1/\sqrt{|q|}$ and $r = \text{Re}(n/q)$, or equivalently,

$$z = \frac{n}{q} + \frac{i}{|q|} \quad \text{as desired.}$$

For the second statement, suppose Q and R are distinct rational parabolic subgroups. Let \hat{X} denote the first contact for $\{Q, R\}$. There exists $\gamma \in \bar{\Gamma}$ such that ${}^\gamma Q = P_0$, ${}^\gamma R = P$ for some P . Note that $\gamma \cdot \hat{X}$ is the first contact for $\{P_0, P\}$. By the $\bar{\Gamma}$ -invariance of the exhaustion functions, it follows that

$$f_Q(\hat{X})^2 = f_{P_0}(\gamma \cdot \hat{X})^2 = \frac{1}{\sqrt{|D|}|\mathcal{Q}(v_{P_0}, v_P)|}.$$

Since $\bar{\Gamma}$ preserves \mathcal{Q} , $|\mathcal{Q}(v_{P_0}, v_P)| = |\mathcal{Q}(v_Q, v_R)|$ and the result follows. \square

3.3. Candidates

We will now compute a finite list subsets of rational parabolic subgroups from which a complete set of $\bar{\Gamma}$ -representative of strongly admissible sets can be chosen.

First, compute all the possible admissible sets of order 2 as follows. Since $h(\bar{\Gamma}) = 1$, every admissible set is $\bar{\Gamma}$ -conjugate to an admissible set containing P_0 . Let $\mathcal{I} = \{P_0, P\}$ be such an admissible set. Then by Proposition 3.2, if P has associated isotropic vector $v_P = (n, p, q)^t$ then $|q| \leq 1/\mu^2$. Since $q \in \mathcal{O}$, this is a finite list of possibilities for q . Furthermore, using $\bar{\Gamma}_{P_0} = \bar{\Gamma} \cap P_0$, we can move a point $X = (z, u)$ to a point $X' = (z', u')$, where u' lies in a fundamental domain for \mathbb{C} modulo translation by \mathcal{O} and $f_0(X) = f_0(X')$. In particular, we can arrange that the u -component of the first contact of \mathcal{I} lies in a fundamental domain for \mathbb{C} modulo translation by \mathcal{O} . Since the first contact of $\{P_0, P\}$ has $u = p/q$, it follows that there are finitely many choices for p . Finally, since v_P is isotropic and there are only finitely many choices for p and q , (3) implies there are finitely many choices for n .

Once we have a finite list of candidates for $\bar{\Gamma}$ -representatives of admissible sets of order two, we check if it is possible to find candidates of order greater than two that contain these candidates. We repeatedly use [Proposition 3.2](#). In particular, if \mathcal{I} is admissible, then

$$|\mathcal{Q}(v_Q, v_R)| \leq \frac{1}{\sqrt{|D|\mu^2}} \quad \text{for every } Q, R \in \mathcal{I}.$$

Given such an \mathcal{I} , we can check if it is admissible by seeing if $\mathcal{D}(\mathcal{I}) \neq \emptyset$.

We see that the isotropic vectors provide a convenient way to study admissible sets. If the value of $|\mathcal{Q}(v_P, v_Q)|$ is small enough, this invariant is enough to classify the $\bar{\Gamma}$ -conjugacy class of $\{P, Q\}$.

Proposition 3.3. *If $\mathcal{I} = \{P, Q\}$ is a subset of rational parabolic subgroups such that $|\mathcal{Q}(v_P, v_Q)|^2 = 1/|D|$, then \mathcal{I} is $\bar{\Gamma}$ -equivalent to $\{P_0, P_w\}$, where P_w is the rational parabolic subgroup that stabilizes the line through $(0, 0, 1)^t$.*

Proof. Since the class number is 1, every rational parabolic subgroup is $\bar{\Gamma}$ -conjugate to P_0 . Without loss of generality, we can assume that $v_P = (1, 0, 0)^t$. Then since $|\mathcal{Q}(v_P, v_Q)|^2 = 1/|D|$, v_Q is of the form $v = (n, p, 1)$ for some $n, p \in \mathcal{O}$. We can multiply by any matrix in $\bar{\Gamma}_{N_0} = \bar{\Gamma} \cap N_0$ without changing $(1, 0, 0)^t$. In particular,

$$\begin{pmatrix} 1 & -\bar{p}\sqrt{D} & |p|^2\omega \\ 0 & 1 & -p \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} n \\ p \\ 1 \end{pmatrix} = \begin{pmatrix} n' \\ 0 \\ 1 \end{pmatrix}.$$

Note that since $(n', 0, 1)^t$ is isotropic, [\(3\)](#) implies that $n' \in \mathbb{R}$. It follows that $n' \in \mathbb{Z}$ and

$$\begin{pmatrix} 1 & 0 & -n' \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} \in \bar{\Gamma}_{N_0}.$$

Thus we can arrange that $\{u, v\}$ is a $\bar{\Gamma}$ -translate of a set of the form

$$\{(1, 0, 0)^t, (0, 0, 1)^t\}. \quad \square$$

4. Cohomology

We follow [\[2\]](#) closely and refer to that paper for the details. Many of the complications arising from orientation issues disappear in our case since we subdivide to get a cell complex such that the stabilizer of each cell fixes the cell pointwise. For the remainder of the paper, let Γ be a congruence group of level N ($\Gamma(N)$, $\Gamma_1(N)$, or $\Gamma_0(N)$) and let $\bar{\Gamma} = \text{SU}(2, 1; \mathcal{O})$.

4.1. Cell complex

The decomposition of the spine has a refinement W into a regular cell complex such that the stabilizer of each cell fixes the cell pointwise. We use W to compute the integral cohomology of $\Gamma \backslash \mathcal{D}$.

Definition 4.1. Let σ be a cell of W . Then the *type* of σ is the $\bar{\Gamma}$ -conjugacy class of σ .

4.2. Orientation

Let ϕ be an oriented cell of W and let $\gamma \in \bar{\Gamma}$. Since $\bar{\Gamma}$ acts by diffeomorphisms, γ takes the orientation of ϕ to some orientation of $\gamma\phi$. Let $\gamma_*\phi$ denote the cell $\gamma\phi$ with this choice of orientation.

Let W_T denote the $\bar{\Gamma}$ -orbit of cells of type T . For each type T , fix a representative cell $\phi_T \in W_T$. Fix orientations on the standard cells ϕ_T . We use these orientations to fix orientations of all the cells of W . In particular, if ϕ is a cell of type T and $\phi = \gamma\phi_T$, then we give ϕ the orientation $\gamma_*\phi$. Note that because the stabilizer of a cell fixes the cell pointwise, this gives a well-defined orientation to each cell of W .

4.3. Incidence function

We now define the incidence function on W . If ψ is a facet of ϕ , then the orientation of ϕ induces an orientation of ψ . This orientation of ψ may or may not agree with the orientation determined above. Let $[\phi, \psi]$ be ± 1 depending on whether this orientation on ϕ does or does not agree with the orientation of ψ . This is precisely the (ϕ, ψ) -component of the cellular boundary operator on W . The incidence function $[\cdot, \cdot]$ is $\bar{\Gamma}$ -invariant in the sense that $[\gamma_*\phi, \gamma_*\psi] = [\phi, \psi]$ for all $\gamma \in \bar{\Gamma}$.

For every subgroup $\Gamma \subset \bar{\Gamma}$, the quotient $\Gamma \backslash W$ has the structure of a regular cell complex with identifications. Since the incidence function defined above is $\bar{\Gamma}$ -invariant, it is in particular Γ -invariant. Thus we can use $[\cdot, \cdot]$ to compute the cellular cohomology of $\Gamma \backslash W$. If ϕ is a cell of W , we will write $\bar{\phi}$ to denote its Γ -conjugacy class, or equivalently the corresponding cell in $\Gamma \backslash W$.

4.4. Cells as orbits

Let X denote the homogeneous space $\Gamma \backslash \bar{\Gamma}$. The following proposition allows us to identify cells of the quotient $\Gamma \backslash W$ with right $\bar{\Gamma}_T$ -orbits in X .

Proposition 4.2. (See [2, Proposition 3.3].) *There is a one-to-one correspondence between*

- (i) Cells in W_T and cosets $\bar{\Gamma} / \bar{\Gamma}_T$.
- (ii) Cells of the quotient $\Gamma \backslash W_T$ and Γ -orbits of W_T .
- (iii) Cells of the quotient $\Gamma \backslash W_T$ and double cosets in $\Gamma \backslash \bar{\Gamma} / \bar{\Gamma}_T$.

Proof. The first statement is a standard fact about stabilizers. Since the stabilizer of a cell fixes the cell pointwise, the cell structure on W descends to give a cell structure on $\Gamma \backslash W$, and the second statement follows. Combining the two statements gives that the cells of $\Gamma \backslash W_T$ are parametrized by $\Gamma \backslash \bar{\Gamma} / \bar{\Gamma}_T$. \square

In light of this proposition, we will now think of a cell of $\Gamma \backslash W$ as a Γ_T -orbits of X or a double coset in $\Gamma \backslash \bar{\Gamma} / \bar{\Gamma}_T$.

The facets of ϕ_T , denoted \mathcal{F}_{ϕ_T} , can be expressed as translates of the various representative cells $\phi_{T'}$. In particular, there exist finite subsets $\mathcal{B}(T, T') \subset \bar{\Gamma}$ such that

$$\mathcal{F}_{\phi_T} = \bigcup_{T'} \bigcup_{\alpha \in \mathcal{B}(T, T')} \alpha \phi_{T'}.$$

Note that the α 's are not uniquely determined, but the cosets $\alpha \bar{\Gamma}_{T'} \in \bar{\Gamma} / \bar{\Gamma}_{T'}$ are determined for each pair (T, T') .

The cell $\bar{\phi}_T$ corresponds to the double coset $\Gamma e \bar{\Gamma}_T$, and the face $\overline{\alpha \phi_{T'}}$ corresponds to the double coset $\Gamma \alpha \bar{\Gamma}_{T'}$. In terms of orbits, if O is the $\bar{\Gamma}_T$ -orbit corresponding to $\bar{\phi}_T$, then the orbit corresponding to $\overline{\alpha \phi_{T'}}$ is the $\bar{\Gamma}_{T'}$ -orbit of X that contains $O \cdot \alpha$. By translation, we can understand the boundary faces of an arbitrary cell $\bar{\phi}$ of $\Gamma \backslash W$ in terms of the α defined above.

Proposition 4.3. (See [2, Proposition 3.20].) *Let ϕ be a cell of type T and let O denote the Γ_T -orbit of X corresponding to $\bar{\phi}$. Then the faces of $\bar{\phi}$ of type T' correspond exactly to the $\bar{\Gamma}_{T'}$ -orbits of X containing $O \cdot \alpha$ as α ranges over $\mathcal{B}(T, T')$.*

From Propositions 4.2 and 4.3, it becomes clear that to compute cohomology in this way, we need the cellular structure of W , the stabilizers $\bar{\Gamma}_T$, the elements $\alpha \in \mathcal{B}(T, T')$, and the incidence numbers $[\phi_T, \psi]$. This is carried out for $k = \mathbb{Q}(i)$ and $k = \mathbb{Q}(\sqrt{-3})$ in the following sections.

5. Examples

5.1. $k = \mathbb{Q}(i)$

In [10], we compute the space W for $k = \mathbb{Q}(i)$. The cells of W fall into twenty-four equivalence classes modulo $\bar{\Gamma}$ consisting of two 3-cells, seven 2-cells, nine 1-cells, and six 0-cells. Representatives of the 3-cells and their boundary faces are shown in Fig. 1. We refer to [10] for the cell data. The results of the cohomology computation are tabulated in Section 6.




Fig. 1. The 3-cells for $SU(2, 1; \mathbb{Z}[i])$.

5.2. $k = \mathbb{Q}(\sqrt{-3})$

In this section we apply the previous results in the case $k = \mathbb{Q}(\sqrt{-3})$. Let $\zeta = (1 + \sqrt{-3})/2$. Let

$$w = \begin{pmatrix} 0 & 0 & -1 \\ 0 & 1 & 0 \\ 1 & 0 & 0 \end{pmatrix}, \quad \tau = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}, \tag{7}$$

$$\gamma_1 = \begin{pmatrix} \zeta^5 & 0 & \zeta^2 \\ -1 & \zeta^2 & 1 \\ \zeta^4 & \sqrt{-3}\zeta & 1 \end{pmatrix}, \quad \gamma_2 = \begin{pmatrix} -1 & \sqrt{-3} & \zeta^5 \\ 0 & 1 & -1 \\ -1 & \sqrt{-3} & \zeta^4 \end{pmatrix}, \tag{8}$$

$$\epsilon = \begin{pmatrix} \zeta & 0 & 0 \\ 0 & \zeta^{-2} & 0 \\ 0 & 0 & \zeta \end{pmatrix}, \quad \text{and} \quad \sigma = \begin{pmatrix} 1 & \sqrt{-3} & \zeta \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{pmatrix}. \tag{9}$$

A weak lower bound for the exhaustion functions is found by examining the values of various exhaustion functions on a set $\Omega \subset \mathcal{D}$ whose $\bar{\Gamma}$ translates cover \mathcal{D} . Then Section 3.2 is then used to compute the possible admissible sets. Each was either rejected or verified using MAPLE. The results are given below.

Proposition 5.1. *Every cell of \mathcal{D}_0 is Γ -conjugate to exactly one of the following:*

- (i) $\mathcal{I}^2 = \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix} \right\}$.
- (ii) $\mathcal{I}_1^3 = \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix} \right\}$.
- (iii) $\mathcal{I}_2^3 = \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} \zeta \\ 1 \\ 1 \end{pmatrix} \right\}$.
- (iv) $\mathcal{I}^4 = \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} \zeta \\ 1 \\ 1 \end{pmatrix} \right\}$.
- (v) $\mathcal{I}^8 = \left\{ \begin{pmatrix} 1 \\ 0 \\ 0 \end{pmatrix}, \begin{pmatrix} 0 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} 1 \\ 0 \\ 1 \end{pmatrix}, \begin{pmatrix} \zeta \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} \zeta^2 \\ 1 \\ 1 \end{pmatrix}, \begin{pmatrix} \zeta \\ \zeta^5 \\ 1 \end{pmatrix}, \begin{pmatrix} \sqrt{-3}\zeta^5 \\ \zeta^5 \\ 1 \end{pmatrix}, \begin{pmatrix} \zeta^2 \\ \zeta \\ 1 \end{pmatrix} \right\}$.

Table 1
Incidence types.

	\mathcal{I}^2	\mathcal{I}_1^3	\mathcal{I}_2^3	\mathcal{I}^4	\mathcal{I}^8
\mathcal{I}_1^2	*	3	3	6	28
\mathcal{I}_1^3	2	*	*	1	8
\mathcal{I}_2^3	12	*	*	3	48
\mathcal{I}^4	24	6	3	*	16
\mathcal{I}^8	12	6	3	2	*

The incidence table is given in Table 1, where the entry below the diagonal means that each column cell has that many row cells in its boundary, and the entry above the diagonal means the column cell appears in the boundary of this many row cells. The entries below the diagonal can be read off from Fig. 2. The entries above the diagonal can be easily computed from Proposition 5.1, since the Γ -conjugacy class of a strongly admissible can be distinguished by the pairwise \mathcal{Q} -inner products of its associated isotropic vectors, except to distinguish \mathcal{I}_1^3 and \mathcal{I}_2^3 we must also compute the dimension of the span of their isotropic vectors.

Since the set of isotropic vectors associated with each piece of \mathcal{D}_0 is known, computing the stabilizers is a finite computation. Let \mathcal{I} be a strongly admissible set and let \mathcal{J} be the associated set of isotropic vectors. If $\gamma \in \text{Stab}(\mathcal{D}(\mathcal{I}))$, then

$$\gamma \cdot \mathcal{D}(\mathcal{I}) = \mathcal{D}({}^\gamma \mathcal{I}) = \mathcal{D}(\mathcal{I}).$$

It follows that γ must take each vector of \mathcal{J} to another vector of \mathcal{J} up to scaling by \mathcal{O}^* .

Proposition 5.2. *The stabilizer of \mathcal{I}^2 is isomorphic to $\mathbb{Z}/12\mathbb{Z}$ and is generated by ϵw . The stabilizer of \mathcal{I}_1^3 is isomorphic to $\mathbb{Z}/3\mathbb{Z} \oplus \mathbb{Z}/6\mathbb{Z}$ and is generated by τw and ϵ . The stabilizer of \mathcal{I}_2^3 is isomorphic to $\mathbb{Z}/3\mathbb{Z}$ and is generated by ϵ^2 . The stabilizer of \mathcal{I}^4 is isomorphic to $\mathbb{Z}/3\mathbb{Z}$ and is generated by ϵ^2 . The stabilizer \mathcal{I}^8 is the order 24 group, Magma small group number 11 and is generated by γ_1 and γ_2 .*

Since the action of $\bar{\Gamma}$ on \mathcal{D}_0 is understood, we can subdivide $\mathcal{D}(\mathcal{I}^2)$ and its boundary into cells in a $\bar{\Gamma}$ -equivariant way as shown in Fig. 2 to yield a cell complex W with the property that the stabilizer of each cell fixes the cell pointwise. The cells of W fall into sixteen equivalence classes modulo $\bar{\Gamma}$ consisting of one 3-cells, seven 2-cells, six 1-cells, and three 0-cells. Representatives are chosen and stabilizers are recomputed. The results are given in Table 2.

6. Numerical results

A computer was used to compute the $\bar{\Gamma}_T$ -orbits for the various congruence subgroups Γ and to create the coboundary matrix. In this process, large sparse matrices were produced, up to $97,542 \times 89,478$ for $\Gamma_0(-2 + 13\sqrt{-3}) \subset \text{SU}(2, 1; \mathbb{Z}[\zeta])$. Then MAGMA




Fig. 2. The 3-cell $\mathcal{D}(\mathcal{I}^2)$ and subdivision for $SU(2, 1; \mathbb{Z}[\zeta])$.

Table 2
Stabilizers of representative cells.

Cell	Dimension	Stabilizer	Generators
X	3	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
A	2	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
B	2	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
C	2	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
E	2	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
F	2	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
G	2	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
a	1	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
b	1	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
c	1	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
d	1	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
e	1	$\mathbb{Z}/3\mathbb{Z}$	ϵ^2
f	1	$\mathbb{Z}/6\mathbb{Z}$	ϵ
m	0	$\mathcal{G}(24, 11)^a$	γ_1, γ_2
n	0	$\mathbb{Z}/12\mathbb{Z}$	ϵw
o	0	$\mathbb{Z}/3\mathbb{Z} \oplus \mathbb{Z}/6\mathbb{Z}$	$\epsilon^2, \tau \epsilon w$

^a This is the order 24 with Magma small group library number 11.

[4] was used to compute the elementary divisors. The computations were only carried out in a few cases for $\Gamma_1(N)$ and $\Gamma(N)$ because the index grows so rapidly. In fact, Holzapfel [8] has an explicit formula for the index $[\Gamma(N) : \bar{\Gamma}]$ for $N \in \mathbb{Z}$, which shows that the index grows like N^8 . Tables 3–5 summarize the numerical results for $k = \mathbb{Q}(i)$ and Table 6 summarizes the results for $k = \mathbb{Q}(\sqrt{-3})$. We remark that although the cohomology in degree 1 is predominately 0 in the range that was computed, by a result of Blasius and Rogawski [3], we expect infinitely many congruence subgroups to yield non-trivial cohomology. Their result, however, is for principal congruence subgroups, not those of the form $\Gamma_0(N)$, and we already see some non-trivial cohomology in H^1 for the former.

Table 3
The integral cohomology for $\Gamma_0(N) \subset \text{SU}(2, 1; \mathbb{Z}[i])$.

N	H^1	H^2	H^3
2	0	\mathbb{Z}	\mathbb{Z}^2
3	0	\mathbb{Z}^2	$\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$
4	0	$\mathbb{Z}^2 \oplus \mathbb{Z}/2\mathbb{Z}$	\mathbb{Z}^6
5	0	\mathbb{Z}^7	\mathbb{Z}^5
6	0	$\mathbb{Z}^4 \oplus \mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}^5 \oplus \mathbb{Z}/2\mathbb{Z}$
7	0	$\mathbb{Z}^{10} \oplus \mathbb{Z}/4\mathbb{Z}$	$\mathbb{Z} \oplus \mathbb{Z}/12\mathbb{Z}$
8	0	$\mathbb{Z}^{11} \oplus (\mathbb{Z}/2\mathbb{Z})^3 \oplus \mathbb{Z}/4\mathbb{Z}$	\mathbb{Z}^{12}
9	0	$\mathbb{Z}^{25} \oplus (\mathbb{Z}/3\mathbb{Z})^2$	$\mathbb{Z}^5 \oplus (\mathbb{Z}/2\mathbb{Z})^2$
10	0	\mathbb{Z}^{34}	\mathbb{Z}^{17}
11	0	$\mathbb{Z}^{39} \oplus \mathbb{Z}/15\mathbb{Z}$	$\mathbb{Z} \oplus \mathbb{Z}/30\mathbb{Z}$
12	0	$\mathbb{Z}^{36} \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/8\mathbb{Z}$	$\mathbb{Z}^{13} \oplus \mathbb{Z}/13\mathbb{Z}$
13	0	\mathbb{Z}^{79}	$\mathbb{Z}^5 \oplus (\mathbb{Z}/3\mathbb{Z})^2$
14	0	$\mathbb{Z}^{58} \oplus \mathbb{Z}/12\mathbb{Z}$	$\mathbb{Z}^5 \oplus \mathbb{Z}/12\mathbb{Z}$
15	0	$\mathbb{Z}^{148} \oplus \mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}^{11} \oplus (\mathbb{Z}/2\mathbb{Z})^3$
16	0	$\mathbb{Z}^{94} \oplus (\mathbb{Z}/2\mathbb{Z})^3 \oplus (\mathbb{Z}/4\mathbb{Z})^4 \oplus (\mathbb{Z}/8\mathbb{Z})^2$	$\mathbb{Z}^{23} \oplus \mathbb{Z}/2\mathbb{Z}$
17	0	$\mathbb{Z}^{166} \oplus (\mathbb{Z}/2\mathbb{Z})^2$	$\mathbb{Z}^5 \oplus (\mathbb{Z}/2\mathbb{Z})^2 \oplus (\mathbb{Z}/4\mathbb{Z})^2$
18	0	$\mathbb{Z}^{142} \oplus \mathbb{Z}/3\mathbb{Z} \oplus \mathbb{Z}/6\mathbb{Z}$	$\mathbb{Z}^{17} \oplus (\mathbb{Z}/2\mathbb{Z})^3$
19	0	$\mathbb{Z}^{211} \oplus \mathbb{Z}/15\mathbb{Z}$	$\mathbb{Z} \oplus \mathbb{Z}/90\mathbb{Z}$
20	0	$\mathbb{Z}^{238} \oplus (\mathbb{Z}/2\mathbb{Z})^5$	$\mathbb{Z}^{41} \oplus (\mathbb{Z}/2\mathbb{Z})^2$
21	0	$\mathbb{Z}^{294} \oplus (\mathbb{Z}/2\mathbb{Z})^2 \oplus \mathbb{Z}/16\mathbb{Z}$	$\mathbb{Z}^3 \oplus (\mathbb{Z}/2\mathbb{Z})^2 \oplus \mathbb{Z}/48\mathbb{Z}$
22	0	$\mathbb{Z}^{238} \oplus \mathbb{Z}/30\mathbb{Z}$	$\mathbb{Z}^5 \oplus \mathbb{Z}/30\mathbb{Z}$
23	0	$\mathbb{Z}^{372} \oplus \mathbb{Z}/132\mathbb{Z}$	$\mathbb{Z} \oplus \mathbb{Z}/132\mathbb{Z}$
24	0	$\mathbb{Z}^{312} \oplus (\mathbb{Z}/2\mathbb{Z})^5 \oplus \mathbb{Z}/4\mathbb{Z} \oplus \mathbb{Z}/8\mathbb{Z}$	$\mathbb{Z}^{25} \oplus (\mathbb{Z}/2\mathbb{Z})^2 \oplus \mathbb{Z}/8\mathbb{Z}$
$1+i$	0	\mathbb{Z}	\mathbb{Z}
$2+i$	0	\mathbb{Z}^3	\mathbb{Z}^2
$3+2i$	0	\mathbb{Z}^7	$\mathbb{Z}^2 \oplus \mathbb{Z}/3\mathbb{Z}$
$4+i$	0	$\mathbb{Z}^9 \oplus \mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}$
$5+2i$	0	$\mathbb{Z}^{21} \oplus \mathbb{Z}/7\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/7\mathbb{Z}$
$6+i$	0	$\mathbb{Z}^{37} \oplus \mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/9\mathbb{Z}$
$5+4i$	0	$\mathbb{Z}^{45} \oplus \mathbb{Z}/5\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/10\mathbb{Z}$
$7+2i$	0	$\mathbb{Z}^{75} \oplus \mathbb{Z}/13\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/13\mathbb{Z}$
$6+5i$	0	$\mathbb{Z}^{103} \oplus \mathbb{Z}/5\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/15\mathbb{Z}$
$8+3i$	0	$\mathbb{Z}^{151} \oplus \mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/18\mathbb{Z}$
$8+5i$	0	$\mathbb{Z}^{225} \oplus \mathbb{Z}/11\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/22\mathbb{Z}$
$9+4i$	0	$\mathbb{Z}^{271} \oplus \mathbb{Z}/4\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/24\mathbb{Z}$
$10+i$	0	$\mathbb{Z}^{291} \oplus \mathbb{Z}/25\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/25\mathbb{Z}$
$10+3i$	0	$\mathbb{Z}^{343} \oplus \mathbb{Z}/9\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/27\mathbb{Z}$
$8+7i$	0	$\mathbb{Z}^{369} \oplus \mathbb{Z}/14\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/2\mathbb{Z} \oplus \mathbb{Z}/28\mathbb{Z}$

Table 4
The integral cohomology of $\Gamma_1(N) \subset \text{SU}(2, 1; \mathbb{Z}[i])$.

N	H^1	H^2	H^3
2	0	$\mathbb{Z} \oplus \mathbb{Z}/2\mathbb{Z}$	\mathbb{Z}^2
3	0	$\mathbb{Z}^{13} \oplus (\mathbb{Z}/2\mathbb{Z})^2 \oplus \mathbb{Z}/4\mathbb{Z}$	\mathbb{Z}^3
4	\mathbb{Z}^2	$\mathbb{Z}^{24} \oplus (\mathbb{Z}/2\mathbb{Z})^3 \oplus (\mathbb{Z}/4\mathbb{Z})^2 \oplus \mathbb{Z}/8\mathbb{Z}$	\mathbb{Z}^{11}
5	0	$\mathbb{Z}^{115} \oplus (\mathbb{Z}/2\mathbb{Z})^5 \oplus (\mathbb{Z}/10\mathbb{Z})^2 \oplus \mathbb{Z}/20\mathbb{Z}$	\mathbb{Z}^{23}
6	0	$\mathbb{Z}^{102} \oplus (\mathbb{Z}/2\mathbb{Z})^5 \oplus \mathbb{Z}/6\mathbb{Z} \oplus (\mathbb{Z}/12\mathbb{Z})^2$	\mathbb{Z}^{19}
7	0	$\mathbb{Z}^{538} \oplus (\mathbb{Z}/2\mathbb{Z})^2 \oplus (\mathbb{Z}/14\mathbb{Z})^2 \oplus \mathbb{Z}/28\mathbb{Z}$	\mathbb{Z}^{23}
8	\mathbb{Z}^6	$\mathbb{Z}^{460} \oplus (\mathbb{Z}/2\mathbb{Z})^5 \oplus (\mathbb{Z}/4\mathbb{Z})^4 \oplus \mathbb{Z}/8\mathbb{Z} \oplus \mathbb{Z}/16\mathbb{Z}$	$\mathbb{Z}^{71} \oplus \mathbb{Z}/2\mathbb{Z}$
$2+i$	0	$\mathbb{Z}^{10} \oplus \mathbb{Z}/2\mathbb{Z}$	\mathbb{Z}^3
$3+2i$	0	$\mathbb{Z}^{91} \oplus \mathbb{Z}/2\mathbb{Z}$	\mathbb{Z}^{11}
$4+i$	0	$\mathbb{Z}^{184} \oplus (\mathbb{Z}/2\mathbb{Z})^2$	$\mathbb{Z}^{15} \oplus \mathbb{Z}/2\mathbb{Z}$

Table 5

The integral cohomology for $\Gamma(N) \subset \text{SU}(2, 1; \mathbb{Z}[i])$.

N	H^1	H^2	H^3
$1 + i$	0	\mathbb{Z}	\mathbb{Z}^2
2	0	$\mathbb{Z}^2 \oplus (\mathbb{Z}/2\mathbb{Z})^2$	\mathbb{Z}^5
3	0	$\mathbb{Z}^{243} \oplus (\mathbb{Z}/2\mathbb{Z})^7 \oplus (\mathbb{Z}/6\mathbb{Z})^7 \oplus \mathbb{Z}/12\mathbb{Z}$	\mathbb{Z}^{55}
4	\mathbb{Z}^6	$\mathbb{Z}^{484} \oplus (\mathbb{Z}/2\mathbb{Z})^7 \oplus (\mathbb{Z}/4\mathbb{Z})^{12} \oplus (\mathbb{Z}/8\mathbb{Z})^2$	$\mathbb{Z}^{95} \oplus \mathbb{Z}/2\mathbb{Z}$

Table 6

The integral cohomology for $\Gamma_0(N) \subset \text{SU}(2, 1; \mathbb{Z}[\zeta])$.

N	H^1	H^2	H^3
5	0	$\mathbb{Z}^8 \oplus \mathbb{Z}/6\mathbb{Z}$	$\mathbb{Z} \oplus \mathbb{Z}/4\mathbb{Z}$
11	0	$\mathbb{Z}^{56} \oplus (\mathbb{Z}/2\mathbb{Z})^2 \oplus \mathbb{Z}/60\mathbb{Z}$	$\mathbb{Z} \oplus \mathbb{Z}/20\mathbb{Z}$
17	0	$\mathbb{Z}^{208} \oplus \mathbb{Z}/3\mathbb{Z} \oplus \mathbb{Z}/24\mathbb{Z}$	$\mathbb{Z} \oplus \mathbb{Z}/48\mathbb{Z}$
23	0	$\mathbb{Z}^{508} \oplus (\mathbb{Z}/2\mathbb{Z})^2 \oplus \mathbb{Z}/264\mathbb{Z}$	$\mathbb{Z} \oplus \mathbb{Z}/88\mathbb{Z}$
$2 + \sqrt{-3}$	0	\mathbb{Z}^7	\mathbb{Z}^2
$3 + \sqrt{-3}$	0	\mathbb{Z}^{15}	$\mathbb{Z}^2 \oplus \mathbb{Z}/2\mathbb{Z}$
$-2 + 5\sqrt{-3}$	0	$\mathbb{Z}^{21} \oplus \mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/3\mathbb{Z}$
$5 + \sqrt{-3}$	0	$\mathbb{Z}^{47} \oplus \mathbb{Z}/5\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/5\mathbb{Z}$
$-3 + 7\sqrt{-3}$	0	$\mathbb{Z}^{67} \oplus \mathbb{Z}/3\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/6\mathbb{Z}$
$6 + \sqrt{-3}$	0	$\mathbb{Z}^{85} \oplus \mathbb{Z}/7\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/7\mathbb{Z}$
$-4 + 9\sqrt{-3}$	0	$\mathbb{Z}^{167} \oplus \mathbb{Z}/5\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/10\mathbb{Z}$
$-2 + 9\sqrt{-3}$	0	$\mathbb{Z}^{197} \oplus \mathbb{Z}/11\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/11\mathbb{Z}$
$8 + \sqrt{-3}$	0	$\mathbb{Z}^{235} \oplus \mathbb{Z}/6\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/12\mathbb{Z}$
$3 + 7\sqrt{-3}$	0	$\mathbb{Z}^{271} \oplus \mathbb{Z}/13\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/13\mathbb{Z}$
$-3 + 11\sqrt{-3}$	0	$\mathbb{Z}^{408} \oplus \mathbb{Z}/8\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/16\mathbb{Z}$
$2 + 9\sqrt{-3}$	0	$\mathbb{Z}^{405} \oplus \mathbb{Z}/17\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/17\mathbb{Z}$
$7 + 5\sqrt{-3}$	0	$\mathbb{Z}^{511} \oplus \mathbb{Z}/9\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/18\mathbb{Z}$
$-6 + 13\sqrt{-3}$	0	$\mathbb{Z}^{687} \oplus \mathbb{Z}/21\mathbb{Z}$	$\mathbb{Z}^2 \oplus \mathbb{Z}/21\mathbb{Z}$

Acknowledgments

I would like to thank Paul Gunnells for many helpful comments and patiently explaining his paper to me. I also thank T.N. Venkataramana, who helped with understanding [3].

References

- [1] A. Ash, Small-dimensional classifying spaces for arithmetic subgroups of general linear groups, *Duke Math. J.* 51 (2) (1984) 459–468.
- [2] A. Ash, P.E. Gunnells, M. McConnell, Cohomology of congruence subgroups of $\text{SL}_4(\mathbb{Z})$, *J. Number Theory* 94 (1) (2002) 181–212.
- [3] D. Blasius, J. Rogawski, Cohomology of congruence subgroups of $\text{SU}(2, 1)^p$ and Hodge cycles on some special complex hyperbolic surfaces, in: *Regulators in Analysis, Geometry and Number Theory*, in: *Progr. Math.*, vol. 171, Birkhäuser Boston, Boston, MA, 2000, pp. 1–15.
- [4] W. Bosma, J. Cannon, C. Playoust, The Magma algebra system. I. The user language, in: *Computational Algebra and Number Theory*, London, 1993, *J. Symbolic Comput.* 24 (3–4) (1997) 235–265.
- [5] L. Dembélé, Examples of automorphic forms on the unitary group $U(3)$, preprint, 2007.
- [6] E. Falbel, J.R. Parker, The geometry of the Eisenstein–Picard modular group, *Duke Math. J.* 131 (2) (2006) 249–289.
- [7] J. Franke, Harmonic analysis in weighted L_2 -spaces, *Ann. Sci. Ecole Norm. Sup.* (4) 31 (2) (1998) 181–279.

- [8] R.-P. Holzapfel, Zeta dimension formula for Picard modular cusp forms of neat natural congruence subgroups, *Abh. Math. Semin. Univ. Hambg.* 68 (1998) 169–192.
- [9] D. Yasaki, On the existence of spines for \mathbb{Q} -rank 1 groups, *Selecta Math. (N.S.)* 12 (3–4) (2006) 541–564.
- [10] D. Yasaki, An explicit spine for the Picard modular group over the Gaussian integers, *J. Number Theory* 128 (1) (2008) 207–234.
- [11] T. Zink, Über die Anzahl der Spitzen einiger arithmetischer Untergruppen unitärer Gruppen, *Math. Nachr.* 89 (1979) 315–320.