



ELSEVIER

Contents lists available at ScienceDirect

Journal of Number Theory

www.elsevier.com/locate/jnt



On Hecke L -functions attached to half-integral weight modular forms

YoungJu Choie^{a,*}, Winfried Kohnen^b

^a Department of Mathematics, Pohang Institute of Science and Technology, POSTECH, Pohang 790-784, Republic of Korea

^b Mathematisches Institut der Universität, INF 205, D-69120 Heidelberg, Germany

ARTICLE INFO

Article history:

Received 2 April 2017

Accepted 1 November 2017

Available online xxxx

Communicated by A. Folsom

MSC:

primary 11F37

secondary 11F66

Keywords:

Hecke eigenform

Half integral weight

L -function

ABSTRACT

We investigate non-vanishing properties of $L(f, s)$ on the real line, when f is a Hecke eigenform of half-integral weight $k + \frac{1}{2}$ on $\Gamma_0(4)$.

© 2018 Elsevier Inc. All rights reserved.

1. Introduction

If f is a cusp form of half-integral weight $k + \frac{1}{2}$ on $\Gamma_0(4)$ with Fourier coefficients $a(n)$ ($n \geq 1$), one can as usual attach to f the Hecke L -series

$$L(f, s) = \sum_{n \geq 1} a(n)n^{-s} \quad (\sigma := \Re(s) \gg 1). \quad (1)$$

* Corresponding author.

E-mail addresses: yjc@postech.ac.kr (Y. Choie), winfried@mathi.uni-heidelberg.de (W. Kohnen).

<https://doi.org/10.1016/j.jnt.2017.11.016>

0022-314X/© 2018 Elsevier Inc. All rights reserved.

It is known [9] that $L(f, s)$ has holomorphic continuation to \mathbf{C} and satisfies a functional equation under $s \mapsto k + \frac{1}{2} - s$, relating $L(f, s)$ and $L(f|W_4, s)$ where W_4 is the Fricke involution.

We note that even if f is a Hecke eigenform, in general $L(f, s)$ has no Euler product. Though $L(f, s)$ has turned out to be useful in the investigation of sign changes of the coefficients $\Re(a(n))$ resp. $\Im(a(n))$ [4], in general otherwise its meaning remains a bit mysterious.

In this paper we will investigate non-vanishing properties of $L(f, s)$ on the real line, when f is a Hecke eigenform (Thm., sect. 3). The proofs which will be given in sect. 4 use the Mellin integral representation and rely on the existence of some special half-integral weight cusp forms with non-vanishing properties on the positive imaginary axis.

As will be clear from the proof, similar (in fact, somewhat easier) arguments also work in the case of cuspidal Hecke eigenforms of integral weight on $\Gamma_1 := SL_2(\mathbf{Z})$. We leave it to the reader to carry out the details and formulate the corresponding statements.

We would like to recall that in the case of Hecke eigenforms on Γ_1 non-vanishing results for their Hecke L -functions at an *arbitrary* point s_0 in the critical strip (not on the critical line) have been proved in [5] (cf. also [7]), using holomorphic kernel functions. This method was carried over to the case of half-integral weight in [8], for arbitrary level. However, in this approach for given s_0 the weight in general has to be large depending on s_0 .

We also would like to point out that in [10] L -functions associated to modular forms of half-integral weight were studied. In particular, certain half-integral weight cusp forms were investigated and the location of their zeros off the critical line were calculated.

Finally, addressing the reader interested in Siegel modular forms, we would like to mention the papers [3, 6] where corresponding non-vanishing results for Koecher–Maass series are given.

Notations. For $z \in \mathbf{C}^*$ we let $z^{\frac{1}{2}} := e^{\frac{1}{2} \log z}$, where $\log z$ is the principal branch of the complex logarithm. For z in the complex upper half-plane \mathcal{H} we put $q = e^{2\pi iz}$.

We denote by $\Gamma_0(4)$ the subgroup of Γ_1 consisting of matrices $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ with $4|c$.

The letter k always denotes an integer. We write $M_k(4)$ for the space of modular forms of weight k for $\Gamma_0(4)$, with trivial character if k is even and non-trivial character ($-\frac{4}{k}$) if k is odd. We let $M_{k+\frac{1}{2}}(4)$ be the space of modular forms of weight $k + \frac{1}{2}$ for $\Gamma_0(4)$ and write $S_{k+\frac{1}{2}}(4)$ for the subspace of cusp forms.

2. Basic facts on modular forms of half-integral weight of level 4

For basic facts on modular forms of half-integral weight we refer to [9] and in the special case of level 4 also to [1, 2].

The group $\Gamma_0(4)$ has three cusps, represented by 0, $\frac{1}{2}$ and $i\infty$. The cusp $\frac{1}{2}$ is $(k + \frac{1}{2})$ -irregular, so a modular form of weight $k + \frac{1}{2}$ on $\Gamma_0(4)$ automatically vanishes at $\frac{1}{2}$.

As is well-known, one has

$$\dim S_{k+\frac{1}{2}}(4) = \sup \{0, [\frac{k}{2}] - 1\}. \quad (2)$$

Recall that on $M_{k+\frac{1}{2}}(4)$ (resp. $S_{k+\frac{1}{2}}(4)$) the Fricke involution W_4 acts by

$$f(z) \mapsto (f|W_4)(z) := (-2iz)^{-k-1/2} f(-\frac{1}{4z}) \quad (z \in \mathcal{H}).$$

The Hecke operators $T(p^2)$ ($p > 2$ prime) commute with W_4 , hence the spaces

$$S_{k+\frac{1}{2}}^{(\pm)}(4) := \{f \in S_{k+\frac{1}{2}}(4) \mid f|W_4 = (\pm)f\}$$

have a basis of Hecke eigenforms of all the $T(p^2)$.

For $f \in S_{k+\frac{1}{2}}(4)$ we put

$$L^*(f, s) := (2\pi)^{-s} \cdot 2^s \cdot \Gamma(s) \cdot L(f, s) \quad (\sigma \gg 1)$$

where $L(f, s)$ is the L -series defined by (1). Then $L^*(f, s)$ has holomorphic continuation to \mathbf{C} and satisfies the functional equation

$$L^*(f, k + \frac{1}{2} - s) = L^*(f|W_4, s). \quad (3)$$

In particular, if $f|W_4 = -f$, then $L^*(f, \frac{k}{2} + \frac{1}{4}) = 0$.

In the rest of this section, we are concerned with certain special modular forms of level 4 which will play a role in sect. 4.

We let

$$\theta(z) = \sum_{n \in \mathbf{Z}} q^{n^2} \quad (z \in \mathcal{H})$$

be the basic theta function which is in $M_{\frac{1}{2}}(4)$. One has

$$\theta|W_4 = \theta. \quad (4)$$

Let

$$P(z) = 1 - 24 \sum_{n \geq 1} \sigma_1(n) q^n \quad (z \in \mathcal{H})$$

be the quasi-Eisenstein series of weight 2 on Γ_1 (where $\sigma_1(n) = \sum_{d|n} d$) and put

$$F_2(z) := \frac{1}{24} (-P(z) + 3P(2z) - 2P(4z)).$$

Then

$$F_2(z) = \sum_{n \geq 1, n \equiv 1 \pmod{2}} \sigma_1(n) q^n$$

and $F_2 \in M_2(4)$. The transformation formula

$$(2z)^{-2} F_2\left(-\frac{1}{4z}\right) = F_2(z) - \frac{1}{16} \theta^4(z) \quad (5)$$

holds.

Further, put

$$\Delta_4(z) := F_2(z) (\theta^4(z) - 16F_2(z)) \quad (z \in \mathcal{H}).$$

Then $\Delta_4 \in M_4(4)$ and one has

$$(2z)^{-4} \Delta_4\left(-\frac{1}{4z}\right) = \Delta_4(z) \quad (6)$$

as follows from (4) and (5).

We also note that Δ_4 vanishes at the cusps 0 and $i\infty$, hence it follows that $\theta\Delta_4$ is a cusp form of weight $\frac{9}{2}$ on $\Gamma_0(4)$. Using (2) we find that the map

$$f \mapsto f \cdot \theta\Delta_4$$

gives an isomorphism of $M_k(4)$ onto $S_{k+\frac{9}{2}}(4)$.

Finally, we define

$$D_2(z) := \theta^4(z) - 32F_2(z) \quad (z \in \mathcal{H}).$$

Then D_2 is in $M_2(4)$ and using (4) and (5) one checks that

$$(2z)^{-2} D_2\left(-\frac{1}{4z}\right) = D_2(z). \quad (7)$$

3. Statement of results

It easily follows from the discussions in sect. 2 that $S_{k+\frac{1}{2}}^{(+)}(4) = \{0\}$ for $k < 4$ (in fact, $S_{k+\frac{1}{2}}(4) = \{0\}$ in this case) and $S_{k+\frac{1}{2}}^{(-)}(4) = \{0\}$ for $k < 6$. We shall prove

Theorem. *Let $\sigma \in \mathbf{R}$ be fixed. Then the following assertions hold:*

- i) *If $k \geq 4$, then there is a Hecke eigenform $f \in S_{k+\frac{1}{2}}^{(+)}(4)$ with $L^*(f, \sigma) \neq 0$.*
- ii) *If $k \geq 6$ and $\sigma \neq \frac{k}{2} + \frac{1}{4}$, then there is a Hecke eigenform $f \in S_{k+\frac{1}{2}}^{(-)}(4)$ with $L^*(f, \sigma) \neq 0$.*

Remarks.

- i) In the proof of the theorem we explicitly construct a cusp form f in the space $S_{k+\frac{1}{2}}^{(+)}(4)$ (resp. in the space $S_{k+\frac{1}{2}}^{(-)}(4)$) such that $L^*(f, \sigma) \neq 0$, for any real σ .
- ii) Let $k = 4$. It follows from sect. 2 that $S_{9/2}^{(+)}(4) = \mathbf{C}\theta\Delta_4$, hence $f_1 := \theta\Delta_4$ is a Hecke eigenform and $L^*(f_1, \sigma) \neq 0$ for all $\sigma \in \mathbf{R}$. It will in fact follow from the proof given in the next sect. that $L^*(f_1, \sigma) > 0$ for all $\sigma \in \mathbf{R}$. Similarly, in the case $k = 6$ one checks that $S_{13/2}^{(-)}(4) = \mathbf{C}\theta D_2\Delta_4$, so $f_2 := \theta D_2\Delta_4$ is a Hecke eigenform and $L^*(f_2, \sigma) \neq 0$ for all $\sigma \in \mathbf{R}$, $\sigma \neq \frac{13}{4}$. It will follow from our proof that $L^*(f_2, \sigma)$ is positive for $\sigma > \frac{13}{4}$ and negative for $\sigma < \frac{13}{4}$.

4. Proofs

We start with the proof of i) whose first part is standard. If $f \in S_{k+\frac{1}{2}}(4)$, then in the usual way by Mellin transform one has

$$L^*(f, s) = \int_0^\infty f(it)(2t)^s \frac{dt}{t} \quad (\sigma \gg 1).$$

Splitting up the integral into the sum of the integral from 0 to $\frac{1}{2}$ and the integral from $\frac{1}{2}$ to ∞ , and substituting $t \mapsto \frac{1}{4t}$ in the former one, we obtain

$$L^*(f, s) = \int_{\frac{1}{2}}^\infty f\left(\frac{i}{4t}\right)\left(\frac{1}{2t}\right)^s \frac{dt}{t} + \int_{\frac{1}{2}}^\infty f(it)(2t)^s \frac{dt}{t}.$$

If $f|W_4 = f$, then $f\left(\frac{i}{4t}\right) = (2t)^{k+1/2}f(it)$, hence we get the identity

$$L^*(f, s) = \int_{\frac{1}{2}}^\infty f(it) \left((2t)^s + (2t)^{k+1/2-s} \right) \frac{dt}{t} \quad (8)$$

which is valid for all $s \in \mathbf{C}$, since f is a cusp form.

We now put $f := \Delta_4 \theta^{2k-7}$. By (4) and (6) we see that $f \in S_{k+\frac{1}{2}}^{(+)}(4)$.

Recall that the valence formula asserts that the sum of the orders (counted with multiplicities) of a non-zero modular form g of weight ℓ on $\Gamma_0(4)$ on the compact modular curve $X_0(4)$ of level 4 is equal to $\frac{\ell}{12}[\Gamma_1 : \Gamma_0(4)]$. We note that $[\Gamma_1 : \Gamma_0(4)] = 6$ and apply the valence formula with $g = \Delta_4 \in M_4(4)$. Since Δ_4 vanishes at the cusps 0 and $i\infty$, it follows that Δ_4 has no zeros on \mathcal{H} , in particular $\Delta_4(it) \neq 0$ for all $t > 0$. Since by definition $\Delta_4(it)$ is real for $t > 0$, by continuity we must have $\Delta_4(it) > 0$ for all t or $\Delta_4(it) < 0$ for all t . We claim that the first alternative holds. Indeed, evaluating (5) at $z = \frac{i}{2}$ we find that

$$\theta^4\left(\frac{i}{2}\right) = 32F_2\left(\frac{i}{2}\right),$$

hence

$$\Delta_4\left(\frac{i}{2}\right) = F_2\left(\frac{i}{2}\right)\left(\theta^4\left(\frac{i}{2}\right) - 16F_2\left(\frac{i}{2}\right)\right) > 0.$$

(Alternatively, we could also have used the convergent q -product expansion

$$\Delta_4(z) = q \prod_{n \geq 1, n \equiv 0, \pm 1 \pmod{4}} (1 - q^n)^8,$$

cf. [2, p. 25].)

By definition, $\theta(z)$ is real and positive on the positive imaginary axis. Thus $f(it) > 0$ for $t > 0$ and so in particular the integrand in (8) is positive for all $t \geq \frac{1}{2}$ if $s = \sigma$ is real, hence $L^*(f, \sigma)$ is positive and so non-zero for all σ . Writing f in terms of a Hecke basis of $S_{k+\frac{1}{2}}^{(+)}(4)$ we deduce the assertion of i).

We now prove ii). If $f|W_4 = -f$, then similarly as in i) we find that

$$L^*(f, s) = \int_{\frac{1}{2}}^{\infty} f(it) \left((2t)^s - (2t)^{k+1/2-s} \right) \frac{dt}{t}.$$

We let $f := \Delta_4 D_2 \theta^{2k-11}$. By (4), (6) and (7) then $f \in S_{k+\frac{1}{2}}^{(-)}(4)$.

We want to show that $L^*(f, \sigma) \neq 0$ for σ real, $\sigma \neq \frac{k}{2} + \frac{1}{4}$. By the functional equation (3) we may assume that $\sigma > \frac{k}{2} + \frac{1}{4}$. In this range clearly

$$(2t)^\sigma - (2t)^{k+1/2-\sigma} \geq 0$$

whenever $t \geq \frac{1}{2}$, and the inequality is strict for $t > \frac{1}{2}$.

As mentioned above, $\theta(it) > 0$ and we proved that $\Delta_4(it) > 0$, for $t > 0$. Thus to prove our claim, it will suffice to show that $D_2(it) > 0$ for $t > \frac{1}{2}$.

We apply the valence formula with $g = D_2 \in M_2(4)$. Observe that $D_2(z)$ by (7) vanishes at $z = \frac{i}{2}$. We conclude that any zero of D_2 in \mathcal{H} must be equivalent under $\Gamma_0(4)$ to $\frac{i}{2}$. Thus to see that $D_2(it) \neq 0$ for $t > \frac{1}{2}$ we have to show that for $t > \frac{1}{2}$ the point it cannot be equivalent to $\frac{i}{2}$ under $\Gamma_0(4)$.

Assume on the contrary that there exists $M \in \Gamma_0(4)$ with $M \circ \frac{i}{2} = it$ and $t > \frac{1}{2}$. Let \mathcal{F} be the standard fundamental domain for Γ_1 consisting of $z \in \mathcal{H}$ with $|z| \geq 1$ and $|x| \leq \frac{1}{2}$. Let $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$. Then $S^2 = -E$. Suppose first that $t \leq 1$. Then $S \circ it \in \mathcal{F}$. Also $S \circ \frac{i}{2} = 2i \in \text{int}\mathcal{F}$ and $SMS \circ 2i = S \circ it$. It follows that $SMS = \pm E$ and so $2i = \frac{i}{t}$, i.e. $t = \frac{1}{2}$, a contradiction.

Now suppose that $t > 1$. Then $M \circ \frac{i}{2} \in \mathcal{F}$. Since $M \circ \frac{i}{2} = MS \circ (S \circ \frac{i}{2}) = MS \circ 2i$ and $2i \in \text{int}\mathcal{F}$, it follows that $MS = \pm E$, i.e. $M = \pm S$, a contradiction since S is not in $\Gamma_0(4)$.

Since D_2 has the value 1 at infinity, by continuity we finally find that indeed $D_2(it) > 0$ for all $t > \frac{1}{2}$.

It follows that $L^*(f, \sigma) \neq 0$ as claimed. Writing f in terms of a Hecke basis as before, we derive the assertion of ii).

Acknowledgment

The first author was partially supported by NRF 2017R1A2B2001807.

References

- [1] H. Cohen, Sums involving the values at negative integers of L -functions of quadratic characters, *Math. Ann.* 217 (3) (1975) 271–285.
- [2] H. Cohen, *Formes modulaires à une et deux variables*, Thèse, Univ. Bordeaux I, 1976.
- [3] S. Das, W. Kohnen, Non-vanishing of Koecher–Maass series attached to cusp forms, *Adv. Math.* 281 (2015) 624–669.
- [4] M. Knopp, W. Kohnen, W. Pribitkin, On the signs of Fourier coefficients of cusp forms, *Ramanujan J.* 7 (1–3) (2003) 269–277.
- [5] W. Kohnen, Nonvanishing of Hecke L -functions associated to cusp forms inside the critical strip, *J. Number Theory* 67 (2) (1977) 182–189.
- [6] W. Kohnen, Non-vanishing of Koecher–Maass series attached to Siegel cusp forms on the real line, *Abh. Math. Semin. Univ. Hambg.* 87 (1) (2017) 39–41.
- [7] A. Raghuram, Non-vanishing of L -functions of cusp forms inside the critical strip, in: *Number Theory*, in: Ramanujan Math. Soc. Lect. Notes Ser., vol. 1, Ramanujan Math. Soc., Mysore, 2005, pp. 97–105.
- [8] B. Ramakrishnan, Karam Deo Shankhadhar, Non-vanishing of L -functions associated to cusp forms of half-integral weight, in: *Automorphic Forms*, in: Springer Proc. Math. Stat., vol. 115, 2014, pp. 223–231.
- [9] G. Shimura, On modular forms of half-integral weight, *Ann. of Math.* 97 (1973) 440–481.
- [10] H. Yoshida, On calculations of zeros of various L -functions, *J. Math. Kyoto Univ.* 35 (4) (1995) 663–696.