



ELSEVIER

Contents lists available at ScienceDirect

Journal of Number Theory

www.elsevier.com/locate/jnt



General Section

Asymptotics for the Fourier coefficients of eta-quotients

Shane Chern

Department of Mathematics, The Pennsylvania State University, University Park,
PA 16802, USA

ARTICLE INFO

Article history:

Received 2 June 2018

Received in revised form 24

September 2018

Accepted 27 November 2018

Available online 17 December 2018

Communicated by F. Pellarin

MSC:

11P55

11P82

Keywords:

Eta-quotients

Asymptotics

Circle method

ABSTRACT

We study the asymptotics for the Fourier coefficients of a broad class of eta-quotients,

$$\prod_{r=1}^R \left(\prod_{k \geq 1} (1 - q^{m_r k}) \right)^{\delta_r},$$

where m_1, \dots, m_R are R distinct positive integers and $\delta_1, \dots, \delta_R$ are R non-zero integers with $\sum_{r=1}^R \delta_r \geq 0$.

© 2018 Elsevier Inc. All rights reserved.

1. Introduction

1.1. Background

Since the pioneering work of Hardy and Ramanujan [12], and Rademacher [20,21], there has been an intensive study of the asymptotic behaviors of the Fourier coefficients of eta-quotients. Here what Hardy and Ramanujan as well as Rademacher treated has important combinatorial background:

E-mail address: shanechern@psu.edu.

A partition of a positive integer n is a non-increasing sequence of positive integers whose sum equals n . Let $p(n)$ count the number of partitions of n with the convention that $p(0) = 1$. It is well known that

$$\sum_{n \geq 0} p(n)q^n = \frac{1}{(q; q)_{\infty}}. \quad (1.1)$$

Here and in the sequel, we adopt the standard q -series notation:

$$(a; q)_{\infty} := \prod_{k \geq 0} (1 - aq^k).$$

It is necessary to mention that the generating function (1.1) is holomorphic on the open unit disk $\mathbb{D} \subset \mathbb{C}$, namely, $|q| < 1$.

Rademacher showed that

$$p(n) = \frac{1}{2\sqrt{2}\pi} \sum_{k \geq 1} A_k(n) \sqrt{k} \frac{d}{dn} \left(\frac{2}{\sqrt{n - \frac{1}{24}}} \sinh \left(\frac{\pi}{k} \sqrt{\frac{2}{3} \left(n - \frac{1}{24} \right)} \right) \right), \quad (1.2)$$

where

$$A_k(n) = \sum_{\substack{0 \leq h < k \\ \gcd(h, k) = 1}} e^{\pi i(s(h, k) - 2nh/k)}$$

with $s(h, k)$ being the Dedekind sum defined in (1.7).

Rademacher's approach is straightforward in essence, however delicate in detail. The basic idea is merely Cauchy's integral formula, but we need various techniques including Ford circles, Farey sequences, modular symmetry and the Dedekind eta-function.

One may naturally consider the following general family of holomorphic functions on the open unit disk \mathbb{D} :

$$G(q) = G(e^{2\pi i\tau}) := \prod_{r=1}^R (q^{m_r}; q^{m_r})_{\infty}^{\delta_r}, \quad (1.3)$$

where $\mathbf{m} = (m_1, \dots, m_R)$ is a sequence of R distinct positive integers and $\boldsymbol{\delta} = (\delta_1, \dots, \delta_R)$ is a sequence of R non-zero integers. This can be treated as an eta-quotient as the Dedekind eta-function is defined by (with $q = e^{2\pi i\tau}$)

$$\eta(\tau) := q^{\frac{1}{24}} (q; q)_{\infty}.$$

If we write

$$G(q) = \sum_{n \geq 0} g(n)q^n, \quad (1.4)$$

then $g(n)$ may be related to certain combinatorial quantity in the theory of partitions. We notice that Rademacher's approach for the asymptotic formula of $p(n)$ can be easily adapted to study the asymptotics of $g(n)$ provided that $\sum_{r=1}^R \delta_r < 0$. For some particular $G(q)$, the interested readers may refer to the work of Grosswald [8], Hagis Jr. [9–11], Iseki [14,15] and many others. On the other hand, when $\sum_{r=1}^R \delta_r \geq 0$, a variance of Rademacher's method provided by O-Y. Chan [6] works. Chan used this method to prove several conjectures of Andrews and Lewis [3] concerning rank/crank differences modulo 3 and 4. We refer the readers to [7,18,19] for other related papers.

For general $G(q)$ with $\sum_{r=1}^R \delta_r < 0$, the recent work of Sussman [23] presented a Rademacher-type formula. Sussman's result can in some sense be treated as a special case of the work of Bringmann and Ono [5], in which the coefficients of harmonic Maass forms are studied. On the other hand, Sills [22] provided an automatic algorithm when $\sum_{r=1}^R \delta_r = 0$. When $\sum_{r=1}^R \delta_r = 1$, B. Kim [17] studied a subclass of such $G(q)$ with the aid of Chan's method.

The purpose of this paper is to obtain asymptotic formulas of Fourier coefficients of a broad class of eta-quotients $G(q)$ with $\sum_{r=1}^R \delta_r \geq 0$.

1.2. The main result and some remarks

Assuming that k and h are positive integers with $\gcd(h, k) = 1$, we put

$$\begin{aligned} \Delta_1 &= -\frac{1}{2} \sum_{r=1}^R \delta_r, & \Delta_2 &= \sum_{r=1}^R m_r \delta_r, \\ \Delta_3(k) &= -\sum_{r=1}^R \frac{\delta_r \gcd^2(m_r, k)}{m_r}, & \Delta_4(k) &= \prod_{r=1}^R \left(\frac{m_r}{\gcd(m_r, k)} \right)^{-\frac{\delta_r}{2}}, \end{aligned} \quad (1.5)$$

and

$$\omega_{h,k} = \exp \left(-\pi i \sum_{r=1}^R \delta_r \cdot s \left(\frac{m_r h}{\gcd(m_r, k)}, \frac{k}{\gcd(m_r, k)} \right) \right), \quad (1.6)$$

where $s(d, c)$ is the Dedekind sum defined by

$$s(d, c) := \sum_{n \bmod c} \left(\left(\frac{dn}{c} \right) \right) \left(\left(\frac{n}{c} \right) \right) \quad (1.7)$$

with

$$((x)) := \begin{cases} x - [x] - 1/2 & \text{if } x \notin \mathbb{Z}, \\ 0 & \text{if } x \in \mathbb{Z}. \end{cases}$$

Let $L = \text{lcm}(m_1, \dots, m_R)$. We divide the set $\{1, 2, \dots, L\}$ into two disjoint subsets:

$$\begin{aligned}\mathcal{L}_{>0} &:= \{1 \leq \ell \leq L : \Delta_3(\ell) > 0\}, \\ \mathcal{L}_{\leq 0} &:= \{1 \leq \ell \leq L : \Delta_3(\ell) \leq 0\}.\end{aligned}$$

Now we are ready to present the main result.

Theorem 1.1. *If $\Delta_1 \leq 0$ and the inequality*

$$\min_{1 \leq r \leq R} \left(\frac{\gcd^2(m_r, \ell)}{m_r} \right) \geq \frac{\Delta_3(\ell)}{24} \quad (1.8)$$

holds for all $1 \leq \ell \leq L$, then for positive integers $n > -\Delta_2/24$, we have

$$\begin{aligned}g(n) = E(n) + \sum_{\ell \in \mathcal{L}_{>0}} 2\pi \Delta_4(\ell) \left(\frac{24n + \Delta_2}{\Delta_3(\ell)} \right)^{-\frac{\Delta_1+1}{2}} \\ \times \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \frac{I_{-\Delta_1-1} \left(\frac{\pi}{6k} \sqrt{\Delta_3(\ell)(24n + \Delta_2)} \right)}{k} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} \omega_{h,k} e^{-\frac{2\pi i n h}{k}},\end{aligned} \quad (1.9)$$

where

$$N = \left\lfloor \sqrt{2\pi \left(n + \frac{\Delta_2}{24} \right)} \right\rfloor, \quad (1.10)$$

$$|E(n)| \ll_{\mathbf{m}, \delta} \begin{cases} 1 & \text{if } \Delta_1 = 0, \\ \left(n + \frac{\Delta_2}{24} \right)^{1/4} & \text{if } \Delta_1 = -\frac{1}{2}, \\ \left(n + \frac{\Delta_2}{24} \right)^{1/2} \log \left(n + \frac{\Delta_2}{24} \right) & \text{if } \Delta_1 = -1, \\ \left(n + \frac{\Delta_2}{24} \right)^{-\Delta_1-1/2} & \text{if } \Delta_1 \leq -\frac{3}{2}, \end{cases} \quad (1.11)$$

and $I_s(x)$ is the modified Bessel function of the first kind.

We have several remarks to make.

1. The main term of $g(n)$ may vanish for certain families of eta-quotients. One example arises from Gauß square exponent theorem [2, Corollary 2.10]:

$$\frac{(q; q)_\infty^2}{(q^2; q^2)_\infty} = \sum_{k=-\infty}^{\infty} (-1)^k q^{k^2}, \quad (1.12)$$

from which we see that the Fourier coefficients of the left-hand side are in $\{0, 1, \pm 2\}$. In this case, we have $L = 2$, and hence we can compute

$$\Delta_3(1) = -\frac{3}{2} \quad \text{and} \quad \Delta_3(2) = 0.$$

This implies that the set $\mathcal{L}_{>0}$ is empty. Consequently, the main term vanishes for all n .

2. Recalling that the asymptotic expansion of $I_s(x)$ (cf. [1, p. 377, (9.7.1)]) states that, for fixed s , when $|\arg x| < \frac{\pi}{2}$,

$$I_s(x) \sim \frac{e^x}{\sqrt{2\pi x}} \left(1 - \frac{4s^2 - 1}{8x} + \frac{(4s^2 - 1)(4s^2 - 9)}{2!(8x)^2} - \dots \right), \quad (1.13)$$

one may say more about the main term. Let $\ell \in \mathcal{L}_{>0}$ be fixed. One thing we observe is that for given k , the sum

$$\sum_{\substack{0 \leq h < k \\ \gcd(h, k) = 1}} \omega_{h, k} e^{-\frac{2\pi i n h}{k}}$$

has period k for $n \in \mathbb{Z}_{>0}$. On the other hand, letting us fix some $k_0 \equiv \ell \pmod{L}$, the following estimate of the tail gives useful information:

$$\begin{aligned} \text{Tail} &:= \sum_{\substack{k_0 < k \leq N \\ k \equiv_L \ell}} \frac{I_{-\Delta_1-1} \left(\frac{\pi}{6k} \sqrt{\Delta_3(\ell)(24n + \Delta_2)} \right)}{k} \sum_{\substack{0 \leq h < k \\ \gcd(h, k) = 1}} \omega_{h, k} e^{-\frac{2\pi i n h}{k}} \\ &\ll \sum_{\substack{k_0 < k \leq N \\ k \equiv_L \ell}} I_{-\Delta_1-1} \left(\frac{\pi}{6k} \sqrt{\Delta_3(\ell)(24n + \Delta_2)} \right) \\ &\ll \sum_{\substack{k_0 < k \leq N \\ k \equiv_L \ell}} \frac{e^{\frac{\pi}{6k} \sqrt{\Delta_3(\ell)(24n + \Delta_2)}}}{k^{-\frac{1}{2}} (24n + \Delta_2)^{\frac{1}{4}}} \\ &\ll N^{\frac{3}{2}} (24n + \Delta_2)^{-\frac{1}{4}} e^{\frac{\pi}{6(k_0+L)} \sqrt{\Delta_3(\ell)(24n + \Delta_2)}} \\ &\ll (24n + \Delta_2)^{\frac{1}{2}} e^{\frac{\pi}{6(k_0+L)} \sqrt{\Delta_3(\ell)(24n + \Delta_2)}} \\ &= o \left(e^{\frac{\pi}{6k_0} \sqrt{\Delta_3(\ell)(24n + \Delta_2)}} \right). \end{aligned} \quad (1.14)$$

3. An explicit bound of $E(n)$ is given in (3.10). It would be necessary if one wants to study the exact sign pattern of $g(n)$.
4. It is helpful to record Sussman's result [23]:

Theorem (Sussman). *If $\Delta_1 > 0$ and the inequality*

$$\min_{1 \leq r \leq R} \left(\frac{\gcd^2(m_r, \ell)}{m_r} \right) \geq \frac{\Delta_3(\ell)}{24}$$

holds for all $1 \leq \ell \leq L$, then for positive integers $n > -\Delta_2/24$, we have

$$\begin{aligned} g(n) = & \sum_{\ell \in \mathcal{L}_{>0}} 2\pi \Delta_4(\ell) \left(\frac{24n + \Delta_2}{\Delta_3(\ell)} \right)^{-\frac{\Delta_1+1}{2}} \\ & \times \sum_{\substack{k \geq 1 \\ k \equiv_L \ell}} \frac{I_{\Delta_1+1} \left(\frac{\pi}{6k} \sqrt{\Delta_3(\ell)(24n + \Delta_2)} \right)}{k} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} \omega_{h,k} e^{-\frac{2\pi i n h}{k}}. \end{aligned} \quad (1.15)$$

One may notice that Sussman's asymptotic formula is almost the same as our main term. His proof is mainly based on Rademacher's original version [20] of circle method, whereas ours relies on O-Y. Chan's version [6]. One major difference arises from the contours chosen for Cauchy's integral formula.

1.3. Notation

Let \mathbb{Z} , \mathbb{R} and \mathbb{C} be the set of integers, real numbers and complex numbers, respectively. Let \mathbb{H} be the upper half complex plane. For a given set \mathcal{S} , we use $\mathcal{S}_{\text{condition}}$ to denote the subset of \mathcal{S} with the given condition (e.g. $\mathbb{Z}_{>0}$ denotes the set of positive integers).

The big- O notation is defined in the usual way: $f(x) = O(g(x))$ means that $|f(x)| \leq Cg(x)$ where C is an absolute constant. If the constant C depends on some variables, then we write $f(x) = O_{\text{variables}}(g(x))$. We may also omit the subscript if these variables are explicit. Furthermore, $f(x) \ll g(x)$ means that $f(x) = O(g(x))$. On the other hand, if $\lim f(x)/g(x) = 0$, then we say $f(x) = o(g(x))$. If $\lim f(x)/g(x) = 1$, then we write $f(x) \sim g(x)$.

We use $u \equiv_m v$ to denote $u \equiv v \pmod{m}$, and likewise $u \not\equiv_m v$ to denote $u \not\equiv v \pmod{m}$.

For $\Delta \in \frac{1}{2}\mathbb{Z}_{\leq 0}$ and $x \in \mathbb{R}_{\geq 1}$, we define

$$\Xi_{\Delta}(x) := \begin{cases} 1 & \text{if } \Delta = 0, \\ 2x^{1/2} & \text{if } \Delta = -\frac{1}{2}, \\ x(\log x + 1) & \text{if } \Delta = -1, \\ \zeta(-\Delta)x^{-2\Delta-1} & \text{otherwise,} \end{cases} \quad (1.16)$$

where $\zeta(\cdot)$ is the Riemann zeta-function.

2. The circle method

Our main ingredient is O-Y. Chan's version of circle method [6] with some slight modifications. For the sake of completeness, we shall reproduce the necessary details.

2.1. Cauchy's integral formula

Given a holomorphic function

$$G(q) = \sum_{n \geq 0} g(n) q^n$$

on a simply connected domain containing the origin, we know from Cauchy's integral formula that

$$g(n) = \frac{1}{2\pi i} \oint_{|q|=r} \frac{G(q)}{q^{n+1}} dq,$$

where the contour integral is taken counter-clockwise.

We take $r = e^{-2\pi\varrho}$ with $\varrho = 1/N^2$ where N is a positive integer to be determined later.

Let h/k be a Farey fraction of order N . We use $\xi_{h,k}$ to denote the interval $[-\theta'_{h,k}, \theta''_{h,k}]$ with $-\theta'_{h,k}$ and $\theta''_{h,k}$ being the positive distances from h/k to its neighboring mediants. Now we dissect the circle $|q| = r$ by Farey arcs and obtain

$$g(n) = \sum_{1 \leq k \leq N} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} e^{-\frac{2\pi i n h}{k}} \int_{\xi_{h,k}} G(e^{2\pi i(h/k + i\varrho + \phi)}) e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi.$$

Letting $z = k(\varrho - i\phi)$ and $\tau = (h + iz)/k$, we may rewrite the integral as

$$g(n) = \sum_{1 \leq k \leq N} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} e^{-\frac{2\pi i n h}{k}} \int_{\xi_{h,k}} G(e^{2\pi i \tau}) e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi. \quad (2.1)$$

2.2. Transformation formulas

Let

$$F(q) = F(e^{2\pi i \tau}) := \frac{1}{(q; q)_\infty}$$

with $\tau \in \mathbb{H}$. We also define the fractional linear transformation

$$\gamma(\tau) := \frac{a\tau + b}{c\tau + d}, \quad \text{with } \gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in SL_2(\mathbb{Z}).$$

We know from the transformation formula of Dedekind eta-function [4, pp. 52–61] that

$$F(e^{2\pi i\tau}) = \exp\left(2\pi i\left(\frac{\tau - \gamma(\tau)}{24} - \frac{s(d, c)}{2} + \frac{a + d}{24c}\right)\right) \sqrt{-i(c\tau + d)} F(e^{2\pi i\gamma(\tau)}), \quad (2.2)$$

where the square root is taken on the principal branch, with $\sqrt{x} > 0$ for real $x > 0$.

Given a positive integer m , to transform $F(e^{2\pi im\tau})$ properly for each choice of k and h (with $\gcd(h, k) = 1$), we proceed as follows.

Let $d = \gcd(m, k)$. We also write $m = dm'$ and $k = dk'$. Choosing an integer $\tilde{h}_{m'}$ such that $\tilde{h}_{m'}m'h \equiv -1 \pmod{k'}$ (this is possible since $\gcd(h, k) = 1$) and putting $b_{m'} = (\tilde{h}_{m'}m'h + 1)/k'$, we obtain the following matrix in $SL_2(\mathbb{Z})$:

$$\gamma_{(m, k)} = \begin{pmatrix} \tilde{h}_{m'} & -b_{m'} \\ k' & -m'h \end{pmatrix}. \quad (2.3)$$

Note that our $\gamma_{(m, k)}$ is different to Chan's definition. However, they are identical if $\gcd(m, k) = 1$.

Recalling that

$$\tau = \frac{h + iz}{k} = \frac{h + iz}{dk'},$$

we have

$$\begin{aligned} \gamma_{(m, k)}(m\tau) &= \frac{\tilde{h}_{m'} \cdot m \frac{h+iz}{dk'} - b_{m'}}{k' \cdot m \frac{h+iz}{dk'} - m'h} = \frac{\tilde{h}_{m'}m'h + \tilde{h}_{m'}(im'z) - (\tilde{h}_{m'}m'h + 1)}{m'hk' + k'(im'z) - m'hk'} \\ &= \frac{\tilde{h}_{m'}}{k'} + \frac{1}{m'k'z}i. \end{aligned}$$

This implies that

$$\gamma_{(m, k)}(m\tau) = \frac{\tilde{h}_{m'} \gcd(m, k)}{k} + \frac{\gcd^2(m, k)z^{-1}}{mk}i. \quad (2.4)$$

On the other hand, since $\gamma_{(m, k)} \in SL_2(\mathbb{Z})$, it follows from (2.2) that

$$\begin{aligned} F(e^{2\pi im\tau}) &= e^{\frac{\pi}{12k}\left(\frac{\gcd^2(m, k)}{mz} - mz\right)} \exp\left(\pi i \cdot s\left(\frac{mh}{\gcd(m, k)}, \frac{k}{\gcd(m, k)}\right)\right) \\ &\quad \times \sqrt{\frac{mz}{\gcd(m, k)}} F(e^{2\pi i\gamma_{(m, k)}(m\tau)}). \end{aligned} \quad (2.5)$$

Let $G(q)$ be defined as in (1.3). Then

$$G(e^{2\pi i\tau}) = \prod_{r=1}^R F(e^{2\pi i m_r \tau})^{-\delta_r}. \quad (2.6)$$

The following result is an immediate consequence of (2.5).

Lemma 2.1. *For any positive integers k and h with $\gcd(h, k) = 1$, let $\tau = (h + iz)/k$ be defined as in §2.1. We have,*

$$G(e^{2\pi i\tau}) = e^{\frac{\pi}{12k} \left(\frac{\Delta_3(k)}{z} + \Delta_2 z \right)} z^{\Delta_1} \omega_{h,k} \Delta_4(k) \prod_{r=1}^R F(e^{2\pi i \gamma_{(m_r, k)}(m_r \tau)})^{-\delta_r}. \quad (2.7)$$

2.3. Some bounds

We now present several necessary bounds, most of which are given by Chan.

A standard result on Farey fractions states that (cf. [13, Chap. 3])

$$\frac{1}{2kN} \leq \theta'_{h,k}, \theta''_{h,k} \leq \frac{1}{kN}. \quad (2.8)$$

Consequently,

$$\frac{1}{kN} \leq |\xi_{h,k}| \leq \frac{2}{kN}. \quad (2.9)$$

We next notice that $z = k(\varrho - i\phi)$. Hence

$$\Re(z) = k\varrho = \frac{k}{N^2}. \quad (2.10)$$

This implies that

$$|z| \geq \frac{k}{N^2}. \quad (2.11)$$

On the other hand, we have

$$\Re\left(\frac{1}{z}\right) \geq \frac{k}{2}. \quad (2.12)$$

This is because

$$\Re\left(\frac{1}{z}\right) = \frac{1}{k} \frac{\varrho}{\varrho^2 + \phi^2} \geq \frac{1}{k} \frac{N^{-2}}{N^{-4} + k^{-2}N^{-2}} = \frac{k}{k^2N^{-2} + 1} \geq \frac{k}{1+1} = \frac{k}{2},$$

where we use the fact $k \leq N$ in the last inequality.

In the author's recent work with Tang and Wang [7], it is shown that

Lemma 2.2. Let $\tau = (h + iz)/k$ be defined as in §2.1. We have

$$\left| F(e^{2\pi i \gamma_{(m,k)}(m\tau)}) \right| \leq \exp \left(\frac{e^{-\pi \gcd^2(m,k)/m}}{(1 - e^{-\pi \gcd^2(m,k)/m})^2} \right), \quad (2.13)$$

$$\left| \frac{1}{F(e^{2\pi i \gamma_{(m,k)}(m\tau)})} \right| \leq \exp \left(\frac{e^{-\pi \gcd^2(m,k)/m}}{(1 - e^{-\pi \gcd^2(m,k)/m})^2} \right). \quad (2.14)$$

Proof. We write in this proof $q = e^{2\pi i \varpi}$ with $\varpi \in \mathbb{H}$. Recall that $p(n)$ denotes the number of partitions of n with the convention that $p(0) = 1$. It is clear that $p(n) > 0$ for all $n \geq 0$.

We first have

$$|F(q)| \leq \sum_{n \geq 0} p(n) |q|^n = F(|q|).$$

On the other hand,

$$\left| \frac{1}{F(q)} \right| = \prod_{k \geq 1} |1 - q^k| \leq \prod_{k \geq 1} (1 + |q|^k) \leq \prod_{k \geq 1} \frac{1}{1 - |q|^k} = F(|q|).$$

For real $0 \leq x < 1$, we have

$$\begin{aligned} \log F(x) &= - \sum_{k \geq 1} \log(1 - x^k) = \sum_{k \geq 1} \sum_{m \geq 1} \frac{x^{km}}{m} = \sum_{n \geq 1} x^n \sum_{d|n} \frac{1}{d} \\ &\leq \sum_{n \geq 1} n x^n = \frac{x}{(1 - x)^2}. \end{aligned}$$

Noting that $|q| = |e^{2\pi i \varpi}| = e^{-2\pi \Im(\varpi)}$, we have

$$F(|q|) \leq \exp \left(\frac{e^{-2\pi \Im(\varpi)}}{(1 - e^{-2\pi \Im(\varpi)})^2} \right). \quad (2.15)$$

We remark that the right-hand side can be treated as a decreasing function of $\Im(\varpi)$.

Finally, we see from (2.4) that

$$\Im(\gamma_{(m,k)}(m\tau)) = \frac{\gcd^2(m,k)}{mk} \Re\left(\frac{1}{z}\right) \geq \frac{\gcd^2(m,k)}{mk} \frac{k}{2} = \frac{\gcd^2(m,k)}{2m},$$

where we use (2.12). Hence the left-hand sides of (2.13) and (2.14) are both bounded from above by

$$\exp \left(\frac{e^{-2\pi\Im \left(e^{2\pi i \gamma(m,k)(m\tau)} \right)}}{\left(1 - e^{-2\pi\Im \left(e^{2\pi i \gamma(m,k)(m\tau)} \right)} \right)^2} \right) \leq \exp \left(\frac{e^{-\pi \gcd^2(m,k)/m}}{\left(1 - e^{-\pi \gcd^2(m,k)/m} \right)^2} \right). \quad \square$$

We also extend a bound given in [7, Lemma 3.3].

Lemma 2.3. *Let η_1, \dots, η_R be R integers, all non-zero, and let $\varpi_1, \dots, \varpi_R$ be R complex numbers in \mathbb{H} . Then*

$$\left| \prod_{r=1}^R F(e^{2\pi i \varpi_r})^{\eta_r} - 1 \right| \leq \exp \left(\sum_{r=1}^R \frac{|\eta_r| e^{-2\pi\Im(\varpi_r)}}{(1 - e^{-2\pi\Im(\varpi_r)})^2} \right) - 1. \quad (2.16)$$

Proof. Let $q_r = e^{2\pi i \varpi_r}$ for $r = 1, \dots, R$. Note that $F(q_r)^{\eta_r}$ can be treated as the generating function of $|\eta_r|$ -colored partitions when η_r is a positive integer, while when η_r is a negative integer, it is the generating function of weighted $|\eta_r|$ -colored distinct partitions where the weight is given by $(-1)^\sharp$ with \sharp counting the total number of parts.

Hence, for $r = 1, \dots, R$, if we write

$$F(e^{2\pi i \varpi_r})^{\eta_r} = \sum_{n \geq 0} a_r(n) q_r^n \quad \text{and} \quad F(e^{2\pi i \varpi_r})^{|\eta_r|} = \sum_{n \geq 0} \tilde{a}_r(n) q_r^n,$$

then $|a_r(n)| \leq \tilde{a}_r(n)$ for $n \geq 0$ with $a_r(0) = \tilde{a}_r(0) = 1$.

Hence

$$\begin{aligned} \left| \prod_{r=1}^R F(e^{2\pi i \varpi_r})^{\eta_r} - 1 \right| &= \left| \sum_{n_1 \geq 0} \cdots \sum_{n_R \geq 0} a_1(n_1) \cdots a_R(n_R) q_1^{n_1} \cdots q_R^{n_R} - 1 \right| \\ &= \left| \sum_{\substack{n_1 \geq 0 \\ \dots \\ n_R \geq 0 \\ (n_1, \dots, n_R) \neq (0, \dots, 0)}} a_1(n_1) \cdots a_R(n_R) q_1^{n_1} \cdots q_R^{n_R} \right| \\ &\leq \sum_{\substack{n_1 \geq 0 \\ \dots \\ n_R \geq 0 \\ (n_1, \dots, n_R) \neq (0, \dots, 0)}} \tilde{a}_1(n_1) \cdots \tilde{a}_R(n_R) |q_1|^{n_1} \cdots |q_R|^{n_R} \\ &= \sum_{n_1 \geq 0} \cdots \sum_{n_R \geq 0} \tilde{a}_1(n_1) \cdots \tilde{a}_R(n_R) |q_1|^{n_1} \cdots |q_R|^{n_R} - 1 \\ &= \prod_{r=1}^R F(|q_r|)^{|\eta_r|} - 1 \leq \exp \left(\sum_{r=1}^R \frac{|\eta_r| e^{-2\pi\Im(\varpi_r)}}{(1 - e^{-2\pi\Im(\varpi_r)})^2} \right) - 1, \end{aligned}$$

where we use (2.15) in the last inequality. \square

2.4. An integral

The last task in this section is to evaluate a useful integral.

Lemma 2.4. *Let $a \in \mathbb{R}_{>0}$, $b \in \mathbb{R}$ and $c \in \frac{1}{2}\mathbb{Z}_{\leq 0}$. Let $\gcd(h, k) = 1$. Define*

$$I := \int_{\xi_{h,k}} e^{\frac{\pi}{12k}(\frac{a}{z} + bz)} z^c e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi. \quad (2.17)$$

Then, for those positive integers n with $24n + b > 0$, we have

$$I = \frac{2\pi}{k} \left(\frac{24n + b}{a} \right)^{-\frac{c+1}{2}} I_{-c-1} \left(\frac{\pi}{6k} \sqrt{a(24n + b)} \right) + E(I), \quad (2.18)$$

where

$$|E(I)| \leq \frac{2^{-c} \pi^{-1} e^{\frac{\pi a}{3}} N^{-c}}{n + \frac{b}{24}} e^{2\pi \varrho(n + \frac{b}{24})}. \quad (2.19)$$

Proof. We first put $w = z/k = \varrho - i\phi$ and reverse the integral order to obtain

$$I = \frac{1}{2\pi i} \int_{\varrho - i\theta''_{h,k}}^{\varrho + i\theta'_{h,k}} 2\pi e^{\frac{\pi a}{12k^2 w}} e^{2\pi w(n + \frac{b}{24})} (kw)^c dw.$$

One may separate the integral into three parts

$$\begin{aligned} I &= \frac{1}{2\pi i} \left(\int_{\Gamma} - \int_{-\infty - i\theta''_{h,k}}^{\varrho - i\theta''_{h,k}} + \int_{-\infty + i\theta'_{h,k}}^{\varrho + i\theta'_{h,k}} \right) 2\pi e^{\frac{\pi a}{12k^2 w}} e^{2\pi w(n + \frac{b}{24})} (kw)^c dw \\ &=: J_1 - J_2 + J_3, \end{aligned}$$

where

$$\Gamma := (-\infty - i\theta''_{h,k}) \rightarrow (\varrho - i\theta''_{h,k}) \rightarrow (\varrho + i\theta'_{h,k}) \rightarrow (-\infty + i\theta'_{h,k})$$

is a Hankel contour.

To compute the main term J_1 , we make the following change of variables $t = wk\sqrt{(24n + b)/a}$ to obtain

$$J_1 = \frac{2\pi}{k} \left(\frac{24n + b}{a} \right)^{-\frac{c+1}{2}} \frac{1}{2\pi i} \int_{\tilde{\Gamma}} e^{\frac{\pi}{12k} \sqrt{a(24n+b)}(t + \frac{1}{t})} t^c dt.$$

Note that the new contour $\tilde{\Gamma}$ is still a Hankel contour. Recalling the contour integral representation of $I_s(x)$:

$$I_s(x) = \frac{1}{2\pi i} \int_{\Gamma} t^{-s-1} e^{\frac{x}{2}} e^{(t+\frac{1}{t})} dt \quad (\Gamma \text{ is a Hankel contour}),$$

we conclude

$$J_1 = \frac{2\pi}{k} \left(\frac{24n+b}{a} \right)^{-\frac{c+1}{2}} I_{-c-1} \left(\frac{\pi}{6k} \sqrt{a(24n+b)} \right).$$

For the error term $E(I)$, which comes from J_2 and J_3 , we put $w = x + i\theta$ with $-\infty \leq x \leq \varrho$ and $\theta \in \{\theta'_{h,k}, -\theta''_{h,k}\}$. Following Chan [6, p. 120], we have

$$\left| e^{2\pi w(n+\frac{b}{24})} \right| = e^{2\pi x(n+\frac{b}{24})},$$

and

$$\left| e^{\frac{\pi a}{12k^2 w}} \right| = e^{\frac{\pi a}{12k^2} \Re(\frac{1}{w})} = e^{\frac{\pi a}{12k^2} \frac{x}{x^2+\theta^2}} \leq e^{\frac{\pi a}{12k^2} \frac{x}{\theta^2}} \leq e^{\frac{\pi a}{12k^2} \varrho(2kN)^2} = e^{\frac{\pi a}{3}},$$

$$|(kw)^c| = (|kw|^{-1})^{-c} \leq \left(\frac{1}{k\sqrt{x^2+\theta^2}} \right)^{-c} \leq \left(\frac{1}{k|\theta|} \right)^{-c} \leq (2N)^{-c},$$

where we use the bound $\frac{1}{2kN} \leq |\theta| \leq \frac{1}{kN}$. Hence for $j = 2$ and 3 , we have

$$\begin{aligned} |J_j| &\leq \frac{1}{2\pi} \int_{-\infty}^{\varrho} 2\pi e^{\frac{\pi a}{3}} e^{2\pi x(n+\frac{b}{24})} (2N)^{-c} dx \\ &= \frac{2^{-c-1} \pi^{-1} e^{\frac{\pi a}{3}} N^{-c}}{n + \frac{b}{24}} e^{2\pi \varrho(n+\frac{b}{24})}. \end{aligned}$$

This implies that

$$|E(I)| = |-J_2 + J_3| \leq |J_2| + |J_3| \leq \frac{2^{-c} \pi^{-1} e^{\frac{\pi a}{3}} N^{-c}}{n + \frac{b}{24}} e^{2\pi \varrho(n+\frac{b}{24})}. \quad \square$$

3. Asymptotics

We rewrite (2.1) as follows:

$$g(n) = \sum_{1 \leq k \leq N} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} e^{-\frac{2\pi i n h}{k}} \int_{\xi_{h,k}} G(e^{2\pi i \tau}) e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi$$

$$\begin{aligned}
&= \sum_{\ell=1}^L \sum_{\substack{1 \leq k \leq N \\ k \equiv \ell \pmod L}} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} e^{-\frac{2\pi i n h}{k}} \int_{\xi_{h,k}} G(e^{2\pi i \tau}) e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi \\
&=: \sum_{\ell=1}^L S_{\ell}.
\end{aligned} \tag{3.1}$$

Recalling that $L = \text{lcm}(m_1, \dots, m_R)$, we have, for those k with $k \equiv \ell \pmod L$, the following two identities hold:

$$\Delta_3(k) = \Delta_3(\ell) \quad \text{and} \quad \Delta_4(k) = \Delta_4(\ell),$$

since $\gcd(m_r, k) = \gcd(m_r, \ell)$ for all $r = 1, \dots, R$.

Throughout this section, we always assume that $24n + \Delta_2 > 0$.

3.1. Estimating S_{ℓ} with $\ell \in \mathcal{L}_{\leq 0}$

We first assume $\ell \in \mathcal{L}_{\leq 0}$. In this case, the contribution of S_{ℓ} is relatively small.

Recalling the bounds (2.10)–(2.14) and the assumption $\Delta_1 \leq 0$, it follows from (2.7) that, if $k \equiv \ell \pmod L$, then

$$\begin{aligned}
&|G(e^{2\pi i \tau}) e^{-2\pi i n \phi} e^{2\pi n \varrho}| \\
&= \left| \prod_{r=1}^R F(e^{2\pi i \gamma(m_r, k)}(m_r \tau))^{-\delta_r} \right| \left| e^{\frac{\pi}{12k} \left(\frac{\Delta_3(\ell)}{z} + \Delta_2 z \right)} \right| \Delta_4(\ell) |z|^{\Delta_1} e^{2\pi n \varrho} \\
&= \left| \prod_{r=1}^R F(e^{2\pi i \gamma(m_r, k)}(m_r \tau))^{-\delta_r} \right| e^{\frac{\pi}{12k} (\Delta_3(\ell) \Re(\frac{1}{z}) + \Delta_2 \Re(z))} \Delta_4(\ell) |z|^{\Delta_1} e^{2\pi n \varrho} \\
&\leq \Delta_4(\ell) k^{\Delta_1} N^{-2\Delta_1} e^{2\pi \varrho \left(n + \frac{\Delta_2}{24} \right)} \exp \left(\frac{\pi \Delta_3(\ell)}{24} + \sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right).
\end{aligned}$$

Consequently,

$$\begin{aligned}
|S_{\ell}| &= \left| \sum_{\substack{1 \leq k \leq N \\ k \equiv \ell \pmod L}} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} e^{-\frac{2\pi i n h}{k}} \int_{\xi_{h,k}} G(e^{2\pi i \tau}) e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi \right| \\
&\leq \sum_{\substack{1 \leq k \leq N \\ k \equiv \ell \pmod L}} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} \int_{\xi_{h,k}} \Delta_4(\ell) k^{\Delta_1} N^{-2\Delta_1} e^{2\pi \varrho \left(n + \frac{\Delta_2}{24} \right)} \\
&\quad \times \exp \left(\frac{\pi \Delta_3(\ell)}{24} + \sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) d\phi
\end{aligned}$$

$$\begin{aligned}
&\leq \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \sum_{\substack{0 \leq h < k \\ \gcd(h, k) = 1}} \Delta_4(\ell) k^{\Delta_1} N^{-2\Delta_1} e^{2\pi \varrho \left(n + \frac{\Delta_2}{24}\right)} \\
&\quad \times \exp \left(\frac{\pi \Delta_3(\ell)}{24} + \sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) \frac{2}{kN} \\
&\hspace{15em} \text{(by (2.9))} \\
&\leq \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \Delta_4(\ell) k^{\Delta_1} N^{-2\Delta_1} e^{2\pi \varrho \left(n + \frac{\Delta_2}{24}\right)} \\
&\quad \times \exp \left(\frac{\pi \Delta_3(\ell)}{24} + \sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) \frac{2}{kN} k \\
&= 2\Delta_4(\ell) N^{-2\Delta_1-1} e^{2\pi \varrho \left(n + \frac{\Delta_2}{24}\right)} \left(\sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} k^{\Delta_1} \right) \\
&\quad \times \exp \left(\frac{\pi \Delta_3(\ell)}{24} + \sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) \\
&\leq 2\Delta_4(\ell) e^{2\pi \varrho \left(n + \frac{\Delta_2}{24}\right)} \Xi_{\Delta_1}(N) \exp \left(\frac{\pi \Delta_3(\ell)}{24} + \sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right).
\end{aligned}$$

Here we use the trivial bound

$$N^{-2\Delta_1-1} \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} k^{\Delta_1} \leq \Xi_{\Delta_1}(N),$$

where $\Xi_{\Delta_1}(N)$ is defined in (1.16).

To summarize, we have

Lemma 3.1. *For $\ell \in \mathcal{L}_{\leq 0}$, it holds that*

$$|S_\ell| \leq 2\Delta_4(\ell) e^{2\pi \varrho \left(n + \frac{\Delta_2}{24}\right)} \Xi_{\Delta_1}(N) \exp \left(\frac{\pi \Delta_3(\ell)}{24} + \sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right). \quad (3.2)$$

3.2. Estimating S_ℓ with $\ell \in \mathcal{L}_{>0}$

Assume that $\ell \in \mathcal{L}_{>0}$. Here $g(n)$ is dominated by these S_ℓ .

We deduce from (2.7) that

$$\begin{aligned}
 S_\ell &= \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \sum_{\substack{0 \leq h < k \\ \gcd(h, k) = 1}} e^{-\frac{2\pi i n h}{k}} \int_{\xi_{h, k}} G(e^{2\pi i \tau}) e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi \\
 &= \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \sum_{\substack{0 \leq h < k \\ \gcd(h, k) = 1}} e^{-\frac{2\pi i n h}{k}} \int_{\xi_{h, k}} \prod_{r=1}^R F(e^{2\pi i \gamma_{(m_r, k)}(m_r \tau)})^{-\delta_r} \\
 &\quad \times \omega_{h, k} \Delta_4(\ell) e^{\frac{\pi}{12k} \left(\frac{\Delta_3(\ell)}{z} + \Delta_2 z \right)} z^{\Delta_1} e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi \\
 &= \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \sum_{\substack{0 \leq h < k \\ \gcd(h, k) = 1}} e^{-\frac{2\pi i n h}{k}} \int_{\xi_{h, k}} \omega_{h, k} \Delta_4(\ell) \\
 &\quad \times e^{\frac{\pi}{12k} \left(\frac{\Delta_3(\ell)}{z} + \Delta_2 z \right)} z^{\Delta_1} e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi \\
 &\quad + \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \sum_{\substack{0 \leq h < k \\ \gcd(h, k) = 1}} e^{-\frac{2\pi i n h}{k}} \int_{\xi_{h, k}} \left(\prod_{r=1}^R F(e^{2\pi i \gamma_{(m_r, k)}(m_r \tau)})^{-\delta_r} - 1 \right) \omega_{h, k} \Delta_4(\ell) \\
 &\quad \times e^{\frac{\pi}{12k} \left(\frac{\Delta_3(\ell)}{z} + \Delta_2 z \right)} z^{\Delta_1} e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi \\
 &=: T_{\ell, 1} + T_{\ell, 2}.
 \end{aligned} \tag{3.3}$$

We first deal with $T_{\ell, 2}$, which is relatively small comparing with $T_{\ell, 1}$. The following bound is necessary.

Lemma 3.2. *If $\Delta_3(\ell) > 0$, we have, for $k \equiv \ell \pmod{L}$,*

$$\begin{aligned}
 &\left| \left(\prod_{r=1}^R F(e^{2\pi i \gamma_{(m_r, k)}(m_r \tau)})^{-\delta_r} - 1 \right) e^{\frac{\pi}{12k} \frac{\Delta_3(\ell)}{z}} \right| \\
 &\leq e^{\frac{\pi \Delta_3(\ell)}{24}} \left(\exp \left(\sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) - 1 \right).
 \end{aligned} \tag{3.4}$$

Proof. It follows from Lemma 2.3 that

$$\left| \prod_{r=1}^R F(e^{2\pi i \gamma_{(m_r, k)}(m_r \tau)})^{-\delta_r} - 1 \right| \leq \exp \left(\sum_{r=1}^R \frac{|\delta_r| e^{-2\pi \Im(\gamma_{(m_r, k)}(m_r \tau))}}{(1 - e^{-2\pi \Im(\gamma_{(m_r, k)}(m_r \tau))})^2} \right) - 1.$$

We also know from (2.4) that

$$\Im(\gamma_{(m_r, k)}(m_r \tau)) = \frac{\gcd^2(m_r, \ell)}{m_r k} \Re\left(\frac{1}{z}\right).$$

Hence

$$\begin{aligned}
 & \left| \left(\prod_{r=1}^R F \left(e^{2\pi i \gamma_{(m_r, k)}(m_r \tau)} \right)^{-\delta_r} - 1 \right) e^{\frac{\pi}{12k} \frac{\Delta_3(\ell)}{z}} \right| \\
 &= e^{\frac{\pi \Delta_3(\ell)}{12k} \Re(\frac{1}{z})} \left| \prod_{r=1}^R F \left(e^{2\pi i \gamma_{(m_r, k)}(m_r \tau)} \right)^{-\delta_r} - 1 \right| \\
 &\leq e^{\frac{\pi \Delta_3(\ell)}{12k} \Re(\frac{1}{z})} \left(\exp \left(\sum_{r=1}^R \frac{|\delta_r| e^{\frac{-2\pi \gcd^2(m_r, \ell)}{m_r k} \Re(\frac{1}{z})}}{\left(1 - e^{\frac{-2\pi \gcd^2(m_r, \ell)}{m_r k} \Re(\frac{1}{z})} \right)^2} \right) - 1 \right). \quad (3.5)
 \end{aligned}$$

Note that $\Re(\frac{1}{z}) \geq \frac{k}{2}$. If we put

$$x = e^{-\frac{\pi \Delta_3(\ell)}{12k} \Re(\frac{1}{z})},$$

then $x \in (0, e^{-\pi \Delta_3(\ell)/24}] \subseteq (0, 1)$ since $\Delta_3(\ell) > 0$. Letting

$$u_r = \frac{24 \gcd^2(m_r, \ell)}{\Delta_3(\ell) m_r}$$

and recalling (1.8)

$$\frac{\gcd^2(m_r, \ell)}{m_r} \geq \frac{\Delta_3(\ell)}{24} > 0,$$

we have $u_r \geq 1$ for all $r = 1, \dots, R$.

Now (3.5) becomes

$$\left| \left(\prod_{r=1}^R F \left(e^{2\pi i \gamma_{(m_r, k)}(m_r \tau)} \right)^{-\delta_r} - 1 \right) e^{\frac{\pi}{12k} \frac{\Delta_3(\ell)}{z}} \right| \leq \frac{1}{x} \left(\exp \left(\sum_{r=1}^R \frac{|\delta_r| x^{u_r}}{(1 - x^{u_r})^2} \right) - 1 \right).$$

Let

$$W(x) = \frac{1}{x} \left(\exp \left(\sum_{r=1}^R \frac{|\delta_r| x^{u_r}}{(1 - x^{u_r})^2} \right) - 1 \right).$$

To prove (3.4), it suffices to show that $W(x)$ is a non-decreasing function of x on $(0, e^{-\pi \Delta_3(\ell)/24}]$ since the right-hand side of (3.4) is exactly $W(e^{-\pi \Delta_3(\ell)/24})$. It is equivalent to show that $W'(x) \geq 0$ on this interval. We have

$$W'(x) = -\frac{1}{x^2} \left(\exp \left(\sum_{r=1}^R \frac{|\delta_r| x^{u_r}}{(1-x^{u_r})^2} \right) - 1 \right) \\ + \frac{1}{x} \exp \left(\sum_{r=1}^R \frac{|\delta_r| x^{u_r}}{(1-x^{u_r})^2} \right) \sum_{r=1}^R \left(\frac{2|\delta_r| u_r x^{-1+2u_r}}{(1-x^{u_r})^3} + \frac{|\delta_r| u_r x^{-1+u_r}}{(1-x^{u_r})^2} \right).$$

It suffices to show that

$$\exp \left(-\sum_{r=1}^R \frac{|\delta_r| x^{u_r}}{(1-x^{u_r})^2} \right) + \sum_{r=1}^R \left(\frac{2|\delta_r| u_r x^{2u_r}}{(1-x^{u_r})^3} + \frac{|\delta_r| u_r x^{u_r}}{(1-x^{u_r})^2} \right) \geq 1.$$

We next observe that

$$\sum_{r=1}^R \left(\frac{2|\delta_r| u_r x^{2u_r}}{(1-x^{u_r})^3} + \frac{|\delta_r| u_r x^{u_r}}{(1-x^{u_r})^2} \right) \geq \sum_{r=1}^R \frac{|\delta_r| x^{u_r}}{(1-x^{u_r})^2}.$$

This is valid as for $r = 1, \dots, R$, we have

$$\frac{2|\delta_r| u_r x^{2u_r}}{(1-x^{u_r})^3} + \frac{|\delta_r| u_r x^{u_r}}{(1-x^{u_r})^2} - \frac{|\delta_r| x^{u_r}}{(1-x^{u_r})^2} = \frac{|\delta_r| x^{u_r} (-1 + u_r + x^{u_r} + u_r x^{u_r})}{(1-x^{u_r})^3} \geq 0$$

since $u_r \geq 1$ and $x \in (0, e^{-\pi\Delta_3(\ell)/24}]$.

Letting

$$y = \sum_{r=1}^R \frac{|\delta_r| x^{u_r}}{(1-x^{u_r})^2},$$

we see that $y > 0$ when $x \in (0, e^{-\pi\Delta_3(\ell)/24}]$.

Now it suffices to show that $e^{-y} + y \geq 1$, which is obvious as $e^{-y} + y$ is increasing for $y > 0$. We therefore arrive at the desired result. \square

Remark 3.1. It is helpful to mention more about the necessity of the assumption (1.8). Suppose that there exist some m_r and ℓ such that

$$\frac{\Delta_3(\ell)}{24} > \frac{\gcd^2(m_r, \ell)}{m_r} > 0,$$

then in this case $0 < u_r < 1$. It is not hard to compute that

$$\lim_{x \rightarrow 0} W(x) = \infty.$$

This means that $W(x)$ is not bounded on the interval $(0, e^{-\pi\Delta_3(\ell)/24}]$, which is not what we hope.

Recalling again the bound (2.11) and the assumption $\Delta_1 \leq 0$, it follows from Lemma 3.2 that

$$\begin{aligned}
 |T_{\ell,2}| &\leq \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} \int_{\xi_{h,k}} \Delta_4(\ell) k^{\Delta_1} N^{-2\Delta_1} e^{2\pi\varrho\left(n+\frac{\Delta_2}{24}\right)} e^{\frac{\pi\Delta_3(\ell)}{24}} \\
 &\quad \times \left(\exp \left(\sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) - 1 \right) d\phi \\
 &\leq \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} \Delta_4(\ell) k^{\Delta_1} N^{-2\Delta_1} e^{2\pi\varrho\left(n+\frac{\Delta_2}{24}\right)} e^{\frac{\pi\Delta_3(\ell)}{24}} \\
 &\quad \times \left(\exp \left(\sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) - 1 \right) \frac{2}{kN} \\
 &\leq \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \Delta_4(\ell) k^{\Delta_1} N^{-2\Delta_1} e^{2\pi\varrho\left(n+\frac{\Delta_2}{24}\right)} e^{\frac{\pi\Delta_3(\ell)}{24}} \\
 &\quad \times \left(\exp \left(\sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) - 1 \right) \frac{2}{kN} k \\
 &= 2\Delta_4(\ell) N^{-2\Delta_1-1} e^{2\pi\varrho\left(n+\frac{\Delta_2}{24}\right)} e^{\frac{\pi\Delta_3(\ell)}{24}} \left(\sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} k^{\Delta_1} \right) \\
 &\quad \times \left(\exp \left(\sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) - 1 \right) \\
 &\leq 2\Delta_4(\ell) e^{\frac{\pi\Delta_3(\ell)}{24}} e^{2\pi\varrho\left(n+\frac{\Delta_2}{24}\right)} \Xi_{\Delta_1}(N) \\
 &\quad \times \left(\exp \left(\sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) - 1 \right).
 \end{aligned}$$

To summarize, we have

Lemma 3.3. For $\ell \in \mathcal{L}_{>0}$, it holds that

$$\begin{aligned}
 |T_{\ell,2}| &\leq 2\Delta_4(\ell) e^{\frac{\pi\Delta_3(\ell)}{24}} e^{2\pi\varrho\left(n+\frac{\Delta_2}{24}\right)} \Xi_{\Delta_1}(N) \\
 &\quad \times \left(\exp \left(\sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) - 1 \right). \tag{3.6}
 \end{aligned}$$

At last, we estimate $T_{\ell,1}$.

Lemma 3.4. For $\ell \in \mathcal{L}_{>0}$, it holds that

$$T_{\ell,1} = D_{\ell} + 2\pi \Delta_4(\ell) \left(\frac{24n + \Delta_2}{\Delta_3(\ell)} \right)^{-\frac{\Delta_1+1}{2}} \\ \times \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \frac{I_{-\Delta_1-1} \left(\frac{\pi}{6k} \sqrt{\Delta_3(\ell)(24n + \Delta_2)} \right)}{k} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} \omega_{h,k} e^{-\frac{2\pi i n h}{k}}, \quad (3.7)$$

where

$$|D_{\ell}| \leq \frac{2^{-\Delta_1} \pi^{-1} e^{\frac{\pi \Delta_3(\ell)}{3}} N^{-\Delta_1+2}}{n + \frac{\Delta_2}{24}} e^{2\pi \varrho \left(n + \frac{\Delta_2}{24} \right)}. \quad (3.8)$$

Proof. It follows from Lemma 2.4 that

$$T_{\ell,1} = \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} e^{-\frac{2\pi i n h}{k}} \int_{\xi_{h,k}} \omega_{h,k} \Delta_4(\ell) \\ \times e^{\frac{\pi}{12k} \left(\frac{\Delta_3(\ell)}{z} + \Delta_2 z \right)} z^{\Delta_1} e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi \\ = \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} e^{-\frac{2\pi i n h}{k}} \omega_{h,k} \Delta_4(\ell) \\ \times \int_{\xi_{h,k}} e^{\frac{\pi}{12k} \left(\frac{\Delta_3(\ell)}{z} + \Delta_2 z \right)} z^{\Delta_1} e^{-2\pi i n \phi} e^{2\pi n \varrho} d\phi \\ = D_{\ell} + 2\pi \Delta_4(\ell) \left(\frac{24n + \Delta_2}{\Delta_3(\ell)} \right)^{-\frac{\Delta_1+1}{2}} \\ \times \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \frac{I_{-\Delta_1-1} \left(\frac{\pi}{6k} \sqrt{\Delta_3(\ell)(24n + \Delta_2)} \right)}{k} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} \omega_{h,k} e^{-\frac{2\pi i n h}{k}}.$$

Here

$$|D_{\ell}| \leq \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} \frac{2^{-\Delta_1} \pi^{-1} e^{\frac{\pi \Delta_3(\ell)}{3}} N^{-\Delta_1}}{n + \frac{\Delta_2}{24}} e^{2\pi \varrho \left(n + \frac{\Delta_2}{24} \right)} \\ \leq \frac{2^{-\Delta_1} \pi^{-1} e^{\frac{\pi \Delta_3(\ell)}{3}} N^{-\Delta_1+2}}{n + \frac{\Delta_2}{24}} e^{2\pi \varrho \left(n + \frac{\Delta_2}{24} \right)}. \quad \square$$

3.3. The asymptotic formula of $g(n)$

We know from Lemma 3.4 that the main term of $g(n)$ is

$$\sum_{\ell \in \mathcal{L}_{>0}} 2\pi \Delta_4(\ell) \left(\frac{24n + \Delta_2}{\Delta_3(\ell)} \right)^{-\frac{\Delta_1+1}{2}} \\ \times \sum_{\substack{1 \leq k \leq N \\ k \equiv_L \ell}} \frac{I_{-\Delta_1-1} \left(\frac{\pi}{6k} \sqrt{\Delta_3(\ell)(24n + \Delta_2)} \right)}{k} \sum_{\substack{0 \leq h < k \\ \gcd(h,k)=1}} \omega_{h,k} e^{-\frac{2\pi i n h}{k}}. \quad (3.9)$$

Furthermore, the total error term is

$$|E(n)| \leq \sum_{\ell \in \mathcal{L}_{\leq 0}} |S_\ell| + \sum_{\ell \in \mathcal{L}_{>0}} (|T_{\ell,2}| + |D_\ell|) \\ \leq \frac{2^{-\Delta_1} \pi^{-1} N^{-\Delta_1+2}}{n + \frac{\Delta_2}{24}} e^{2\pi \varrho \left(n + \frac{\Delta_2}{24} \right)} \sum_{\ell \in \mathcal{L}_{>0}} e^{\frac{\pi \Delta_3(\ell)}{3}} \\ + 2e^{2\pi \varrho \left(n + \frac{\Delta_2}{24} \right)} \Xi_{\Delta_1}(N) \\ \times \left(\sum_{1 \leq \ell \leq L} \Delta_4(\ell) \exp \left(\frac{\pi \Delta_3(\ell)}{24} + \sum_{r=1}^R \frac{|\delta_r| e^{-\pi \gcd^2(m_r, \ell)/m_r}}{(1 - e^{-\pi \gcd^2(m_r, \ell)/m_r})^2} \right) \right. \\ \left. - \sum_{\ell \in \mathcal{L}_{>0}} \Delta_4(\ell) e^{\frac{\pi \Delta_3(\ell)}{24}} \right), \quad (3.10)$$

where we use Lemmas 3.1, 3.3 and 3.4.

At last, we set

$$N = \left\lfloor \sqrt{2\pi \left(n + \frac{\Delta_2}{24} \right)} \right\rfloor.$$

It is easy to check that

$$|E(n)| \ll_{\mathbf{m}, \delta} \begin{cases} 1 & \text{if } \Delta_1 = 0, \\ \left(n + \frac{\Delta_2}{24} \right)^{1/4} & \text{if } \Delta_1 = -\frac{1}{2}, \\ \left(n + \frac{\Delta_2}{24} \right)^{1/2} \log \left(n + \frac{\Delta_2}{24} \right) & \text{if } \Delta_1 = -1, \\ \left(n + \frac{\Delta_2}{24} \right)^{-\Delta_1-1/2} & \text{if } \Delta_1 \leq -\frac{3}{2}. \end{cases}$$

We therefore arrive at Theorem 1.1.

Table 1The values of $\Delta_3(\ell)$ and $\Delta_4(\ell)$ for $1 \leq \ell \leq L$.

ℓ	1	2	3	4	5	6	7	8	9	10
$\Delta_3(\ell)$	$\frac{3}{5}$	$-\frac{18}{5}$	$\frac{3}{5}$	$-\frac{18}{5}$	3	$-\frac{18}{5}$	$\frac{3}{5}$	$-\frac{18}{5}$	$\frac{3}{5}$	6
$\Delta_4(\ell)$	$\frac{\sqrt{5}}{2}$	$\sqrt{5}$	$\frac{\sqrt{5}}{2}$	$\sqrt{5}$	$\frac{1}{2}$	$\sqrt{5}$	$\frac{\sqrt{5}}{2}$	$\sqrt{5}$	$\frac{\sqrt{5}}{2}$	1

Table 2The values of $\frac{\pi}{6k} \sqrt{\Delta_3(\ell)}$.

k	1	3	5	7	9	10
$\frac{\pi}{6k} \sqrt{\Delta_3(\ell)}$	$\frac{\pi}{2\sqrt{15}}$	$\frac{\pi}{6\sqrt{15}}$	$\frac{\pi}{10\sqrt{3}}$	$\frac{\pi}{14\sqrt{15}}$	$\frac{\pi}{18\sqrt{15}}$	$\frac{\pi}{10\sqrt{6}}$

4. An application

We end this paper with an application of the main result. Here we will study the asymptotic behavior of the following eta-quotient:

$$G(q) = \frac{(q^2; q^2)_\infty^3}{(q; q)_\infty^2 (q^{10}; q^{10})_\infty} = \sum_{n \geq 0} g(n) q^n. \quad (4.1)$$

This eta-quotient is closely related to the rank statistics for cubic partition pairs. We refer the readers to [16] for details. In particular, (4.1) appears on p. 5 of [16] (the third line of (3.2) therein).

Theorem 4.1. *We have, as $n \rightarrow \infty$,*

$$g(n) \sim 3^{\frac{3}{4}} 5^{\frac{1}{4}} (24n - 6)^{-\frac{3}{4}} \exp \left(\frac{\pi}{2\sqrt{15}} \sqrt{24n - 6} \right). \quad (4.2)$$

Proof. We have $\mathbf{m} = (1, 2, 10)$ and $\delta = (-2, 3, -1)$. It is straightforward to compute that $\Delta_1 = 0$ and $\Delta_2 = -6$. We also have $L = 10$. The values of $\Delta_3(\ell)$ and $\Delta_4(\ell)$ for $1 \leq \ell \leq L$ are listed in Table 1. Hence $\mathcal{L}_{>0} = \{1, 3, 5, 7, 9, 10\}$.

We know from Theorem 1.1 along with (1.13) and (1.14) that $g(n)$ is dominated by the largest

$$I_{-\Delta_1-1} \left(\frac{\pi}{6k} \sqrt{\Delta_3(\ell)(24n + \Delta_2)} \right)$$

provided that this term does not vanish. From Table 2, we see that when $k = 1$, the aforementioned modified Bessel function of the first kind has the largest order.

For $k = 1$ (and hence $\ell = 1$), we further compute that

$$\frac{1}{k} \sum_{\substack{0 \leq h < k \\ \gcd(h, k) = 1}} \omega_{h, k} e^{-\frac{2\pi i n h}{k}} = \cos(2\pi n) = 1$$

for all $n \in \mathbb{Z}_{>0}$. We also have

$$2\pi \Delta_4(\ell) \left(\frac{24n + \Delta_2}{\Delta_3(\ell)} \right)^{-\frac{\Delta_1+1}{2}} = 3^{\frac{1}{2}} \pi (24n - 6)^{-\frac{1}{2}}.$$

We therefore deduce that, as $n \rightarrow \infty$,

$$\begin{aligned} g(n) &\sim 3^{\frac{1}{2}} \pi (24n - 6)^{-\frac{1}{2}} I_{-1} \left(\frac{\pi}{2\sqrt{15}} \sqrt{24n - 6} \right) \\ &\sim 3^{\frac{3}{4}} 5^{\frac{1}{4}} (24n - 6)^{-\frac{3}{4}} \exp \left(\frac{\pi}{2\sqrt{15}} \sqrt{24n - 6} \right), \end{aligned}$$

where we use (1.13). \square

Acknowledgments

I want to thank the referee for careful reading and useful comments.

References

- [1] M. Abramowitz, I.A. Stegun (Eds.), *Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables*, 10th printing, United States Department of Commerce, National Bureau of Standards, 1972.
- [2] G.E. Andrews, *The Theory of Partitions*, Reprint of the 1976 original, Cambridge Math. Lib., Cambridge University Press, Cambridge, 1984, xvi+255 pp.
- [3] G.E. Andrews, R. Lewis, The ranks and cranks of partitions moduli 2, 3, and 4, *J. Number Theory* 85 (1) (2000) 74–84.
- [4] T.M. Apostol, *Modular Functions and Dirichlet Series in Number theory*, second edition, Grad. Texts in Math., vol. 41, Springer-Verlag, New York, 1990, x+204 pp.
- [5] K. Bringmann, K. Ono, Coefficients of harmonic Maass forms, in: *Partitions, q -Series, and Modular Forms*, in: Dev. Math., vol. 23, Springer, New York, 2012, pp. 23–38.
- [6] O-Y. Chan, Some asymptotics for cranks, *Acta Arith.* 120 (2) (2005) 107–143.
- [7] S. Chern, D. Tang, L. Wang, Some inequalities for Garvan’s bicrank function of 2-colored partitions, Preprint, available at arXiv:1805.06575, 2018.
- [8] E. Grosswald, Some theorems concerning partitions, *Trans. Amer. Math. Soc.* 89 (1958) 113–128.
- [9] P. Hagis Jr., Partitions into odd summands, *Amer. J. Math.* 85 (1963) 213–222.
- [10] P. Hagis Jr., On the partitions of an integer into distinct odd summands, *Amer. J. Math.* 87 (1965) 867–873.
- [11] P. Hagis Jr., Some theorems concerning partitions into odd summands, *Amer. J. Math.* 88 (1966) 664–681.
- [12] G.H. Hardy, S. Ramanujan, Asymptotic formulae in combinatory analysis [Proc. London Math. Soc. (2) 16 (1917), Records for 1 March 1917], in: *Collected Papers of Srinivasa Ramanujan*, vol. 244, AMS Chelsea Publ., Providence, RI, 2000.
- [13] G.H. Hardy, E.M. Wright, *An Introduction to the Theory of Numbers*, fifth edition, The Clarendon Press, Oxford University Press, New York, 1979, xvi+426 pp.
- [14] S. Iseki, On some partition functions, *J. Math. Soc. Japan* 12 (1960) 81–88.
- [15] S. Iseki, Partitions in certain arithmetic progressions, *Amer. J. Math.* 83 (1961) 243–264.
- [16] B. Kim, Partition statistics for cubic partition pairs, *Electron. J. Combin.* 18 (1) (2011) 128.
- [17] B. Kim, Periodicity of signs of Fourier coefficients of eta-quotients, *J. Math. Anal. Appl.* 385 (2) (2012) 998–1004.
- [18] B. Kim, H. Nam, On a conjecture of Soon-Yi Kang on a certain partition rank difference, *Ramanujan J.* 35 (3) (2014) 467–477.

- [19] E. Kim, On a sign pattern for the crank function of cubic partition pairs, *J. Number Theory* 190 (2018) 229–240.
- [20] H. Rademacher, On the partition function $p(n)$, *Proc. London Math. Soc.* (2) 43 (4) (1937) 241–254.
- [21] H. Rademacher, On the expansion of the partition function in a series, *Ann. of Math.* (2) 44 (1943) 416–422.
- [22] A.V. Sills, Towards an automation of the circle method, in: *Gems in Experimental Mathematics*, in: *Contemp. Math.*, vol. 517, Amer. Math. Soc., Providence, RI, 2010, pp. 321–338.
- [23] E. Sussman, Rademacher series for η -quotients, Preprint, available at [arXiv:1710.03415](https://arxiv.org/abs/1710.03415), 2017.