



Contents lists available at ScienceDirect

Journal of Number Theory

[www.elsevier.com/locate/jnt](http://www.elsevier.com/locate/jnt)



# Hadamard grade of power series

J.-P. Allouche<sup>a,\*</sup>, M. Mendès France<sup>b</sup>

<sup>a</sup> CNRS, Math., Équipe Combinatoire et Optimisation, Université Pierre et Marie Curie, Case 189, 4 Place Jussieu, F-75252 Paris Cedex 05, France

<sup>b</sup> Université Bordeaux I, Mathématiques, F-33405 Talence Cedex, France

## ARTICLE INFO

### Article history:

Received 4 January 2011

Revised 25 March 2011

Accepted 4 April 2011

Available online xxxx

Communicated by David Goss

À Vera Sós pour son 80<sup>e</sup> anniversaire, avec admiration et amitié

### MSC:

11R58

11D88

13F25

11B85

### Keywords:

Hadamard product

Algebraic power series

Hadamard grade

Automatic sequences

## ABSTRACT

The Hadamard product of two power series  $\sum a_n z^n$  and  $\sum b_n z^n$  is the power series  $\sum a_n b_n z^n$ . We define the (Hadamard) grade of a power series  $A$  to be the least number (finite or infinite) of algebraic power series, the Hadamard product of which equals  $A$ . We study and discuss this notion.

© 2011 Elsevier Inc. All rights reserved.

## 1. Introduction

The Hadamard product of two power series is simply the power series obtained by termwise multiplication of their coefficients. Since the seminal paper of Hadamard [19] this product occurred in various, sometimes unexpected, fields (look for example at the introduction of [5]). We revisit the question of what can be said about the Hadamard product of “simple” series. We introduce in particular the closure under finite Hadamard products of the set of algebraic power series.

\* Corresponding author.

E-mail addresses: [allouche@math.jussieu.fr](mailto:allouche@math.jussieu.fr) (J.-P. Allouche), [michel.mendes-france@math.u-bordeaux1.fr](mailto:michel.mendes-france@math.u-bordeaux1.fr) (M. Mendès France).

## 2. The early work of Hadamard

**Definition 1.** Let  $A(z) := \sum_{n \geq 0} a_n z^n$  and  $B(z) := \sum_{n \geq 0} b_n z^n$  be two power series with coefficients in some commutative field  $K$ . Their *Hadamard product* (sometimes also called *child product*, see [26, Section 8, p. 251], *Schur product*, or *quasi-inner product*, see [5, p. 36]) is the power series  $A * B(z)$  defined by

$$A * B(z) := \sum_{n \geq 0} a_n b_n z^n.$$

Note that this definition makes sense either by considering series  $A$  and  $B$  as formal power series, or, if  $K$  is a subfield of the field of complex numbers  $\mathbb{C}$ , as series possibly converging in a neighborhood of 0 (if  $A$  and  $B$  converge in a neighborhood of 0, then obviously  $A * B$  also converges in some neighborhood of 0). If  $A$  and  $B$  are two functions, we let  $A * B$  denote the Hadamard product of their respective power series, provided they exist.

### 2.1. Singularities of Hadamard products

Starting from the easy observation that for  $a$  and  $b$  nonzero

$$\frac{1}{a-z} * \frac{1}{b-z} = \frac{1}{ab-z}$$

one may ask whether the singularities of the Hadamard product of two series with coefficients in  $\mathbb{C}$  are related to the product of singularities of these series. Hadamard realized that this is indeed the case.

**Theorem 1.** (See Hadamard [19].) Let  $A(z) = \sum a_n z^n$  and  $B(z) = \sum b_n z^n$  be two power series holomorphic in a neighborhood of 0. Let  $\sigma_A$  and  $\sigma_B$  be their respective sets of singularities. Then the singularities of  $A * B$  are among the values  $\alpha\beta$ , where  $\alpha \in \sigma_A$  and  $\beta \in \sigma_B$ . In other words  $\sigma_{A*B} \subset \sigma_A \cdot \sigma_B := \{\alpha\beta, \alpha \in \sigma_A, \beta \in \sigma_B\}$ .

The proof, which is by no means easy, involves the study of the integral

$$\oint A(u) B\left(\frac{z}{u}\right) \frac{du}{u} = \sum a_n b_n z^n$$

extended to a circle around the origin. The holomorphy of  $A(z)$  and  $B(z)$  in a neighborhood of 0 is therefore an essential hypothesis for the result to hold. And indeed in Section 2.2 below we show what may happen if say  $\limsup_{n \rightarrow \infty} |a_n|^{1/n} = \infty$ .

**Remark 1.** Other papers on singularities of Hadamard product may be of interest for the reader: see in particular [4, 11, 12].

### 2.2. A counter-example: the Euler series

The hypothesis that both  $A$  and  $B$  converge in some neighborhood of the origin is necessary for the validity of Theorem 1. Indeed consider the Euler  $E(z)$  formal power series defined by

$$E(z) := \sum_{n \geq 0} (-1)^n n! z^n.$$

This series, considered as a complex series, diverges everywhere except at 0. As will be explained below,  $E(z)$  has the integral representation  $I(z)$  where

$$(\#) \quad I(z) = \int_0^{\infty} \frac{e^{-u}}{1+zu} du = -\frac{1}{z} e^{1/z} \log \frac{1}{z} + S\left(\frac{1}{z}\right)$$

with

$$S(y) := -ye^y \left( \gamma + \sum_{n \geq 1} \frac{(-1)^n y^n}{n.n!} \right)$$

and  $\gamma$  is the Euler constant. The integral above is a function well-defined in the complex plane except on the real negative axis  $\{z = s + it, t = 0, s < 0\}$ .

Now let  $B(z) := e^z$ . We have  $E * B(z) = \sum_{n \geq 0} (-1)^n z^n = \frac{1}{1+z}$  which has only one singularity at  $z = -1$ , while  $e^z$  does not have any singularity.

In order to demystify the integral representation  $E(z) = I(z)$ , compute the derivatives of  $I(z)$ :

$$I^{(n)}(z) = (-1)^n \int_0^{\infty} \frac{n! e^{-u}}{(1+zu)^{n+1}} du$$

which reduces at  $z = 0$  to  $I^{(n)}(0) = (-1)^n (n!)^2$ . Therefore the formal Taylor series for  $I(z)$  is

$$\sum_{n \geq 0} (-1)^n n! z^n$$

which indeed coincides with  $E(z)$ .

The last part of identity (#) is an illuminating computation (see Hardy's book [20, pp. 26–27]) that describes precisely the singularities of  $E(z)$ .

**Theorem 2.** (See [20].) *The integral representation of the Euler series  $\sum_{n \geq 0} (-1)^n n! z^n$  can be continued for all  $z$  to a many-valued function with an infinity of branches differing by integral multiples of  $2i\pi z^{-1} e^{1/z}$ . One branch tends to 1 as  $z$  tends to 0 through positive values. More precisely*

$$I(z) = -\frac{1}{z} e^{1/z} \log \frac{1}{z} + S\left(\frac{1}{z}\right)$$

where

$$S(y) := -ye^y \left( \gamma + \sum_{n \geq 1} \frac{(-1)^n y^n}{n.n!} \right)$$

and  $\gamma$  is the Euler constant.  $I(z)$  lives on the same Riemann surface as  $\log z$ . In particular, for  $z = 1$  [Euler and Hardy write]

$$\sum_{n \geq 0} (-1)^n n! = -e \left( \gamma + \sum_{n \geq 1} \frac{(-1)^n}{nn!} \right) = .5963 \dots$$

**Proof.** (This proof is attributed to Euler by Hardy in [20]).

Suppose  $z = x$  is a positive real number. Performing the change of variable  $t = \frac{x}{1+xu}$ , we have

$$I(x) = \int_0^{\infty} \frac{e^{-u}}{1+xu} du = -\frac{1}{x} e^{1/x} \operatorname{li}(e^{-1/x})$$


where  $\operatorname{li}(\xi) := \int_0^{\xi} \frac{ds}{\log s}$ , for  $\xi \in [0, 1)$ , is the *logarithmic integral* (*logarithm-integral* in Hardy's terminology). Now

$$\begin{aligned} -\operatorname{li}(e^{-y}) &= \int_y^{\infty} \frac{e^{-u}}{u} du = \int_1^{\infty} \frac{e^{-u}}{u} du - \int_0^1 \frac{1-e^{-u}}{u} du - \int_1^y \frac{du}{u} + \int_0^y \frac{1-e^{-u}}{u} du \\ &= -\gamma - \log y - \sum_{n \geq 1} \frac{(-1)^n y^n}{n \cdot n!} \end{aligned}$$

which concludes the proof.  $\square$

### 3. Digression: elementary optics

Consider  $N$  transparent plates one against the other with respective thicknesses  $a_1, a_2, \dots, a_N$  ( $a_n > 0$ ) and indices of refraction  $n_1, n_2, \dots, n_N$  ( $n_k > 1$ ).



An incoming ray hits the system at the origin with an incident angle  $\theta$  and reappears after  $N$  refractions at  $M$  on the opposite side. Let  $\theta_k$  be the angle at which the luminous path meets the  $k$ th plate. The Snell–Descartes law states that  $n_k \sin \theta_k$  is independent of  $k$ . An easy computation shows that the length  $AM = \varphi(\theta)$  is given by

$$\varphi(\theta) = \sum_{1 \leq k \leq N} \frac{a_k \sin \theta}{\sqrt{n_k^2 - \sin^2 \theta}}.$$

Put  $z := \sin \theta$  and  $\psi(z) := \varphi(\theta)$ , thus

$$\psi(z) = \sum_{1 \leq k \leq N} \frac{a_k z}{\sqrt{n_k^2 - z^2}}.$$

In [23] the second named author and A. Sebbar notice that

$$\psi(z) = \frac{z}{\sqrt{1-z^2}} * \sum_{1 \leq k \leq N} \frac{a_k n_k z}{n_k^2 - z^2}.$$

If none of the  $a_k$  vanish and if the  $n_k$  are pairwise distinct, then  $\psi$  is an algebraic function of degree  $2^N$ . The above Hadamard representation is somewhat simpler than the original  $\psi$  since one factor is rational whereas the other is quadratic.

The above identity is even valid for  $N = \infty$ ,  $a_k$  and  $n_k \in \mathbb{C}$ , provided the series converges. The second factor can often be summed, especially if  $a_k$  and  $n_k$  are rational functions of  $k$ . Choose for example  $n_k = (2k+1)$  and  $a_k = (2k+1)^{-1}$ . Then the identity reads:

$$\sum_{k \geq 0} \frac{z}{(2k+1)\sqrt{(2k+1)^2 - z^2}} = \frac{\pi}{4} \frac{z}{\sqrt{1-z^2}} * \tan \frac{\pi z}{2}.$$

This could be thought of as a quasi-summation of the left hand side series.

Let us test the formula. For  $z \rightarrow 0$  it reduces to

$$\sum_{k \geq 0} \frac{1}{(2k+1)^2} = \frac{\pi^2}{8}.$$

More generally, put  $\tan X := \sum_{j \geq 0} t_{2j+1} X^{2j+1}$ . Observe that

$$\frac{1}{(2k+1)\sqrt{(2k+1)^2 - z^2}} = \frac{1}{(2k+1)^2} \sum_{j \geq 0} (-1)^j \binom{-\frac{1}{2}}{j} \frac{z^{2j}}{(2k+1)^{2j}}.$$

Then the above Hadamard product leads to the well-known equality

$$\sum_{k \geq 0} \frac{1}{(2k+1)^{2j+2}} = \frac{\pi^{2j+2}}{2^{2j+3}} t_{2j+1}.$$

The example  $a_k = (2k+1)^{-1}$  and  $n_k = (2k+1)$  corresponds to an optical system with infinitely many thinner and thinner plates but with no physical meaning since  $n_k = (2k+1)$  tends to infinity. In real materials the highest indices seem not to go beyond 4 or 5.

One last remark before closing this section. The identity showing that  $\psi$  is a Hadamard product of two simple functions can easily be extended. Let  $H(z) := \sum_{j \geq 0} h_j z^{2j+1}$  be any odd function holomorphic around the origin ( $|z| < R$ ). Then, for all complex nonzero  $a_k$  and  $n_k$ , and for  $N \leq +\infty$ ,

$$\sum_{1 \leq k \leq N} a_k H\left(\frac{z}{n_k}\right) = H(z) * \sum_{1 \leq k \leq N} \frac{a_k n_k z}{n_k^2 - z^2}$$

provided  $|z| < R \inf_k |n_k|$ . The proof consists in expanding both sides of the identity into Taylor series, just as in the case  $H(z) = z(1-z^2)^{-1/2}$ . Needless to say that similar identities hold if  $H$  is even, or neither odd nor even.

#### 4. Recalling some definitions and results

In what follows we will need the definition of *D-finite* power series and the definition of *automatic* sequences.

**Definition 2.** (See [24].) A power series is *D-finite* (or *differentially finite* or *holonomic*) if it satisfies a linear differential equation with polynomial coefficients.

**Example 1.** The power series  $F := \sum_{n \geq 0} \frac{z^n}{n!}$  is *D-finite*, since it satisfies  $F' - F = 0$ . The series  $\sum_{n \geq 0} \sqrt{n} z^n$  is not *D-finite* [16]. The series  $\sum_{n \geq 1} \log n z^n$  and  $\sum_{n \geq 1} p_n z^n$  where  $p_n$  is the  $n$ th prime are not *D-finite* [13]. A nice example of a non-*D-finite* power series is  $\sum_{n \geq 1} \zeta(2n+1) z^n$  where  $\zeta$  is the Riemann function (see [3]).

**Definition 3.** Let  $q$  be an integer  $\geq 2$ . A sequence  $(a_n)_{n \geq 0}$  is *q-automatic* if the  $q$ -kernel of  $(a_n)_{n \geq 0}$ , i.e., the set of subsequences  $\{(a_{q^k n + j})_{n \geq 0}, k \geq 0, j \in [0, q^k - 1]\}$ , is finite.

#### 5. The Hadamard grade of a power series

**Definition 4.** A power series  $A(z) := \sum a_n z^n$  with coefficients in a commutative field  $K$  is said to have *finite Hadamard grade* over  $K$  if there exist finitely many power series  $B_1(z), B_2(z), \dots, B_k(z)$  that are algebraic (over  $K(z)$ ) such that  $A = B_1 * B_2 * \dots * B_k$ . The least  $k$  having this property is called the *Hadamard grade* over  $K$  of  $A$ .

We give some first properties of power series with finite Hadamard grade.

##### Theorem 3.

- The Hadamard product of two power series having finite Hadamard grade has finite Hadamard grade.
- If  $A$  is a power series that is algebraic over  $K(z)$ , then  $A$  has Hadamard grade 1.
- If  $A$  is a power series with finite Hadamard grade, then  $A$  is *D-finite*.

**Proof.** The first two assertions are clear. For the third, recall that a power series is *D-finite* (or *differentially finite* or *holonomic*) if it satisfies a linear differential equation with polynomial coefficients (see [24]); also recall that any algebraic power series is *D-finite* and that the Hadamard product of two *D-finite* power series is also *D-finite* (see [24]).  $\square$

**Remark 2.** The *D-finiteness* of the Hadamard product of two *D-finite* power series can be found in the paper [24], but as noted by Stanley this is already in the paper of Jungen [22] who indicates that this result can be found in the unpublished manuscripts of Hurwitz (available at École Polytechnique de Zurich). Jungen also gives the same characterization as Stanley's for the coefficients of *D-finite* power series.

The next result shows that in positive characteristic the definition of the grade is not really pertinent (that is why in the next sections we will focus on the case where  $K$  is the field of complex numbers  $\mathbb{C}$  or the field of rational numbers  $\mathbb{Q}$ ).

**Theorem 4.** A power series on a field of positive characteristic has finite Hadamard grade if and only if it is algebraic.

**Proof.** Only one direction need to be proven. This was done by Furstenberg [15] for a finite field. Fliess [14] noted that the proof still works for a perfect field of positive characteristic. Deligne [8] gave a general proof.  $\square$

**Remark 3.** For more concerning algebraic power series the reader can look for example at [1,17,18,26] and the references therein.

## 6. Hadamard grade of formal power series with complex or with rational coefficients

### 6.1. The base field is $\mathbb{C}$

From now on we will restrict ourselves to the case where the field  $K$  is either  $\mathbb{C}$  or  $\mathbb{Q}$ . We begin with a theorem stating that a minimal Hadamard decomposition of an irrational power series into finitely many algebraic power series only involves algebraic irrational power series.

**Theorem 5.** *Let  $A(z)$  be a power series with complex coefficients. If  $A$  is irrational and has finite Hadamard grade  $d$ , then in any Hadamard decomposition  $A = B_1 * B_2 * \cdots * B_d$  with algebraic  $B_i$ 's, all the  $B_i$ 's are irrational.*

**Proof.** It suffices to use the minimality of  $d$  and a result of Jungen [22] stating that the Hadamard product of a rational power series and an algebraic power series is algebraic.  $\square$

The following result is classical.

**Theorem 6.** *Let  $F = \sum_{n \geq 0} a_n z^n$  be a formal power series with coefficients in the complex field  $\mathbb{C}$  that is algebraic over  $\mathbb{C}(z)$ . Then the radius of convergence of  $F$  is  $> 0$  and  $F$  has finitely many singularities on the circle of convergence.*

Using this theorem and Hadamard's Theorem 1 yields the following corollary.

**Corollary 1.** *Let  $A(z)$  be a formal power series with coefficients in  $\mathbb{C}$  which has finite Hadamard grade. Then  $A$  has a positive radius of convergence, and  $A$  has only finitely many singularities on the circle of convergence.*

There is a simple necessary condition for a complex power series to be of finite Hadamard grade in terms of diagonals of rational functions.

**Theorem 7.** *If  $A$  is a complex power series with finite Hadamard grade  $d$ , then  $A$  is the (complete) diagonal of a complex rational power series with  $2d$  variables.*

**Proof.** The proof is done by induction on  $d$ . The case  $d = 1$  can be found in the paper of Furstenberg [15]. Now if  $A_1, A_2, \dots, A_d$  are complex algebraic power series, then there exist a two-variable rational power series  $B(X, Y) := \sum b_{k,\ell} X^k Y^\ell$  and a rational power series  $C(Z_1, Z_2, \dots, Z_{2d-2}) := \sum c_{j_1, j_2, \dots, j_{2d-2}} Z_1^{j_1} Z_2^{j_2} \cdots Z_{2d-2}^{j_{2d-2}}$  such that  $A_d(z) = \sum b_{k,k} z^k$  and  $A_1(z) * A_2(z) * \cdots * A_{d-1}(z) = \sum c_{j,j,\dots,j} z^j$ . The (ordinary Cauchy) product of the series  $B$  and  $C$ , i.e.,  $B(X, Y)C(Z_1, Z_2, \dots, Z_{2d-2}) = \sum b_{k,\ell} c_{j_1, j_2, \dots, j_{2d-2}} X^k Y^\ell Z_1^{j_1} Z_2^{j_2} \cdots Z_{2d-2}^{j_{2d-2}}$  is a rational power series whose complete diagonal  $\sum b_{j,j} c_{j,j,\dots,j} z^j$  is equal to  $A_1(z) * A_2(z) * \cdots * A_d(z)$ .  $\square$

### 6.2. The base field is $\mathbb{Q}$

What can be said of a power series with rational coefficients and finite Hadamard grade over  $\mathbb{Q}$ ? We begin with a simple remark.

**Remark 4.** The family of power series with rational coefficients and with finite Hadamard grade over  $\mathbb{Q}$  is countable.

Our next result links finite Hadamard grade over  $\mathbb{Q}$  and automaticity for the case of power series with rational coefficients (for more about *automatic sequences* the reader can look at [2]).

**Theorem 8.** Let  $A(z) := \sum a_n z^n$  be a power series with rational coefficients. If  $A$  has finite Hadamard grade over the field of rational numbers  $\mathbb{Q}$ , then there are only finitely many primes dividing at least one of the denominators of the  $a_n$ 's. Furthermore for all primes  $p$  outside these finitely many primes, the series  $\bar{A}(z) := \sum (a_n \bmod p) z^n$  is algebraic over  $\mathbb{F}_p(z)$ . In particular the sequence  $(a_n \bmod p)_{n \geq 0}$  is  $p$ -automatic. Even more, for these primes  $p$  and for any  $r \geq 1$ , the sequence  $(a_n \bmod p^r)_{n \geq 0}$  is  $p$ -automatic.

**Proof.** Write  $A = B_1 * B_2 * \cdots * B_d$  where the  $B_i$ 's are algebraic over  $\mathbb{C}(z)$ . From Eisenstein's theorem [10] (see Remark 5 below) the denominators of the coefficients of each  $B_i$  have only finitely many distinct prime factors. Let  $\mathcal{P}$  be the (finite) set of primes that divide the denominator of at least one  $B_i$ . For each prime  $p \notin \mathcal{P}$  one can write  $(A \bmod p) = (B_1 \bmod p) * (B_2 \bmod p) * \cdots * (B_d \bmod p)$ . Each  $(B_i \bmod p)$  is algebraic over  $\mathbb{F}_p(z)$ , so is their Hadamard product (Theorem 4). Actually for any  $r \geq 1$  the sequence  $(a_n \bmod p^r)_{n \geq 0}$  is  $p$ -automatic: this was proved by Christol (see [6,7], see also [9]).  $\square$

### Remark 5.

- The Eisenstein theorem states: Let  $\sum a_n z^n$  be an algebraic power series with rational coefficients. Then there exist two nonzero integers  $A$  and  $C$  such that, for all  $n \geq 0$ , the number  $CA^n a_n$  is an integer. This theorem was actually proved by Heine [21].
- Christol (see, e.g., [7]) calls *globally bounded* any series  $f = \sum a_n z^n$  with rational coefficients such that  $f$  has a positive radius of convergence (when it is seen as a power series on  $\mathbb{C}$ ) and such that there exist  $\alpha, \beta \in \mathbb{Q}^*$  such that  $\alpha f(\beta z)$  belongs to  $\mathbb{Z}[[z]]$ . He shows that any algebraic power series  $\sum a_n z^n$  is *globally automatic*, i.e., for all but finitely many primes  $p$  and for all integers  $r \geq 2$ , the sequence  $(a_n \bmod p^r)_{n \geq 0}$  is  $p$ -automatic.

## 7. Examples, counterexamples, and questions

Using the results of the previous sections we can give examples and counterexamples of power series having finite Hadamard grade: we will restrict to power series **with rational coefficients**, and consider their **Hadamard grade over  $\mathbb{Q}$** , although some of the statements and questions below could be formulated for the Hadamard grade over  $K$  of power series with coefficients in  $K$ , where  $K$  is any field of characteristic zero.

(a) The power series  $e^z$ ,  $\log(1+z)$ ,  $\sum \pm z^n$  where the sequence  $\pm 1$  is not ultimately periodic, all have infinite Hadamard grade.

**Proof.** The denominators of the power series  $e^z$  and  $\log(1+z)$  are divisible by infinitely many primes. If the sequence  $\pm$  is not ultimately periodic, then the series  $\sum \pm z^n$  admits the unit circle as natural boundary (this was proved by Szegő in [25]).  $\square$

(b) There exist power series with finite Hadamard grade  $\geq 2$  (i.e., nonalgebraic series with finite Hadamard grade, since all series with Hadamard grade  $\geq 2$  are transcendental).

**Proof.** There is a classical example of a nonalgebraic Hadamard product of two algebraic power series (see [22]). Namely the Hadamard square of  $\sum \binom{2n}{n} z^n = (1-4z)^{-1/2}$ , is equal to  $\sum \binom{2n}{n}^2 z^n = \frac{2}{\pi} \int_0^{\pi/2} \frac{d\theta}{(1-16z \sin^2 \theta)^{1/2}}$  which is transcendental.  $\square$

(c) There exist  $D$ -finite power series that have infinite Hadamard grade. There even exist  $D$ -finite power series with integer coefficients that have infinite Hadamard grade.



**Proof.** First the exponential series  $\sum_{n \geq 0} z^n/n!$  has infinite Hadamard grade since the denominators of its coefficients are divisible by infinitely many primes (actually *all* the primes); it is of course  $D$ -finite. Second the Euler series  $E(z) := \sum_{n \geq 0} (-1)^n n! z^n$  is clearly  $D$ -finite since  $z^2 E'(z) + (1+z)E(z) = 1$ , while it has infinite Hadamard grade (its radius of convergence is 0).  $\square$

**Question.** Is there a  $D$ -finite power series with integer coefficients and positive radius of convergence which is algebraic on  $\mathbb{F}_p(z)$  for all primes  $p$  when reduced modulo  $p$ , but has infinite Hadamard grade?

**Question.** Is it true that algebraic power series can be decomposed into finite Hadamard products of quadratic power series?

(d) *There exist two power series  $A$  and  $B$  having infinite Hadamard grade, such that  $A * B$  has finite Hadamard grade.*

**Proof.** Of course if  $A$  is even and  $B$  odd, then  $A * B = 0$ . But, from Section 2.2, we have even more than the claim above: take  $A(z) := E(z)$  the Euler series and  $B(z) := e^z$ . These power series both have infinite Hadamard grade, and their Hadamard product is rational, hence of Hadamard grade 1. Another example is given by the Hadamard square of  $\sum \pm z^n$ , where the sequence of  $\pm$ 's is not ultimately periodic.  $\square$

## Questions.

- Is it true that for all  $k > 0$  there exists a power series whose grade is  $k$ ?
- Is it true that if  $B$  is a nonzero rational power series, then for every power series  $A$  the Hadamard grade of  $A + B$  and of  $AB$  are both equal to the Hadamard grade of  $A$ ?
- Is it true that if the Hadamard grade of a power series  $A$  is infinite, then for any nonzero power series  $B$  of finite Hadamard grade, the Hadamard grades of  $A + B$  and  $AB$  are both infinite?
- Let  $a$  be an irrational number (real or complex). Is it true that the grade of  $(1+z)^a$  is infinite?
- Let  $\sum a_n z^n$  be algebraic irrational. Is it true that for all integers  $k \geq 1$  the Hadamard grade of  $\sum a_n^k z^n$  is equal to  $k$ ? Is it the case if  $a_n := \binom{2n}{n}$ ?
- Let  $A(z) := \sum a_n z^n$ . Let  $\mathcal{H}_k$  denote the Hadamard grade of the Hadamard product of  $k$  factors equal to  $A$ . If there exists  $C > 0$  such that  $\mathcal{H}_k < C$  for all  $k$ , is it true that  $A(z)$  must be a rational function? What if  $A$  satisfies the weaker condition  $\mathcal{H}_k = o(k)$ ? Or even the weaker condition that there exists an integer  $k \geq 2$  such that  $\mathcal{H}_k < k$ ? (Note that conversely if  $A$  is rational then  $\mathcal{H}_k = 1$  for all  $k \geq 1$ .)

**Remark 6.** More examples (where  $\mathbb{Q}$  is replaced by  $\mathbb{C}$ ) can be obtained from results of [3] on non- $D$ -finite power series.

## Acknowledgments

We want to thank warmly J.-P. Bézivin, M. Bousquet-Mélou, and J.-M. Deshouillers for comments and discussions.

## References

- [1] J.-P. Allouche, Note sur un article de Sharif and Woodcock, J. Théor. Nombres Bordeaux 1 (1989) 163–187.
- [2] J.-P. Allouche, J. Shallit, Automatic Sequences. Theory, Applications, Generalizations, Cambridge University Press, Cambridge, 2003.
- [3] J.P. Bell, S. Gerhold, M. Klazar, F. Luca, Non-holonomicity of sequences defined via elementary functions, Ann. Comb. 12 (2008) 1–16.
- [4] É. Borel, Sur les singularités des séries de Taylor, Bull. Soc. Math. France 26 (1898) 238–248.
- [5] L.R. Bragg, Trigonometric integrals and Hadamard products, Amer. Math. Monthly 106 (1999) 36–42.

- [6] G. Christol, Limites uniformes  $p$ -adiques de fonctions algébriques, vol. 6, Thèse Sciences Math., Université Paris 6, 1977.
- [7] G. Christol, Globally bounded solutions of differential equations, in: *Lecture Notes in Math.*, vol. 1434, 1990, pp. 45–64.
- [8] P. Deligne, Intégration sur un cycle évanescant, *Invent. Math.* 76 (1983) 129–143.
- [9] J. Denef, L. Lipshitz, Algebraic power series and diagonals, *J. Number Theory* 26 (1987) 46–67.
- [10] G. Eisenstein, Über eine allgemeine Eigenschaft der Reihenentwicklungen aller algebraischen Funktionen, *Berlin. Sitzber.* (1852) 441–443.
- [11] G. Faber, Bemerkungen zu einem funktionentheoretischen Satze des Herrn Hadamard, *Jahresber. Deutsch. Math.-Verein.* 16 (1892/1907) 285–298.
- [12] J. Fill, P. Flajolet, N. Kapur, Singularity analysis, Hadamard products, and tree recurrences, *J. Comput. Appl. Math.* 174 (2005) 271–313.
- [13] P. Flajolet, S. Gerhold, B. Salvy, On the non-holonomic character of logarithms, powers, and the  $n$ th prime function, *Electron. J. Combin.* 11 (2005), Article #A2, 16 pp.
- [14] M. Fliess, Sur certaines familles de séries formelles, Thèse, Paris, VII, 1972.
- [15] H. Furstenberg, Algebraic functions over finite fields, *J. Algebra* 7 (1967) 271–277.
- [16] S. Gerhold, On some non-holonomic sequences, *Electron. J. Combin.* 11 (2004), Article #R87, 7 pp.
- [17] D. Griffiths, Power series expansions of algebraic functions, PhD Thesis, Macquarie University, 1992.
- [18] D. Griffiths, Series expansions of algebraic functions, in: W. Bosma, A. van der Poorten (Eds.), *Computational Algebra and Number Theory*, Kluwer, 1995, pp. 267–277.
- [19] J. Hadamard, Théorème sur les séries entières, *Acta Math.* 22 (1899) 55–63.
- [20] G.H. Hardy, *Divergent Series*, 2nd edition, Chelsea Publishing Company, 1991.
- [21] E. Heine, Der Eisensteinsche Satz über Reihen-Entwicklung algebraischer Functionen, *J. Reine Angew. Math.* 45 (1853) 285–302.
- [22] R. Jungen, Sur les séries de Taylor n'ayant que des singularités algébrique-logarithmiques sur leur cercle de convergence, *Comment. Math. Helv.* 3 (1931) 266–306.
- [23] M. Mendès France, A. Sebbar, Reflections on refraction (geometrical optics), *Ann. Henri Poincaré* 9 (2008) 625–638.
- [24] R.P. Stanley, Differentiability finite power series, *European J. Combin.* 1 (1980) 175–188.
- [25] G. Szegő, Über Potenzreihen mit endlich vielen verschiedenen Koeffizienten, in: *Sitzungsberichte der Preussischen Akademie der Wissenschaften, Phys.-Math. Klasse* (1922) 88–91. Reprinted in collected papers: R. Askey (Ed.), *Contemp. Math.*, vol. 1, Birkhäuser, Boston, Basel, Stuttgart, 1982, pp. 557–561.
- [26] A.J. van der Poorten, Power series representing algebraic functions, in: *Séminaire de Théorie des Nombres*, Paris, 1990–1991, pp. 241–262.