

Exact multiplicity of solutions and S-shaped bifurcation curves for the p -Laplacian perturbed Gelfand problem in one space variable [☆]

Shin-Hwa Wang ^{a,*}, Tzung-Shin Yeh ^a

^a Department of Mathematics, National Tsing Hua University, Hsinchu, Taiwan 300, Republic of China

^b Department of Applied Mathematics, Hsuan Chuang University, Hsinchu, Taiwan 300, Republic of China

Received 3 October 2006

Available online 28 January 2008

Submitted by H.W. Broer

Abstract

We study exact multiplicity of positive solutions and the bifurcation curve of the p -Laplacian perturbed Gelfand problem from combustion theory

$$\begin{cases} (\varphi_p(u'(x)))' + \lambda \exp\left(\frac{au}{a+u}\right) = 0, & -1 < x < 1, \\ u(-1) = u(1) = 0, \end{cases}$$

where $p > 1$, $\varphi_p(y) = |y|^{p-2}y$, $(\varphi_p(u'))'$ is the one-dimensional p -Laplacian, $\lambda > 0$ is the Frank–Kamenetskii parameter, $u(x)$ is the dimensionless temperature, and the reaction term $f(u) = \exp(\frac{au}{a+u})$ is the temperature dependence obeying the Arrhenius reaction-rate law. We find explicitly $\tilde{a} = \tilde{a}(p) > 0$ such that, if the activation energy $a \geq \tilde{a}$, then the bifurcation curve is S-shaped in the $(\lambda, \|u\|_\infty)$ -plane. More precisely, there exist $0 < \lambda_* < \lambda^* < \infty$ such that the problem has exactly three positive solutions for $\lambda_* < \lambda < \lambda^*$, exactly two positive solutions for $\lambda = \lambda_*$ and $\lambda = \lambda^*$, and a unique positive solution for $0 < \lambda < \lambda_*$ and $\lambda^* < \lambda < \infty$.
© 2007 Elsevier Inc. All rights reserved.

Keywords: Exact multiplicity; Positive solution; S-shaped bifurcation curve; p -Laplacian; Perturbed Gelfand problem; Time map

1. Introduction

We study exact multiplicity of positive solutions and the bifurcation curve of the p -Laplacian perturbed Gelfand problem

$$\begin{cases} (\varphi_p(u'(x)))' + \lambda \exp\left(\frac{au}{a+u}\right) = 0, & -1 < x < 1, \\ u(-1) = u(1) = 0, \end{cases} \quad (1.1)$$

[☆] Work partially supported by the National Science Council, Republic of China.

* Corresponding author. Fax: +886 3 5723888.

E-mail addresses: shwang@math.nthu.edu.tw (S.-H. Wang), tsyeh@hcu.edu.tw (T.-S. Yeh).




Fig. 1. S-shaped bifurcation curve \bar{S} of (1.1) for $p > 1$ and $a \geq \tilde{a}$.

where $p > 1$, $\varphi_p(y) = |y|^{p-2}y$, $(\varphi_p(u'))'$ is the one-dimensional p -Laplacian. This is the one-dimensional case of a problem arising in the study of (steady state) solid fuel ignition models in thermal combustion theory, and it was discussed in [1] and many references cited within for the case when $p = 2$ (Laplacian case). See also [7, Section 4]. In this context, the quantity p is a characteristic of the medium, $\lambda > 0$ is the Frank–Kamenetskii parameter, $u(x)$ is the dimensionless temperature, the reaction term $f(u) = \exp(\frac{au}{a+u})$ is the temperature dependence obeying the Arrhenius reaction-rate law, and $a > 0$ is the activation energy.

It is known that, for any given $\alpha > 0$, there exists a unique $\lambda = \lambda(\alpha) > 0$ such that (1.1) admits a unique positive solution u with $\|u\|_\infty = \alpha$. We define the bifurcation curve of (1.1)

$$\bar{S} = \{(\lambda, \|u\|_\infty) : \lambda > 0 \text{ and } u \text{ is a positive solution of (1.1)}\}.$$

In Theorem 2.1 stated below, which is the main result in this paper, we find explicitly $\tilde{a} = \tilde{a}(p) > 0$ such that, if the activation energy $a \geq \tilde{a}$, then the bifurcation curve \bar{S} is S-shaped in the $(\lambda, \|u\|_\infty)$ -plane; that is, \bar{S} has *exactly two* turning points at some points $(\lambda^*, \|u_{\lambda^*}\|_\infty)$ and $(\lambda_*, \|u_{\lambda_*}\|_\infty)$ such that

- (i) $\lambda_* < \lambda^*$ and $\|u_{\lambda_*}\|_\infty < \|u_{\lambda^*}\|_\infty$,
- (ii) at $(\lambda^*, \|u_{\lambda^*}\|_\infty)$ the bifurcation curve \bar{S} turns to the left,
- (iii) at $(\lambda_*, \|u_{\lambda_*}\|_\infty)$ the bifurcation curve \bar{S} turns to the right.

More precisely, problem (1.1) has exactly three positive solutions for $\lambda_* < \lambda < \lambda^*$, exactly two positive solutions for $\lambda = \lambda_*$ and $\lambda = \lambda^*$, and a unique positive solution for $0 < \lambda < \lambda_*$ and $\lambda^* < \lambda < \infty$. See Fig. 1.

We recall some results on S-shaped bifurcation curves for bifurcation problems related to (1.1). When $p = 2$, problem (1.1) reduces to


$$\begin{cases} u''(x) + \lambda \exp\left(\frac{au}{a+u}\right) = 0, & -1 < x < 1, \\ u(-1) = u(1) = 0. \end{cases} \quad (1.2)$$

It is well known that, if $0 < a \leq 4$, the bifurcation curve \bar{S} for (1.2) is a monotone curve in the $(\lambda, \|u\|_\infty)$ -plane, see, e.g., [2, p. 482]. That is, \bar{S} has no turning point. In [12], using the quadrature method, Wang proved the next S-shaped bifurcation curve theorem for (1.2) for $a > a^* \approx 4.4967$ with some constant a^* defined in [12, Eq. (2.21)].

Theorem 1.1. (See Wang [12], Fig. 1.) Let $p = 2$. Consider (1.2). There exists $a^* \approx 4.4967$ such that if $a > a^*$, then

$$\lim_{\alpha \rightarrow 0^+} \lambda(\alpha) = 0, \quad \lim_{\alpha \rightarrow \infty} \lambda(\alpha) = \infty,$$

and the bifurcation curve \bar{S} is S-shaped in the $(\lambda, \|u\|_\infty)$ -plane. More precisely, there exist $0 < \lambda_* < \lambda^* < \infty$ such that (1.2) has exactly three positive solutions for $\lambda_* < \lambda < \lambda^*$, exactly two positive solutions for $\lambda = \lambda_*$ and $\lambda = \lambda^*$, and a unique positive solution for $0 < \lambda < \lambda_*$ and $\lambda^* < \lambda < \infty$.

Fig. 2. Graphs of $\tilde{a} > 4(p-1)$ for $1 < p \leq 3$.

Note that the result in Theorem 1.1 was improved by Korman and Li [8, Theorem 3.1] for $a > a^{**} \approx 4.35$ by applying a bifurcation theorem of Crandall and Rabinowitz [4]. When $\Omega = B_1$ is the unit ball in \mathbb{R}^2 , Du and Lou [6] proved that the bifurcation curve $\{(\lambda, \|u\|_\infty)\}$ of the perturbed Gelfand problem

$$\begin{cases} \Delta u + \lambda \exp\left(\frac{au}{a+u}\right) = 0 & \text{in } B_1, \\ u = 0 & \text{on } \partial B_1 \end{cases}$$

is S-shaped if a is large enough.

In [13], Wang studied problem (1.2) with nonlinearity $f(u) = \exp(\frac{au}{a+u})$ replaced by $\tilde{f}(u) = (1 + u/a)^m \exp(\frac{au}{a+u})$, in which \tilde{f} is the temperature dependence of m th ($m < 1$) order reaction rate obeying the *general* Arrhenius reaction-rate law. Du [5] extended the results of [13] for the one-dimensional case to cover both dimensions one and two, and extended the results of [6] for the special case $m = 0$ to the general case $0 \leq m < 1$.

Recently, Ramaswamy and Shivaji [11, Section 4] studied existence of multiple positive solutions of the p -Laplacian perturbed Gelfand problem

$$\begin{cases} \Delta_p u + \lambda \exp\left(\frac{au}{a+u}\right) = 0 & \text{in } \Omega, \\ u = 0 & \text{on } \partial\Omega, \end{cases} \quad (1.3)$$

where $\Delta_p u = \operatorname{div}(|\nabla u|^{p-2} \nabla u)$, $\lambda > 0$, and Ω is a general bounded domain in \mathbb{R}^N , $N \geq 2$ with $\partial\Omega$ of class C^2 and connected. (If $N = 1$, they assumed that Ω is a bounded open interval.) They showed that problem (1.3) has *at least three* positive solutions for a certain range of λ if a is large enough by applying the sub-super solution techniques. Note that it is well known that a necessary condition for multiplicity for (1.3) and also for (1.1) is $a > 4(p-1)$, see, e.g., [11, Section 4]. See also Jacobsen and Schmitt [7, Section 4] for precise existence and multiplicity results for radial solutions of the p -Laplacian Gelfand problem

$$\begin{cases} \Delta_p u + \lambda \exp(u) = 0 & \text{in } B_1, \\ u = 0 & \text{on } \partial B_1, \end{cases}$$

where B_1 is the unit ball in \mathbb{R}^N ($N \geq 1$).

The paper is organized as follows. Section 2 contains statements of Theorem 2.1, Lemmas 2.2–2.4 used to prove Theorem 2.1, and the proof of Theorem 2.1. Section 3 contains proofs of Lemmas 2.2–2.4.

2. Main result

Theorem 2.1. (See Figs. 1 and 2.) Let $p > 1$. Consider (1.1). If

$$a \geq \tilde{a} \equiv \begin{cases} \max\{4, (4p-2)(\ln p) + 0.7\} & \text{if } 1 < p < 2, \\ (4p-2)(\ln p) + \eta & \text{if } p \geq 2, \end{cases} \quad (2.1)$$

where

$$\eta \approx 0.622$$

is the positive zero of

$$M(u) \equiv 4u \exp(u/2) - u - 4 \ln 2$$

satisfying $M(u) < 0$ for $0 < u < \eta$, then the bifurcation curve \bar{S} is S-shaped in the $(\lambda, \|u\|_\infty)$ -plane. More precisely, there exist $0 < \lambda_* < \lambda^* < \infty$ such that (1.1) has exactly three positive solutions for $\lambda_* < \lambda < \lambda^*$, exactly two positive solutions for $\lambda = \lambda_*$ and $\lambda = \lambda^*$, and a unique positive solution for $0 < \lambda < \lambda_*$ and $\lambda^* < \lambda < \infty$.

Remark 1. In (2.1),

(i) For $1 < p < 2$,

$$\tilde{a} \equiv \max \left\{ 4, (4p-2)(\ln p) + 0.7 \right\} = \begin{cases} 4 & \text{for } 1 < p \leq p^*, \\ (4p-2)(\ln p) + 0.7 & \text{for } p^* < p < 2, \end{cases}$$

where we define $p^* \approx 1.846$ is the unique root of the equation $(4p-2)(\ln p) + 0.7 = 4$ in the interval $(1, 2)$.

(ii) For $p = 2$, $\tilde{a} = 6(\ln 2) + \eta \approx 4.781 > 4.4967 \approx a^*$, in which a^* stated in Theorem 1.1, is a lower bound of activation energy a such that the bifurcation curve \bar{S} is S-shaped in the $(\lambda, \|u\|_\infty)$ -plane. Hence Theorem 2.1 contains no new result for the case $p = 2$.

(iii) For $p > 1$,

$$\tilde{a} > 4(p-1) \tag{2.2}$$

by a numerical simulation given in Fig. 2.

We first consider the bifurcation curve of positive solutions of the p -Laplacian Dirichlet problem

$$\begin{cases} (\varphi_p(u'(x)))' + \lambda f(u(x)) = 0, & -1 < x < 1, \\ u(-1) = u(1) = 0, \end{cases} \tag{2.3}$$

where $f \in C[0, \infty) \cap C^2(0, \infty)$, and $\lambda > 0$ is a bifurcation parameter. We define

$$F(u) = \int_0^u f(t) dt \quad \text{and} \quad \theta(u) = pF(u) - uf(u), \tag{2.4}$$

and assume that function f satisfies the following hypotheses (H1)–(H3):

(H1) $f(u) > 0$ for $u \in [0, \infty)$, and $m_\infty \equiv \lim_{u \rightarrow \infty} f(u)/u^{p-1} = 0$.

(H2) There exist numbers $0 < C_1 < D_1 < \gamma < C_2 < D_2 < \infty$ such that

$$\theta(D_1) = \theta(D_2) = \theta'(C_1) = \theta'(C_2) = 0, \tag{2.5}$$

and either

$$\theta''(u) = (p-2)f'(u) - uf''(u) \begin{cases} < 0 & \text{on } (0, \gamma), \\ = 0 & \text{for } u = \gamma, \\ > 0 & \text{on } (\gamma, \infty), \end{cases} \tag{2.6}$$

or there exist a number $\tilde{\gamma} \in (0, C_1)$ such that

$$\theta''(u) = (p-2)f'(u) - uf''(u) \begin{cases} > 0 & \text{on } (0, \tilde{\gamma}), \\ = 0 & \text{for } u = \tilde{\gamma}, \\ < 0 & \text{on } (\tilde{\gamma}, \gamma), \\ = 0 & \text{for } u = \gamma, \\ > 0 & \text{on } (\gamma, \infty), \end{cases} \quad (2.7)$$

and $\theta(\tilde{\gamma}) - \tilde{\gamma}\theta'(\tilde{\gamma}) - \theta(C_2) \geq 0$.

(H3) $uf'(u)/f(u) \geq -1/(p+1)$ on $(0, C_1)$ and $uf'(u)/f(u)$ is increasing on (C_1, D_1) .

The time map formula which we apply to study p -Laplacian problem (2.3) takes the form as follows:

$$T(\alpha) \equiv \left(\frac{p-1}{p}\right)^{1/p} \int_0^\alpha \frac{1}{[F(\alpha) - F(u)]^{1/p}} du = \lambda^{1/p} \quad \text{for } 0 < \alpha < \infty, \quad (2.8)$$

see, e.g., [3, Lemmas 2.1 and 2.2] and [10, Lemma 2.4] for the derivation of the time map formula $T(\alpha)$ for problem (2.3). So positive solutions u of (2.3) correspond to $\|u\|_\infty = \alpha$ and $T(\alpha) = \lambda^{1/p}$. Thus to study the number of positive solutions of (2.3) is equivalent to study the shape of the time map $T(\alpha)$ on $(0, \infty)$.

To prove Theorem 2.1, we need the next Lemmas 2.2–2.4 whose proofs are given in the next section.

Lemma 2.2. Let $p > 1$. Consider (2.3). Suppose $f \in C[0, \infty) \cap C^2(0, \infty)$ satisfies (H1)–(H3), then

$$\lim_{\alpha \rightarrow 0^+} T(\alpha) = 0, \quad \lim_{\alpha \rightarrow \infty} T(\alpha) = \infty, \quad (2.9)$$

and $T(\alpha)$ has exactly two positive critical points, $\alpha^* < \alpha_*$, on $(0, \infty)$, such that $T(\alpha^*)$ is a local maximum on $(0, \infty)$ and $T(\alpha_*)$ is a local minimum on $(0, \infty)$. More precisely, there exist

$$0 < \lambda_* \equiv (T(\alpha_*))^p < \lambda^* \equiv (T(\alpha^*))^p < \infty$$

such that (2.3) has exactly three positive solutions for $\lambda_* < \lambda < \lambda^*$, exactly two positive solutions for $\lambda = \lambda_*$ and $\lambda = \lambda^*$, and a unique positive solution for $0 < \lambda < \lambda_*$ and $\lambda^* < \lambda < \infty$.

Lemma 2.3. Let $p > 1$. If

$$a \geq \tilde{a} = \begin{cases} \max\{4, (4p-2)(\ln p) + 0.7\} & \text{if } 1 < p < 2, \\ (4p-2)(\ln p) + \eta & \text{if } p \geq 2, \end{cases}$$

then

(i) $\theta(a) < 0$.

(ii) If $1 < p \leq 2$, then there exist numbers $0 < C_1 < D_1 < \gamma < C_2 < D_2 < \infty$ such that (2.5) and (2.6) hold. Moreover,

$$C_1 = \frac{a}{2(p-1)} [a - 2p + 2 - \sqrt{a^2 + 4a - 4pa}], \quad (2.10)$$

$$\gamma = \frac{a}{2p} [a - 2p + 2 + \sqrt{a^2 + 4a - 4pa + 4}], \quad (2.11)$$

$$C_2 = \frac{a}{2(p-1)} [a - 2p + 2 + \sqrt{a^2 + 4a - 4pa}]. \quad (2.12)$$

(iii) If $p \geq 2$, then there exist numbers $0 < \tilde{\gamma} < C_1 < D_1 < \gamma < C_2 < D_2 < \infty$ such that (2.5) and (2.7) hold. In addition,

$$\theta(\tilde{\gamma}) - \tilde{\gamma}\theta'(\tilde{\gamma}) - \theta(C_2) > 0. \quad (2.13)$$

Moreover, C_1 , γ , and C_2 are given in (2.10)–(2.12) respectively, and

$$\tilde{\gamma} = \frac{a}{2p} [a - 2p + 2 - \sqrt{a^2 + 4a - 4pa + 4}]. \quad (2.14)$$

Lemma 2.4. Let $p \geq 2$. If $a \geq \tilde{a} = (4p - 2)(\ln p) + \eta$, then

- (i) $\tilde{\gamma} < \frac{3}{20}a$,
- (ii) $C_1 < \frac{43}{100}a$,
- (iii) $\gamma > \frac{139}{100}a$.

We are now in a position to prove Theorem 2.1 by applying Lemma 2.2.

Proof of Theorem 2.1. We show that, for $p > 1$, $f(u) = \exp(\frac{au}{a+u})$ satisfies (H1)–(H3) if $a \geq \tilde{a}$. The proof is divided into two cases:

- (A) $1 < p < 2$, and
- (B) $p \geq 2$.

Proof of Case (A) $1 < p < 2$. First, for $1 < p < 2$, for $f(u) = \exp(\frac{au}{a+u})$, it is easy to see that $f \in C[0, \infty) \cap C^2(0, \infty)$ satisfies (H1). We then compute that, for $\theta(u) = pF(u) - uf(u)$ in (2.4),

$$\theta'(u) = (p-1)f(u) - uf'(u) = \left[p - 1 - \frac{a^2u}{(a+u)^2} \right] \exp\left(\frac{au}{a+u}\right) \quad (2.15)$$

and

$$\theta''(u) = (p-2)f'(u) - uf''(u) = \left[\frac{pu^2 + (2pa - a^2 - 2a)u - a^2(2-p)}{(a+u)^4} \right] a^2 \exp\left(\frac{au}{a+u}\right). \quad (2.16)$$

So, it is easy to see that $\theta(0) = 0$, $\theta'(0) = p - 1 > 0$. Also $\lim_{u \rightarrow \infty} \theta(u) = \infty$ since $\lim_{u \rightarrow \infty} \theta'(u) = (p-1)\exp(a) > 0$. For

$$1 < p < 2 \quad \text{and} \quad a \geq \tilde{a} = \max\{4, (4p-2)(\ln p) + 0.7\},$$

by Lemma 2.3(ii), there exist numbers $0 < C_1 < D_1 < \gamma < C_2 < D_2 < \infty$, where C_1 , γ , and C_2 are given in (2.10)–(2.12) respectively, such that (2.5) and (2.6) hold. So f satisfies (H2).

We finally prove that f satisfies (H3). It is clear that $uf'(u) = \frac{a^2u}{(a+u)^2} \exp(\frac{au}{a+u}) > 0$ on $(0, \infty)$. So $uf'(u)/f(u) > 0 > -1/(p+1)$ on $(0, C_1)$. In addition, we compute that

$$\left(\frac{uf'(u)}{f(u)} \right)' = \frac{a^2(a-u)}{(a+u)^3} > 0 \quad \text{on } (0, a) \supset (C_1, D_1)$$

since $(0 < C_1 <) D_1 < a$ by applying Lemma 2.3(i) and (ii). Hence f satisfies (H3) if $a \geq \tilde{a}$.

We summarize above results and we conclude that, for $1 < p < 2$, $f(u) = \exp(\frac{au}{a+u})$ satisfies (H1)–(H3) if $a \geq \tilde{a}$. \square

Proof of Case (B) $p \geq 2$. Parts of the proof of Case (B) $p \geq 2$ are similar to, or the same as, those of Case (A) $1 < p < 2$.

First, for $p \geq 2$, for $f(u) = \exp(\frac{au}{a+u})$, it is easy to see that $f \in C[0, \infty) \cap C^2(0, \infty)$ satisfies (H1). We then compute that, for $\theta(u) = pF(u) - uf(u)$ in (2.4), we obtain (2.15) and (2.16) for $\theta'(u)$ and $\theta''(u)$, respectively. So it is easy to see that $\theta(0) = 0$, $\theta'(0) = p - 1 > 0$. Also $\lim_{u \rightarrow \infty} \theta(u) = \infty$ since $\lim_{u \rightarrow \infty} \theta'(u) = (p-1)\exp(a) > 0$. For

$$p \geq 2 \quad \text{and} \quad a \geq \tilde{a} = (4p-2)(\ln p) + \eta,$$

by Lemma 2.3(iii), there exist numbers $0 < \tilde{\gamma} < C_1 < D_1 < \gamma < C_2 < D_2 < \infty$, where $\tilde{\gamma}$, C_1 , γ , and C_2 are given in (2.14), (2.10)–(2.12) respectively, such that (2.5) and (2.7) hold. In addition, $\theta(\tilde{\gamma}) - \tilde{\gamma}\theta'(\tilde{\gamma}) - \theta(C_2) > 0$. So f satisfies (H2).

Finally, applying Lemma 2.3(i) and (iii), we can prove that f satisfies (H3) for $a \geq \tilde{a}$ by using the same argument used in the proof of Case (A) for $1 < p < 2$. We conclude that, for $p \geq 2$, $f(u) = \exp(\frac{au}{a+u})$ satisfies (H1)–(H3) if $a \geq \tilde{a}$. \square

By above, it follows from Lemma 2.2 that the bifurcation curve \bar{S} is S-shaped in the $(\lambda, \|u\|_\infty)$ -plane. Moreover, there exist $0 < \lambda_* < \lambda^* < \infty$ such that (1.1) has exactly three positive solutions for $\lambda_* < \lambda < \lambda^*$, exactly two positive solutions for $\lambda = \lambda_*$ and $\lambda = \lambda^*$, and a unique positive solution for $0 < \lambda < \lambda_*$ and $\lambda^* < \lambda < \infty$.

The proof of Theorem 2.1 is complete. \square

3. Proofs of Lemmas 2.2–2.4

Proof of Lemma 2.2. Eq. (2.9) follows by a slight generalization of [9, Theorems 2.6 and 2.9]. By (2.8) for $T(\alpha)$, we compute that

$$T'(\alpha) = \left(\frac{p-1}{p^{p+1}} \right)^{1/p} \frac{1}{\alpha} \int_0^\alpha \frac{\Delta\theta}{(\Delta F)^{(p+1)/p}} du \quad (3.1)$$

and

$$T''(\alpha) = \left(\frac{p-1}{p^{p+1}} \right)^{1/p} \frac{1}{\alpha^2} \int_0^\alpha \frac{-\frac{p+1}{p}(\Delta\theta)(\Delta\tilde{f}) + \Delta F(\Delta\tilde{\theta}')}{(\Delta F)^{(2p+1)/p}} du, \quad (3.2)$$

where $\Delta F = F(\alpha) - F(u)$, $\Delta\theta = \theta(\alpha) - \theta(u)$, $\Delta\tilde{f} = \alpha f(\alpha) - u f(u)$, and $\Delta\tilde{\theta}' = \alpha\theta'(\alpha) - u\theta'(u)$. By (3.1) and (3.2), we obtain that

$$\begin{aligned} T''(\alpha) + \frac{p}{\alpha} T'(\alpha) &= \left(\frac{p-1}{p^{p+1}} \right)^{1/p} \frac{1}{\alpha^2} \int_0^\alpha \frac{(\Delta F)(\Delta\phi) + \frac{p+1}{p}(\Delta\theta)^2}{(\Delta F)^{(2p+1)/p}} du \\ &\geq \left(\frac{p-1}{p^{p+1}} \right)^{1/p} \frac{1}{\alpha^2} \int_0^\alpha \frac{\Delta\phi}{(\Delta F)^{(p+1)/p}} du, \end{aligned} \quad (3.3)$$

where $\phi(u) = u\theta'(u) - \theta(u)$ and $\Delta\phi = \phi(\alpha) - \phi(u)$.

We know that $\theta(0) = 0$, $\theta'(0) = (p-1)f(0) > 0$. By (H1) and (H2), we obtain $\lim_{u \rightarrow \infty} \theta(u) = \infty$. In addition, there exists a positive number $D_3 \in (D_2, \infty)$ such that

$$\begin{cases} \theta(u) > 0 & \text{on } (0, D_1), \quad \theta(u) < 0 & \text{on } (D_1, D_2), \\ 0 < \theta(u) < \theta(C_1) & \text{on } (D_2, D_3), \quad \theta(u) \geq \theta(C_1) & \text{on } [D_3, \infty), \\ \theta(D_1) = \theta(D_2) = 0 \end{cases} \quad (3.4)$$

and

$$\begin{cases} \theta'(u) > 0 & \text{on } (0, C_1), \quad \theta'(u) < 0 & \text{on } (C_1, C_2), \quad \theta'(u) > 0 & \text{on } (C_2, \infty), \\ \theta'(C_1) = \theta'(C_2) = 0. \end{cases} \quad (3.5)$$

By (3.1), (3.4) and (3.5), we obtain that

$$T'(\alpha) > 0 \quad \text{for } \alpha \in (0, C_1], \quad T'(\alpha) < 0 \quad \text{for } \alpha \in [D_1, C_2] \quad \text{and} \quad T'(\alpha) > 0 \quad \text{for } \alpha \in [D_3, \infty).$$

Hence $T(\alpha)$ has at least two critical points, a local maximum on (C_1, D_1) and a local minimum on (C_2, D_3) . We then prove that $T(\alpha)$ has exactly one critical point, a local maximum on (C_1, D_1) , and $T(\alpha)$ has exactly one critical point, a local minimum on (C_2, D_3) , respectively.

First, we are able to prove that $T(\alpha)$ has exactly one critical point at some α^* , a local maximum, on (C_1, D_1) . We show this by showing that

$$T''(\alpha) + \frac{M_1}{p\alpha} T'(\alpha) < 0 \quad \text{for } \alpha \in (C_1, D_1)$$

for some positive function

$$M_1 = \max_{0 \leq u \leq \alpha} \frac{\alpha f(\alpha) - u f(u)}{F(\alpha) - F(u)}.$$

This proof is similar to that of [14, Theorem 1.1], and consequently, we omit it.

Next, we prove that $T(\alpha)$ has exactly one critical point, a local minimum, on (C_2, D_3) . If there exist numbers $0 < C_1 < \gamma < \infty$ such that (2.6) holds, since $\phi'(u) = u\theta''(u) = u[(p-2)f'(u) - uf''(u)]$, we obtain that

$$\phi'(u) < 0 \quad \text{on } (0, \gamma), \quad \phi'(\gamma) = 0, \quad \phi'(u) > 0 \quad \text{on } (\gamma, \infty).$$

In addition, it is easy to compute that $\phi(0) = 0$ and $\phi(C_2) = -\theta(C_2) > 0$. So we obtain that $\phi(u)$ is strictly increasing on (C_2, D_3) and $\phi(u) < \phi(C_2)$ for $u \in (0, C_2)$. The above imply that $\phi(u) < \phi(\alpha)$ for $\alpha \in (C_2, D_3)$, $u \in (0, \alpha)$. By (3.3), we obtain $T''(\alpha) + (p/\alpha)T'(\alpha) > 0$ for $\alpha \in (C_2, D_3)$. That is, if $\alpha_* \in (C_2, D_3)$ is a critical point of $T(\alpha)$, then $T(\alpha_*)$ must be a local minimum. Thus $T(\alpha)$ has exactly one critical point at α_* , a local minimum, on (C_2, D_3) .

If there exist numbers $0 < \tilde{\gamma} < C_1 < \gamma < \infty$ such that (2.7) holds, since $\phi'(u) = u\theta''(u) = u[(p-2)f'(u) - uf''(u)]$, similarly, we obtain that

$$\begin{cases} \phi'(u) > 0 & \text{on } (0, \tilde{\gamma}), & \phi'(u) < 0 & \text{on } (\tilde{\gamma}, \gamma), & \phi'(u) > 0 & \text{on } (\gamma, \infty), \\ \phi'(\tilde{\gamma}) = 0, & \phi'(\gamma) = 0. \end{cases}$$

In addition, it is easy to compute that $\phi(0) = 0$, $\phi(C_1) = -\theta(C_1) < 0$ and $\phi(C_2) = -\theta(C_2) > 0$. So we obtain that $\phi(u)$ is strictly increasing on (C_2, D_3) and $\phi(u) \leq \phi(C_2)$ for $u \in (0, C_2)$ by the assumption in (H2) that $\theta(\tilde{\gamma}) - \tilde{\gamma}\theta'(\tilde{\gamma}) - \theta(C_2) \geq 0$. By (3.3), we obtain $T''(\alpha) + (p/\alpha)T'(\alpha) > 0$ for $\alpha \in (C_2, D_3)$. That is, if $\alpha_* \in (C_2, D_3)$ is a critical point of $T(\alpha)$, then $T(\alpha_*)$ must be a local minimum. Thus $T(\alpha)$ has exactly one critical point at α_* , a local minimum, on (C_2, D_3) .

We summarize above results and we obtain that $T(\alpha)$ has exactly two positive critical points, $\alpha^* < \alpha_*$, on $(0, \infty)$, such that $T(\alpha^*)$ is a local maximum on $(0, \infty)$ and $T(\alpha_*)$ is a local minimum on $(0, \infty)$.

So there exist

$$0 < \lambda_* \equiv (T(\alpha_*))^p < \lambda^* \equiv (T(\alpha^*))^p < \infty$$

such that (2.3) has exactly three positive solutions for $\lambda_* < \lambda < \lambda^*$, exactly two positive solutions for $\lambda = \lambda_*$ and $\lambda = \lambda^*$, and a unique positive solution for $0 < \lambda < \lambda_*$ and $\lambda^* < \lambda < \infty$.

The proof of Lemma 2.2 is complete. \square

Proof of Lemma 2.3. First, we know that $\theta(0) = 0$, $\theta'(0) = p - 1 > 0$, and $\lim_{u \rightarrow \infty} \theta(u) = \infty$.

(i) We prove $\theta(a) < 0$. The proof is divided into two cases:

(A) $1 < p < 2$, and

(B) $p \geq 2$.

Proof of Case (A) $1 < p < 2$. Let $a \geq \tilde{a} = \max\{4, (4p-2)(\ln p) + 0.7\}$.

We first consider the graph of the function

$$g(u) \equiv pf(u) = p \exp\left(\frac{au}{a+u}\right),$$

on $(0, a)$. It is easy to see that $g(u)$ satisfies

$$g(u) > 0 \quad \text{on } (0, a), \tag{3.6}$$

$$g'(u) = \frac{pa^2}{(a+u)^2} \exp\left(\frac{au}{a+u}\right) > 0 \quad \text{on } (0, a), \tag{3.7}$$

and

$$g''(u) = \frac{pa^2[a(a-2) - 2u]}{(a+u)^4} \exp\left(\frac{au}{a+u}\right) > 0 \quad \text{on } (0, a) \tag{3.8}$$

for $a \geq \tilde{a} \geq 4$.

Secondly, define $a^* \in (0, a)$ by

$$a^* = \frac{a(a-2\ln p)}{a+2\ln p} = a - \frac{4a\ln p}{a+2\ln p},$$




Fig. 3. Graph of function $g(u) \equiv pf(u) = p \exp\left(\frac{au}{a+u}\right)$ on $(0, a)$ for $a \geq \tilde{a}$ and $1 < p < 2$. Note that $g''(u) > 0$ on $(0, a)$.

which satisfies

$$g(a^*) = pf(a^*) = p \exp\left(\frac{aa^*}{a+a^*}\right) = \exp(a/2).$$

We then compute that

$$\begin{aligned} \theta(a) &= pF(a) - af(a) \\ &= \int_0^a p \exp\left(\frac{at}{a+t}\right) dt - a \exp(a/2) \\ &= \int_{a^*}^a \left[p \exp\left(\frac{at}{a+t}\right) - \exp(a/2) \right] dt + \int_0^{a^*} \left[p \exp\left(\frac{at}{a+t}\right) - \exp(a/2) \right] dt \\ &= \int_{a^*}^a \left[p \exp\left(\frac{at}{a+t}\right) - \exp(a/2) \right] dt - \int_0^{a^*} \left[\exp(a/2) - p \exp\left(\frac{at}{a+t}\right) \right] dt \\ &< \frac{1}{2} \left\{ \frac{4a \ln p}{a+2 \ln p} (p-1) \exp(a/2) - [\exp(a/2) - p] \frac{a(a-2 \ln p)}{a+2 \ln p} \right\} \end{aligned}$$

since

$$\int_{a^*}^a \left[p \exp\left(\frac{at}{a+t}\right) - \exp(a/2) \right] dt < \frac{1}{2} (p-1) \exp(a/2) (a-a^*) = \frac{1}{2} \left\{ \frac{4a \ln p}{a+2 \ln p} (p-1) \exp(a/2) \right\}$$

and

$$-\int_0^{a^*} \left[\exp(a/2) - p \exp\left(\frac{at}{a+t}\right) \right] dt < -\frac{1}{2} [\exp(a/2) - p] a^* = -\frac{1}{2} [\exp(a/2) - p] \frac{a(a-2 \ln p)}{a+2 \ln p}$$

by (3.6)–(3.8), see Fig. 3. So we have

$$\begin{aligned} \theta(a) &< \frac{1}{2} \left\{ \frac{4a \ln p}{a+2 \ln p} (p-1) \exp(a/2) - [\exp(a/2) - p] \frac{a(a-2 \ln p)}{a+2 \ln p} \right\} \\ &= -\frac{a}{2(a+2 \ln p)} \{ (a-2 \ln p) [\exp(a/2) - p] - (4p-4)(\ln p) \exp(a/2) \} \end{aligned}$$




Fig. 4. Graph of function $h(p, 4)$ for $1 < p < 2$. $h(\tilde{p}, 4) = 0$ where $\tilde{p} \approx 1.847$.

$$\begin{aligned}
 &= -\frac{a}{2(a+2\ln p)} \{ [a - (4p-2)\ln p] \exp(a/2) - p(a-2\ln p) \} \\
 &\equiv -\frac{a}{2(a+2\ln p)} h(p, a)
 \end{aligned} \tag{3.9}$$

where

$$h(p, a) = [a - (4p-2)\ln p] \exp(a/2) - p(a-2\ln p).$$

So

$$\theta(a) < 0 \quad \text{if } h(p, a) > 0.$$

For fixed $1 < p < 2$ and for $a \geq \tilde{a} \equiv \max\{4, (4p-2)(\ln p) + 0.7\}$, we compute that

$$\begin{aligned}
 \frac{\partial}{\partial a} h(p, a) &= \left[\frac{a}{2} - (2p-1)(\ln p) + 1 \right] \exp(a/2) - p \\
 &\geq [(2p-1)(\ln p) + 0.35 - (2p-1)(\ln p) + 1]p - p
 \end{aligned}$$

since $a \geq \tilde{a} \geq (4p-2)(\ln p) + 0.7$ and $\exp(a/2) \geq \exp((2p-1)\ln p) > \exp(\ln p) = p$. So

$$\frac{\partial}{\partial a} h(p, a) \geq [(2p-1)(\ln p) + 0.35 - (2p-1)(\ln p) + 1]p - p = 0.35p > 0.$$

In addition,

(I) For $1 < p \leq p^* (\approx 1.846)$, $\tilde{a}(p) = 4$. Letting $a = \tilde{a} = 4$, we obtain

$$h(p, 4) = [4 - (4p-2)\ln p] \exp(2) - p(4-2\ln p) > 0,$$


see Fig. 4. (Note that the first positive zero for $h(p, 4)$ is $\tilde{p} \approx 1.847$.)

(II) For $(1.846 \approx) p^* < p < 2$, $\tilde{a}(p) = (4p-2)(\ln p) + 0.7$. Letting $a = \tilde{a}(p) = (4p-2)(\ln p) + 0.7$, we compute that

$$\begin{aligned}
 h(p, \tilde{a}(p)) &= [a - (4p-2)\ln p] \exp(a/2) - p(a-2\ln p) \\
 &= [(4p-2)(\ln p) + 0.7 - (4p-2)(\ln p)] \exp[(2p-1)(\ln p) + 0.35] \\
 &\quad - p[(4p-2)(\ln p) + 0.7 - 2\ln p] \\
 &= 0.7p^{2p-1} \exp(0.35) - 4(p-1)p(\ln p) - 0.7p \\
 &> 0,
 \end{aligned}$$

see Fig. 5. (Note that $h(\tilde{p}, \tilde{a}(\tilde{p})) = 0$ for $\tilde{p} \approx 1.828$ and $h(p, \tilde{a}(p)) > 0$ for $p > p^* \approx 1.846$.)

So by above, we obtain that, if $1 < p < 2$ and $a \geq \tilde{a}$, then $h(p, a) > 0$, and hence $\theta(a) < 0$. So Lemma 2.3(i) holds for Case (A) $1 < p < 2$. \square

Fig. 5. Graph of function $h(p, \tilde{a}(p))$ for $(1.846 \approx) p^* < p < 2$.

Proof of Case (B) $p \geq 2$. Let $a \geq \tilde{a} = (4p - 2)(\ln p) + \eta$.

Similarly as before, we first consider the graph of the function

$$g(u) \equiv pf(u) = p \exp\left(\frac{au}{a+u}\right)$$

on $(0, a)$. It is easy to see that $g(u)$ satisfies (3.6)–(3.8) for $a \geq \tilde{a} > 4$ by (2.2).

Secondly, define $a^* \in (0, a)$ by

$$a^* = \frac{a(a - 2 \ln p)}{a + 2 \ln p} = a - \frac{4a \ln p}{a + 2 \ln p},$$

which satisfies

$$g(a^*) = pf(a^*) = p \exp\left(\frac{aa^*}{a+a^*}\right) = \exp(a/2).$$

Then, by the same arguments as we did in the proof of Lemma 2.3(i), we obtain (3.9)

$$\theta(a) < -\frac{a}{2(a + 2 \ln p)} h(p, a)$$

where

$$h(p, a) = [a - (4p - 2) \ln p] \exp(a/2) - p(a - 2 \ln p).$$

So

$$\theta(a) < 0 \quad \text{if } h(p, a) > 0.$$

For fixed $p \geq 2$ and for $a \geq \tilde{a} = (4p - 2)(\ln p) + \eta$, we compute that

$$\begin{aligned} \frac{\partial}{\partial a} h(p, a) &= \left[\frac{a}{2} - (2p - 1)(\ln p) + 1 \right] \exp(a/2) - p \\ &\geq \left[(2p - 1)(\ln p) + \frac{\eta}{2} - (2p - 1)(\ln p) + 1 \right] p - p \end{aligned}$$

since $a \geq \tilde{a} = (4p - 2)(\ln p) + \eta$ and $\exp(a/2) \geq \exp((2p - 1) \ln p) > \exp(\ln p) = p$. So

$$\frac{\partial}{\partial a} h(p, a) \geq \left[(2p - 1)(\ln p) + \frac{\eta}{2} - (2p - 1)(\ln p) + 1 \right] p - p = \frac{\eta p}{2} > 0. \quad (3.10)$$

In addition, for $p \geq 2$, letting $a = \tilde{a} = (4p - 2)(\ln p) + \eta$, we obtain

$$\begin{aligned} h(p, (4p - 2)(\ln p) + \eta) &= \eta \exp((2p - 1)(\ln p) + \eta/2) - p[(4p - 2)(\ln p) + \eta - 2 \ln p] \\ &= \eta \exp((2p - 1)(\ln p) + \eta/2) - p[(4p - 4)(\ln p) + \eta] \\ &= p[\eta \exp(\eta/2) p^{2p-2} - (4p - 4)(\ln p) + \eta] \\ &\equiv pR(p), \end{aligned}$$

where

$$R(p) = \eta \exp(\eta/2) p^{2p-2} - (4p - 4)(\ln p) + \eta.$$

We compute that

$$\begin{aligned} R'(p) &= 2\eta \exp(\eta/2) p^{2p-2} \left[\frac{p-1}{p} + \ln p \right] - \frac{4p-4}{p} - 4(\ln p) \\ &= 2[\eta \exp(\eta/2) p^{2p-2} - 2] \left[\frac{p-1}{p} + \ln p \right] \\ &> 0 \quad \text{on } [2, \infty) \end{aligned}$$

since for $p \geq 2$,

- (I) $\eta \exp(\eta/2) p^{2p-2} - 2 > 0$ since $\eta \exp(\eta/2) p^{2p-2} - 2$ is strictly increasing in $p > 2$ and $\eta \exp(\eta/2) 2^2 - 2 \approx 1.394 > 0$, and
 (II) $\frac{p-1}{p} + \ln p > 0$.

In addition,

$$R(2) = 4\eta \exp(\eta/2) - 4(\ln 2) + \eta \approx 1.244 > 0.$$

So by above, we conclude that, for $p \geq 2$, $R(p) > 0$ and hence $h(p, (4p - 2)(\ln p) + \eta) > 0$. Thus, by (3.10), for $a \geq \tilde{a} = (4p - 2)(\ln p) + \eta$, we have $h(p, a) > 0$ and hence $\theta(a) < 0$. So Lemma 2.3(i) holds for Case (B) $p \geq 2$. \square

By above, Lemma 2.3(i) holds for $p > 1$.

(ii) For $1 < p < 2$, by (2.15), $\theta(u)$ has exactly two positive critical points at $C_1 < C_2$, and we compute that

$$\begin{aligned} C_1 &= \frac{a}{2(p-1)} [a - 2p + 2 - \sqrt{a^2 + 4a - 4pa}], \\ C_2 &= \frac{a}{2(p-1)} [a - 2p + 2 + \sqrt{a^2 + 4a - 4pa}]. \end{aligned}$$

Also, by (2.16), it is easy to see that $\theta(u)$ has exactly one positive inflection point at γ , and we compute that

$$\gamma = \frac{a}{2p} [a - 2p + 2 + \sqrt{a^2 + 4a - 4pa + 4}] > \frac{a}{2p} [(4(p-1) - 2p) + 2 + 2]$$

since $a^2 + 4a - 4pa + 4 = a[a - 4(p-1)] + 4 > 4$ and $a > 4(p-1)$ by (2.2). So

$$\gamma > \frac{a}{2p} [(4(p-1) - 2p) + 2 + 2] = a.$$

Since $\theta(a) < 0$ and by above analysis for $\theta(u)$, we know that $\theta(u)$ has exactly two positive zeros at $D_1 < D_2$ such that $0 < C_1 < D_1 < \gamma < C_2 < D_2 < \infty$. So Lemma 2.3(ii) holds.

(iii) For $p \geq 2$, similarly, by (2.15), $\theta(u)$ has exactly two positive critical points at $C_1 < C_2$, and we compute that

$$\begin{aligned} C_1 &= \frac{a}{2(p-1)} [a - 2p + 2 - \sqrt{a^2 + 4a - 4pa}], \\ C_2 &= \frac{a}{2(p-1)} [a - 2p + 2 + \sqrt{a^2 + 4a - 4pa}]. \end{aligned}$$




Fig. 6. Graph of function $\theta'(u)$ on $(0, C_2)$. Note that $-C_1\theta'(\tilde{\gamma}) - (\gamma - a)\theta'(a) > 0$ for $a \geq \tilde{a}$ and $p \geq 2$.

Also, by (2.16), it is easy to see that $\theta(u)$ has exactly two positive inflection points at $\tilde{\gamma} < \gamma$, and we compute that

$$\begin{aligned}\tilde{\gamma} &= \frac{a}{2p} \left[a - 2p + 2 - \sqrt{a^2 + 4a - 4pa + 4} \right], \\ \gamma &= \frac{a}{2p} \left[a - 2p + 2 + \sqrt{a^2 + 4a - 4pa + 4} \right] > \frac{a}{2p} \left[(4(p-1) - 2p) + 2 + 2 \right]\end{aligned}$$

since $a^2 + 4a - 4pa + 4 = a[a - 4(p-1)] + 4 > 4$ and $a > 4(p-1)$ by (2.2). So

$$\gamma > \frac{a}{2p} \left[(4(p-1) - 2p) + 2 + 2 \right] = a.$$

Since $\theta(a) < 0$ and by above analysis for $\theta(u)$, we know that $\theta(u)$ has exactly two positive zeros at $D_1 < D_2$ such that $0 < \tilde{\gamma} < C_1 < D_1 < \gamma < C_2 < D_2 < \infty$.

Finally, we prove (2.13) $\theta(\tilde{\gamma}) - \tilde{\gamma}\theta'(\tilde{\gamma}) - \theta(C_2) > 0$ by applying Lemma 2.4, see Fig. 6.

Let $p \geq 2$ and $a \geq \tilde{a} = (4p-2)(\ln p) + \eta$. Suppose Lemma 2.4 holds. We then observe the graph of $\theta'(u)$ on $(0, C_2)$ as in Fig. 6 and we compute that

$$\begin{aligned}\theta(\tilde{\gamma}) - \tilde{\gamma}\theta'(\tilde{\gamma}) - \theta(C_2) &= \int_0^{\tilde{\gamma}} \theta'(u) du - \tilde{\gamma}\theta'(\tilde{\gamma}) - \int_0^{C_2} \theta'(u) du \\ &= \int_0^{\tilde{\gamma}} \theta'(u) du - \tilde{\gamma}\theta'(\tilde{\gamma}) - \int_0^{C_1} \theta'(u) du - \int_{C_1}^{C_2} \theta'(u) du \\ &= -\tilde{\gamma}\theta'(\tilde{\gamma}) - \int_{\tilde{\gamma}}^{C_1} \theta'(u) du - \int_{C_1}^{C_2} \theta'(u) du \\ &> -C_1\theta'(\tilde{\gamma}) - (\gamma - a)\theta'(a)\end{aligned}$$

since $-\tilde{\gamma}\theta'(\tilde{\gamma}) - \int_{\tilde{\gamma}}^{C_1} \theta'(u) du > -C_1\theta'(\tilde{\gamma})$ and $-\int_{C_1}^{C_2} \theta'(u) du > -(\gamma - a)\theta'(a)$. So we have

$$\begin{aligned}\theta(\tilde{\gamma}) - \tilde{\gamma}\theta'(\tilde{\gamma}) - \theta(C_2) &> -C_1\theta'(\tilde{\gamma}) - (\gamma - a)\theta'(a) \\ &= -C_1 \left[(p-1)f(\tilde{\gamma}) - \tilde{\gamma}f'(\tilde{\gamma}) \right] - (\gamma - a)\theta'(a) \\ &> -C_1(p-1)f(\tilde{\gamma}) - (\gamma - a)\theta'(a) \quad (\text{since } \tilde{\gamma}f'(\tilde{\gamma}) > 0) \\ &> -\frac{43}{100}a(p-1)f\left(\frac{3}{20}a\right) - \frac{39}{100}a\theta'(a)\end{aligned}$$




Fig. 7. Graph of function $K(p, \tilde{a}(p))$ for $2 \leq p \leq 2.5$. $K(2, \tilde{a}(2)) \approx 6.183 > 0$.

by Lemma 2.4(i)–(iii) and since f is increasing on $(0, a)$. So

$$\begin{aligned} \theta(\tilde{\gamma}) - \tilde{\gamma}\theta'(\tilde{\gamma}) - \theta(C_2) &> -\frac{43}{100}a(p-1)f\left(\frac{3}{20}a\right) - \frac{39}{100}a\theta'(a) \\ &= -\frac{43}{100}a(p-1)\exp(3a/23) + \frac{39}{100}a\left(\frac{a}{4} - p + 1\right)\exp(a/2) \\ &\equiv \frac{1}{400}a\exp(3a/23)K(p, a), \end{aligned} \quad (3.11)$$

where

$$K(p, a) = 39(a - 4p + 4)[\exp(17a/46)] - 172(p - 1).$$

We easily compute that, for $a \geq \tilde{a} = (4p - 2)(\ln p) + \eta$,

$$\frac{\partial}{\partial a}K(p, a) = \frac{39}{46}(17a - 68p + 114)[\exp(17a/46)] > 0$$

since $17a - 68p + 114 \geq 17[(4p - 2)(\ln p) + \eta] - 68p + 114 > 0$ for $p \geq 2$, we omit the proof. In addition,

$$K(p, \tilde{a}(p)) = 39[(4p - 2)(\ln p) + \eta - 4p + 4][\exp(17((4p - 2)(\ln p) + \eta)/46)] - 172(p - 1) > 0$$

by some numerical simulation given in Fig. 7, which shows that $K(p, \tilde{a}(p))$ is a strictly increasing function of $p \geq 2$ and $K(2, \tilde{a}(2)) \approx 6.183 > 0$.

We summarize above results and we conclude that, for $p \geq 2$ and $a \geq \tilde{a} = (4p - 2)(\ln p) + \eta$, $K(p, a) > 0$, and hence $\theta(\tilde{\gamma}) - \tilde{\gamma}\theta'(\tilde{\gamma}) - \theta(C_2) > 0$ by (3.11). This completes the proof of Lemma 2.3(iii).

The proof of Lemma 2.3 is complete. \square

Proof of Lemma 2.4. Let $p \geq 2$.

(i) We take

$$R_1(p, a) \equiv \frac{\tilde{\gamma}}{a} = \frac{1}{2p}[a - 2p + 2 - \sqrt{a^2 + 4a - 4pa + 4}].$$

If $a \geq \tilde{a} = (4p - 2)(\ln p) + \eta$, then it is easy to compute that

$$\frac{\partial}{\partial a}R_1(p, a) = \frac{-a + 2p - 2 + \sqrt{a^2 + 4a - 4pa + 4}}{2p\sqrt{a^2 + 4a - 4pa + 4}} \begin{cases} < 0 & \text{if } p > 2, \\ = 0 & \text{if } p = 2 \end{cases}$$

since $a > 2p - 2$ by (2.2), and

$$(-a + 2p - 2)^2 - (a^2 + 4a - 4pa + 4) = 4p(p - 2) \begin{cases} > 0 & \text{if } p > 2, \\ = 0 & \text{if } p = 2. \end{cases}$$




Fig. 8. Graph of function $R_1(p, \tilde{a}(p))$ for $2 \leq p \leq 20$. $\max_{p \geq 2} R_1(p, \tilde{a}(p)) = R_1(p_1, \tilde{a}(p_1)) \approx 0.149$ for $p = p_1 \approx 3.838$.




Fig. 9. Graph of function $R_2(p, \tilde{a}(p))$ for $2 \leq p \leq 20$. $\max_{p \geq 2} R_2(p, \tilde{a}(p)) = R_2(2, \tilde{a}(2)) \approx 0.4243$.

Then, for all $p \geq 2$ and $a \geq \tilde{a}(p)$, we obtain that

$$R_1(p, a) \leq R_1(p, \tilde{a}(p)) \leq \max_{p \geq 2} R_1(p, \tilde{a}(p)) = R_1(p_1, \tilde{a}(p_1)) \approx 0.149 < 0.150 = \frac{3}{20}$$

for $p_1 \approx 3.838$, by some numerical simulation given in Fig. 8. So part (i) follows.

(ii) We take

$$R_2(p, a) \equiv \frac{C_1}{a} = \frac{1}{2(p-1)} [a - 2p + 2 - \sqrt{a^2 + 4a - 4pa}].$$

If $a \geq \tilde{a} = (4p - 2)(\ln p) + \eta$, then it is easy to compute that

$$\frac{\partial}{\partial a} R_2(p, a) = \frac{-a + 2p - 2 + \sqrt{a^2 + 4a - 4pa}}{2(p-1)\sqrt{a^2 + 4a - 4pa}} < 0$$

since $a > 2p - 2$ by (2.2), and

$$(-a + 2p - 2)^2 - (a^2 + 4a - 4pa) = 4(p-1)^2 > 0.$$

Then, for all $p \geq 2$ and $a \geq \tilde{a}(p)$, we obtain that

$$R_2(p, a) \leq R_2(p, \tilde{a}(p)) \leq \max_{p \geq 2} R_2(p, \tilde{a}(p)) = R_2(2, \tilde{a}(2)) \approx 0.424 < 0.430 = \frac{43}{100}$$

by some numerical simulation given in Fig. 9. (Numerical simulation shows that $R_2(p, \tilde{a}(p))$ is a strictly decreasing function of $p \geq 2$.) So part (ii) follows.




Fig. 10. Graph of function $R_3(p, \tilde{a}(p))$ for $2 \leq p \leq 20$. $\min_{p \geq 2} R_3(p, \tilde{a}(p)) = R_3(2, \tilde{a}(2)) \approx 1.390$.

(iii) We take

$$R_3(p, a) \equiv \frac{\gamma}{a} = \frac{1}{2p} [a - 2p + 2 + \sqrt{a^2 + 4a - 4pa + 4}].$$

If $a \geq \tilde{a} = (4p - 2)(\ln p) + \eta$, then it is easy to compute that

$$\frac{\partial}{\partial a} R_3(p, a) = \frac{a - 2p + 2 + \sqrt{a^2 + 4a - 4pa + 4}}{2p\sqrt{a^2 + 4a - 4pa + 4}} > 0 \quad \text{if } p \geq 2$$

since $a > 2p - 2$ by (2.2). Then, for all $p \geq 2$ and $a \geq \tilde{a}(p)$, we obtain that

$$R_3(p, a) \geq R_3(p, \tilde{a}(p)) \geq \min_{p \geq 2} R_3(p, \tilde{a}(p)) = R_3(2, \tilde{a}(2)) \approx 1.3904 > 1.390 = \frac{139}{100}$$

by some numerical simulation given in Fig. 10. (Numerical simulation shows that $R_3(p, \tilde{a}(p))$ is a strictly increasing function of $p \geq 2$.) So part (iii) follows.

The proof of Lemma 2.4 is complete. \square

Acknowledgments

The authors thank the referee for giving so many illuminating comments and suggestions on the original manuscript, upon which this paper is completely revised. The authors also thank Dr. Idris Addou for giving many valuable comments. Much of the computation in this paper has been checked using the symbolic manipulator *Mathematica 5.0*.

References

- [1] J. Bebernes, D. Eberly, *Mathematical Problems from Combustion Theory*, Springer-Verlag, New York, 1989.
- [2] K.J. Brown, M.M.A. Ibrahim, R. Shivaji, S-shaped bifurcation curves, *Nonlinear Anal.* 5 (1981) 475–486.
- [3] J.G. Cheng, Uniqueness results for the one-dimensional p -Laplacian, *J. Math. Anal. Appl.* 311 (2005) 381–388.
- [4] M.G. Crandall, P.H. Rabinowitz, Bifurcation, perturbation of simple eigenvalues and linearized stability, *Arch. Ration. Mech. Anal.* 52 (1973) 161–180.
- [5] Y. Du, Exact multiplicity and S-shaped bifurcation curve for some semilinear elliptic problems from combustion theory, *SIAM J. Math. Anal.* 32 (2000) 707–733.
- [6] Y. Du, Y. Lou, Proof of a conjecture for the perturbed Gelfand equation from combustion theory, *J. Differential Equations* 173 (2001) 213–230.
- [7] J. Jacobsen, K. Schmitt, The Liouville–Bratu–Gelfand problem for radial operators, *J. Differential Equations* 184 (2002) 283–298.
- [8] P. Korman, Y. Li, On the exactness of an S-shaped bifurcation curve, *Proc. Amer. Math. Soc.* 127 (1999) 1011–1020.
- [9] T. Laetsch, The number of solutions of a nonlinear two point boundary value problem, *Indiana Univ. Math. J.* 20 (1970) 1–13.
- [10] A. Lakmeche, A. Hammoudi, Multiple positive solutions of the one-dimensional p -Laplacian, *J. Math. Anal. Appl.* 317 (2006) 43–49.
- [11] M. Ramaswamy, R. Shivaji, Multiple positive solutions for classes of p -Laplacian equations, *Differential Integral Equations* 17 (2004) 1255–1261.

- [12] S.-H. Wang, On S -shaped bifurcation curves, *Nonlinear Anal.* 22 (1994) 1475–1485.
- [13] S.-H. Wang, Rigorous analysis and estimates of S -shaped bifurcation curves in a combustion problem with general Arrhenius reaction-rate laws, *Proc. R. Soc. Lond. Ser. A* 454 (1998) 1031–1048.
- [14] S.-H. Wang, T.-S. Yeh, Exact multiplicity and ordering properties of positive solutions of a p -Laplacian Dirichlet problem and their applications, *J. Math. Anal. Appl.* 287 (2003) 380–398.