



Available online at www.sciencedirect.com

SCIENCE @ DIRECT®

J. Math. Anal. Appl. 300 (2004) 343–350

Journal of
MATHEMATICAL
ANALYSIS AND
APPLICATIONS

www.elsevier.com/locate/jmaa

Hyers–Ulam–Rassias stability of Cauchy equation in the space of Schwartz distributions [☆]

Jaeyoung Chung

Department of Mathematics, Kunsan National University, Kunsan 573-701, South Korea

Received 8 March 2004

Submitted by T. Krisztin

Abstract

We reformulate and prove the Hyers–Ulam–Rassias stability of Cauchy equation in the space of Schwartz tempered distributions and Fourier hyperfunctions.

© 2004 Elsevier Inc. All rights reserved.

Keywords: Gauss transforms; Heat kernel; Distributions; Hyperfunctions; Heat polynomial; Stability; Cauchy equation

1. Introduction

Generalizing the stability theorem of D.H. Hyers [10,11] which was motivated by S.M. Ulam [17], Th.M. Rassias [15] and Z. Gajda [7] showed the following stability theorem for the Cauchy equation:

Theorem 1.1 [7,15]. *Let $f : E_1 \rightarrow E_2$ with E_1, E_2 Banach spaces be an approximately additive, that is, f satisfies*

$$\|f(x+y) - f(x) - f(y)\| \leq \epsilon(\|x\|^p + \|y\|^p), \quad p \neq 1, \quad (1.1)$$

[☆] This work was supported by Korea Research Foundation Grant (KRF-2003-002-C00012).
E-mail address: jychung@kunsan.ac.kr.

for all $x, y \in E_1$ ($x \neq 0$ and $y \neq 0$ if $p < 0$). Then there exists a unique mapping $g : E_1 \rightarrow E_2$ such that

$$g(x + y) - g(x) - g(y) = 0$$

and

$$\|f(x) - g(x)\| \leq \frac{2\epsilon}{|2^p - 2|} \|x\|^p \quad (1.2)$$

for all $x \in E_1$. Here the inequality (1.1) holds for $x, y \neq 0$ and (1.2) for $x \neq 0$ if $p < 0$.

The above stability theorem was firstly proved for the case $p < 1$ by Th.M. Rassias [15] and later it was proved for the case $p > 1$ by Z. Gajda [7].

In this paper, we reformulate and prove the above stability theorem in the spaces of generalized functions such as the space \mathcal{S}' of Schwartz tempered distributions and the space \mathcal{F}' of Fourier hyperfunctions for the case that p is even integer greater than 2. Note that the above inequality (1.1) makes no sense if f is a tempered distribution or Fourier hyperfunction. As in [1,2,5,6,13] making use of the pullbacks of generalized function we extend the inequality (1.1) to the spaces of tempered distributions and Fourier hyperfunctions as follows:

$$\|u \circ A - u \circ P_1 - u \circ P_2\| \leq \epsilon(\|x\|^{2p} + \|y\|^{2p}). \quad (1.3)$$

Here $u \circ A$, $u \circ P_1$ and $u \circ P_2$ are the pullbacks of u in \mathcal{S}' or \mathcal{F}' by A , P_1 and P_2 , respectively, where A , P_1 and P_2 are the functions $A(x, y) = x + y$, $P_1(x, y) = x$ and $P_2(x, y) = y$, $x, y \in \mathbb{R}^n$. Also the inequality $\|v\| \leq \epsilon(\|x\|^{2p} + \|y\|^{2p})$ in (1.3) means that $|\langle v, \varphi \rangle| \leq \epsilon \int (\|x\|^{2p} + \|y\|^{2p}) |\varphi(x, y)| dx dy$ for all test functions $\varphi \in \mathcal{S}$ (respectively \mathcal{F}).

For the pullback of tempered distributions we refer to [9, Chapters V–VI]. As a matter of fact, the pullbacks $u \circ A$, $u \circ P_1$, $u \circ P_2$ can be written in a transparent way as

$$\begin{aligned} \langle u \circ A, \varphi(x, y) \rangle &= \left\langle u, \int \varphi(x - y, y) dy \right\rangle, \\ \langle u \circ P_1, \varphi(x, y) \rangle &= \left\langle u, \int \varphi(x, y) dy \right\rangle, \\ \langle u \circ P_2, \varphi(x, y) \rangle &= \left\langle u, \int \varphi(x, y) dx \right\rangle \end{aligned}$$

for all test functions $\varphi \in \mathcal{S}(\mathbb{R}^{2n})$.

As a results, we prove that every solution of the inequality (1.3) can be approximated by a linear function in the sense that there exists a unique $a \in \mathbb{C}^n$ such that

$$\|u - a \cdot x\| \leq \frac{2\epsilon}{4^p - 2} \|x\|^{2p}.$$

2. Distributions and hyperfunctions

We first introduce briefly some spaces of generalized functions such as the space \mathcal{S}' of tempered distributions and the space \mathcal{F}' of Fourier hyperfunctions which is a natural

generalization of \mathcal{S}' . Here we use the multi-index notations for $x = (x_1, \dots, x_n) \in \mathbb{R}^n$, $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}_0^n$,

$$|\alpha| = \alpha_1 + \dots + \alpha_n, \quad \alpha! = \alpha_1! \dots \alpha_n!,$$

$$x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}, \quad \partial^\alpha = \partial_1^{\alpha_1} \dots \partial_n^{\alpha_n},$$

where \mathbb{N}_0 is the set of non-negative integers and $\partial_j = \partial/\partial x_j$.

Definition 2.1 [3,8,9,16]. We denote by \mathcal{S} or $\mathcal{S}(\mathbb{R}^n)$ the Schwartz space of all infinitely differentiable functions φ in \mathbb{R}^n such that

$$\|\varphi\|_{\alpha,\beta} = \sup_x |x^\alpha \partial^\beta \varphi(x)| < \infty \tag{2.1}$$

for all $\alpha, \beta \in \mathbb{N}_0^n$, equipped with the topology defined by the seminorms $\|\cdot\|_{\alpha,\beta}$. The elements of \mathcal{S} are called rapidly decreasing functions and the elements of the dual space \mathcal{S}' are called *tempered distributions*.

As a matter of fact, it is known in [3] that (2.1) is equivalent to

$$\sup_{x \in \mathbb{R}^n} |x^\alpha \varphi(x)| < \infty, \quad \sup_{\xi \in \mathbb{R}^n} |\xi^\beta \hat{\varphi}(\xi)| < \infty \tag{2.1'}$$

for all $\alpha, \beta \in \mathbb{N}_0^n$.

Imposing growth conditions on $\|\cdot\|_{\alpha,\beta}$ in (2.1) Sato and Kawai introduced the space \mathcal{F} of test functions for the Fourier hyperfunctions as follows:

Definition 2.2 [4,8,16]. We denote by \mathcal{F} or $\mathcal{F}(\mathbb{R}^n)$ the Sato space of all infinitely differentiable functions φ in \mathbb{R}^n such that

$$\|\varphi\|_{A,B} = \sup_{x,\alpha,\beta} \frac{|x^\alpha \partial^\beta \varphi(x)|}{A^{|\alpha|} B^{|\beta|} \alpha! \beta!} < \infty \tag{2.2}$$

for some positive constants A, B .

We say that $\varphi_j \rightarrow 0$ as $j \rightarrow \infty$ if $\|\varphi_j\|_{A,B} \rightarrow 0$ as $j \rightarrow \infty$ for some $A, B > 0$, and denote by \mathcal{F}' the strong dual of \mathcal{F} and call its elements *Fourier hyperfunctions*.

It is known in [4] that the inequality (2.2) is equivalent to

$$\sup_{x \in \mathbb{R}^n} |\varphi(x)| \exp k|x| < \infty, \quad \sup_{\xi \in \mathbb{R}^n} |\hat{\varphi}(\xi)| \exp h|\xi| < \infty \tag{2.2'}$$

for some $h, k > 0$.

It is easy to see the following topological inclusions:

$$\mathcal{F} \hookrightarrow \mathcal{S}, \quad \mathcal{S}' \hookrightarrow \mathcal{F}'.$$

From now on a *test function* means an element in the Schwartz space \mathcal{S} or the Sato space \mathcal{F} and a *generalized function* means a *tempered distribution* or a *Fourier hyperfunction*.

3. Main theorems

We employ the n -dimensional heat kernel, that is, the fundamental solution $E_t(x)$ of the heat operator $\partial_t - \Delta_x$ in $\mathbb{R}_x^n \times \mathbb{R}_t^+$ given by

$$E_t(x) = \begin{cases} (4\pi t)^{-n/2} \exp(-|x|^2/4t), & t > 0, \\ 0, & t \leq 0. \end{cases}$$

Here the semigroup property

$$(E_t * E_s)(x) = E_{t+s}(x) \quad (3.1)$$

of the heat kernel will be very useful later. Now let a tempered distribution u be given. Then its *Gauss transform*

$$Gu(x, t) = (u * E_t)(x) = \langle u_y, E_t(x - y) \rangle, \quad x \in \mathbb{R}^n, t > 0, \quad (3.2)$$

is a C^∞ -function in $\mathbb{R}^n \times (0, \infty)$. As a matter of fact we can represent tempered distributions via some solutions of the heat equation as follows:

Proposition 3.1 [14]. *Let $u \in \mathcal{S}'(\mathbb{R}^n)$. Then its Gauss transform $Gu(x, t)$ is a C^∞ -solution of heat equation satisfying:*

(i) *There exist positive constants C , M and N such that*

$$|Gu(x, t)| \leq Ct^{-M} (1 + |x|)^N \quad \text{in } \mathbb{R}^n \times (0, \delta); \quad (3.3)$$

(ii) *$Gu(x, t) \rightarrow u$ as $t \rightarrow 0^+$ in the sense that for every $\varphi \in \mathcal{S}$,*

$$\langle u, \varphi \rangle = \lim_{t \rightarrow 0^+} \int Gu(x, t) \varphi(x) dx.$$

Conversely, every C^∞ -solution $U(x, t)$ of heat equation satisfying the growth condition (3.3) can be uniquely expressed as $U(x, t) = Gu(x, t)$ for some $u \in \mathcal{S}'$.

Similarly we can represent Fourier hyperfunctions as initial values of solutions of heat equation as a special case of the results in [12]. In this case, the estimate (3.3) is replaced by the following: For every $\epsilon > 0$ there exists a positive constant C_ϵ such that

$$|Gu(x, t)| \leq C_\epsilon \exp(\epsilon(|x| + 1/t)) \quad \text{in } \mathbb{R}^n \times (0, \delta). \quad (3.3')$$

Definition 3.2. Let v be in \mathcal{S}' or \mathcal{F}' . Then we denote by $\|v\| \leq \psi$ if

$$|\langle v, \varphi \rangle| \leq \|\psi \varphi\|_{L^1} \quad (3.4)$$

for all test functions φ .

Now we prove main theorems.

Theorem 3.3. *Let u in \mathcal{S}' or \mathcal{F}' satisfy the inequality*

$$\|u \circ A - u \circ P_1 - u \circ P_2\| \leq \epsilon(x^{2\gamma} + y^{2\gamma}) \quad (3.5)$$

for some $\gamma \in \mathbb{N}_0^n$, $|\gamma| > 1$. Then there exists a unique $a \in \mathbb{C}^n$ such that

$$\|u - a \cdot x\| \leq \frac{2\epsilon}{4^{|\gamma|} - 2} x^{2\gamma}. \tag{3.6}$$

Proof. Convolving in each side of (3.5) the tensor product $E_t(x)E_s(y)$ of n -dimensional heat kernels as a function of x, y the left-hand side of (3.5) can be written as

$$\begin{aligned} [(u \circ A) * (E_t(x)E_s(y))](\xi, \eta) &= \langle u \circ A, E_t(\xi - x)E_s(\eta - y) \rangle \\ &= \left\langle u_x, \int E_t(\xi - x + y)E_s(\eta - y) dy \right\rangle \\ &= \left\langle u_x, \int E_t(\xi + \eta - x - y)E_s(y) dy \right\rangle \\ &= \langle u_x, (E_t * E_s)(\xi + \eta - x) \rangle \\ &= \langle u_x, E_{t+s}(\xi + \eta - x) \rangle \\ &= Gu(\xi + \eta, t + s), \end{aligned}$$

and similarly

$$\begin{aligned} [(u \circ P_1) * (E_t(x)E_s(y))](\xi, \eta) &= Gu(\xi, t), \\ [(u \circ P_2) * (E_t(x)E_s(y))](\xi, \eta) &= Gu(\eta, s), \end{aligned}$$

where $Gu(\xi, t)$ is the Gauss transform of u .

Also the right-hand side of (3.5) can be written as

$$[\epsilon(x^{2\gamma} + y^{2\gamma}) * (E_t(x)E_s(y))](\xi, \eta) = \epsilon(H_{2\gamma}(\xi, t) + H_{2\gamma}(\eta, s)),$$

where $H_{2\gamma}$ is the heat polynomial of degree 2γ which is given by

$$H_{2\gamma}(\xi, t) = [x^{2\gamma} * E_t(x)](\xi) = (2\gamma)! \sum_{0 \leq \alpha \leq \gamma} \frac{t^{|\alpha|} \xi^{2\gamma - 2\alpha}}{\alpha!(2\gamma - 2\alpha)!}.$$

Thus the inequality (3.5) is converted to the following stability problem involving the Gauss transform of u :

$$|Gu(\xi + \eta, t + s) - Gu(\xi, t) - Gu(\eta, s)| \leq \epsilon(H_{2\gamma}(\xi, t) + H_{2\gamma}(\eta, s)) \tag{3.7}$$

for all $\xi, \eta \in \mathbb{R}^n, t, s > 0$.

Now we follow the same method as in [15]. Replacing both ξ and η by $\xi/2$, both t and s by $t/2$ in (3.7) we have

$$|Gu(\xi, t) - 2Gu(2^{-1}\xi, 2^{-1}t)| \leq 2\epsilon H_{2\gamma}(2^{-1}\xi, 2^{-1}t)$$

for all $\xi \in \mathbb{R}^n, t > 0$. Making use of the induction argument and triangle inequality we have

$$\begin{aligned} |Gu(\xi, t) - 2^n Gu(2^{-n}\xi, 2^{-n}t)| &\leq \epsilon \sum_{j=1}^n 2^j H_{2\gamma}(2^{-j}\xi, 2^{-j}t) \\ &\leq \epsilon (2\gamma)! \sum_{0 \leq \alpha \leq \gamma} a_{n,\alpha} \frac{t^{|\alpha|} \xi^{2\gamma - 2\alpha}}{\alpha!(2\gamma - 2\alpha)!} \end{aligned} \tag{3.8}$$

for all $n \in \mathbb{N}$, $\xi \in \mathbb{R}^n$, $t > 0$, where $a_{n,\alpha} = 2^{|\alpha|+1}(1 - 2^{(|\alpha|-2\gamma+1)n})/(2^{2\gamma} - 2^{|\alpha|+1})$.

Replacing ξ, t by $2^{-m}\xi, 2^{-m}t$, respectively in (3.8) and multiplying 2^m in the result it follows easily from the fact $|\gamma| > 1$ that

$$A_m(\xi, t) := 2^n Gu(2^{-m}\xi, 2^{-m}t)$$

is a Cauchy sequence which converges locally uniformly. Now let

$$A(\xi, t) = \lim_{m \rightarrow \infty} A_m(\xi, t).$$

Then $A(\xi, t)$ is the unique mapping in $\mathbb{R}^n \times (0, \infty)$ satisfying

$$|Gu(\xi, t) - A(\xi, t)| \leq \epsilon(2\gamma)! \sum_{0 \leq \alpha \leq \gamma} a_\alpha \frac{t^{|\alpha|} \xi^{2\gamma-2\alpha}}{\alpha!(2\gamma - 2\alpha)!}, \tag{3.9}$$

$$A(\xi + \eta, t + s) - A(\xi, t) - A(\eta, s) = 0 \tag{3.10}$$

for all $\xi, \eta \in \mathbb{R}^n$, $t, s > 0$, where $a_\alpha = 2^{|\alpha|+1}/(2^{2\gamma} - 2^{|\alpha|+1})$. Indeed, the inequality (3.9) follows immediately from (3.8). To prove (3.10), replacing ξ, η, t, s by $2^{-m}\xi, 2^{-m}\eta, 2^{-m}t, 2^{-m}s$ in (3.7), respectively, multiplying 2^m and letting $m \rightarrow \infty$ it follows immediately from the fact $|\gamma| > 1$. To prove the uniqueness of $A(\xi, t)$, let $B(\xi, t)$ be another function satisfying (3.9) and (3.10). Then it follows from (3.9), (3.10) and the triangle inequality that for all $n \in \mathbb{N}$,

$$\begin{aligned} |A(\xi, t) - B(\xi, t)| &\leq n \left| A\left(\frac{x}{n}, \frac{t}{n}\right) - B\left(\frac{x}{n}, \frac{t}{n}\right) \right| \\ &\leq 2\epsilon(2\gamma)! n^{1-|\gamma|} \sum_{0 \leq \alpha \leq \gamma} a_\alpha \frac{t^{|\alpha|} \xi^{2\gamma-2\alpha}}{\alpha!(2\gamma - 2\alpha)!}. \end{aligned} \tag{3.11}$$

Letting $n \rightarrow \infty$, we have $A(\xi, t) = B(\xi, t)$ for all $\xi \in \mathbb{R}^n$, $t > 0$. This proves the uniqueness.

Now it is easy to see that every continuous solution $A(\xi, t)$ of the Cauchy equation (3.10) has the form

$$A(\xi, t) = a \cdot \xi + bt$$

for some $a \in \mathbb{C}^n$, $b \in \mathbb{C}$. Letting $t \rightarrow 0^+$ in (3.9) we have

$$\|u - a \cdot \xi\| \leq \frac{2\epsilon}{4^{|\gamma|} - 2} \xi^{2\gamma}.$$

This completes the proof. \square

Now we consider the inequality

$$\|u \circ A - u \circ P_1 - u \circ P_2\| \leq \epsilon(\|x\|^{2p} + \|y\|^{2p})$$

for some integer $p > 1$.

Theorem 3.4. *Let u in \mathcal{S}' or \mathcal{F}' satisfy the inequality*

$$\|u \circ A - u \circ P_1 - u \circ P_2\| \leq \epsilon(\|x\|^{2p} + \|y\|^{2p}) \tag{3.12}$$

for some integer $p > 1$. Then there exists a unique $a \in \mathbb{C}^n$ such that

$$\|u - a \cdot x\| \leq \frac{2\epsilon}{4^p - 2} \|x\|^{2p}. \tag{3.13}$$

Proof. Note that we can write

$$\|x\|^{2p} = \sum_{|\gamma|=p} \frac{p!}{\gamma!} x^{2\gamma}.$$

Thus convolving in each side of (3.12) the tensor product $E_t(x)E_s(y)$ of n -dimensional heat kernels as a function of x, y the inequality (3.12) is converted to the following inequality as in the proof of Theorem 3.3:

$$\begin{aligned} & |Gu(\xi + \eta, t + s) - Gu(\xi, t) - Gu(\eta, s)| \\ & \leq \epsilon \sum_{|\gamma|=p} \frac{p!}{\gamma!} (H_{2\gamma}(\xi, t) + H_{2\gamma}(\eta, s)) \end{aligned} \tag{3.14}$$

for all $\xi, \eta \in \mathbb{R}^n, t, s > 0$.

Now making use of the same approach as in the proof of above theorem we have

$$\|u - a \cdot \xi\| \leq \sum_{|\gamma|=p} \frac{p!}{\gamma!} \left(\frac{2\epsilon}{4^{|\gamma|} - 2} \xi^{2\gamma} \right) = \frac{2\epsilon}{4^p - 2} \|\xi\|^{2p}.$$

This completes the proof. \square

References

- [1] J.A. Baker, Distributional methods for functional equations, *Aequationes Math.* 62 (2001) 136–142.
- [2] J. Chung, Stability of functional equations in the spaces of distributions and hyperfunctions, *J. Math. Anal. Appl.* 286 (2003) 177–186.
- [3] J. Chung, S.-Y. Chung, D. Kim, Une caractérisation de l'espace de Schwartz, *C. R. Acad. Sci. Paris Sér. I Math.* 316 (1993) 23–25.
- [4] J. Chung, S.-Y. Chung, D. Kim, A characterization for Fourier hyperfunctions, *Publ. Res. Inst. Math. Sci.* 30 (1994) 203–208.
- [5] J. Chung, S.Y. Lee, Some functional equations in the spaces of generalized functions, *Aequationes Math.* 65 (2003) 267–279.
- [6] S.-Y. Chung, Reformulation of some functional equations in the space of Gevrey distributions and regularity of solutions, *Aequationes Math.* 59 (2000) 108–123.
- [7] Z. Gajda, On stability of additive mappings, *Internat. J. Math. Math. Sci.* 14 (1991) 431–434.
- [8] I.M. Gelfand, G.E. Shilov, *Generalized Functions I, II*, Academic Press, New York, 1968.
- [9] L. Hörmander, *The Analysis of Linear Partial Differential Operators I*, Springer-Verlag, Berlin, 1983.
- [10] D.H. Hyers, On the stability of the linear functional equations, *Proc. Nat. Acad. Sci. USA* 27 (1941) 222–224.
- [11] D.H. Hyers, G. Isac, Th.M. Rassias, *Stability of Functional Equations in Several Variables*, Birkhäuser, 1998.
- [12] G. Isac, Th.M. Rassias, On the Hyers–Ulam stability of ψ -additive mappings, *J. Approx. Theory* 72 (1993) 131–137.
- [13] E.L. Koh, The Cauchy functional equation in distributions, *Proc. Amer. Math. Soc.* 106 (1989) 641–647.
- [14] T. Matsuzawa, A calculus approach to hyperfunctions III, *Nagoya Math. J.* 118 (1990) 133–153.

- [15] Th.M. Rassias, On the stability of linear mapping in Banach spaces, Proc. Amer. Math. Soc. 72 (1978) 297–300.
- [16] L. Schwartz, Théorie des distributions, Hermann, Paris, 1966.
- [17] S.M. Ulam, Problems in Modern Mathematics, Wiley, New York, 1964, Chapter VI.