



NLS ground states on metric graphs with localized nonlinearities



Lorenzo Tentarelli

Dipartimento di Scienze Matematiche “G.L. Lagrange”, Politecnico di Torino, Corso Duca degli Abruzzi, 24, 10129 Torino, Italy

ARTICLE INFO

Article history:

Received 27 February 2015
Available online 3 August 2015
Submitted by V. Radulescu

Keywords:

Minimization
Metric graphs
NLS energy
Nonlinear Schrödinger equation
Localized nonlinearity

ABSTRACT

We investigate the existence of ground states for the focusing subcritical NLS energy on metric graphs with localized nonlinearities. In particular, we find two thresholds on the measure of the region where the nonlinearity is localized that imply, respectively, existence or nonexistence of ground states. In order to obtain these results we adapt to the context of metric graphs some classical techniques from the Calculus of Variations.

© 2015 Elsevier Inc. All rights reserved.

1. Introduction

In this paper we discuss the existence of a ground state for the NLS energy functional with a *localized nonlinearity*

$$E(u, \mathcal{G}) := \frac{1}{2} \|u'\|_{L^2(\mathcal{G})}^2 - \frac{1}{p} \|u\|_{L^p(\mathcal{K})}^p, \quad (1)$$

where $p \in (2, 6)$ is a real parameter and \mathcal{G} and \mathcal{K} are two connected *metric graphs*, such that \mathcal{K} is a *compact* subgraph of \mathcal{G} . Namely, we investigate the minimization problem

$$\min E(u, \mathcal{G}), \quad u \in H^1(\mathcal{G}), \quad \|u\|_{L^2(\mathcal{G})}^2 = \mu \quad (2)$$

where $\mu > 0$ is a given number.

Here we present a rather informal description of the problem and of the main results of the paper, whereas a precise setting and formal definitions are given in Sections 2 and 3.

It is well known that, when the graph \mathcal{G} is compact, the minimization problem is trivial. The aim of this paper, therefore, is the study of existence and nonexistence of solutions to (2) when \mathcal{G} is made up of a compact graph \mathcal{K} where the nonlinearity is localized and a finite number of *half-lines* \mathcal{R}_i incident at some

E-mail address: lorenzo.tentarelli@polito.it.

vertices of \mathcal{K} . For the sake of completeness, we recall that the case of *non-localized* nonlinearities (where the *lack of compactness* is stronger) is investigated in [3].

The main results we are going to present are the following. In [Theorem 3.1](#) we adapt to the context of metric graphs a classical *level argument* (see e.g. [9,22,7]) in order to overcome the lack of compactness. More in detail, first we show that under general assumptions

$$\inf_{\substack{u \in H^1(\mathcal{G}) \\ \|u\|_{L^2(\mathcal{G})}^2 = \mu}} E(u, \mathcal{G}) \leq 0$$

and further that, if

$$\inf_{\substack{u \in H^1(\mathcal{G}) \\ \|u\|_{L^2(\mathcal{G})}^2 = \mu}} E(u, \mathcal{G}) < 0, \quad \text{then the infimum is attained.}$$

Consequently, we exploit these two properties to prove that existence and nonexistence of ground states strongly depend on the exponent p of the nonlinearity and on the metric properties of the compact graph \mathcal{K} . In particular, [Theorem 3.3](#) states that, if $p \in (2, 4)$, then there always exists a minimizer for E , whereas [Theorem 3.4](#) states that, when $p \in [4, 6)$, the existence of minimizers depends on the measure of \mathcal{K} ; precisely, there is a threshold for $\text{meas}(\mathcal{K})$ over which the minimum is always attained and another one under which the minimum cannot be attained. In [Corollary 3.6](#) we slightly improve the nonexistence threshold, by adding some further assumptions on the inner structure of \mathcal{G} .

Finally we highlight that all the functions we consider are real-valued. Indeed, as pointed out in [3], this is not restrictive since $E(|u|, \mathcal{G}) \leq E(u, \mathcal{G})$ and any ground state is in fact real-valued (and then, without loss of generality, nonnegative), up to multiplication by a constant phase $e^{i\vartheta}$.

Among the physical motivations for the investigation of this kind of problem, nowadays the most topical ones come from the study of *quantum graphs*. This subject has gained popularity in recent years (as extensively pointed out in [18,15]) not only because graphs emulate successfully complex mesoscopic and optical networks, but also because they manage to reproduce universal properties of quantum chaotic systems. In particular, in [16] (and the references therein) two main applications are exhibited to motivate the study of NLS energy functionals with localized nonlinearities. Indeed, they can be of interest both in the analysis of the effects of nonlinearity on transmission through a complex network of nonlinear one-dimensional leads (for instance, *optical fibers*, e.g. [5]) and in the analysis of the properties of *Bose–Einstein condensates* in non-regular traps (see also [3,12]). In other words, even though these are idealized models, they actually describe systems made up by several long leads (where the nonlinearity is, in principle, negligible) linked to a “dense” sub-network (where the nonlinearity is strong) and reproduce accurately the effects that the latter induces on the whole network.

It is also worth recalling that the investigation of the NLS equation with localized nonlinearities in standard domains has been the object of several papers. In the one-dimensional case we first mention [20], that introduces a pioneering model of nonlinearities concentrated at a finite number of points. Then, [4] contains a rigorous definition of the problem with potentials modeled by *Dirac delta functions* with amplitude depending nonlinearly on the wave function. Furthermore, [11] shows that such nonlinearities can be seen as *point-like* limits of spatially concentrated nonlinearities, whereas [10] deals with the associated issue of the asymptotic stability. Finally, for a rigorous setting of the case of a *point-interaction* with strength depending nonlinearly on the wave function in dimension three, we refer the reader to [1], while in [17,21] one can find several models of concentrated nonlinearities in dimension three or higher.

In this framework, our work can be seen as a slight extension of the one-dimensional case (although we only focus on ground states of the NLS energy) in that a nonlinear potential is spread over a compact graph in place of a Dirac delta based at single point.




Fig. 1. A compact metric graph \mathcal{K} with 14 vertices and 24 bounded edges. In particular there are: 1 self-loop, 1 multiple edge, 2 vertices of degree one and 2 vertices of degree two.

The paper is organized as follows. In Sections 2 and 3 we present, respectively, the precise setting of the problem and the statements of the main results. Section 4 contains a discussion of some preliminary and auxiliary facts and techniques that may have interest in themselves. Finally, in Section 5 we prove the results stated in Section 3.

2. Setting and notation

Although it is a central notion in this paper, we do not present a deep and detailed discussion on metric graphs. For a modern account on the subject we refer the reader to [3,14,19] and references therein.

We start with a brief recall on metric graphs and, in particular, on the class of metric graphs that we focus on in this paper. First of all, the graphs that we consider are multigraphs, namely, graphs with possibly multiple edges and self-loops. Moreover we only deal with connected metric graphs, that is, connected graphs with the “representation” introduced in [3], which associates each edge e of length l_e with an interval $I_e := [0, l_e]$ parametrized by a variable x_e ($x_e = 0$ representing either the starting or the endpoint of e , depending on the orientation choice). In addition, if $l_e = +\infty$, then e is called half-line and $I_e := [0, +\infty)$ (now $x_e = 0$ representing the starting point of the half-line).

Throughout this paper we denote by $\mathcal{K} = (V_{\mathcal{K}}, E_{\mathcal{K}})$ a (non-trivial) compact connected metric graph, i.e. a connected metric graph without half-lines (or, equivalently, without vertices at infinity). We stress the fact that there are no constraints on the degree of the vertices. This means that there are possibly vertices of degree one or two (see, for instance, Fig. 1).

Furthermore, we consider N distinct half-lines $\mathcal{R}_1, \dots, \mathcal{R}_N$ starting from vertices (possibly not distinct) in \mathcal{K} and ending, respectively, into N (distinct) vertices at infinity v_1, \dots, v_N . Then we define the graph

$$\begin{aligned} \mathcal{G} &:= (V, E), \\ V &:= V_{\mathcal{K}} \cup \{v_1, \dots, v_N\}, \quad E := E_{\mathcal{K}} \cup \{\mathcal{R}_1, \dots, \mathcal{R}_N\} \end{aligned} \tag{3}$$

(a typical case is depicted in Fig. 2).

Clearly the measure of the compact graph \mathcal{K} , defined by

$$\text{meas}(\mathcal{K}) := \sum_{e \in E_{\mathcal{K}}} \text{meas}(I_e), \tag{4}$$

is finite, while that of \mathcal{G} is equal to ∞ (and so \mathcal{G} is not compact). Consequently, a function $u : \mathcal{G} \rightarrow \mathbb{R}$ can be regarded as a family of functions $(u_e)_{e \in E}$, where $u_e : I_e \rightarrow \mathbb{R}$ is the restriction of u to the edge I_e , and L^p spaces can be defined over \mathcal{G} in the natural way, with norm

$$\|u\|_{L^p(\mathcal{G})}^p = \sum_{e \in E} \|u_e\|_{L^p(I_e)}^p.$$




Fig. 2. A noncompact metric graph \mathcal{G} obtained attaching 5 half-lines (and 5 vertices at infinity) to the graph \mathcal{K} of Fig. 1. Note that \mathcal{R}_1 is attached to a vertex of degree one, \mathcal{R}_4 and \mathcal{R}_5 to a vertex of degree three and \mathcal{R}_2 and \mathcal{R}_3 to distinct vertices of degree two.

The Sobolev space $H^1(\mathcal{G})$ can be defined as the set of those functions $u : \mathcal{G} \rightarrow \mathbb{R}$ such that $u = (u_e)_{e \in E}$ is continuous on \mathcal{G} and $u_e \in H^1(I_e)$ for every edge $e \in E$, with the natural norm

$$\|u\|_{H^1(\mathcal{G})}^2 := \|u'\|_{L^2(\mathcal{G})}^2 + \|u\|_{L^2(\mathcal{G})}^2.$$

Fix, now, the graphs \mathcal{K} and \mathcal{G} as above and let μ and p be such that

$$\mu > 0 \quad \text{and} \quad 2 < p < 6. \tag{5}$$

Then the functional in (1) is well defined over $H^1(\mathcal{G})$ and takes the form

$$E(u, \mathcal{G}) = \frac{1}{2} \sum_{e \in E} \int_{I_e} |u'_e(x_e)|^2 dx_e - \frac{1}{p} \sum_{e \in E} \int_{I_e} |u_e(x_e)|^p dx_e.$$

Moreover, defining

$$H^1_\mu(\mathcal{G}) := \{u \in H^1(\mathcal{G}) : \|u\|_{L^2(\mathcal{G})}^2 = \mu\}, \tag{6}$$

the minimization problem (2) can be written in a compact form as

$$\min_{u \in H^1_\mu(\mathcal{G})} E(u, \mathcal{G}). \tag{7}$$

Remark 2.1. Note that if $u \in H^1_\mu(\mathcal{G})$ and λ is positive, then the function

$$w(x) := \lambda^{\frac{2}{6-p}} u \left(\lambda^{\frac{p-2}{6-p}} x \right)$$

belongs to $H^1_{\lambda\mu}(\mathcal{G}')$, where \mathcal{G}' is a graph obtained from \mathcal{G} by means of a homothety of factor $\lambda^{\frac{2-p}{6-p}}$. Moreover a change of variable shows that

$$E(w, \mathcal{G}') = \lambda^{\frac{2+p}{6-p}} E(u, \mathcal{G}),$$

and hence the problem with mass constraint $\lambda\mu$ over the graph \mathcal{G}' is equivalent to problem (7).

3. Main results

In this section we present the main results on *existence* and *nonexistence* of a solution for problem (7).

We start with a general result.

Theorem 3.1. *Let \mathcal{G} be a graph as in (3) and let μ, p satisfy (5). Then*

$$\inf_{u \in H^1_\mu(\mathcal{G})} E(u, \mathcal{G}) \leq 0. \tag{8}$$

Moreover, if

$$\inf_{u \in H^1_\mu(\mathcal{G})} E(u, \mathcal{G}) < 0, \tag{9}$$

then the infimum is attained and, up to a change of sign, every minimizer is strictly positive.

It is easy to construct an example of graph \mathcal{G} for which the infimum is achieved.

Example 3.2. Let p, μ satisfy (5) and define the function

$$\varphi_\mu(x) := \mu^\alpha \varphi_1(\mu^\beta x), \quad \alpha = \frac{2}{6-p}, \quad \beta = \frac{p-2}{6-p},$$

where

$$\varphi_1(x) := \Gamma_p \operatorname{sech}(\gamma_p x)^{\alpha/\beta}, \quad \text{with } \Gamma_p, \gamma_p > 0.$$

It is well known that φ_μ (usually called a *soliton of mass μ*) is a minimizer for (7) when $\mathcal{G} = \mathbb{R}$ and the nonlinearity is **not** localized and, in particular,

$$\frac{1}{2} \int_{\mathbb{R}} |\varphi'_\mu(x)|^2 dx - \frac{1}{p} \int_{\mathbb{R}} |\varphi_\mu(x)|^p dx = \mathcal{E} < 0$$

(for more details, see [2,3]). Now let \mathcal{K} be a segment of length L and assume, in addition, that $N = 2$ and that the two half-lines are incident at the endpoints of \mathcal{K} . Hence the graph \mathcal{G} can be seen as a straight line and the energy functional reads

$$E(u, \mathcal{G}) := \frac{1}{2} \int_{\mathbb{R}} |u'(x)|^2 dx - \frac{1}{p} \int_{-L/2}^{L/2} |u(x)|^p dx.$$

Hence, if \tilde{L} is a positive constant such that

$$\int_{|x| > \tilde{L}/2} |\varphi_\mu(x)|^p dx < |\mathcal{E}|,$$

then we see that

$$E(\varphi_\mu, \mathcal{G}) < 0 \quad \text{whenever} \quad L > \tilde{L}$$

and, from Theorem 3.1, E has a minimizer in $H^1_\mu(\mathcal{G})$.

In fact, in more general cases, existence or nonexistence of minimizers for E strongly depends on the exponent of the nonlinearity and on the measure of the compact graph \mathcal{K} . This is described in the following two theorems.




Fig. 3. (a) A *double bridge* graph; (b) a partition where each \mathcal{G}_i contains a half-line and a bounded edge; (c) a partition where \mathcal{G}_1 contains a half-line and all the bounded edges and \mathcal{G}_2 only a half-line.

Theorem 3.3. *Let \mathcal{G} be as in (3) and $\mu > 0$. Then for every $p \in (2, 4)$ the minimum in (7) is attained.*

Theorem 3.4. *Let \mathcal{G} be as in (3) and $\mu > 0$. If $p \in [4, 6)$, then there exist two positive constants L_1, L_2 such that*

$$\text{meas}(\mathcal{K}) > L_1 \quad \Rightarrow \quad \text{the minimum in (7) is attained}, \tag{10}$$

$$\text{meas}(\mathcal{K}) < L_2 \quad \Rightarrow \quad \text{the minimum in (7) is not attained}. \tag{11}$$

In some cases we can improve the threshold that guarantees nonexistence. To this aim, we give the following definition.

Definition 3.5. Let $N \geq 2$ and \mathcal{G} be a graph satisfying (3). In addition, let $(\mathcal{G}_i)_{i=1}^\nu$, with $2 \leq \nu \leq N$, be a family of non-trivial subgraphs of \mathcal{G} (each consisting of edges and vertices of \mathcal{G}). We say that $(\mathcal{G}_i)_{i=1}^\nu$ is a *partition* of \mathcal{G} if:

1. $\bigcup_{i=1}^\nu \mathcal{G}_i = \mathcal{G}$;
2. $\text{meas}(\mathcal{G}_i \cap \mathcal{G}_l) = 0$ whenever $i \neq l$;
3. for every $i \in \{1, \dots, \nu\}$, there exists $j \in \{1, \dots, N\}$ such that $\mathcal{R}_j \subset \mathcal{G}_i$.

In other words, a partition of a graph is the union of “essentially” pairwise disjoint subgraphs, each containing at least a half-line. We also stress the fact that every graph admits, in principle, several distinct partitions. Two examples of partitions are given in Fig. 3 in the case of a *double bridge* graph.

Now we can extend (11) as follows.

Corollary 3.6. *Let \mathcal{G} , μ and p satisfy the assumptions of Theorem 3.4. In addition, let $(\mathcal{G}_i)_{i=1}^\nu$ be a partition of \mathcal{G} and $\mathcal{K}_i = \mathcal{G}_i \cap \mathcal{K}$, for $i = 1, \dots, \nu$. If L_2 is the same of Theorem 3.4 and*

$$\text{meas}(\mathcal{K}_i) < L_2 \quad \forall i = 1, \dots, \nu,$$

then the minimum in (7) is not attained.

Remark 3.7. In order to understand how [Corollary 3.6](#) “improves” [Theorem 3.4](#), consider again the double bridge graph. If one chooses the partition of [Fig. 3\(b\)](#), then

$$\text{meas}(\mathcal{K}_1) = \text{meas}(\mathcal{K}_2) = \frac{\text{meas}(\mathcal{K})}{2}.$$

Hence, although $\text{meas}(\mathcal{K}) > L_2$, one can claim that ground states do not exist (provided $\text{meas}(\mathcal{K}_1) = \text{meas}(\mathcal{K}_2) < L_2$).

Remark 3.8. The bounds L_1 and L_2 deserve some further comment. As we will show in [Section 5](#), we can compute them explicitly:

$$\begin{aligned} \mu L_1 &= \frac{N^2}{2} && \text{if } p = 4 \\ \mu^{\frac{p-2}{6-p}} L_1 &= c_p N^{\frac{4}{6-p}} && \text{if } p \in (4, 6) \\ \mu^{\frac{p-2}{6-p}} L_2 &= C_p^{\frac{4-p}{6-p}} C_\infty^{-p} && \text{if } p \in [4, 6), \end{aligned} \tag{12}$$

where c_p is defined by [\(31\)](#), while C_p and C_∞ come, respectively, from inequalities [\(13\)](#) and [\(14\)](#). In addition one can check that they are invariant under the transformations

$$\mu \rightarrow \lambda\mu \quad \mathcal{G} \rightarrow \lambda^{\frac{2-p}{6-p}} \mathcal{G},$$

coherently with [Remark 2.1](#). This entails that the results we stated in [Theorems 3.3 and 3.4](#) can be equivalently formulated in terms of the mass μ in place of $\text{meas}(\mathcal{K})$. Precisely, for $\text{meas}(\mathcal{K}) > 0$ fixed, if $p \in (2, 4)$, then there always exists a solution to [\(7\)](#); whereas, if $p \in [4, 6)$, then there exist $\mu_1, \mu_2 > 0$ such that

$$\begin{aligned} \mu > \mu_1 &\Rightarrow \text{the minimum in (7) is attained,} \\ \mu < \mu_2 &\Rightarrow \text{the minimum in (7) is not attained.} \end{aligned}$$

4. Preliminary and auxiliary results

In this section we introduce some tools that we will use in the proofs of the main results in [Section 5](#). The first one is a version of the Gagliardo–Nirenberg inequalities over a graph \mathcal{G} .

Proposition 4.1. *Let \mathcal{G} be as in [\(3\)](#). For every $p \in [2, \infty]$ there exist two positive constants C_p (depending only on p) and C_∞ such that*

$$\|u\|_{L^p(\mathcal{G})}^p \leq C_p \|u\|_{L^2(\mathcal{G})}^{\frac{p}{2}+1} \|u'\|_{L^2(\mathcal{G})}^{\frac{p}{2}-1} \quad \forall u \in H^1(\mathcal{G}), \quad \text{if } p < \infty, \tag{13}$$

$$\|u\|_{L^\infty(\mathcal{G})} \leq C_\infty \|u\|_{L^2(\mathcal{G})}^{\frac{1}{2}} \|u'\|_{L^2(\mathcal{G})}^{\frac{1}{2}} \quad \forall u \in H^1(\mathcal{G}). \tag{14}$$

Proof. Consider a nonnegative function $u \in H^1(\mathcal{G})$. We define its *decreasing rearrangement* as the function $u^* : \mathbb{R}^+ \rightarrow \mathbb{R}$ such that

$$u^*(x) := \inf\{t \geq 0 : \rho(t) \leq x\},$$

where

$$\rho(t) := \sum_{e \in \mathbb{E}} \text{meas}(\{x_e \in I_e : u_e(x_e) > t\}), \quad t \geq 0,$$

is the *distribution function* of u (for more details, see [3,14]). Then, clearly,

$$\int_{\mathcal{G}} |u(x)|^r dx = \int_{\mathbb{R}^+} |u^*(x)|^r dx \quad \text{and} \quad \sup_{\mathcal{G}} u = \sup_{\mathbb{R}^+} u^* \tag{15}$$

for every $r \geq 1$. Moreover $u^* \in H^1(\mathbb{R}^+)$ and, by Proposition 3.1 in [3],

$$\int_{\mathbb{R}^+} |(u^*)'(x)|^2 dx \leq \int_{\mathcal{G}} |u'(x)|^2 dx. \tag{16}$$

Then, from the Gagliardo–Nirenberg inequality on \mathbb{R}^+ (cf. [6,13])

$$\begin{aligned} \|u\|_{L^p(\mathcal{G})}^p &= \|u^*\|_{L^p(\mathbb{R}^+)}^p \leq C_p \|u^*\|_{L^2(\mathbb{R}^+)}^{\frac{p}{2}+1} \|(u^*)'\|_{L^2(\mathbb{R}^+)}^{\frac{p}{2}-1} \\ &\leq C_p \|u\|_{L^2(\mathcal{G})}^{\frac{p}{2}+1} \|u'\|_{L^2(\mathcal{G})}^{\frac{p}{2}-1}. \end{aligned}$$

In the very same way it is possible to prove (14). \square

Remark 4.2. It is well known that the *best constant* of (14) when $\mathcal{G} = \mathbb{R}$ is equal to 1. Indeed, assume that $u \in H^1(\mathbb{R})$ satisfies $|u(0)| = \max_{\mathbb{R}} |u(x)|$ and let \bar{u} be the *even part* of u . Then

$$u^2(0) = \bar{u}^2(0) = 2 \int_{-\infty}^0 \bar{u}(x)\bar{u}'(x)dx \leq \|\bar{u}\|_{L^2(\mathbb{R})} \|\bar{u}'\|_{L^2(\mathbb{R})},$$

so that

$$\|u\|_{L^\infty(\mathbb{R})} \leq \|u\|_{L^2(\mathbb{R})}^{\frac{1}{2}} \|u'\|_{L^2(\mathbb{R})}^{\frac{1}{2}}.$$

Since equality holds for $u = e^{-|x|}$, the claim follows. With a slight further effort, one can also check that when $\mathcal{G} = \mathbb{R}^+$ the best constant is $\sqrt{2}$.

According to these remarks, we see that for a generic graph \mathcal{G} the best constant in (14) satisfies $C_\infty \leq \sqrt{2}$. In addition, if the graph contains at least two half-lines, then one can repeat the proof of Proposition 4.1 using the *symmetric decreasing* rearrangement on \mathbb{R} in place of the *decreasing* one on \mathbb{R}^+ (see again [3,14]), thus obtaining that (in this case) $C_\infty \leq 1$.

Next we deal with the minimization problem (7) on a straight line with an extra Dirichlet condition.

Proposition 4.3. *Let $m > 0$, $a > 0$. Then the function $u(x) = ae^{-\frac{a^2|x|}{m}}$ satisfies*

$$\|u'\|_{L^2(\mathbb{R})}^2 = \min_{v \in H_{m,a}^1(\mathbb{R})} \|v'\|_{L^2(\mathbb{R})}^2 \tag{17}$$

where $H_{m,a}^1(\mathbb{R}) := \{v \in H_m^1(\mathbb{R}) : v(0) = a\}$.

Proof. Consider inequality (14). From Remark 4.2 we know that, when $\mathcal{G} = \mathbb{R}$, the best constant is $C_\infty = 1$. Hence, for a non-identically zero function v in $H^1(\mathbb{R})$, we find that

$$\|v'\|_{L^2(\mathbb{R})}^2 \geq \frac{\|v\|_{L^\infty(\mathbb{R})}^4}{\|v\|_{L^2(\mathbb{R})}^2}$$

and, assuming also that $v \in H^1_{m,a}(\mathbb{R})$,

$$\|v'\|_{L^2(\mathbb{R})}^2 \geq \frac{a^4}{m}. \tag{18}$$

Since $u \in H^1_{m,a}(\mathbb{R})$ and

$$\|u'\|_{L^2(\mathbb{R})}^2 = \frac{a^4}{m},$$

we see that the infimum is attained by u . \square

The following is an analogous result over the half-line.

Corollary 4.4. *Let $m > 0, a > 0$. Then the function $u(x) := ae^{-\frac{a^2x}{2m}}$, satisfies*

$$\|u'\|_{L^2(\mathbb{R}^+)}^2 = \min_{v \in H^1_{m,a}(\mathbb{R}^+)} \|v'\|_{L^2(\mathbb{R}^+)}^2. \tag{19}$$

Finally, we recall a standard result about the optimality conditions satisfied by any solution of (7) (for a proof, see [3]).

Proposition 4.5. *Let \mathcal{G} be as in (3) and let μ, p satisfy (5). Suppose, in addition, that u is a solution of (7). Then*

(i) *there exists a real constant λ such that, for every edge $e \in E$*

$$u''_e + \kappa_e u_e |u_e|^{p-2} = \lambda u_e \quad (\text{Euler-Lagrange equation}) \tag{20}$$

where $\kappa_e = 1$ if $e \in E_{\mathcal{K}}$ and $\kappa_e = 0$ if $e \in E \setminus E_{\mathcal{K}}$;

(ii) *for every vertex $v \in V_{\mathcal{K}}$*

$$\sum_{e \succ v} \frac{du_e}{dx_e}(v) = 0 \quad (\text{Kirchhoff conditions}) \tag{21}$$

where “ $e \succ v$ ” means that the edge e is incident at v ;

(iii) *up to a change of sign, $u > 0$ on \mathcal{G} .*

Remark 4.6. The symbol $\frac{du_e}{dx_e}(v)$ is a shorthand notation for $u'_e(0)$ or $-u'_e(l_e)$, according to the fact that x_e is equal to 0 or l_e at v .

5. Proof of the main results

Throughout this section, it is convenient to identify each $u \in H^1(\mathcal{G})$ with a $(N + 1)$ -ple of functions $\psi, \varphi_1, \dots, \varphi_N$ such that

$$\psi \in H^1(\mathcal{K}), \quad \varphi_i \in H^1(\mathcal{R}_i) \quad \text{for } i = 1, \dots, N, \tag{22}$$

and

$$\psi(w) = \varphi_i(w) \quad \text{for } i = 1, \dots, N, \tag{23}$$

where w denotes the starting vertex of \mathcal{R}_i . Then (7) is equivalent to the minimization problem

$$\min \left(\frac{1}{2} \|\psi'\|_{L^2(\mathcal{K})}^2 - \frac{1}{p} \|\psi\|_{L^p(\mathcal{K})}^p + \frac{1}{2} \sum_{i=1}^N \|\varphi'_i\|_{L^2(\mathcal{R}_i)}^2 \right),$$

with $(\psi, \varphi_1, \dots, \varphi_N)$ varying in the set of the $(N + 1)$ -ples satisfying (22)–(23) and subject to the mass constraint

$$\|\psi\|_{L^2(\mathcal{K})}^2 + \sum_{i=1}^N \|\varphi_i\|_{L^2(\mathcal{R}_i)}^2 = \mu. \tag{24}$$

Proof of Theorem 3.1. We prove the two statements separately.

Part (i): proof of (8). Consider a function $\xi \in C_0^\infty(\mathbb{R}^+)$ with $\|\xi\|_{L^2(\mathbb{R}^+)}^2 = \mu$. Then, for every $\lambda > 0$ we define a function

$$u_\lambda(x) = \begin{cases} \sqrt{\lambda} \xi(\lambda x) & \text{if } x \in \mathcal{R}_1 \\ 0 & \text{elsewhere.} \end{cases}$$

As a consequence, $u_\lambda \in H_\mu^1(\mathcal{G})$ and $E(u_\lambda, \mathcal{G}) = \frac{\lambda^2}{2} \|\xi'\|_{L^2(\mathbb{R}^+)}^2$. Hence

$$\inf_{u \in H_\mu^1(\mathcal{G})} E(u, \mathcal{G}) \leq \lim_{\lambda \rightarrow 0} E(u_\lambda, \mathcal{G}) = 0$$

and (8) is proved.

Part (ii): (9) entails existence. Note that from (13)

$$E(u, \mathcal{G}) \geq \frac{1}{2} \|u'\|_{L^2(\mathcal{G})}^2 - \frac{C_p \mu^{\frac{p+2}{4}}}{p} \|u'\|_{L^2(\mathcal{G})}^{\frac{p}{2}-1} \quad \forall u \in H_\mu^1(\mathcal{G}). \tag{25}$$

Then, since $p/2 - 1 < 2$ by (5), there results

$$\|u\|_{H^1(\mathcal{G})}^2 \leq C + CE(u, \mathcal{G}) \tag{26}$$

(here C depends on μ). Consider now a minimizing sequence u_k for problem (7). By (26) it is bounded in $H^1(\mathcal{G})$ and therefore (up to subsequences)

$$u_k \rightharpoonup u \quad \text{in } H^1(\mathcal{G}).$$

Recalling that $u_k = (\psi_k, \varphi_{1k}, \dots, \varphi_{Nk})$ with $\psi_k, \varphi_{1k}, \dots, \varphi_{Nk}$ satisfying (22)–(24), we see that there exist $\psi \in H^1(\mathcal{K})$ and $\varphi_i \in H^1(\mathcal{R}_i)$, satisfying (23), such that

$$\psi_k \rightharpoonup \psi \quad \text{in } H^1(\mathcal{K}), \quad \varphi_{ik} \rightharpoonup \varphi_i \quad \text{in } H^1(\mathcal{R}_i) \quad \text{for } i = 1, \dots, N.$$

Moreover, since \mathcal{K} is compact, we also have

$$\psi_k \rightarrow \psi \quad \text{in } L^p(\mathcal{K}).$$

Setting $u = (\psi, \varphi_1, \dots, \varphi_N)$, by weak lower semicontinuity,

$$E(u, \mathcal{G}) \leq \liminf_k E(u_k, \mathcal{G}) = \inf_{v \in H_\mu^1(\mathcal{G})} E(v, \mathcal{G}) < 0 \tag{27}$$

and

$$\|u\|_{L^2(\mathcal{G})}^2 \leq \mu.$$

If we prove, in addition, that the latter inequality is in fact an equality, then the function u is a minimizer. Note that, by (27), $u \not\equiv 0$. Suppose now that $0 < \|u\|_{L^2(\mathcal{G})}^2 < \mu$. Hence there exists $\sigma > 1$ such that $\|\sigma u\|_{L^2(\mathcal{G})}^2 = \mu$. Consequently

$$\begin{aligned} E(\sigma u, \mathcal{G}) &= \frac{\sigma^2}{2} \int_{\mathcal{G}} |u'(x)|^2 dx - \frac{\sigma^p}{p} \int_{\mathcal{K}} |u(x)|^p dx \\ &< \sigma^2 E(u, \mathcal{G}) < E(u, \mathcal{G}) \leq \inf_{v \in H_{\mu}^1(\mathcal{G})} E(v, \mathcal{G}), \end{aligned}$$

contradicting (27). Then $\|u\|_{L^2(\mathcal{G})}^2 = \mu$ and this proves (9). Finally, $u > 0$ immediately follows from Proposition 4.5. \square

Proof of Theorem 3.3. Let $L = \text{meas}(\mathcal{K})$ and consider the function $u = (\psi, \varphi_1, \dots, \varphi_N)$ defined by

$$\psi \equiv a, \quad \varphi_i(x) = ae^{-\frac{a^2 x}{2m}} \quad \text{for } i = 1, \dots, N, \tag{28}$$

with

$$a \in \left(0, \sqrt{\mu/L}\right) \quad \text{and} \quad m = \frac{\mu - a^2 L}{N}. \tag{29}$$

Then (22)–(24) are satisfied and the energy functional reads

$$E(u, \mathcal{G}) = \frac{a^4 N^2}{8(\mu - a^2 L)} - \frac{a^p L}{p}. \tag{30}$$

Now, since $p \in (2, 4)$, when a is sufficiently small we see that $E(u, \mathcal{G}) < 0$ and hence existence of minimizers for (7) follows from Theorem 3.1. \square

Proof of Theorem 3.4. We break the proof in two parts.

Part (i): proof of (10). Let $L = \text{meas}(\mathcal{K})$ and let u be the function defined in (28)–(29), so that the energy functional reads as in (30). When $p = 4$ one can easily check that, if

$$L > \frac{N^2}{2\mu},$$

then there exists $a_0 \in \left(0, \sqrt{\mu/L}\right)$ such that $E(u, \mathcal{G}) < 0$. On the other hand, when $p \in (4, 6)$ we claim that, if

$$L > c_p \frac{N^{\frac{4}{6-p}}}{\mu^{\frac{p-2}{6-p}}}$$

with

$$c_p = \left[\left(\frac{p(p-4)}{16} \right)^{\frac{2}{p-2}} + \frac{p}{8} \left(\frac{p(p-4)}{16} \right)^{\frac{4-p}{p-2}} \right]^{\frac{p-2}{6-p}}, \tag{31}$$

then there exists again $a_0 \in (0, \sqrt{\mu/L})$ such that $E(u, \mathcal{G}) < 0$. Indeed, $E(u, \mathcal{G}) < 0$ is equivalent to $g(a) < \mu$ where

$$g(a) = a^2L + \frac{N^2p}{8L}a^{4-p}.$$

Easy computations show that g has a unique critical point given by

$$\bar{a} = \left(\frac{N^2p(p-4)}{16L^2} \right)^{\frac{1}{p-2}}.$$

As g is strictly convex, this is a global minimizer for g in $(0, \sqrt{\mu/L})$ and, in addition,

$$g(\bar{a}) = L^{-\frac{6-p}{p-2}} \left[\left(\frac{N^2p(p-4)}{16} \right)^{\frac{2}{p-2}} + \frac{N^2p}{8} \left(\frac{N^2p(p-4)}{16} \right)^{\frac{4-p}{p-2}} \right].$$

Now, if

$$\bar{a} < \sqrt{\mu/L} \quad \text{and} \quad g(\bar{a}) < \mu,$$

then the claim is proved setting $a_0 = \bar{a}$. The former inequality explicitly reads

$$\mu^{\frac{p-2}{6-p}}L > \left(\frac{N^2p(p-4)}{16} \right)^{\frac{2}{6-p}}, \tag{32}$$

whereas the latter reads

$$\begin{aligned} \mu^{\frac{p-2}{6-p}}L &> \left[\left(\frac{N^2p(p-4)}{16} \right)^{\frac{2}{p-2}} + \frac{N^2p}{8} \left(\frac{N^2p(p-4)}{16} \right)^{\frac{4-p}{p-2}} \right]^{\frac{p-2}{6-p}} \\ &= N^{\frac{4}{6-p}}c_p. \end{aligned} \tag{33}$$

Now, observing that (33) entails (32), letting

$$L_1 = \begin{cases} \frac{1}{2}\mu^{-1}N^2 & \text{if } p = 4 \\ c_p\mu^{\frac{2-p}{6-p}}N^{\frac{4}{6-p}} & \text{if } p \in (4, 6) \end{cases} \tag{34}$$

we see that (10) follows from Theorem 3.1.

Part (ii): proof of (11). Suppose that there exists a function $u \in H_\mu^1(\mathcal{G})$ such that

$$E(u, \mathcal{G}) \leq 0. \tag{35}$$

We claim that

$$\|u'\|_{L^2(\mathcal{G})}^2 \leq \frac{1}{C_\infty^4\mu} \|u\|_{L^\infty(\mathcal{G})}^{4\left(\frac{p}{4}\right)^{n+1}} (C_\infty^4\mu L)^{\sum_{i=0}^n \left(\frac{p}{4}\right)^i} \quad \forall n \geq 0 \tag{36}$$

(once again $L = \text{meas}(\mathcal{K})$). To check this by induction, we first note that from (35) we have

$$\|u'\|_{L^2(\mathcal{G})}^2 \leq \frac{2}{p} \|u\|_{L^p(\mathcal{K})}^p \leq L \|u\|_{L^\infty(\mathcal{G})}^p, \tag{37}$$

since $2/p < 1$, which corresponds to (36) when $n = 0$. Moreover, for fixed $n > 0$, assume that

$$\|u'\|_{L^2(\mathcal{G})}^2 \leq \frac{1}{C_\infty^4 \mu} \|u\|_{L^\infty(\mathcal{G})}^{4\left(\frac{p}{4}\right)^n} (C_\infty^4 \mu L)^{\sum_{i=0}^{n-1} \left(\frac{p}{4}\right)^i}. \tag{38}$$

Combining with (14), we see that

$$\|u\|_{L^\infty(\mathcal{G})}^p \leq \|u\|_{L^\infty(\mathcal{G})}^{4\left(\frac{p}{4}\right)^{n+1}} (C_\infty^4 \mu L)^{\sum_{i=1}^n \left(\frac{p}{4}\right)^i} \tag{39}$$

and plugging into (37) we recover (36). Furthermore, combining (35) with (25) and (14), we find that

$$\|u\|_{L^\infty(\mathcal{G})} \leq C_\infty C_p^{\frac{1}{6-p}} \mu^{\frac{2}{6-p}}$$

and thus, putting into inequality (36), we obtain

$$\begin{aligned} \|u'\|_{L^2(\mathcal{G})}^2 &\leq C_p^2 \mu^3 (C_\infty^4 \mu L)^{n+1}, & \text{if } p = 4, \\ \|u'\|_{L^2(\mathcal{G})}^2 &\leq C_p^{\frac{4}{6-p}} \mu^{\frac{p+2}{6-p}} \left(C_\infty^{\frac{4p}{p-4}} C_p^{\frac{4}{6-p}} \mu^{\frac{4(p-2)}{(p-4)(6-p)}} L^{\frac{4}{p-4}} \right)^{\left(\frac{p}{4}\right)^{n+1} - 1}, & \text{if } p \in (4, 6), \end{aligned}$$

for every $n \geq 0$. Now, when the terms in brackets in the two inequalities are strictly smaller than 1, letting $n \rightarrow \infty$ we obtain that $\|u'\|_{L^2(\mathcal{G})}^2 = 0$, which is a contradiction since $\|u\|_{L^2(\mathcal{G})}^2 = \mu > 0$ (see [8]). Consequently, if we choose

$$L_2 = C_p^{\frac{4-p}{6-p}} \mu^{\frac{2-p}{6-p}} C_\infty^{-p},$$

then we see that $E(u, \mathcal{G}) > 0$ whenever $L < L_2$ and hence (11) follows from Theorem 3.1. \square

Proof of Corollary 3.6. Consider a function u in $H_\mu^1(\mathcal{G})$. According to the partition $(\mathcal{G}_i)_{i=1}^\nu$, the energy functional reads

$$E(u, \mathcal{G}) = \sum_{i=1}^\nu E(u|_{\mathcal{G}_i}, \mathcal{G}_i). \tag{40}$$

Now if $u|_{\mathcal{G}_i} \equiv 0$ for some i , then $E(u|_{\mathcal{G}_i}, \mathcal{G}_i) = 0$. On the other hand, if $u|_{\mathcal{G}_i} \not\equiv 0$, since

$$\|u|_{\mathcal{G}_i}\|_{L^2(\mathcal{G}_i)}^2 \leq \mu,$$

then there exists $\sigma \geq 1$ satisfying

$$\int_{\mathcal{G}_i} \sigma^2 |u(x)|^2 dx = \mu,$$

so that

$$E(\sigma u|_{\mathcal{G}_i}, \mathcal{G}_i) \leq \sigma^2 E(u|_{\mathcal{G}_i}, \mathcal{G}_i).$$

Therefore, since $\text{meas}(\mathcal{K}_i) < L_2$, (11) entails that $E(\sigma u|_{\mathcal{G}_i}, \mathcal{G}_i) > 0$, whence $E(u|_{\mathcal{G}_i}, \mathcal{G}_i) > 0$. Now, as $u \in H_\mu^1(\mathcal{G})$, then $u|_{\mathcal{G}_i} \not\equiv 0$ for at least one i . As a consequence, $E(u, \mathcal{G}) > 0$, which concludes the proof.

Note that we omit the case of a function $u \in H_\mu^1(\mathcal{G})$ with support contained in a subgraph of the partition having trivial compact part, since this immediately entails that $E(u, \mathcal{G}) > 0$. \square

Acknowledgments

The author is grateful to Enrico Serra for many helpful suggestions during the preparation of this work and to Paolo Tilli and Riccardo Adami for several enlightening discussions.

References

- [1] R. Adami, G. Dell'Antonio, R. Figari, A. Teta, The Cauchy problem for the Schrödinger equation in dimension three with concentrated nonlinearity, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 20 (3) (2003) 477–500.
- [2] R. Adami, D. Noja, N. Visciglia, Constrained energy minimization and ground states for NLS with point defects, *Discrete Contin. Dyn. Syst. Ser. B* 18 (5) (2013) 1155–1188.
- [3] R. Adami, E. Serra, P. Tilli, NLS ground states on graphs, *Calc. Var. Partial Differential Equations* (2015), <http://dx.doi.org/10.1007/s00526-014-0804-z>, in press.
- [4] R. Adami, A. Teta, A class of nonlinear Schrödinger equations with concentrated nonlinearity, *J. Funct. Anal.* 180 (1) (2001) 148–175.
- [5] G.P. Agrawal, *Nonlinear Fiber Optics*, 4th ed., Academic Press, 2007.
- [6] M. Agueh, Sharp Gagliardo–Nirenberg inequalities and mass transport theory, *J. Dynam. Differential Equations* 18 (4) (2006) 1069–1093.
- [7] J. Bellazzini, N. Visciglia, On the orbital stability for a class of nonautonomous NLS, *Indiana Univ. Math. J.* 59 (3) (2010) 1211–1230.
- [8] H. Brezis, *Functional Analysis, Sobolev Spaces and Partial Differential Equations*, Universitext, Springer, New York, 2011.
- [9] H. Brezis, L. Nirenberg, Positive solutions of nonlinear elliptic equations involving critical Sobolev exponents, *Comm. Pure Appl. Math.* 36 (4) (1983) 437–477.
- [10] V.S. Buslaev, A.I. Komech, E.A. Kopylova, D. Stuart, On asymptotic stability of Solitary waves in Schrödinger equation coupled to nonlinear oscillator, *Comm. Partial Differential Equations* 33 (4) (2008) 669–705.
- [11] C. Cacciapuoti, D. Finco, D. Noja, A. Teta, The NLS equation in dimension one with spatially concentrated nonlinearities: the pointlike limit, *Lett. Math. Phys.* 104 (12) (2014) 1557–1570.
- [12] F. Dalfovo, S. Giorgini, L.P. Pitaevskii, S. Stringari, Theory of Bose–Einstein condensation in trapped gases, *Rev. Modern Phys.* 71 (3) (1999) 463–511.
- [13] J. Dolbeault, M.J. Esteban, A. Laptev, M. Loss, One-dimensional Gagliardo–Nirenberg–Sobolev inequalities: remarks on duality and flows, *J. Lond. Math. Soc.* 90 (2) (2014) 525–550.
- [14] L. Friedlander, Extremal properties of eigenvalues for a metric graph, *Ann. Inst. Fourier* 55 (1) (2005) 199–211.
- [15] S. Gnutzmann, U. Smilansky, Quantum graphs: applications to quantum chaos and universal spectral statistics, *Adv. Phys.* 55 (5–6) (2006) 527–625.
- [16] S. Gnutzmann, U. Smilansky, S. Derevyanko, Stationary scattering from a nonlinear network, *Phys. Rev. A* 83 (3) (2011). Article number 033831.
- [17] G. Jona-Lasinio, C. Presilla, J. Sjöstrand, On Schrödinger equation with concentrated nonlinearities, *Ann. Phys.* 240 (1) (1995) 1–21.
- [18] T. Kottos, U. Smilansky, Periodic orbit theory and spectral statistics for quantum graphs, *Ann. Phys.* 274 (1) (1999) 76–124.
- [19] P. Kuchment, Quantum graphs I. Some basic structures, *Waves Random Media* 14 (1) (2004) 107–128.
- [20] B.A. Malomed, M.Ya. Azbel, Modulational instability of a wave scattered by a nonlinear center, *Phys. Rev. B* 47 (16) (1993) 10402–10406.
- [21] F. Nier, The dynamics of some quantum open systems with short range nonlinearities, *Nonlinearity* 11 (4) (1998) 1127–1172.
- [22] F. Prinari, N. Visciglia, On a minimization problem involving the critical Sobolev exponent, *Adv. Nonlinear Stud.* 7 (4) (2007) 551–564.