



Doubly nonlocal system with Hardy–Littlewood–Sobolev critical nonlinearity

J. Giacomoni^{a,*}, T. Mukherjee^b, K. Sreenadh^b

^a Université de Pau et des Pays de l'Adour, CNRS, E2S, LMAP UMR 5142, avenue de l'université, 64013 Pau cedex, France

^b Department of Mathematics, Indian Institute of Technology Delhi, Hauz Khaz, New Delhi 110016, India

ARTICLE INFO

Article history:

Received 14 May 2018

Available online 24 July 2018

Submitted by V. Radulescu

Keywords:

Nonlocal operator

Fractional Laplacian

Choquard equation

Hardy–Littlewood–Sobolev critical exponent

ABSTRACT

This article concerns about the existence and multiplicity of weak solutions for the following nonlinear doubly nonlocal problem with critical nonlinearity in the sense of Hardy–Littlewood–Sobolev inequality

$$\begin{cases} (-\Delta)^s u = \lambda |u|^{q-2} u + \left(\int_{\Omega} \frac{|v(y)|^{2_{\mu}^*}}{|x-y|^{\mu}} dy \right) |u|^{2_{\mu}^*-2} u & \text{in } \Omega \\ (-\Delta)^s v = \delta |v|^{q-2} v + \left(\int_{\Omega} \frac{|u(y)|^{2_{\mu}^*}}{|x-y|^{\mu}} dy \right) |v|^{2_{\mu}^*-2} v & \text{in } \Omega \\ u = v = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

where Ω is a smooth bounded domain in \mathbb{R}^n , $n > 2s$, $s \in (0, 1)$, $(-\Delta)^s$ is the well known fractional Laplacian, $\mu \in (0, n)$, $2_{\mu}^* = \frac{2n-\mu}{n-2s}$ is the upper critical exponent in the Hardy–Littlewood–Sobolev inequality, $1 < q < 2$ and $\lambda, \delta > 0$ are real parameters. We study the fibering maps corresponding to the functional associated with $(P_{\lambda, \delta})$ and show that minimization over suitable subsets of Nehari manifold renders the existence of at least two non trivial solutions of $(P_{\lambda, \delta})$ for suitable range of λ and δ .

© 2018 Elsevier Inc. All rights reserved.

1. Introduction

Let $\Omega \subset \mathbb{R}^n$ be a bounded domain with smooth boundary $\partial\Omega$ (at least C^2), $n > 2s$ and $s \in (0, 1)$. We consider the following nonlinear doubly nonlocal system with critical nonlinearity:

* Corresponding author.

E-mail addresses: jacques.giacomoni@univ-pau.fr (J. Giacomoni), tulimukh@gmail.com (T. Mukherjee), sreenadh@gmail.com (K. Sreenadh).

$$(P_{\lambda,\delta}) \begin{cases} (-\Delta)^s u = \lambda |u|^{q-2} u + \left(\int_{\Omega} \frac{|v(y)|^{2_\mu^*}}{|x-y|^\mu} dy \right) |u|^{2_\mu^*-2} u & \text{in } \Omega \\ (-\Delta)^s v = \delta |v|^{q-2} v + \left(\int_{\Omega} \frac{|u(y)|^{2_\mu^*}}{|x-y|^\mu} dy \right) |v|^{2_\mu^*-2} v & \text{in } \Omega \\ u = v = 0 & \text{in } \mathbb{R}^n \setminus \Omega, \end{cases}$$

where Ω is a smooth bounded domain in \mathbb{R}^n , $n > 2s$, $s \in (0, 1)$, $\mu \in (0, n)$, $2_\mu^* = \frac{2n-\mu}{n-2s}$ is the upper critical exponent in the Hardy–Littlewood–Sobolev inequality, $1 < q < 2$, $\lambda, \delta > 0$ are real parameters and $(-\Delta)^s$ is the fractional Laplace operator defined as

$$(-\Delta)^s u(x) = 2C_s^n \text{P.V.} \int_{\mathbb{R}^n} \frac{u(x) - u(y)}{|x-y|^{n+2s}} dy$$

where P.V. denotes the Cauchy principal value and $C_s^n = \pi^{-\frac{n}{2}} 2^{2s-1} s \frac{\Gamma(\frac{n+2s}{2})}{\Gamma(1-s)}$, Γ being the Gamma function. The fractional Laplacian is the infinitesimal generator of Lévy stable diffusion process and arise in anomalous diffusion in plasma, population dynamics, geophysical fluid dynamics, flames propagation, chemical reactions in liquids and American options in finance, see [3] for instance.

In the local case, authors in [5] studied the existence of ground states for the nonlinear Choquard equation

$$-\Delta u + V(x)u = \left(\int_{\Omega} \frac{|u(y)|^p}{|x-y|^\mu} dy \right) |u|^{p-2} u \quad \text{in } \mathbb{R}^n, \quad (1.1)$$

where $p > 1$ and $n \geq 3$. Recently, Ghimenti, Moroz and Schaftingen [16] proved the existence of least action nodal solution for the problem

$$-\Delta u + u = (I_\alpha * |u|^2)u \quad \text{in } \mathbb{R}^n,$$

where $*$ denotes the convolution and I_α denotes the Riesz potential. Further results related to Choquard equations can be found in the survey paper [24] and the references therein. Alves, Figueiredo and Yang [1] proved existence of a nontrivial solution via penalization method for the following Choquard equation

$$-\Delta u + V(x)u = (|x|^{-\mu} * F(u))f(u) \quad \text{in } \mathbb{R}^n,$$

where $0 < \mu < N$, $N = 3$, V is a continuous real valued function and F is the primitive of function f . In the nonlocal case, Choquard equations involving fractional Laplacian is a recent topic of research. Authors in [9] obtained regularity, existence, nonexistence, symmetry as well as decays properties for the problem

$$(-\Delta)^s u + \omega u = (|x|^{\alpha-n} * |u|^p) |u|^{p-2} u \quad \text{in } \mathbb{R}^n,$$

where $\omega > 0$, $p > 1$ and $s \in (0, 1)$. Fractional Choquard equations also known as nonlinear fractional Schrödinger equations with Hartree-type nonlinearity arise in the study of mean field limit of weakly interacting molecules, physics of multi particle systems and the quantum mechanical theory, etc. These are recently studied by some authors in [6,10,22]. We also refer [2] and [4] for the qualitative behaviour of Choquard type problems in the semiclassical limit.

Concerning the boundary value problems involving the Choquard nonlinearity, the Brezis–Nirenberg type problem that is

$$-\Delta u = \lambda u + \left(\int_{\Omega} \frac{|u|^{2^*_\mu}}{|x-y|^\mu} dy \right) |u|^{2^*_\mu-2} u \text{ in } \Omega, \quad u = 0 \text{ in } \mathbb{R}^n \setminus \Omega$$

where Ω is bounded domain in \mathbb{R}^n , was studied by Gao and Yang in [14]. They proved the existence, multiplicity and nonexistence results for a range of λ . Moreover, in [13] authors proved the existence results for a class of critical Choquard equations in critical case. Among the very recent works, we cite [29] where Shen, Gao and Yang obtained the existence of multiple solutions for non-homogenous critical Choquard equation using the variational methods when $0 < \lambda < \lambda_1$, where λ_1 denotes the first eigenvalue of $-\Delta$ with Dirichlet boundary condition.

Coming to the system of equations, elliptic systems involving fractional Laplacian and homogeneous nonlinearity have been studied in [17,20,11] using variational methods. We also refer [23] to readers for a detailed study on variational methods for fractional elliptic problems. Guo et al. in [19] studied a nonlocal system involving fractional Sobolev critical exponent and fractional Laplacian. We also cite [8,12,32] as some very recent works on the study of fractional elliptic systems and [33] and [34] for Schrödinger type nonlocal equations and [18] for problems involving singular nonlinearities with critical growth. However there is a very few literature available on fractional elliptic system involving Choquard type nonlinearity and fractional elliptic system with critical Choquard inequality has not been studied yet, to the best of our knowledge.

In this present paper, we discuss the existence and multiplicity result for the problem $(P_{\lambda,\delta})$. We seek help of the Nehari manifold techniques where minimization over suitable components of Nehari manifold provide the weak solution to the problem. We divide the problem into two cases that is $0 < \mu \leq 4s$ and $\mu > 4s$ and show existence of at least two solutions while bounding the parameters λ and δ optimally. The existence results in the first case are completely new for the system case and optimal in the sense of obtaining the constant Θ (defined in Lemma 3.3). We also reach the expected first critical level that is

$$I_{\lambda,\delta}(u_1, v_1) + \frac{n - \mu + 2s}{2n - \mu} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2n-\mu}{n-\mu+2s}}$$

where (u_1, v_1) denote the first solution of $(P_{\lambda,\delta})$, in this case (see Lemma 4.9) analogously to the local setting case (refer Lemma 2.4 in [13]). Whereas in the latter case, we obtain the multiplicity for a smaller range of λ and δ that is Θ_0 (defined in Theorem 4.13) as compared to Θ . We use an accurate blow up analysis involving the minimizers of the embeddings to achieve the goal. In the case $0 < \mu \leq 4s$, our results are sharp in the sense that the restrictions on the parameters λ and δ are used only to show that Nehari set is a manifold. Moreover using an iterative scheme, regularity results known for nonlocal problems involving fractional laplacian and strong maximum principle, we show the existence of a positive solution (see Proposition 4.8).

Theorem 1.1. *Assume $1 < q < 2$ and $0 < \mu < n$ then there exist positive constants Θ and Θ_0 such that*

1. *if $\mu \leq 4s$ and $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$, the system $(P_{\lambda,\delta})$ admits at least two nontrivial solutions,*
2. *if $\mu > 4s$ and $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta_0$, the system $(P_{\lambda,\delta})$ admits at least two nontrivial solutions.*

Moreover, there exists a positive solution for $(P_{\lambda,\delta})$.

Remark 1.2. We remark that the solution obtained for $(P_{\lambda,\delta})$ (other than the positive solution) is not even semi trivial. The proof follows along the same line as section 5 (p. 841) of [7].

Our paper is organized as follows: Section 2 contains the functional setting and various asymptotic estimates involving minimizers of best constants. We analyse the fibering maps associated to the Nehari

manifold in section 3. Lastly, section 4 contains the proof of main result where we show the existence of at least two non trivial solutions.

2. Function spaces and some asymptotic estimates

Consider the function space $H^s(\Omega)$ as the usual fractional Sobolev space $W^{s,2}(\Omega)$ defined by

$$H^s(\Omega) = \left\{ u \in L^2(\Omega) : \int_{\Omega} \int_{\Omega} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy < +\infty \right\}.$$

Setting $Q := \mathbb{R}^{2n} \setminus (\mathcal{C}\Omega \times \mathcal{C}\Omega)$ where $\mathcal{C}\Omega = \mathbb{R}^n \setminus \Omega$, we define the Banach space

$$X := \left\{ u : \mathbb{R}^n \rightarrow \mathbb{R} \text{ measurable} : u \in L^2(\Omega), \int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy < +\infty \right\}$$

with the norm defined as

$$\|u\|_X := \|u\|_{L^2(\Omega)} + \left(\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy \right)^{\frac{1}{2}} = \|u\|_{L^2(\Omega)} + \left(\frac{1}{C_s^n} \int_{\Omega} u(-\Delta)^s u \, dx dy \right)^{\frac{1}{2}}.$$

If we set $X_0 := \{u \in X : u = 0 \text{ in } \mathbb{R}^n \setminus \Omega\}$, then it can be shown that X_0 forms a Hilbert space with the inner product

$$\langle u, v \rangle = \int_Q \frac{(u(x) - u(y))(v(x) - v(y))}{|x - y|^{n+2s}} dx dy$$

for $u, v \in X_0$ and thus the corresponding norm is

$$\|u\|_{X_0} = \|u\| := \left(\int_Q \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy \right)^{\frac{1}{2}}.$$

Then X_0 can be equivalently considered as completion of $C_0^\infty(\Omega)$ under the norm $\|\cdot\|_X$. It holds that $X_0 \hookrightarrow L^r(\Omega)$ continuously for $r \in [1, 2_s^*]$ and compactly for $r \in [1, 2_s^*)$, where $2_s^* = \frac{2n}{n-2s}$. Now consider the product space $Y := X_0 \times X_0$ endowed with the norm $\|(u, v)\|^2 := \|u\|^2 + \|v\|^2$. Before defining the weak solution for $(P_{\lambda,\delta})$, we need to certify that whenever $u \in X_0$, the term

$$\int_{\Omega} (|x|^{-\mu} * |u|^{2_\mu^*}) |u|^{2_\mu^*} dx = \int_{\Omega} \int_{\Omega} \frac{|u(x)|^{2_\mu^*} |v(y)|^{2_\mu^*}}{|x - y|^\mu} dx dy$$

is well defined. This is certified by the following well known Hardy–Littlewood–Sobolev inequality.

Proposition 2.1. (Hardy–Littlewood–Sobolev inequality) [p. 106, Theorem 4.3, [21]] *Let $t, r > 1$ and $0 < \mu < n$ with $1/t + \mu/n + 1/r = 2$, $f \in L^t(\mathbb{R}^n)$ and $h \in L^r(\mathbb{R}^n)$. There exists a sharp constant $C(t, n, \mu, r)$, independent of f, h such that*

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{f(x)h(y)}{|x-y|^\mu} dx dy \leq C(t, n, \mu, r) \|f\|_{L^t(\mathbb{R}^n)} \|h\|_{L^r(\mathbb{R}^n)}. \quad (2.1)$$

If $t = r = \frac{2n}{2n-\mu}$ then

$$C(t, n, \mu, r) = C(n, \mu) = \pi^{\frac{\mu}{2}} \frac{\Gamma(\frac{n}{2} - \frac{\mu}{2})}{\Gamma(n - \frac{\mu}{2})} \left\{ \frac{\Gamma(\frac{n}{2})}{\Gamma(n)} \right\}^{-1 + \frac{\mu}{n}}.$$

In this case there is equality in (2.1) if and only if $f \equiv (\text{constant})h$ and

$$h(x) = A(\gamma^2 + |x - a|^2)^{-\frac{(2n-\mu)}{2}}$$

for some $A \in \mathbb{C}$, $0 \neq \gamma \in \mathbb{R}$ and $a \in \mathbb{R}^n$.

Remark 2.2. For $u \in H^s(\mathbb{R}^n)$, if we let $f = h = |u|^p$ then by Hardy–Littlewood–Sobolev inequality,

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x)|^p |u(y)|^p}{|x-y|^\mu} dx dy$$

is well defined for all p satisfying

$$2_\mu := \left(\frac{2n-\mu}{n} \right) \leq p \leq \left(\frac{2n-\mu}{n-2s} \right) := 2_\mu^*.$$

Next result is a basic inequality whose proof can be worked out in a similar manner as proof of Proposition 3.2(3.3) of [15].

Lemma 2.3. For $u, v \in L^{\frac{2n}{2n-\mu}}(\mathbb{R}^n)$, we have

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x)|^p |v(y)|^p}{|x-y|^\mu} dx dy \leq \left(\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x)|^p |u(y)|^p}{|x-y|^\mu} dx dy \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|v(x)|^p |v(y)|^p}{|x-y|^\mu} dx dy \right)^{\frac{1}{2}},$$

where $\mu \in (0, n)$ and $p \in [2_\mu, 2_\mu^*]$.

Proof. We recall the semigroup property of the Riesz potential which states that if $I_\alpha : \mathbb{R}^n \rightarrow \mathbb{R}$ denotes the Riesz potential given by

$$I_\alpha(x) = \frac{A_\alpha}{|x|^{n-\alpha}}, \text{ where } x \in \mathbb{R}^n \setminus \{0\} \text{ and } A_\alpha = \frac{\Gamma(\frac{n-\alpha}{2})}{\Gamma(\frac{\alpha}{2}) \pi^{n/2} 2^\alpha}.$$

Then I_α satisfies $I_\alpha = I_{\alpha/2} * I_{\alpha/2}$. Using this along with Hölder's inequality we obtain

$$\begin{aligned}
& \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x)|^p |v(y)|^p}{|x-y|^\mu} dx dy \\
&= \frac{1}{A_{n-\mu}} \int_{\mathbb{R}^n} (I_{n-\mu} * |u|^p) |v|^p dx = \frac{1}{A_{n-\mu}} \int_{\mathbb{R}^n} (I_{\frac{n-\mu}{2}} * |u|^p) (I_{\frac{n-\mu}{2}} * |v|^p) dx \\
&\leq \frac{1}{A_{n-\mu}} \left(\int_{\mathbb{R}^n} (I_{\frac{n-\mu}{2}} * |u|^p)^2 dx \right)^{1/2} \left(\int_{\mathbb{R}^n} (I_{\frac{n-\mu}{2}} * |v|^p)^2 dx \right)^{1/2} \\
&= \left(\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|u(x)|^p |u(y)|^p}{|x-y|^\mu} dx dy \right)^{\frac{1}{2}} \left(\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|v(x)|^p |v(y)|^p}{|x-y|^\mu} dx dy \right)^{\frac{1}{2}}. \quad \square
\end{aligned}$$

Therefore, it easily follows using Lemma 2.3 that for every $(u, v) \in Y$, $\int_{\Omega} (|x|^{-\mu} * |u|^{2_\mu^*}) |v|^{2_\mu^*} dx < +\infty$.

In the context of Hardy–Littlewood–Sobolev inequality that is Proposition 2.1, for any $u \in X_0$ we get a constant $C > 0$ such that

$$\int_{\Omega} (|x|^{-\mu} * |u|^{2_\mu^*}) |u|^{2_\mu^*} dx = \int_{\Omega} \int_{\Omega} \frac{|u(x)|^{2_\mu^*} |u(y)|^{2_\mu^*}}{|x-y|^\mu} dx dy \leq C \|u\|_{L^{2_\mu^*}(\Omega)}^{22_\mu^*}. \quad (2.2)$$

For notational convenience, if $u, v \in X_0$ we set

$$B(u, v) := \int_{\Omega} (|x|^{-\mu} * |u|^{2_\mu^*}) |v|^{2_\mu^*} dx.$$

Definition 2.4. We say that $(u, v) \in Y$ is a weak solution to $(P_{\lambda, \delta})$ if for every $(\phi, \psi) \in Y$, it satisfies

$$\begin{aligned}
C_s^n(\langle u, \phi \rangle + \langle v, \psi \rangle) &= \int_{\Omega} (\lambda |u|^{q-2} u \phi + \delta |v|^{q-2} v \psi) dx \\
&+ \int_{\Omega} (|x|^{-\mu} * |v|^{2_\mu^*}) |u|^{2_\mu^*-2} u \phi dx + \int_{\Omega} (|x|^{-\mu} * |u|^{2_\mu^*}) |v|^{2_\mu^*-2} v \psi dx.
\end{aligned}$$

Equivalently, if we define the functional $I_{\lambda, \delta} : Y \rightarrow \mathbb{R}$ as

$$I_{\lambda, \delta}(u) := \frac{C_s^n}{2} \|(u, v)\|^2 - \frac{1}{q} \int_{\Omega} (\lambda |u|^q + \delta |v|^q) - \frac{2}{22_\mu^*} B(u, v)$$

then the critical points of $I_{\lambda, \delta}$ correspond to the weak solutions of $(P_{\lambda, \delta})$. A direct computation leads to $I_{\lambda, \delta} \in C^1(Y, \mathbb{R})$ such that for any $(\phi, \psi) \in Y$

$$\begin{aligned}
(I'_{\lambda, \delta}(u, v), (\phi, \psi)) &= C_s^n(\langle u, \phi \rangle + \langle v, \psi \rangle) - \int_{\Omega} (\lambda |u|^{q-2} u \phi + \delta |v|^{q-2} v \psi) dx \\
&- \int_{\Omega} (|x|^{-\mu} * |v|^{2_\mu^*}) |u|^{2_\mu^*-2} u \phi dx - \int_{\Omega} (|x|^{-\mu} * |u|^{2_\mu^*}) |v|^{2_\mu^*-2} v \psi dx.
\end{aligned} \quad (2.3)$$

We define

$$S_s = \inf_{u \in X_0 \setminus \{0\}} \frac{\int_{\mathbb{R}^{2n}} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy}{\left(\int_{\mathbb{R}^n} |u|^{2_s^*} dx \right)^{2/2_s^*}} = \inf_{u \in X_0 \setminus \{0\}} \frac{\|u\|^2}{\|u\|_{L^{2_s^*}(\mathbb{R}^n)}^2}.$$

Consider the family of functions $\{U_\epsilon\}$ defined as

$$U_\epsilon(x) = \epsilon^{-\frac{(n-2s)}{2}} u^*\left(\frac{x}{\epsilon}\right), \quad x \in \mathbb{R}^n \quad (2.4)$$

where $u^*(x) = \bar{u}\left(\frac{x}{S_s^{2s}}\right)$, $\bar{u}(x) = \frac{\tilde{u}(x)}{\|\tilde{u}\|_{L^{2_s^*}(\mathbb{R}^n)}}$ and $\tilde{u}(x) = \alpha(\beta^2 + |x|^2)^{-\frac{n-2s}{2}}$ with $\alpha \in \mathbb{R} \setminus \{0\}$ and $\beta > 0$ are fixed constants. Then for each $\epsilon > 0$, U_ϵ satisfies

$$(-\Delta)^s u = |u|^{2_s^*-2} u \quad \text{in } \mathbb{R}^n$$

and verifies the equality

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|U_\epsilon(x) - U_\epsilon(y)|^2}{|x - y|^{n+2s}} dx dy = \int_{\mathbb{R}^n} |U_\epsilon|^{2_s^*} dx = S_s^{\frac{n}{2s}}. \quad (2.5)$$

For a proof, we refer to [28]. Next, in spirit of the inequality (2.2) we define the best constant

$$S_s^H := \inf_{u \in X_0 \setminus \{0\}} \frac{\int_{\mathbb{R}^{2n}} \frac{|u(x) - u(y)|^2}{|x - y|^{n+2s}} dx dy}{\left(\int_{\mathbb{R}^n} (|x|^{-\mu} * |u|^{2_s^*}) |u|^{2_s^*} dx \right)^{\frac{1}{2_s^*}}} = \inf_{u \in X_0 \setminus \{0\}} \frac{\|u\|^2}{B(u, u)^{\frac{1}{2_s^*}}}.$$

Lemma 2.5. *The constant S_s^H is achieved by u if and only if u is of the form*

$$C \left(\frac{t}{t^2 + |x - x_0|^2} \right)^{\frac{n-2s}{2}}, \quad x \in \mathbb{R}^n$$

for some $x_0 \in \mathbb{R}^n$, $C > 0$ and $t > 0$. Moreover,

$$S_s^H = \frac{S_s}{C(n, \mu)^{\frac{1}{2_s^*}}}. \quad (2.6)$$

Proof. By the Hardy–Littlewood–Sobolev inequality we easily get that

$$S_s^H \geq \frac{S_s}{C(n, \mu)^{\frac{1}{2_s^*}}}.$$

Also from Proposition 2.1 we know that the inequality in (2.1) is an equality if and only if u is of the form

$$C \left(\frac{t}{t^2 + |x - x_0|^2} \right)^{\frac{n-2s}{2}}, \quad x \in \mathbb{R}^n.$$

While we know that if u is of this form then it also forms a minimizer for the constant S_s , thus we obtain the result and (2.6) follows directly. \square

We set

$$\tilde{S}_s^H = \inf_{(u,v) \in Y \setminus \{(0,0)\}} \frac{\|(u,v)\|^2}{\left(\int_{\Omega} (|x|^{-\mu} * |u|^{2_\mu^*}) |v|^{2_\mu^*} dx\right)^{\frac{1}{2_\mu^*}}} = \inf_{(u,v) \in Y \setminus \{(0,0)\}} \frac{\|(u,v)\|^2}{B(u,v)^{\frac{1}{2_\mu^*}}}$$

and show the relation between S_s^H and \tilde{S}_s^H in the following lemma. The argument follows closely the line of Lemma 3.3 of [7] but for sake of completeness, we include it here.

Lemma 2.6. *There holds $\tilde{S}_s^H = 2S_s^H$.*

Proof. Let $\{g_k\} \subset X_0$ be a minimizing sequence for S_s^H . Let $r_1, r_2 > 0$ be specified later and set the sequences $u_k = r_1 g_k$ and $v_k = r_2 g_k$ in X_0 . From the definition of S_s^H we have

$$\tilde{S}_s^H \leq \left(\frac{r_1^2 + r_2^2}{r_1 r_2}\right) \left(\frac{\|g_k\|^2}{B(g_k, g_k)^{\frac{1}{2_\mu^*}}}\right) = \left(\frac{r_1}{r_2} + \frac{r_2}{r_1}\right) \left(\frac{\|g_k\|^2}{B(g_k, g_k)^{\frac{1}{2_\mu^*}}}\right). \quad (2.7)$$

Let us define the function $f: \mathbb{R}^+ \rightarrow \mathbb{R}^+$ by setting $f(x) = x + x^{-1}$. Then it is easy to see that f attains its minimum at $x_0 = 1$ with the minimum value $f(1) = 2$. We choose r_1, r_2 in (2.7) such that $r_1 = r_2$ and letting $k \rightarrow \infty$ in (2.7) we get

$$\tilde{S}_s^H \leq 2S_s^H. \quad (2.8)$$

To prove the reverse inequality we consider the minimizing sequence $\{(u_k, v_k)\} \subset Y \setminus \{(0,0)\}$ for \tilde{S}_s^H . We set $w_k = r_k v_k$ for $r_k > 0$ with $B(u_k, u_k) = B(w_k, w_k)$. This along with Lemma 2.3 gives

$$B(u_k, w_k) \leq B(u_k, u_k)^{\frac{1}{2}} B(w_k, w_k)^{\frac{1}{2}} = B(u_k, u_k) = B(w_k, w_k).$$

Thus we obtain

$$\begin{aligned} \frac{\|(u_k, v_k)\|^2}{B(u_k, v_k)^{\frac{1}{2_\mu^*}}} &= r_k \frac{\|(u_k, v_k)\|^2}{B(u_k, w_k)^{\frac{1}{2_\mu^*}}} \geq r_k \frac{\|u_k\|^2}{B(u_k, u_k)^{\frac{1}{2_\mu^*}}} + r_k r_k^{-2} \frac{\|w_k\|^2}{B(w_k, w_k)^{\frac{1}{2_\mu^*}}} \\ &\geq f(r_k) S_s^H \geq 2S_s^H. \end{aligned}$$

Now passing on the limit as $k \rightarrow \infty$ we get

$$2S_s^H \leq \tilde{S}_s^H. \quad (2.9)$$

Finally from (2.8) and (2.9) it follows that $S_s^H = 2\tilde{S}_s^H$. \square

We recall the definition of U_ϵ from (2.4). Without loss of generality, we assume $0 \in \Omega$ and fix $\delta > 0$ such that $B_{4\delta} \subset \Omega$. Let $\eta \in C^\infty(\mathbb{R}^n)$ be such that $0 \leq \eta \leq 1$ in \mathbb{R}^n , $\eta \equiv 1$ in B_δ and $\eta \equiv 0$ in $\mathbb{R}^n \setminus B_{2\delta}$. For $\epsilon > 0$, we denote by u_ϵ the following function

$$u_\epsilon(x) = \eta(x) U_\epsilon(x),$$

for $x \in \mathbb{R}^n$, where U_ϵ is defined in section 2. We have the following results for u_ϵ from Proposition 21 and 22 of [28].

Proposition 2.7. Let $s \in (0, 1)$ and $n > 2s$. Then, the following estimates hold true as $\epsilon \rightarrow 0$

$$\begin{aligned} \text{(i)} \quad & \int_{\mathbb{R}^{2n}} \frac{|u_\epsilon(x) - u_\epsilon(y)|^2}{|x - y|^{n+2s}} \, dx dy \leq S_s^{\frac{n}{2s}} + O(\epsilon^{n-2s}), \\ \text{(ii)} \quad & \int_{\Omega} |u_\epsilon|^{2^*} \, dx = S_s^{\frac{n}{2s}} + O(\epsilon^n), \\ \text{(iii)} \quad & \end{aligned}$$

$$\int_{\mathbb{R}^n} |u_\epsilon(x)|^2 \, dx \geq \begin{cases} C_s \epsilon^{2s} + O(\epsilon^{n-2s}) & \text{if } n > 4s \\ C_s \epsilon^{2s} |\log \epsilon| + O(\epsilon^{2s}) & \text{if } n = 4s \\ C_s \epsilon^{n-2s} + O(\epsilon^{2s}) & \text{if } n < 4s \end{cases},$$

for some positive constant C_s , depending on s .

Using (2.6), Proposition 2.7(i) can be written as

$$\int_{\mathbb{R}^n} \frac{|u_\epsilon(x) - u_\epsilon(y)|^2}{|x - y|^{n+2s}} \, dx dy \leq S_s^{\frac{n}{2s}} + O(\epsilon^{n-2s}) = \left((C(n, \mu))^{\frac{n-2s}{2n-\mu}} (S_s^H)^{\frac{n}{2s}} \right)^{\frac{n}{2s}} + O(\epsilon^{n-2s}). \quad (2.10)$$

Proposition 2.8. The following estimates hold true:

$$\left(\int_{\Omega} \int_{\Omega} \frac{|u_\epsilon(x)|^{2^*} |u_\epsilon(y)|^{2^*}}{|x - y|^\mu} \, dx dy \right)^{\frac{n-2s}{2n-\mu}} \leq (C(n, \mu))^{\frac{n(n-2s)}{2s(2n-\mu)}} (S_s^H)^{\frac{n-2s}{2s}} + O(\epsilon^n),$$

and

$$\left(\int_{\Omega} \int_{\Omega} \frac{|u_\epsilon(x)|^{2^*} |u_\epsilon(y)|^{2^*}}{|x - y|^\mu} \, dx dy \right)^{\frac{n-2s}{2n-\mu}} \geq \left((C(n, \mu))^{\frac{n}{2s}} (S_s^H)^{\frac{2n-\mu}{2s}} - O(\epsilon^n) \right)^{\frac{n-2s}{2n-\mu}}.$$

Proof. By Hardy–Littlewood–Sobolev inequality, Proposition 2.7(ii) and 2.6, we get

$$\begin{aligned} & \left(\int_{\Omega} \int_{\Omega} \frac{|u_\epsilon(x)|^{2^*} |u_\epsilon(y)|^{2^*}}{|x - y|^\mu} \, dx dy \right)^{\frac{n-2s}{2n-\mu}} \\ & \leq (C(n, \mu))^{\frac{n-2s}{2n-\mu}} \|u_\epsilon\|_{L^{2^*}_s(\Omega)}^2 = (C(n, \mu))^{\frac{n-2s}{2n-\mu}} \left(S_s^{\frac{n}{2s}} + O(\epsilon^n) \right)^{\frac{n-2s}{n}} \\ & = (C(n, \mu))^{\frac{n-2s}{2n-\mu}} \left((C(n, \mu))^{\frac{n(n-2s)}{2s(2n-\mu)}} (S_s^H)^{\frac{n}{2s}} + O(\epsilon^n) \right)^{\frac{n-2s}{n}} \\ & = (C(n, \mu))^{\frac{n(n-2s)}{2s(2n-\mu)}} (S_s^H)^{\frac{n-2s}{2s}} + O(\epsilon^n). \end{aligned}$$

Next, we consider

$$\begin{aligned}
& \int_{\Omega} \int_{\Omega} \frac{|u_{\epsilon}(x)|^{2_{\mu}^*} |u_{\epsilon}(y)|^{2_{\mu}^*}}{|x-y|^{\mu}} \, dx dy \\
& \geq \int_{B_{\delta}} \int_{B_{\delta}} \frac{|u_{\epsilon}(x)|^{2_{\mu}^*} |u_{\epsilon}(y)|^{2_{\mu}^*}}{|x-y|^{\mu}} \, dx dy = \int_{B_{\delta}} \int_{B_{\delta}} \frac{|U_{\epsilon}(x)|^{2_{\mu}^*} |U_{\epsilon}(y)|^{2_{\mu}^*}}{|x-y|^{\mu}} \, dx dy \\
& = \int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|U_{\epsilon}(x)|^{2_{\mu}^*} |U_{\epsilon}(y)|^{2_{\mu}^*}}{|x-y|^{\mu}} \, dx dy - 2 \int_{\mathbb{R}^n \setminus B_{\delta}} \int_{B_{\delta}} \frac{|U_{\epsilon}(x)|^{2_{\mu}^*} |U_{\epsilon}(y)|^{2_{\mu}^*}}{|x-y|^{\mu}} \, dx dy \\
& \quad - \int_{\mathbb{R}^n \setminus B_{\delta}} \int_{\mathbb{R}^n \setminus B_{\delta}} \frac{|U_{\epsilon}(x)|^{2_{\mu}^*} |U_{\epsilon}(y)|^{2_{\mu}^*}}{|x-y|^{\mu}} \, dx dy.
\end{aligned} \tag{2.11}$$

We estimate the integrals in right hand side of (2.11) separately. Firstly to estimate the first integral, by Lemma 2.5 we get that $\{U_{\epsilon}\}$ forms minimizers of S_s^H . Therefore using (2.5) we get

$$\int_{\mathbb{R}^n} \int_{\mathbb{R}^n} \frac{|U_{\epsilon}(x)|^{2_{\mu}^*} |U_{\epsilon}(y)|^{2_{\mu}^*}}{|x-y|^{\mu}} \, dx dy = \left(\frac{\|U_{\epsilon}\|^2}{S_s^H} \right)^{2_{\mu}^*} = \left(\frac{S_s^{n/2s}}{S_s^H} \right)^{2_{\mu}^*} = C(n, \mu)^{n/2s} (S_s^H)^{\frac{2n-\mu}{2s}}. \tag{2.12}$$

Secondly, consider

$$\begin{aligned}
& \int_{\mathbb{R}^n \setminus B_{\delta}} \int_{B_{\delta}} \frac{|U_{\epsilon}(x)|^{2_{\mu}^*} |U_{\epsilon}(y)|^{2_{\mu}^*}}{|x-y|^{\mu}} \, dx dy \\
& \leq C_{2,s} \int_{\mathbb{R}^n \setminus B_{\delta}} \int_{B_{\delta}} \frac{\epsilon^{\mu-2n}}{|x-y|^{\mu} (1+|\frac{x}{\epsilon}|^2)^{\frac{2n-\mu}{2}} (1+|\frac{y}{\epsilon}|^2)^{\frac{2n-\mu}{2}}} \, dx dy \\
& = \epsilon^{2n-\mu} C_{2,s} \int_{\mathbb{R}^n \setminus B_{\delta}} \int_{B_{\delta}} \frac{1}{|x-y|^{\mu} (\epsilon^2 + |x|^2)^{\frac{2n-\mu}{2}} (\epsilon^2 + |y|^2)^{\frac{2n-\mu}{2}}} \, dx dy
\end{aligned}$$

where $C_{2,s}$ is an appropriate positive constant. Let $D := B_{\delta} \times (\mathbb{R}^n \setminus B_{\delta})$ then

$$\begin{aligned}
& \epsilon^{2n-\mu} C_{2,s} \int_{\mathbb{R}^n \setminus B_{\delta}} \int_{B_{\delta}} \frac{1}{|x-y|^{\mu} (\epsilon^2 + |x|^2)^{\frac{2n-\mu}{2}} (\epsilon^2 + |y|^2)^{\frac{2n-\mu}{2}}} \, dx dy \\
& = \epsilon^{2n-\mu} C_{2,s} \left(\int_{D \cap \{|x-y| \leq 1\}} + \int_{D \cap \{|x-y| > 1\}} \right) \frac{1}{|x-y|^{\mu} (\epsilon^2 + |x|^2)^{\frac{2n-\mu}{2}} (\epsilon^2 + |y|^2)^{\frac{2n-\mu}{2}}} \, dx dy.
\end{aligned}$$

Consider

$$\begin{aligned}
& \epsilon^{2n-\mu} C_{2,s} \int_{D \cap \{|x-y| > 1\}} \frac{1}{|x-y|^{\mu} (\epsilon^2 + |x|^2)^{\frac{2n-\mu}{2}} (\epsilon^2 + |y|^2)^{\frac{2n-\mu}{2}}} \, dx dy \\
& \leq \epsilon^{2n-\mu} C_{2,s} \int_{D \cap \{|x-y| > 1\}} \frac{1}{(\epsilon^2 + |x|^2)^{\frac{2n-\mu}{2}} (\epsilon^2 + |y|^2)^{\frac{2n-\mu}{2}}} \, dx dy
\end{aligned}$$

$$\begin{aligned}
&\leq \epsilon^{2n-\mu} C_{2,s} \int_{B_\delta} \frac{dx}{(\epsilon^2 + |x|^2)^{\frac{2n-\mu}{2}}} \int_{\mathbb{R}^n \setminus B_\delta} \frac{dy}{(\epsilon^2 + |y|^2)^{\frac{2n-\mu}{2}}} \\
&\leq \epsilon^{2n-\mu} C_{2,s} \int_0^{\delta/\epsilon} \frac{\epsilon^{\mu-n} t^{n-1} dt}{(1+t^2)^{\frac{2n-\mu}{2}}} \int_{\mathbb{R}^n \setminus B_\delta} \frac{dy}{(|y|^2)^{\frac{2n-\mu}{2}}} = O(\epsilon^n).
\end{aligned}$$

Next we observe that the set $D \cap \{|x-y| < 1\}$ is bounded and if $x, y \in D \cap \{|x-y| < 1\}$ then there exist constants $\alpha, \beta > 0$ such that $\alpha \leq |x|, |y| \leq \beta$. This implies that

$$\begin{aligned}
&\epsilon^{2n-\mu} C_{2,s} \int_{D \cap \{|x-y| < 1\}} \frac{1}{|x-y|^\mu (\epsilon^2 + |x|^2)^{\frac{2n-\mu}{2}} (\epsilon^2 + |y|^2)^{\frac{2n-\mu}{2}}} dx dy \\
&\leq \epsilon^{2n-\mu} C_{2,s} \int_{D \cap \{|x-y| < 1\}} \frac{1}{|x-y|^\mu (|x|^2)^{\frac{2n-\mu}{2}} (|y|^2)^{\frac{2n-\mu}{2}}} dx dy \\
&\leq O(\epsilon^{2n-\mu}) \int_{D \cap \{|x-y| < 1\}} \frac{1}{|x-y|^\mu} dx dy = O(\epsilon^{2n-\mu})
\end{aligned}$$

since $\mu \in (0, n)$. Therefore

$$\int_{\mathbb{R}^n \setminus B_\delta} \int_{B_\delta} \frac{|U_\epsilon(x)|^{2_\mu^*} |U_\epsilon(y)|^{2_\mu^*}}{|x-y|^\mu} dx dy \leq O(\epsilon^n). \quad (2.13)$$

Lastly, in a similar manner we have

$$\begin{aligned}
&\int_{\mathbb{R}^n \setminus B_\delta} \int_{\mathbb{R}^n \setminus B_\delta} \frac{|U_\epsilon(x)|^{2_\mu^*} |U_\epsilon(y)|^{2_\mu^*}}{|x-y|^\mu} dx dy \\
&\leq C'_{2,s} \int_{\mathbb{R}^n \setminus B_\delta} \int_{\mathbb{R}^n \setminus B_\delta} \frac{\epsilon^{\mu-2n}}{|x-y|^\mu (1 + |\frac{x}{\epsilon}|^2)^{\frac{2n-\mu}{2}} (1 + |\frac{y}{\epsilon}|^2)^{\frac{2n-\mu}{2}}} dx dy \\
&= \epsilon^{2n-\mu} C'_{2,s} \int_{\mathbb{R}^n \setminus B_\delta} \int_{\mathbb{R}^n \setminus B_\delta} \frac{1}{|x-y|^\mu (\epsilon^2 + |x|^2)^{\frac{2n-\mu}{2}} (\epsilon^2 + |y|^2)^{\frac{2n-\mu}{2}}} dx dy \\
&\leq \epsilon^{2n-\mu} C'_{2,s} \int_{\mathbb{R}^n \setminus B_\delta} \int_{\mathbb{R}^n \setminus B_\delta} \frac{1}{|x-y|^\mu |x|^{2n-\mu} |y|^{2n-\mu}} dx dy = O(\epsilon^{2n-\mu})
\end{aligned} \quad (2.14)$$

where $C'_{2,s}$ is an appropriate positive constant. Using the estimates (2.12), (2.13) and (2.14) in (2.11), we get

$$\left(\int_{\Omega} \int_{\Omega} \frac{|u_\epsilon(x)|^{2_\mu^*} |u_\epsilon(y)|^{2_\mu^*}}{|x-y|^\mu} dx dy \right)^{\frac{n-2s}{2n-\mu}} \geq \left((C(n, \mu))^{\frac{n}{2s}} (S_s^H)^{\frac{2n-\mu}{2s}} - O(\epsilon^n) \right)^{\frac{n-2s}{2n-\mu}}.$$

This completes the proof. \square

3. Analysis of fibering maps

In this section we study the fibering maps and establish some preliminaries for the Nehari manifold. It is easy to see that the energy functional $I_{\lambda,\delta}$ is not bounded below on the whole domain Y , so we minimize $I_{\lambda,\delta}$ over proper subsets of the Nehari manifold. We define the set

$$\mathcal{N}_{\lambda,\delta} := \{(u, v) \in Y \setminus \{0\} : (I'_{\lambda,\delta}(u, v), (u, v)) = 0\}$$

and find that the functional $I_{\lambda,\delta}$ is bounded below on $\mathcal{N}_{\lambda,\delta}$.

Lemma 3.1. *$I_{\lambda,\delta}$ is coercive and bounded below on $\mathcal{N}_{\lambda,\delta}$ for any $\lambda, \delta > 0$.*

Proof. Let $\lambda, \delta > 0$ and $(u, v) \in \mathcal{N}_{\lambda,\delta}$. Then it holds that

$$\begin{aligned} I_{\lambda,\delta}(u, v) &= C_s^n \left(\frac{1}{2} - \frac{1}{22_\mu^*} \right) \|(u, v)\|^2 - \left(\frac{1}{q} - \frac{1}{22_\mu^*} \right) \int_{\Omega} (\lambda|u|^q + \delta|v|^q) dx \\ &\geq C_s^n \left(\frac{1}{2} - \frac{1}{22_\mu^*} \right) \|(u, v)\|^2 - \left(\frac{1}{q} - \frac{1}{22_\mu^*} \right) |\Omega|^{\frac{2_s^*-q}{2_s^*}} (\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}}) S_s^{-\frac{q}{2}} \|(u, v)\|^q \end{aligned}$$

and this yields the assertion because $1 < q < 2$. \square

From the definition of $\mathcal{N}_{\lambda,\delta}$, it is obvious that $(u, v) \in \mathcal{N}_{\lambda,\delta}$ if and only if $(u, v) \neq (0, 0)$ and

$$C_s^n \|(u, v)\|^2 = \int_{\Omega} (\lambda|u|^q + \delta|v|^q) dx + 2B(u, v).$$

Let us define the fibering map $\varphi_{u,v} : \mathbb{R}^+ \rightarrow \mathbb{R}$ as

$$\varphi_{u,v}(t) = I_{\lambda,\delta}(tu, tv) = \frac{t^2 C_s^n}{2} \|(u, v)\|^2 - \frac{t^q}{q} \int_{\Omega} (\lambda|u|^q + \delta|v|^q) dx - \frac{t^{22_\mu^*}}{2_\mu^*} B(u, v).$$

This gives another characterization of $\mathcal{N}_{\lambda,\delta}$ as follows

$$\mathcal{N}_{\lambda,\delta} = \{(tu, tv) \in Y \setminus \{(0, 0)\} : \varphi'_{u,v}(t) = 0\}$$

because $\varphi'_{u,v}(t) = (I'_{\lambda,\delta}(tu, tv), (u, v))$. An easy computation yields

$$\varphi'_{u,v}(t) = t C_s^n \|(u, v)\|^2 - t^{q-1} \int_{\Omega} (\lambda|u|^q + \delta|v|^q) dx - 2t^{22_\mu^*-1} B(u, v) \quad (3.1)$$

$$\text{and } \varphi''_{u,v}(t) = C_s^n \|(u, v)\|^2 - (q-1)t^{q-2} \int_{\Omega} (\lambda|u|^q + \delta|v|^q) dx - 2(22_\mu^* - 1)t^{22_\mu^*-2} B(u, v). \quad (3.2)$$

If $(u, v) \in \mathcal{N}_{\lambda,\delta}$ then (3.1) and (3.2) gives

$$\begin{aligned} \varphi''_{u,v}(1) &= (2-q)C_s^n \|(u, v)\|^2 + 2(q-22_\mu^*)B(u, v) \\ &= (2-22_\mu^*)C_s^n \|(u, v)\|^2 + (22_\mu^* - q) \int_{\Omega} (\lambda|u|^q + \delta|v|^q) dx. \end{aligned}$$

Naturally, our next step is to divide $\mathcal{N}_{\lambda,\delta}$ into three subsets corresponding to local minima, local maxima and point of inflexion of $\varphi_{u,v}$ namely

$$\mathcal{N}_{\lambda,\delta}^{\pm} := \{(u, v) \in \mathcal{N}_{\lambda,\delta} : \varphi_{u,v}''(1) \gtrless 0\} \text{ and } \mathcal{N}_{\lambda,\delta}^0 := \{(u, v) \in \mathcal{N}_{\lambda,\delta} : \varphi_{u,v}''(1) = 0\}.$$

Our next lemma says that the local minimizers of $I_{\lambda,\delta}$ on the Nehari manifold $\mathcal{N}_{\lambda,\delta}$ are actually its critical points. So it is enough to prove the existence of minimizers of $I_{\lambda,\delta}$ on $\mathcal{N}_{\lambda,\delta}$.

Lemma 3.2. *Let (u_1, v_1) and (u_2, v_2) be minimizers of $I_{\lambda,\delta}$ on $\mathcal{N}_{\lambda,\delta}^+$ and $\mathcal{N}_{\lambda,\delta}^-$ respectively. Then (u_1, v_1) and (u_2, v_2) are nontrivial weak solutions of $(P_{\lambda,\delta})$.*

Proof. Let $(u_1, v_1) \in \mathcal{N}_{\lambda,\delta}^+$ such that $I_{\lambda,\delta}(u_1, v_1) = \inf_{\mathcal{N}_{\lambda,\delta}^+} I_{\lambda,\delta}$ and define $V := \{(u, v) \in Y : (J'_{\lambda,\delta}(u, v), (u, v)) > 0\}$ where $J_{\lambda,\delta}(u, v) = (I'_{\lambda,\delta}(u, v), (u, v))$. So, $\mathcal{N}_{\lambda,\delta}^+ = \{(u, v) \in V : J_{\lambda,\delta}(u, v) = 0\}$ because for each (u, v) such that $J_{\lambda,\delta}(u, v) = 0$, we have $(J'_{\lambda,\delta}(u, v), (u, v)) > 0$ if and only if $\varphi_{u,v}''(1) > 0$. Therefore there exists a Lagrangian multiplier $\rho \in \mathbb{R}$ such that

$$I'_{\lambda,\delta}(u_1, v_1) = \rho J'_{\lambda,\delta}(u_1, v_1).$$

Since $(u_1, v_1) \in \mathcal{N}_{\lambda,\delta}^+$, $(I'_{\lambda,\delta}(u_1, v_1), (u_1, v_1)) = 0$ and $(J'_{\lambda,\delta}(u_1, v_1), (u_1, v_1)) > 0$. This implies $\rho = 0$. Therefore, (u_1, v_1) is a nontrivial weak solution of $(P_{\lambda,\delta})$. Similarly, we can prove that if $(u_2, v_2) \in \mathcal{N}_{\lambda,\delta}^-$ is such that $I_{\lambda,\delta}(u_2, v_2) = \inf_{\mathcal{N}_{\lambda,\delta}^-} I_{\lambda,\delta}$ then (u_2, v_2) is also a nontrivial weak solution of $(P_{\lambda,\delta})$. \square

For fixed $(u, v) \in Y \setminus \{(0, 0)\}$, we write $\varphi'_{u,v}(t) = t^{22_\mu^* - 1}(m_{u,v}(t) - 2B(u, v))$ where

$$m_{u,v}(t) := t^{2-22_\mu^*} C_s^n \|(u, v)\|^2 - t^{q-22_\mu^*} \int_{\Omega} (\lambda|u|^q + \delta|v|^q) dx.$$

Clearly, $\varphi'_{u,v}(t) = 0$ if and only if $m_{u,v}(t) = 2B(u, v)$ if and only if $(tu, tv) \in \mathcal{N}_{\lambda,\delta}$. So in order to understand the fibering maps, we study the map $m_{u,v}$. Since $2 < 22_\mu^*$ and $1 < q < 2$, we get

$$\lim_{t \rightarrow 0^+} m_{u,v}(t) = -\infty \text{ and } \lim_{t \rightarrow +\infty} m_{u,v}(t) = 0.$$

Claim: The map $m_{u,v}(t)$ has a unique critical point at

$$t_{\max}(u, v) := \left(\frac{(22_\mu^* - q) \int_{\Omega} (\lambda|u|^q + \delta|v|^q) dx}{(22_\mu^* - 2) C_s^n \|(u, v)\|^2} \right)^{\frac{1}{2-q}}.$$

This follows from

$$m'_{u,v}(t) = (2 - 22_\mu^*) t^{1-22_\mu^*} C_s^n \|(u, v)\|^2 - (q - 22_\mu^*) t^{q-1-22_\mu^*} \int_{\Omega} (\lambda|u|^q + \delta|v|^q) dx.$$

We can check that $t_{\max}(u, v)$ solves the equation $m'_{u,v}(t) = 0$. Also we can verify that since $1 < q < 2$,

$$m''_{u,v}(t_{\max}(u, v)) = \frac{(q - 2)(22_\mu^* - 2)^{\frac{2+22_\mu^*-q}{2-q}} (C_s^n \|(u, v)\|^2)^{\frac{2+22_\mu^*-q}{2-q}}}{(22_\mu^* - q)^{\frac{22_\mu^*}{2-q}} \left(\int_{\Omega} (\lambda|u|^q + \delta|v|^q) dx \right)^{\frac{22_\mu^*}{2-q}}} < 0$$

implies that $t_{\max}(u, v)$ is the point of maximum for the map $m_{u,v}(t)$. The uniqueness of the critical point of $m_{u,v}$ at $t_{\max}(u, v)$ guarantees that $m_{u,v}(t)$ is strictly increasing in $(0, t_{\max}(u, v))$ and strictly decreasing in $(t_{\max}(u, v), +\infty)$. If $(tu, tv) \in \mathcal{N}_{\lambda, \delta}$ then

$$t^{2^*_\mu-1} m'_{u,v}(t) = \varphi''_{u,v}(t) = t^{-2} \varphi''_{tu,tv}(1)$$

which implies that $(tu, tv) \in \mathcal{N}_{\lambda, \delta}^+$ (or $\mathcal{N}_{\lambda, \delta}^-$) if and only if $m'_{u,v}(t) > 0$ (or $m'_{u,v}(t) < 0$).

Lemma 3.3. For every $(u, v) \in Y \setminus \{(0, 0)\}$ and λ, δ satisfying $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$, where

$$\Theta := \left[\frac{2^{2^*_\mu-1} (C_s^n)^{\frac{2^*_\mu-q}{2-q}}}{C(n, \mu)} \left(\frac{2-q}{2^*_\mu-q} \right) \left(\frac{2^*_\mu-2}{2^*_\mu-q} \right)^{\frac{2^*_\mu-2}{2-q}} S_s^{\frac{q(2^*_\mu-1)}{2-q} + 2^*_\mu} |\Omega|^{-\frac{(2^*_\mu-q)(2^*_\mu-2)}{2^*_\mu(2-q)}} \right]^{\frac{1}{2^*_\mu-1}}, \quad (3.3)$$

there exist unique $t_1, t_2 > 0$ such that $t_1 < t_{\max}(u, v) < t_2$, $(t_1 u, t_1 v) \in \mathcal{N}_{\lambda, \delta}^+$ and $(t_2 u, t_2 v) \in \mathcal{N}_{\lambda, \delta}^-$. Moreover,

$$I_{\lambda, \delta}(t_1 u, t_1 v) = \inf_{t \in [0, t_{\max}(u, v)]} I_{\lambda, \delta}(tu, tv) \text{ and } I_{\lambda, \delta}(t_2 u, t_2 v) = \sup_{t \geq 0} I_{\lambda, \delta}(tu, tv).$$

Proof. Let $(u, v) \in Y \setminus \{(0, 0)\}$. Then we have already seen that

$$m_{u,v}(t) = 2B(u, v) \quad (3.4)$$

if and only if $(tu, tv) \in \mathcal{N}_{\lambda, \delta}$. Since $B(u, v) > 0$, we say that (3.4) can never hold if we choose λ and δ such that $2B(u, v) > m_{u,v}(t_{\max}(u, v))$ and vice-versa. In this case, $(u, v) \notin \mathcal{N}_{\lambda, \delta}$ and hence not a weak solution to $(P_{\lambda, \delta})$. Using Hölder's inequality and the definition of S_s , we get

$$\int_{\Omega} (\lambda |u|^q + \delta |v|^q) dx \leq S_s^{-\frac{q}{2}} |\Omega|^{\frac{2^*_\mu-q}{2^*_\mu}} (\lambda \|u\|^q + \delta \|v\|^q) \leq S_s^{-\frac{q}{2}} |\Omega|^{\frac{2^*_\mu-q}{2^*_\mu}} \|(u, v)\|^q (\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{2-q}{2}}. \quad (3.5)$$

Also from the definition of \tilde{S}_s^H and Lemma 2.6, we get

$$2B(u, v) \leq 2(\tilde{S}_s^H)^{-2^*_\mu} \|(u, v)\|^{2^*_\mu} = 2^{1-2^*_\mu} S_s^{-2^*_\mu} C(n, \mu) \|(u, v)\|^{2^*_\mu}. \quad (3.6)$$

Using (3.5) we can estimate $m_{u,v}(t_{\max})$ as follows

$$\begin{aligned} m_{u,v}(t_{\max}(u, v)) &= \left[\left(\frac{2^*_\mu-2}{2^*_\mu-q} \right)^{\frac{2^*_\mu-2}{2-q}} - \left(\frac{2^*_\mu-2}{2^*_\mu-q} \right)^{\frac{2^*_\mu-q}{2-q}} \right] \frac{(C_s^n \|(u, v)\|^2)^{\frac{2^*_\mu-q}{2-q}}}{\left(\int_{\Omega} (\lambda |u|^q + \delta |v|^q) dx \right)^{\frac{2^*_\mu-2}{2-q}}} \\ &= \left(\frac{2^*_\mu-2}{2^*_\mu-q} \right)^{\frac{2^*_\mu-2}{2-q}} \left(\frac{2-q}{2^*_\mu-q} \right) \frac{(C_s^n \|(u, v)\|^2)^{\frac{2^*_\mu-q}{2-q}}}{\left(\int_{\Omega} (\lambda |u|^q + \delta |v|^q) dx \right)^{\frac{2^*_\mu-2}{2-q}}} \\ &\geq \left(\frac{2^*_\mu-2}{2^*_\mu-q} \right)^{\frac{2^*_\mu-2}{2-q}} \left(\frac{2-q}{2^*_\mu-q} \right) \frac{(C_s^n)^{\frac{2^*_\mu-q}{2-q}} \|(u, v)\|^{2^*_\mu}}{\left(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} \right)^{2^*_\mu-1} (S_s)^{-\frac{q(2^*_\mu-1)}{2-q}} |\Omega|^{\frac{(2^*_\mu-q)(2^*_\mu-2)}{2^*_\mu(2-q)}}}. \end{aligned} \quad (3.7)$$

Now if λ and δ satisfies $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$, where Θ is given in (3.3), then

$$2^{1-2^*_\mu} S_s^{-2^*_\mu} C(n, \mu) \leq \left(\frac{22^*_\mu - 2}{22^*_\mu - q} \right)^{\frac{22^*_\mu - 2}{2-q}} \left(\frac{2-q}{22^*_\mu - q} \right) \frac{(C_s^n)^{\frac{22^*_\mu - q}{2-q}}}{\left(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} \right)^{2^*_\mu - 1} S_s^{-\frac{q(2^*_\mu - 1)}{2-q}} |\Omega|^{\frac{(2^*_s - q)(22^*_\mu - 2)}{2^*_s(2-q)}}} \quad (3.8)$$

which along with (3.7) implies that

$$0 < 2B(u, v) < 2^{1-2^*_\mu} S_s^{-2^*_\mu} C(n, \mu) \|(u, v)\|^{2^*_\mu} < m_{u,v}(t_{\max}(u, v)). \quad (3.9)$$

Therefore there exist unique $t_1, t_2 > 0$ with $t_1 < t_{\max}(u, v) < t_2$ such that

$$m_{u,v}(t_1) = m_{u,v}(t_2) = 2B(u, v)$$

and $m'_{u,v}(t_1) > 0$ and $m'_{u,v}(t_1) < 0$. This implies $(t_1 u, t_1 v) \in \mathcal{N}_{\lambda,\delta}^+$ and $(t_2 u, t_2 v) \in \mathcal{N}_{\lambda,\delta}^-$ and also $\varphi''_{u,v}(t_1) > 0$ and $\varphi''_{u,v}(t_2) < 0$. From the definition of $\varphi_{u,v}$, we get

$$\begin{aligned} I_{\lambda,\delta}(t_2 u, t_2 v) &\geq I_{\lambda,\delta}(t u, t v) \geq I_{\lambda,\delta}(t_1 u, t_1 v) \text{ for each } t \in [t_1, t_2]; \\ I_{\lambda,\delta}(t_1 u, t_1 v) &\leq I_{\lambda,\delta}(t u, t v) \text{ for each } t \in [0, t_1]. \end{aligned}$$

Thus

$$I_{\lambda,\delta}(t_1 u, t_1 v) = \inf_{t \in [0, t_{\max}(u, v)]} I_{\lambda,\delta}(t u, t v) \text{ and } I_{\lambda,\delta}(t_2 u, t_2 v) = \sup_{t \geq 0} I_{\lambda,\delta}(t u, t v),$$

holds true. \square

We end this section with the following important lemma.

Lemma 3.4. *If $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$, where Θ is as in (3.3) then $\mathcal{N}_{\lambda,\delta}^0 = \emptyset$.*

Proof. We prove this by contradiction, so let $(u, v) \in \mathcal{N}_{\lambda,\delta}^0$. By Lemma 3.3 we know that there exist $t_1, t_2 > 0$ such that $\varphi'_{u,v}(t_1) = 0 = \varphi'_{u,v}(t_2)$ and $\varphi''_{u,v}(t_1) > 0 > \varphi''_{u,v}(t_2)$. But $(u, v) \in \mathcal{N}_{\lambda,\delta}^0$ means that $\varphi''_{u,v}(1) = 0 = \varphi'_{u,v}(1)$. This is possible when either $t_1 = 1$ or $t_2 = 1$. But this again implies that $\varphi''_{u,v}(1) > 0$ or $\varphi''_{u,v}(1) < 0$, a contradiction. \square

4. Existence of minimizers on $\mathcal{N}_{\lambda,\delta}^+$ and $\mathcal{N}_{\lambda,\delta}^-$

Lastly, in this section we present the proof of Theorem 1.1. We divide this section into two subsections where we prove existence of first and second solutions respectively.

Lemma 4.1. *Let $\{(u_k, v_k)\} \subset Y$ be a $(PS)_c$ sequence that is*

$$I_{\lambda,\delta}(u_k, v_k) \rightarrow c \text{ in } \mathbb{R} \text{ and } I'_{\lambda,\delta}(u_k, v_k) \rightarrow 0 \text{ in } Y^* \text{ as } k \rightarrow \infty.$$

Then $\{(u_k, v_k)\}$ is bounded in Y .

Proof. Let $\{(u_k, v_k)\} \subset Y$ be a $(PS)_c$ sequence for $I_{\lambda,\delta}$ such that

$$I_{\lambda,\delta}(u_k, v_k) \rightarrow c \text{ in } \mathbb{R} \text{ and } I'_{\lambda,\delta}(u_k, v_k) \rightarrow 0 \text{ in } Y^* \text{ as } k \rightarrow \infty.$$

This can be equivalently written as

$$\frac{C_s^n}{2} \|(u_k, v_k)\|^2 - \frac{1}{q} \int_{\Omega} (\lambda |u_k|^q + \delta |v_k|^q) dx - \frac{1}{2_{\mu}^*} B(u_k, v_k) = c + o_k(1), \quad (4.1)$$

$$C_s^n \|(u_k, v_k)\|^2 - \int_{\Omega} (\lambda |u_k|^q + \delta |v_k|^q) dx - 2B(u_k, v_k) = o_k(\|(u_k, v_k)\|) \quad (4.2)$$

as $k \rightarrow \infty$. We show the boundedness of the sequence $\{(u_k, v_k)\}$ in Y using the method of contradiction. So assume, on contrary, $\|(u_k, v_k)\| \rightarrow \infty$ as $k \rightarrow \infty$ and set

$$w_k := \frac{u_k}{\|(u_k, v_k)\|}, \quad z_k := \frac{v_k}{\|(u_k, v_k)\|}.$$

Clearly, $\|(w_k, z_k)\| = 1$, for all k which implies that there exists a subsequence, still denoted by $\{(w_k, z_k)\}$, such that $(w_k, z_k) \rightharpoonup (w, z)$ weakly in Y as $k \rightarrow \infty$, for some $(w, z) \in Y$. By fractional Sobolev embedding results, we get

$$\int_{\Omega} (\lambda |w_k|^q + \delta |z_k|^q) dx \rightarrow \int_{\Omega} (\lambda |w|^q + \delta |z|^q) dx \text{ as } k \rightarrow \infty. \quad (4.3)$$

Putting $u_k = w_k \|(u_k, v_k)\|$ and $v_k = z_k \|(u_k, v_k)\|$ in (4.1) and (4.2) and solving we get

$$\begin{aligned} \frac{C_s^n}{2} \|(w_k, z_k)\|^2 - \frac{\|(u_k, v_k)\|^{q-2}}{q} \int_{\Omega} (\lambda |w_k|^q + \delta |z_k|^q) dx - \frac{1}{2_{\mu}^*} \|(u_k, v_k)\|^{2_{\mu}^*-2} B(w_k, z_k) &= o_k(1), \\ C_s^n \|(w_k, z_k)\|^2 - \|(u_k, v_k)\|^{q-2} \int_{\Omega} (\lambda |w_k|^q + \delta |z_k|^q) dx - 2 \|(u_k, v_k)\|^{2_{\mu}^*-2} B(w_k, z_k) &= o_k(1). \end{aligned}$$

From above these two equations and (4.3), we get

$$\begin{aligned} C_s^n \|(w_k, z_k)\|^2 &= \frac{(22_{\mu}^* - q)}{q(2_{\mu}^* - 1)} \|(u_k, v_k)\|^{q-2} \int_{\Omega} (\lambda |w_k|^q + \delta |z_k|^q) dx + o_k(1) \\ &= \frac{(22_{\mu}^* - q)}{q(2_{\mu}^* - 1)} \|(u_k, v_k)\|^{q-2} \int_{\Omega} (\lambda |w|^q + \delta |z|^q) dx + o_k(1). \end{aligned}$$

Since $1 < q < 2$ and $\|(u_k, v_k)\| \rightarrow \infty$ we get $\|(w_k, z_k)\|^2 \rightarrow 0$ as $k \rightarrow \infty$ which contradicts $\|(w_k, z_k)\| = 1$ for all k . This completes the proof. \square

Lemma 4.2. *If $\{(u_k, v_k)\}$ is a $(PS)_c$ sequence for $I_{\lambda, \delta}$ with $(u_k, v_k) \rightharpoonup (u, v)$ weakly in Y as $k \rightarrow \infty$, then $I'_{\lambda, \delta}(u, v) = 0$. Moreover there exists a positive constant D_0 depending on μ, q, s, n, S_s and Ω such that*

$$I_{\lambda, \delta}(u, v) \geq -D_0(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}}), \quad (4.4)$$

where

$$D_0 := \frac{(2-q)(22_{\mu}^* - q)}{42_{\mu}^* q} \left[\left(\frac{22_{\mu}^* C_s^n S_s (n - \mu + 2s)}{(2n - \mu)(22_{\mu}^* - q)} \right)^{-\frac{q}{2}} |\Omega|^{\frac{2_{\mu}^* - q}{2_{\mu}^*}} \right]^{\frac{2}{2-q}}.$$

Proof. Let $\{(u_k, v_k)\} \subset Y$ be a $(PS)_c$ sequence for $I_{\lambda, \delta}$ such that $(u_k, v_k) \rightharpoonup (u, v)$ weakly in Y as $k \rightarrow \infty$. This implies $I'_{\lambda, \delta}(u_k, v_k) = o_k(1)$ in Y^* as $k \rightarrow \infty$. Let $(\phi, \psi) \in Y$. From weak convergence it follows that

$$\lim_{k \rightarrow \infty} \langle u_k, \phi \rangle = \langle u, \phi \rangle \text{ and } \lim_{k \rightarrow \infty} \langle v_k, \psi \rangle = \langle v, \psi \rangle. \quad (4.5)$$

For $q' = \frac{q}{q-1}$ we also have

$$|u_k|^{q-2}u_k \rightharpoonup |u|^{q-2}u, \quad |v_k|^{q-2}v_k \rightharpoonup |v|^{q-2}v \text{ in } L^{q'}(\Omega) \text{ and } u_k \rightharpoonup u, \quad v_k \rightharpoonup v \text{ in } L^{2_s^*}(\Omega) \quad (4.6)$$

as $k \rightarrow \infty$, thanks to the embedding of X_0 into $L^m(\Omega)$ for all $1 \leq m \leq 2_s^*$. Since we assumed $\phi, \psi \in X_0$ which is contained in $L^q(\Omega) \cap L^{2_s^*}(\Omega)$, so from (4.6) it follows that as $k \rightarrow \infty$

$$\int_{\Omega} |u_k|^{q-2}u_k \phi \, dx \rightarrow \int_{\Omega} |u|^{q-2}u \phi \, dx. \quad (4.7)$$

Also since $2_{\mu}^* - 1 = \frac{n - \mu + 2s}{n - 2s}$ and $|u_k|^{2_{\mu}^*} \rightharpoonup |u|^{2_{\mu}^*}, |v_k|^{2_{\mu}^*} \rightharpoonup |v|^{2_{\mu}^*}$ in $L^{\frac{2n}{2n-\mu}}(\Omega)$, we get

$$|u_k|^{2_{\mu}^*-2}u_k \rightharpoonup |u|^{2_{\mu}^*-2}u \text{ and } |v_k|^{2_{\mu}^*-2}v_k \rightharpoonup |v|^{2_{\mu}^*-2}v \text{ in } L^{\frac{2n}{n-\mu+2s}}(\Omega).$$

By Hardy–Littlewood–Sobolev inequality, the Riesz potential defines a linear and continuous map from $L^{\frac{2n}{2n-\mu}}(\Omega)$ to $L^{\frac{2n}{\mu}}(\Omega)$ which gives

$$|x|^{-\mu} * |u_k|^{2_{\mu}^*} \rightharpoonup |x|^{-\mu} * |u|^{2_{\mu}^*} \text{ and } |x|^{-\mu} * |v_k|^{2_{\mu}^*} \rightharpoonup |x|^{-\mu} * |v|^{2_{\mu}^*} \text{ in } L^{\frac{2n}{\mu}}(\Omega). \quad (4.8)$$

This implies that the sequences $(|x|^{-\mu} * |u_k|^{2_{\mu}^*})|v_k|^{2_{\mu}^*-2}v_k$ and $(|x|^{-\mu} * |v_k|^{2_{\mu}^*})|u_k|^{2_{\mu}^*-2}u_k$ converges weakly in $L^{\frac{2n}{n+2s}}(\Omega)$. Through Sobolev embedding we know that

$$|u_k|^{2_{\mu}^*-2}u_k \rightarrow |u|^{2_{\mu}^*-2}u \text{ and } |v_k|^{2_{\mu}^*-2}v_k \rightarrow |v|^{2_{\mu}^*-2}v \text{ in } L^{\frac{2n}{2n-\mu}}(\Omega). \quad (4.9)$$

Taking into account (4.8) and (4.9), for any $\tilde{\psi} \in L^{\infty}(\Omega)$ we obtain

$$\begin{aligned} \int_{\Omega} (|x|^{-\mu} * |u_k|^{2_{\mu}^*})|v_k|^{2_{\mu}^*-2}v_k \tilde{\psi} \, dx &\rightarrow \int_{\Omega} (|x|^{-\mu} * |u|^{2_{\mu}^*})|v|^{2_{\mu}^*-2}v \tilde{\psi} \, dx \\ \text{and } \int_{\Omega} (|x|^{-\mu} * |v_k|^{2_{\mu}^*})|u_k|^{2_{\mu}^*-2}u_k \tilde{\psi} \, dx &\rightarrow \int_{\Omega} (|x|^{-\mu} * |v|^{2_{\mu}^*})|u|^{2_{\mu}^*-2}u \tilde{\psi} \, dx. \end{aligned}$$

Therefore the sequences $(|x|^{-\mu} * |u_k|^{2_{\mu}^*})|v_k|^{2_{\mu}^*-2}v_k$ and $(|x|^{-\mu} * |v_k|^{2_{\mu}^*})|u_k|^{2_{\mu}^*-2}u_k$ converges in the distributional sense. Since the weak limit and the distributional limit coincides, for $\phi, \psi \in X_0(\Omega) \subset L^{2_s^*}(\Omega)$, we get that as $k \rightarrow \infty$

$$\begin{aligned} \int_{\Omega} (|x|^{-\mu} * |u_k|^{2_{\mu}^*})|v_k|^{2_{\mu}^*-2}v_k \psi \, dx &\rightarrow \int_{\Omega} (|x|^{-\mu} * |u|^{2_{\mu}^*})|v|^{2_{\mu}^*-2}v \psi \, dx, \\ \int_{\Omega} (|x|^{-\mu} * |v_k|^{2_{\mu}^*})|u_k|^{2_{\mu}^*-2}u_k \phi \, dx &\rightarrow \int_{\Omega} (|x|^{-\mu} * |v|^{2_{\mu}^*})|u|^{2_{\mu}^*-2}u \phi \, dx. \end{aligned} \quad (4.10)$$

So using (2.3), (4.5), (4.7) and (4.10) we get $(I'_{\lambda, \delta}(u_k, v_k) - I'_{\lambda, \delta}(u, v), (\phi, \psi)) \rightarrow 0$ as $k \rightarrow \infty$, for all $(\phi, \psi) \in Y$ which implies that $I'_{\lambda, \delta}(u, v) = 0$. Therefore (u, v) is a weak solution of $(P_{\lambda, \delta})$ and $(u, v) \in \mathcal{N}_{\lambda, \delta}$, that is

$$C_s^n \|(u, v)\|^2 = \int_{\Omega} (\lambda |u|^q + \delta |v|^q) dx + 2B(u, v)$$

which gives

$$I_{\lambda, \delta}(u, v) = \frac{(2_{\mu}^* - 1)C_s^n}{22_{\mu}^*} \|(u, v)\|^2 - \frac{22_{\mu}^* - q}{22_{\mu}^* q} \int_{\Omega} (\lambda |u|^q + \delta |v|^q) dx. \quad (4.11)$$

Let $D = \left[\frac{2}{q} \cdot \frac{n - \mu + 2s}{2(2n - \mu)} \left(\frac{1}{q} - \frac{1}{22_{\mu}^*} \right)^{-1} \right]$. Using the Hölder inequality, the fractional Sobolev inequality, the definition of S_s and the Young's inequality we get the following estimate

$$\begin{aligned} \int_{\Omega} (\lambda |u|^q + \delta |v|^q) dx &\leq |\Omega|^{\frac{2_s^* - q}{2_s^*}} S_s^{-\frac{q}{2}} (\lambda \|u\|^q + \delta \|v\|^q) \\ &= \left(D^{\frac{q}{2}} (C_s^n)^{\frac{q}{2}} \|u\|^q \right) \left(D^{-\frac{q}{2}} \lambda |\Omega|^{\frac{2_s^* - q}{2_s^*}} (C_s^n S_s)^{-\frac{q}{2}} \right) + \left(D^{\frac{q}{2}} (C_s^n)^{\frac{q}{2}} \|v\|^q \right) \left(D^{-\frac{q}{2}} \delta |\Omega|^{\frac{2_s^* - q}{2_s^*}} (C_s^n S_s)^{-\frac{q}{2}} \right) \\ &\leq \frac{n - \mu + 2s}{2(2n - \mu)} \left(\frac{1}{q} - \frac{1}{22_{\mu}^*} \right)^{-1} C_s^n (\|u\|^2 + \|v\|^2) + \tilde{D} \left(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} \right) \\ &= \frac{n - \mu + 2s}{2(2n - \mu)} \left(\frac{1}{q} - \frac{1}{22_{\mu}^*} \right)^{-1} C_s^n \|(u, v)\|^2 + \tilde{D} \left(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} \right), \end{aligned} \quad (4.12)$$

where $\tilde{D} := \frac{2 - q}{2} \left(D^{-\frac{q}{2}} |\Omega|^{\frac{2_s^* - q}{2_s^*}} (C_s^n S_s)^{-\frac{q}{2}} \right)^{\frac{2}{2-q}}$. Using (4.12) in (4.11), we finally obtain (4.4) with $D_0 = \left(\frac{22_{\mu}^* - q}{22_{\mu}^* q} \right) \tilde{D}$. This completes the proof. \square

As a consequence of Lemma 3.4 we infer that for any λ, δ satisfying $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$,

$$\mathcal{N}_{\lambda, \delta} = \mathcal{N}_{\lambda, \delta}^+ \cup \mathcal{N}_{\lambda, \delta}^-.$$

In spirit of Lemma 3.1, we define the following

$$l_{\lambda, \delta} = \inf_{\mathcal{N}_{\lambda, \delta}} I_{\lambda, \delta} \text{ and } l_{\lambda, \delta}^{\pm} = \inf_{\mathcal{N}_{\lambda, \delta}^{\pm}} I_{\lambda, \delta}.$$

Then we have the following result.

Lemma 4.3. *The following holds true:*

- (i) If $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$, then $l_{\lambda, \delta} \leq l_{\lambda, \delta}^+ < 0$,
- (ii) $\inf\{\|(u, v)\| : (u, v) \in \mathcal{N}_{\lambda, \delta}^-, I_{\lambda, \delta}(u, v) \leq M\} < +\infty$ for each $M > 0$.

Proof. (i) Let $(u, v) \in \mathcal{N}_{\lambda, \delta}^+$ implying that $\varphi'_{u, v}(1) = 0$ and $\varphi''_{u, v}(1) > 0$. Therefore

$$\frac{(2 - q)C_s^n}{2(22_{\mu}^* - q)} \|(u, v)\|^2 > B(u, v).$$

Using this we deduce that

$$\begin{aligned} I_{\lambda,\delta}(u, v) &= \left(\frac{1}{2} - \frac{1}{q}\right) C_s^n \|(u, v)\|^2 + \left(\frac{2}{q} - \frac{1}{2_\mu^*}\right) B(u, v) \\ &< \left(\frac{q-2}{2q} + \frac{2-q}{22_\mu^* q}\right) C_s^n \|(u, v)\|^2 = \frac{2-q}{2q} \left(\frac{1}{2_\mu^*} - 1\right) \|(u, v)\|^2 < 0. \end{aligned}$$

This along with the definition of $l_{\lambda,\delta}$ and $l_{\lambda,\delta}^+$ implies that $l_{\lambda,\delta} \leq l_{\lambda,\delta}^+ < 0$.

(ii) Let $(u, v) \in \mathcal{N}_{\lambda,\delta}^-$. Then using Lemma 2.3 and (2.2) we get

$$0 > \varphi_{u,v}''(1) \geq (2-q)C_s^n \|(u, v)\|^2 - 2(22_\mu^* - q)(\tilde{S}_s^H)^{-2_\mu^*} \|(u, v)\|^{22_\mu^*}.$$

This gives

$$\|(u, v)\| \geq \left(\frac{(2-q)C_s^n}{2(22_\mu^* - q)(\tilde{S}_s^H)^{-2_\mu^*}} \right)^{\frac{1}{22_\mu^* - 2}} > 0$$

which implies that $\inf\{\|(u, v)\| : (u, v) \in \mathcal{N}_{\lambda,\delta}^-\} > 0$. Therefore $\inf\{\|(u, v)\| : (u, v) \in \mathcal{N}_{\lambda,\delta}^-\} > 0$. Now let $I_{\lambda,\delta}(u, v) \leq M$ for some $M > 0$ then an easy computation yields

$$\left(\frac{1}{2} - \frac{1}{22_\mu^*}\right) C_s^n \|(u, v)\|^2 - K_{\lambda,\delta} \left(\frac{1}{q} - \frac{1}{22_\mu^*}\right) \|(u, v)\|^q \leq M$$

where $K_{\lambda,\delta} = S_s^{-\frac{q}{2}} |\Omega|^{\frac{2_\mu^* - q}{2s^*}} (\lambda + \delta)$ which completes the proof. \square

Our next result is established by using the implicit function theorem and it plays a crucial role in proving Theorem 4.5.

Proposition 4.4. Assume $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$ and $w = (u, v) \in \mathcal{N}_{\lambda,\delta}$. Then there exist $\epsilon > 0$ and a differentiable function $\zeta : B_\epsilon(0) \subset Y \rightarrow \mathbb{R}^+$ ($B_\epsilon(0)$ denotes ball of radius ϵ with center origin) such that $\zeta(0) = 1$, $\zeta(z)(w - z) \in \mathcal{N}_{\lambda,\delta}$ and

$$(\zeta'(0), z) = -\frac{2(\langle u, z_1 \rangle + \langle v, z_2 \rangle) - T_{\lambda,\delta}(w, z) - 2M(z)}{(2-q)C_s^n \|(u, v)\|^2 - 2(22_\mu^*)B(u, v)} \quad (4.13)$$

for all $z = (z_1, z_2) \in B_\epsilon(0)$, where

$$\begin{aligned} T_{\lambda,\delta}(w, z) &= q \int_{\Omega} (\lambda |u|^{q-2} u z_1 + \delta |v|^{q-2} v z_2) dx, \\ M(z) &= \int_{\Omega} ((|x|^{-\mu} * |v|^{2_\mu^*}) |u|^{2_\mu^* - 2} u z_1 + (|x|^{-\mu} * |u|^{2_\mu^*}) |v|^{2_\mu^* - 2} v z_2) dx. \end{aligned}$$

Proof. For $w = (u, v) \in \mathcal{N}_{\lambda,\delta}$, let us define $\mathfrak{F}_w : \mathbb{R}^+ \times Y \rightarrow \mathbb{R}^n$ by

$$\begin{aligned} \mathfrak{F}_w(\rho, z) &:= (I'_{\lambda,\delta}(\rho(w - z)), (\rho(w - z))) \\ &= \rho^2 C_s^n \|(u - z_1, v - z_2)\|^2 - \rho^q \int_{\Omega} (\lambda |u - z_1|^{q-2} (u - z_1) + \delta |v - z_2|^{q-2} (v - z_2)) dx - 2\rho^{22_\mu^*} B(u - z_1, v - z_2) \end{aligned}$$

where $\rho \in \mathbb{R}^+$ and $z = (z_1, z_2) \in Y$. Then clearly $\mathfrak{F}_w(1, (0, 0)) = (I'_{\lambda, \delta}(w), w) = 0$ since $w \in \mathcal{N}_{\lambda, \delta}$. Also

$$\begin{aligned} \frac{d}{d\rho} \mathfrak{F}_w(1, (0, 0)) &= 2C_s^n \|(u, v)\|^2 - q \int_{\Omega} (\lambda |u|^q + \delta |v|^q) dx - 2(22_{\mu}^*) B(u, v) \\ &= (2 - q) C_s^n \|(u, v)\|^2 - 2(22_{\mu}^* - q) \int_{\Omega} (\lambda |u|^q + \delta |v|^q) dx = \varphi''_{u, v}(1) \neq 0 \end{aligned}$$

because of Lemma 3.4. Therefore we can apply the implicit function theorem to obtain a $\epsilon > 0$ and a differentiable map $\zeta : B_{\epsilon}(0) \subset Y \rightarrow \mathbb{R}^+$ with $\zeta(0) = 1$ and satisfies (4.13). Also $\mathfrak{F}_w(\zeta) = 0$ for all $z \in B_{\epsilon}(0)$ which is equivalent to

$$(I'_{\lambda, \delta}(\zeta(z)(w - z)), \zeta(z)(w - z)) = 0, \text{ for all } z \in B_{\epsilon}(0),$$

that is $\zeta(z)(w - z) \in \mathcal{N}_{\lambda, \delta}$. \square

Theorem 4.5. *If $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$ then there exists a $(PS)_{l_{\lambda, \delta}}$ sequence $\{(u_k, v_k)\} \subset \mathcal{N}_{\lambda, \delta}$ for $I_{\lambda, \delta}$.*

Proof. We use the Ekeland Variational principle to say that there exists a minimizing sequence $\{(u_k, v_k)\} \subset \mathcal{N}_{\lambda, \delta}$ such that

$$I_{\lambda, \delta}(u_k, v_k) < l_{\lambda, \delta} + \frac{1}{k} \quad \text{and} \quad I_{\lambda, \delta}(u_k, v_k) < I_{\lambda, \delta}(w_1, w_2) + \frac{1}{k} \|(w_1, w_2) - (u_k, v_k)\|, \quad (4.14)$$

for each $(w_1, w_2) \in \mathcal{N}_{\lambda, \delta}$. From Lemma 4.3(i) we know that $l_{\lambda, \delta} < 0$, therefore we can find k sufficiently large such that

$$I_{\lambda, \delta}(u_k, v_k) = \left(\frac{1}{2} - \frac{1}{22_{\mu}^*} \right) C_s^n \|(u_k, v_k)\|^2 - \left(\frac{1}{q} - \frac{1}{22_{\mu}^*} \right) \int_{\Omega} (\lambda |u_k|^q + \delta |v_k|^q) dx < \frac{l_{\lambda, \delta}}{2}. \quad (4.15)$$

This gives us

$$-\frac{2_{\mu}^* q}{(22_{\mu}^* - q)} l_{\lambda, \delta} < \int_{\Omega} (\lambda |u_k|^q + \delta |v_k|^q) dx < S_s^{-\frac{q}{2}} |\Omega|^{\frac{2_{\mu}^* - q}{2_{\mu}^*}} (\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{2-q}{2}} \|(u_k, v_k)\|^q. \quad (4.16)$$

Consequently $(u_k, v_k) \neq 0$. From (4.16) we get

$$\|(u_k, v_k)\| > \left(-\frac{2_{\mu}^* q l_{\lambda, \delta}}{22_{\mu}^* - q} S_s^{\frac{q}{2}} |\Omega|^{-\frac{2_{\mu}^* - q}{2_{\mu}^*}} \left(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} \right)^{\frac{q-2}{2}} \right)^{\frac{1}{q}} \quad (4.17)$$

and from (4.15) we get

$$\|(u_k, v_k)\| < \left(\frac{(22_{\mu}^* - q)}{q(2_{\mu}^* - 1)} S_s^{-\frac{q}{2}} |\Omega|^{\frac{2_{\mu}^* - q}{2_{\mu}^*}} \left(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} \right)^{\frac{2-q}{2}} \right)^{\frac{1}{q}}. \quad (4.18)$$

Claim: $I'_{\lambda, \delta}(u_k, v_k) \rightarrow 0$ in Y^* as $k \rightarrow \infty$.

Let us fix $k \in \mathbb{N}$ then by applying Proposition 4.4 to $w_k = (u_k, v_k)$, we get that there exists a function $\zeta_k : B_{\epsilon_k}(0) \rightarrow \mathbb{R}^+$ for some $\epsilon_k > 0$ such that $\zeta_k(h)(w_k - h) \in \mathcal{N}_{\lambda, \delta}$ for $h = (h_1, h_2) \in B_{\epsilon_k}(0)$. Let us take $\tau \in (0, \epsilon_k)$ and $z \in Y$ with $z \neq 0$ in Y . We set

$$\tilde{z} = \frac{\tau z}{\|z\|} \text{ and } h_\tau = \zeta_k(\tilde{z})(w_k - \tilde{z}).$$

Then Lemma 4.4 implies that $\tilde{z} \in \mathcal{N}_{\lambda,\delta}$ and using (4.14) with $(w_1, w_2) = h_\tau$ we get

$$I_{\lambda,\delta}(h_\tau) - I_{\lambda,\delta}(w_k) \geq -\frac{1}{k}\|h_\tau - w_k\|.$$

Now applying the mean value theorem we obtain

$$(I'_{\lambda,\delta}(w_k), h_\tau - w_k) + o(\|h_\tau - w_k\|) \geq -\frac{1}{k}\|h_\tau - w_k\|.$$

Substituting the value of h_τ in this, we get

$$(I'_{\lambda,\delta}(w_k), -\tilde{z}) + (\zeta_k(\tilde{z}) - 1)(I'_{\lambda,\delta}(w_k), w_k - \tilde{z}) \geq -\frac{1}{k}\|h_\tau - w_k\| + o(\|h_\tau - w_k\|).$$

Then using the fact that $\zeta'_k(\tilde{h})(w_k - \tilde{h}) \in \mathcal{N}_{\lambda,\delta}$, we get

$$-\tau \left(I'_{\lambda,\delta}(w_k), \frac{z}{\|z\|} \right) + (\zeta_k(\tilde{z}) - 1)(I'_{\lambda,\delta}(w_k) - I'_{\lambda,\delta}(h_\tau), w_k - \tilde{h}) \geq -\frac{1}{k}\|h_\tau - w_k\| + o(\|h_\tau - w_k\|). \quad (4.19)$$

Since $\|h_\tau - w_k\| \leq \tau|\zeta_k(\tilde{z})| + |\zeta_k(\tilde{z}) - 1|\|w_k\|$ and

$$\lim_{\tau \rightarrow 0} \frac{|\zeta_k(\tilde{z}) - 1|}{\tau} \leq \|\zeta'_k(0)\|,$$

passing to the limit as $\tau \rightarrow 0$ in (4.19), we get for some constant $M > 0$

$$\left(I'_{\lambda,\delta}(w_k), \frac{z}{\|z\|} \right) \leq \frac{M}{k}(1 + \|\zeta'_k(0)\|).$$

This will prove our claim once we are able to show that $\sup_k \|\zeta'_k(0)\|_{Y^*} < +\infty$. Let $w = (w_1, w_2) \in Y$. Then using Hölder's inequality we get

$$\int_{\Omega} (\lambda |u_k|^{q-1} w_1 + \delta |v_k|^{q-1} w_2) dx \leq (\lambda + \delta) C_q^q \| (u_k, v_k) \|^{q-1} \| (w_1, w_2) \|, \quad (4.20)$$

where $C_q = \sup\{\int_{\Omega} |u|^q : \|u_k\| = 1\}$. Again using Hölder inequality, Hardy–Littlewood–Sobolev inequality and fractional Sobolev embeddings, we can estimate the following

$$\begin{aligned} & \int_{\Omega} (|x|^\mu * |u_k|^{2_\mu^*}) |v_k|^{2_\mu^*-1} w_1 \, dx \\ & \leq C(n, \mu) \left(\int_{\Omega} (|v_k|^{2_\mu^*-1} w_1)^{\frac{2n}{2n-\mu}} \right)^{\frac{2n-\mu}{2n}} \left(\int_{\Omega} |u_k|^{2_\mu^* \cdot \frac{2n}{2n-\mu}} \right)^{\frac{2n-\mu}{2n}} \\ & \leq C(n, \mu) \left[\left(\int_{\Omega} |v_k|^{2_s^*} \right)^{\frac{n-\mu+2s}{2n-\mu}} \left(\int_{\Omega} |w_1|^{2_s^*} \right)^{\frac{1}{2_\mu^*}} \right]^{\frac{2n-\mu}{2n}} \left(\int_{\Omega} |u_k|^{2_s^*} \right)^{\frac{2n-\mu}{2n}} \\ & \leq M_1 \|(u_k, v_k)\|^\alpha \|(w_1, w_2)\|, \end{aligned} \quad (4.21)$$

where $\alpha = 2_s^* \left(\frac{3n - 2\mu + 2s}{2n} \right)$ and $M_1 > 0$ is a constant. Similarly we can show that there exists $M_2 > 0$ such that

$$\int_{\Omega} (|x|^{\mu} * |v_k|^{2_s^*}) |u_k|^{2_s^*-1} w_2 \, dx \leq M_2 \| (u_k, v_k) \|^\alpha \| (w_1, w_2) \|. \quad (4.22)$$

Consequently using (4.20), (4.21) and (4.22) in (4.13) we get

$$|(\zeta'_k(0), w)| \leq \frac{M_3 \| (w_1, w_2) \|}{|(2-q)C_s^n \| (u_k, v_k) \|^2 - 2(22_\mu^* - q)B(u_k, v_k)|}$$

where $M_3 > 0$ is a constant independent of (u_k, v_k) , thanks to (4.18).

Claim: There exists a $M_4 > 0$ such that

$$|(2-q)C_s^n \| (u_k, v_k) \|^2 - 2(22_\mu^* - q)B(u_k, v_k)| \geq M_4.$$

On contrary, let us assume that there exists a subsequence still denoted by $\{(u_k, v_k)\} \subset \mathcal{N}_{\lambda, \delta}$ such that

$$|(2-q)C_s^n \| (u_k, v_k) \|^2 - 2(22_\mu^* - q)B(u_k, v_k)| = o_k(1). \quad (4.23)$$

Since $(u_k, v_k) \in \mathcal{N}_{\lambda, \delta}$, we have

$$\begin{aligned} C_s^n \| (u_k, v_k) \|^2 &= \left(\frac{22_\mu^* - q}{22_\mu^* - 2} \right) \int_{\Omega} (\lambda |u_k|^q + \delta |v_k|^q) \, dx + o_k(1) \\ &\leq \left(\frac{22_\mu^* - q}{22_\mu^* - 2} \right) S_s^{-\frac{q}{2}} |\Omega|^{\frac{2_s^* - q}{2_s^*}} (\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{2-q}{2}} \| (u_k, v_k) \|^q + o_k(1) \end{aligned}$$

which implies that

$$C_s^n \| (u_k, v_k) \|^{2-q} \leq \left(\frac{22_\mu^* - q}{22_\mu^* - 2} \right) S_s^{-\frac{q}{2}} |\Omega|^{\frac{2_s^* - q}{2_s^*}} (\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{2-q}{2}} + o_k(1). \quad (4.24)$$

Also (4.23) gives us

$$C_s^n \| (u_k, v_k) \|^2 = \left(\frac{2(22_\mu^* - q)}{2 - q} \right) B(u_k, v_k) + o_k(1) \leq \left(\frac{2(22_\mu^* - q)}{2 - q} \right) (\tilde{S}_s^H)^{-2_\mu^*} \| (u_k, v_k) \|^{22_\mu^*} + o_k(1)$$

which implies that

$$\| (u_k, v_k) \| \geq \left(\frac{C_s^n (2-q) (\tilde{S}_s^H)^{2_\mu^*}}{2(22_\mu^* - q)} \right)^{\frac{1}{22_\mu^* - 2}} + o_k(1) \quad (4.25)$$

where we have used the fact that $\| (u_k, v_k) \| \neq o_k(1)$ because of (4.17). From (4.24) and (4.25), for large k we obtain

$$C_s^n \left(\frac{C_s^n (2-q) (\tilde{S}_s^H)^{2_\mu^*}}{2(22_\mu^* - q)} \right)^{\frac{2-q}{22_\mu^* - 2}} \leq \left(\frac{22_\mu^* - q}{22_\mu^* - 2} \right) S_s^{-\frac{q}{2}} |\Omega|^{\frac{2_s^* - q}{2_s^*}} (\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{2-q}{2}}.$$

Then using Lemma 2.6 and (2.6), the above inequality yields

$$(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}}) \geq \left[\frac{2^{2^*_\mu-1} (C_s^n)^{\frac{22^*_\mu-q}{2-q}}}{C(n, \mu)} \left(\frac{2-q}{22^*_\mu-q} \right) \left(\frac{22^*_\mu-2}{22^*_\mu-q} \right)^{\frac{22^*_\mu-2}{2-q}} S_s^{\frac{q(2^*_\mu-1)}{2-q} + 2^*_\mu} |\Omega|^{-\frac{(2^*_\mu-q)(22^*_\mu-2)}{2^*_\mu(2-q)}} \right]^{\frac{1}{2^*_\mu-1}}.$$

This contradicts the assumption that $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$. Hence the claim holds true and we finally obtain

$$\left(I'_{\lambda, \delta}(w_k), \frac{z}{\|z\|} \right) \leq \frac{M}{k}.$$

This establishes our first claim and completes the proof. \square

4.1. First solution

We now prove the existence of first solution for the problem $(P_{\lambda, \delta})$.

Theorem 4.6. *Let $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$. Then there exists a $(u_1, v_1) \in \mathcal{N}_{\lambda, \delta}^+$ such that (u_1, v_1) is a weak solution of $(P_{\lambda, \delta})$. Moreover, (u_1, v_1) satisfies $I_{\lambda, \delta}(u_1, v_1) = l_{\lambda, \delta} = l_{\lambda, \delta}^+ < 0$.*

Proof. By Theorem 4.5 we know that there exists a $(PS)_{l_{\lambda, \delta}}$ sequence $\{(u_k, v_k)\} \subset \mathcal{N}_{\lambda, \delta}$ for $I_{\lambda, \delta}$ that is

$$\lim_{k \rightarrow \infty} I_{\lambda, \delta}(u_k, v_k) = l_{\lambda, \delta} \leq l_{\lambda, \delta}^+ < 0 \text{ and } \lim_{k \rightarrow \infty} I'_{\lambda, \delta}(u_k, v_k) = 0 \text{ in } Y^*.$$

By Lemma 4.1 we know that this sequence $\{(u_k, v_k)\}$ is bounded in Y . Therefore there exists $(u_1, v_1) \in Y$ such that upto a subsequence, $(u_k, v_k) \rightharpoonup (u_1, v_1)$ weakly in Y and $(u_k, v_k) \rightarrow (u_1, v_1)$ strongly in $L^m(\Omega)$, for $m \in [1, 2^*_s)$ as $k \rightarrow \infty$. Therefore $\lim_{k \rightarrow \infty} \int_{\Omega} (\lambda |u_k|^q + \delta |v_k|^q) dx = \int_{\Omega} (\lambda |u_1|^q + \delta |v_1|^q) dx$. We already know that (u_1, v_1) is a weak solution of $(P_{\lambda, \delta})$, by Lemma 4.2. Since $\{(u_k, v_k)\} \subset \mathcal{N}_{\lambda, \delta}$ we obtain

$$\begin{aligned} I_{\lambda, \delta}(u_k, v_k) &= \left(\frac{1}{2} - \frac{1}{22^*_\mu} \right) C_s^n \|(u_k, v_k)\|^2 - \left(\frac{1}{q} - \frac{1}{22^*_\mu} \right) \int_{\Omega} (\lambda |u_k|^q + \delta |v_k|^q) dx \\ &\geq - \left(\frac{1}{q} - \frac{1}{22^*_\mu} \right) \int_{\Omega} (\lambda |u_k|^q + \delta |v_k|^q) dx. \end{aligned}$$

From Lemma 4.3 we know that $l_{\lambda, \delta} < 0$, so passing on the limit $k \rightarrow \infty$ we get

$$\int_{\Omega} (\lambda |u_1|^q + \delta |v_1|^q) dx \geq - \frac{22^*_\mu}{(22^*_\mu - q)} l_{\lambda, \delta} > 0.$$

This implies that $(u_1, v_1) \in \mathcal{N}_{\lambda, \delta}$ is non-trivial solution of $(P_{\lambda, \delta})$.

Claim: $(u_k, v_k) \rightarrow (u_1, v_1)$ strongly in Y as $k \rightarrow \infty$ and $I_{\lambda, \delta}(u_1, v_1) = l_{\lambda, \delta}^+$.

Using $(u_1, v_1) \in \mathcal{N}_{\lambda, \delta}$ and Fatou's Lemma we have

$$\begin{aligned} l_{\lambda, \delta} &\leq I_{\lambda, \delta}(u_1, v_1) = \left(\frac{2^*_\mu - 1}{22^*_\mu} \right) C_s^n \|(u_1, v_1)\|^2 - \left(\frac{22^*_\mu - q}{22^*_\mu q} \right) \int_{\Omega} (\lambda |u_1|^q + \delta |v_1|^q) dx \\ &\leq \liminf_{k \rightarrow \infty} \left(\left(\frac{2^*_\mu - 1}{22^*_\mu} \right) C_s^n \|(u_k, v_k)\|^2 - \left(\frac{22^*_\mu - q}{22^*_\mu q} \right) \int_{\Omega} (\lambda |u_k|^q + \delta |v_k|^q) dx \right) \\ &= \liminf_{k \rightarrow \infty} I_{\lambda, \delta}(u_k, v_k) = l_{\lambda, \delta}. \end{aligned}$$

This implies that $I_{\lambda,\delta}(u_1, v_1) = l_{\lambda,\delta}$ and $\|(u_k, v_k)\| \rightarrow \|(u_1, v_1)\|$ as $k \rightarrow \infty$. We have

$$\|(u_k - u_1, v_k - v_1)\|^2 = \|(u_k, v_k)\|^2 - \|(u_1, v_1)\|^2 + o_k(1).$$

Therefore $(u_k, v_k) \rightarrow (u_1, v_1)$ strongly in Y as $k \rightarrow \infty$. To establish our claim, it remains to show that $(u_1, v_1) \in \mathcal{N}_{\lambda,\delta}^+$. On the contrary, if $(u_1, v_1) \in \mathcal{N}_{\lambda,\delta}^-$ then by Lemma 3.3, there exist unique $t_2 > t_1 > 0$ such that

$$(t_1 u, t_1 v) \in \mathcal{N}_{\lambda,\delta}^+ \text{ and } (t_2 u_1, t_2 v_1) \in \mathcal{N}_{\lambda,\delta}^-.$$

Particularly, $t_1 < t_2 = 1$. Since $\varphi'_{u,v}(t_1) = 0$ and $\varphi''(t_1) > 0$, so t_1 is local minimum of $\varphi_{u,v}$. Therefore there exists a $\hat{t} \in (t_1, 1]$ such that $I_{\lambda,\delta}(t_1 u_1, t_1 v_1) < I_{\lambda,\delta}(\hat{t} u_1, \hat{t} v_1)$. Hence

$$l_{\lambda,\delta} \leq I_{\lambda,\delta}(t_1 u_1, t_1 v_1) < I_{\lambda,\delta}(\hat{t} u_1, \hat{t} v_1) \leq I_{\lambda,\delta}(u_1, v_1) = l_{\lambda,\delta}$$

which contradicts that $(u_1, v_1) \in \mathcal{N}_{\lambda,\delta}^-$. \square

Lemma 4.7. *There exists a non negative local minimum of $I_{\lambda,\delta}$.*

Proof. Suppose (u_1, v_1) be as obtained in Theorem 4.6. Then it is also a local minimum for $I_{\lambda,\delta}$, the proof follows as [p. 291, [31]]. If $u_1, v_1 \geq 0$ then the lemma is proved. Else consider $(|u_1|, |v_1|)$ then by Lemma 3.3 we know that there exists a t_1 such that $(t_1 |u_1|, t_1 |v_1|) \in \mathcal{N}_{\lambda,\delta}^+$. Since $m_{|u_1|, |v_1|}(1) \leq m_{u_1, v_1}(1) = 2B(u_1, v_1) = 2B(|u_1|, |v_1|) = m_{|u_1|, |v_1|}(t_1)$ and $0 < m'_{u_1, v_1}(1) \leq m'_{|u_1|, |v_1|}(1)$. This implies $t_1 \geq 1$ and thus we have

$$I_{\lambda,\delta}(t_1 |u_1|, t_1 |v_1|) \leq I_{\lambda,\delta}(|u_1|, |v_1|) \leq I_{\lambda,\delta}(u_1, v_1) = \inf I_{\lambda,\delta}(\mathcal{N}_{\lambda,\delta}^+).$$

Hence we obtain a non negative local minimum of $I_{\lambda,\delta}$ over $\mathcal{N}_{\lambda,\delta}^+$. \square

We now prove the positivity of the solution (u_1, v_1) of $(P_{\lambda,\delta})$.

Proposition 4.8. *The non negative weak solution (u_1, v_1) of $(P_{\lambda,\delta})$ obtained in Lemma 4.7 is positive in Ω that is $u_1, v_1 > 0$ in Ω . Moreover for each compact subset K of Ω , there exists a $m_K > 0$ such that $u_1, v_1 \geq m_K$ in K .*

Proof. We divide the proof into two cases. Consider u_1 first and v_1 can be shown to be positive in exactly same way.

Case (1): Let $\frac{2_s^*}{(q-1)} > \frac{n}{2s}$ then there exists a sequence $\{u_\epsilon\}_{\epsilon>0} \subset C_c^\infty(\Omega)$ such that $u_\epsilon \rightarrow u_1$ in $L^{2_s^*}(\Omega)$ as $\epsilon \rightarrow 0$. That means $u_\epsilon^{q-1} \rightarrow u_1^{q-1}$ in $L^{\frac{2_s^*}{(q-1)}}(\Omega)$ as $\epsilon \rightarrow 0$. Now let

$$w_\epsilon := (-\Delta)^{-s}(\lambda u_\epsilon^{q-1}).$$

Then using Proposition 1.4(iii) of [27], we get that $\{w_\epsilon\}$ is a Cauchy sequence in $C^\beta(\mathbb{R}^n)$ where $\beta = \min \left\{ s, 2s - \frac{n(q-1)}{2_s^*} \right\}$ and

$$\|w_\epsilon\|_{C^\beta(\mathbb{R}^n)} \leq C \|u_\epsilon^{q-1}\|_{L^{\frac{2_s^*}{(q-1)}}(\Omega)}. \quad (4.26)$$

We know that there exists a $M_0 > 0$ such that $\sup_\epsilon \|w_\epsilon\|_\infty < M_0$, so by Lebesgue dominated convergence theorem we get

$$\limsup_{\epsilon > 0} \int_{\mathbb{R}^n} ((-\Delta)^s w_\epsilon) w_\epsilon \, dx < +\infty.$$

This implies that $\{w_\epsilon\}$ is bounded in X_0 , hence up to a subsequence, w_ϵ converges weakly to a $w \in X_0$ in X_0 as $\epsilon \rightarrow 0$. Furthermore w satisfies the equation

$$(-\Delta)^s w = \lambda u_1^{q-1} \text{ in } \Omega, \quad w = 0 \text{ in } \mathbb{R}^n \setminus \Omega.$$

In addition we have $w_\epsilon \rightarrow w$ in $C^\beta(\mathbb{R}^n)$ as $\epsilon \rightarrow 0$ and passing to the limit in (4.26) we obtain $w \in C(\bar{\Omega})$. Since (u_1, v_1) solves $(P_{\lambda, \delta})$ it is clear that u_1 satisfies

$$(-\Delta)^s u_1 \geq \lambda u_1^{q-1} \text{ in } \Omega, \quad u_1 = 0 \text{ in } \mathbb{R}^n \setminus \Omega.$$

Therefore $u_1 \geq w$ in Ω , thanks to comparison principle (refer Proposition 4.1 in [26]). Also now by strong maximum principle (refer [30]), we conclude that $w > 0$ in Ω and there exists a $m_K > 0$ for each K compact subset of Ω such that $w > m_K$ in K .

Case (2): Let $\frac{2_s^*}{(q-1)} \leq \frac{n}{2s}$ and consider the following iterative scheme

$$(-\Delta)^s w_k = \lambda w_{k-1}^{q-1} \text{ in } \Omega, \quad w_k = 0 \text{ in } \mathbb{R}^n \setminus \Omega$$

with $w_0 = u_1$. Then take $k = 1$ at first and let $\{w_{0,\epsilon}\} \subset C_c^\infty(\Omega)$ such that $w_{0,\epsilon} \rightarrow w_0 = u_1$ in $L^{2_s^*}(\Omega)$ as $\epsilon \rightarrow 0$ which means $w_{0,\epsilon}^{q-1} \rightarrow u_1^{q-1}$ in $L^{\frac{2_s^*}{q-1}}(\Omega)$ as $\epsilon \rightarrow 0$. We define

$$w_\epsilon^1 := (-\Delta)^{-s} (\lambda w_{0,\epsilon}^{q-1}).$$

Set $q_1 = \frac{2_s^*}{q-1}$ and we get using Proposition 1.4(ii) of [27] that $\{w_\epsilon^1\}$ is a Cauchy sequence in $L^{q_2}(\Omega)$ where $q_2 = \frac{nq_1}{n-2q_1s} > q_1$ and

$$\|w_\epsilon^1\|_{L^{q_2}(\Omega)} \leq C \|w_{0,\epsilon}^{q-1}\|_{L^{q_1}(\Omega)}. \quad (4.27)$$

Necessarily $w_\epsilon^1 \rightarrow w_1$ as $\epsilon \rightarrow 0$ in $L^{q_2}(\Omega)$ so passing on the limit as $\epsilon \rightarrow 0$ in (4.27) we obtain $w_1 \in L^{q_2}(\Omega)$. Proceeding similarly, at each stage we get $w_k \in L^{q_k}(\Omega)$ where $q_k = \frac{nq_{k-1}}{n-2q_{k-1}s}$ and note that $w_k \not\equiv 0$ for each k . Clearly $\{q_k\}$ forms an increasing sequence and the map $t \mapsto \frac{nt}{n-2st}$ has no fixed point. So obviously there exists a $k_0 > 0$ such that $q_{k_0} > \frac{n}{2s}$ and for this k_0 we get $w_{k_0+1} \in C^\beta(\mathbb{R}^n)$, by Proposition 1.4(iii) of [27]. By comparison principle we already know that $\{w_k\}$ forms a non increasing sequence and $u_1 \geq w_1$. Thus arguing same as Case (1) we get

$$u_1 \geq w_1 \geq w_2 \geq \dots \geq w_{k_0+1} > 0 \text{ in } \Omega.$$

Also there exists a $m_K > 0$ for each K compact subset of Ω such that $w_{k_0+1} > m_K$ in K . \square

This result suggests that there is no harm to consider (u_1, v_1) as positive (as this property of the first solution will be used further while proving the existence of second solution in the case $\mu \leq 4s$).

4.2. Second solution

Now, we establish the existence of second solution for $(P_{\lambda, \delta})$. We prove this by showing that minimum of $I_{\lambda, \delta}$ is achieved over $\mathcal{N}_{\lambda, \delta}^-$. We consider two cases separately that is when $\mu \leq 4s$ and when $\mu \geq 4s$. In

the first case we are able to show that when $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$, $(P_{\lambda,\delta})$ has two weak solutions whereas in the other case for $\mu > 4s$ we get another threshold Θ_0 which may be ‘less than or equal to’ Θ such that whenever $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta_0$, $(P_{\lambda,\delta})$ possesses two weak solutions.

Lemma 4.9. *If $\mu \leq 4s$ and $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$, then there exists $(w_0, z_0) \in Y \setminus \{(0, 0)\}$ such that $w_0, z_0 \geq 0$ and*

$$\sup_{t \geq 0} I_{\lambda,\delta}((u_1, v_1) + t(w_0, z_0)) < c_1 := I_{\lambda,\delta}(u_1, v_1) + \frac{n - \mu + 2s}{2n - \mu} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2n - \mu}{n - \mu + 2s}}.$$

Proof. Using (2.10), we can find $r_1 > 0$ such that

$$\int_{\mathbb{R}^{2n}} \frac{|u_\epsilon(x) - u_\epsilon(y)|^2}{|x - y|^{n+2s}} dx dy \leq \left((C(n, \mu))^{\frac{n-2s}{2n-\mu}} S_s^H \right)^{\frac{n}{2s}} + r_1 \epsilon^{n-2s}. \quad (4.28)$$

Also using Proposition 2.8, we can find $r_2 > 0$ such that

$$\int_{\Omega} (|x|^{-\mu} * |u_\epsilon|^{2_\mu^*}) |u_\epsilon|^{2_\mu^*} dx \geq C(n, \mu)^{\frac{n}{2s}} (S_s^H)^{\frac{2n-\mu}{2s}} - r_2 \epsilon^n. \quad (4.29)$$

From proof of Lemma 5.1 of [25], we know that for fixed ρ such that $1 < \rho < \frac{n}{n-2s}$ we have

$$\int_{\Omega} |u_\epsilon|^\rho \leq r_3 \epsilon^{\frac{(n-2s)\rho}{2}}, \quad (4.30)$$

where $r_3 > 0$ is an appropriate constant. Now let $0 < \epsilon < \delta$ then $u_\epsilon = U_\epsilon$ in $B_\epsilon(0)$.

Claim: There exists a constant $r_4 > 0$ such that

$$\int_{|x| \leq \epsilon} \int_{\Omega} \frac{|u_\epsilon(x)|^{2_\mu^*-1} |u_\epsilon(y)|^{2_\mu^*}}{|x - y|^\mu} dx dy \geq r_4 \epsilon^{\frac{n-2s}{2}}. \quad (4.31)$$

To show this, we split the left hand side of (4.31) into two integrals and estimate them separately. We recall the definition of u_ϵ and firstly consider

$$\begin{aligned} & \int_{|x| \leq \epsilon} \int_{|y| \leq \epsilon} \frac{|u_\epsilon(x)|^{2_\mu^*-1} |u_\epsilon(y)|^{2_\mu^*}}{|x - y|^\mu} dy dx \\ &= \frac{\alpha^{2_\mu^*-1}}{\|\tilde{u}\|_{L^{2_\mu^*}(\mathbb{R}^n)}^{2_\mu^*-1}} \int_{|x| \leq \epsilon} \int_{|y| \leq \epsilon} \frac{\epsilon^{\frac{(2s-n)(2_\mu^*-1)}{2}}}{|x - y|^\mu \left(\beta^2 + \left| \frac{x}{\epsilon S_s^{\frac{1}{2s}}} \right|^2 \right)^{\frac{(n-2s)(2_\mu^*-1)}{2}} \left(\beta^2 + \left| \frac{y}{\epsilon S_s^{\frac{1}{2s}}} \right|^2 \right)^{\frac{(n-2s)2_\mu^*}{2}}} dy dx \\ &\geq E_1 \int_{|x| \leq \epsilon} \int_{|y| \leq \epsilon} \frac{\epsilon^{\frac{(2s-n)(2_\mu^*-1)}{2} - \mu}}{(1 + |x|^2)^{\frac{(n-2s)(2_\mu^*-1)}{2}} (1 + |y|^2)^{\frac{(n-2s)2_\mu^*}{2}}} dy dx \\ &= E_1 \int_{|x| \leq 1} \int_{|y| \leq 1} \frac{\epsilon^{\frac{n-2s}{2}}}{(1 + |x|^2)^{\frac{(n-2s)(2_\mu^*-1)}{2}} (1 + |y|^2)^{\frac{(n-2s)2_\mu^*}{2}}} dy dx = O\left(\epsilon^{\frac{n-2s}{2}}\right) \end{aligned}$$

where $E_1 > 0$ is appropriate constant that changes value at each step. Secondly, in a similar manner we get

$$\begin{aligned}
& \int_{|x| \leq \epsilon} \int_{|y| > \epsilon} \frac{|u_\epsilon(y)|^{2_\mu^*} |u_\epsilon(x)|^{2_\mu^*-1}}{|x-y|^\mu} dy dx \\
&= \frac{\alpha^{2_\mu^*-1}}{\|\tilde{u}\|_{L^{2_\mu^*}(\mathbb{R}^n)}^{2_\mu^*-1}} \int_{|x| \leq \epsilon} \int_{|y| > \epsilon} \frac{\epsilon^{\frac{(2s-n)(2_\mu^*-1)}{2}}}{|x-y|^\mu \left(\beta^2 + \left| \frac{x}{\epsilon S_s^{2s}} \right|^2 \right)^{\frac{(n-2s)(2_\mu^*-1)}{2}} \left(\beta^2 + \left| \frac{y}{\epsilon S_s^{2s}} \right|^2 \right)^{\frac{(n-2s)2_\mu^*}{2}}} dy dx \\
&\geq E'_1 \int_{|x| \leq \epsilon} \int_{|y| > \epsilon} \frac{\epsilon^{\frac{(2s-n)(2_\mu^*-1)}{2}-\mu}}{(|y|+\epsilon)^\mu (1+|\frac{x}{\epsilon}|^2)^{\frac{(n-2s)(2_\mu^*-1)}{2}} (1+|\frac{y}{\epsilon}|^2)^{\frac{(n-2s)2_\mu^*}{2}}} dy dx \\
&= E'_1 \int_{|x| \leq 1} \int_{|y| > 1} \frac{\epsilon^{\frac{n-2s}{2}}}{(1+|x|^2)^{\frac{(n-2s)(2_\mu^*-1)}{2}} (1+|y|^2)^{\frac{(n-2s)2_\mu^*}{2}} (1+|y|)^\mu} dy dx = O\left(\epsilon^{\frac{n-2s}{2}}\right)
\end{aligned}$$

where $E'_1 > 0$ is appropriate constant that changes value at each step. This establishes our claim. We can find appropriate constants $\rho_{1,\lambda}, \rho_{1,\delta}, \rho_2 > 0$ such that the following inequalities hold:

(1) $\lambda \left(\frac{(c+d)^q}{q} - \frac{c^q}{q} - dc^{1-q} \right) \geq -\frac{\rho_{1,\lambda} d^\rho}{r_3}$ and $\delta \left(\frac{(c+d)^q}{q} - \frac{c^q}{q} - dc^{1-q} \right) \geq -\frac{\rho_{1,\delta} d^\rho}{r_3}$, for all $c > 0$, $d \geq 0$.

(2) For each $\epsilon > 0$, $m \leq u_1, v_1$ on compact subsets of Ω where $m > 0$ is a constant, we get

$$\begin{aligned}
& \frac{1}{2_\mu^*} B(u_1 + tu_\epsilon, v_1 + tv_\epsilon) - \frac{1}{2_\mu^*} B(u_1, v_1) - \int_{\Omega} \int_{\Omega} \frac{|u_1(y)|^{2_\mu^*} |v_1(x)|^{2_\mu^*-2} v_1(x) t u_\epsilon(x)}{|x-y|^\mu} dy dx \\
& - \int_{\Omega} \int_{\Omega} \frac{|v_1(y)|^{2_\mu^*} |u_1(x)|^{2_\mu^*-2} u_1(x) t v_\epsilon(x)}{|x-y|^\mu} dy dx \\
& \geq \frac{t^{2_\mu^*}}{2_\mu^*} B(u_\epsilon, v_\epsilon) + \frac{\rho_2 t^{2_\mu^*-1}}{(2_\mu^*-1)} \int_{|x| \leq \epsilon} \int_{\Omega} \frac{|u_\epsilon(y)|^{2_\mu^*} |u_\epsilon(x)|^{2_\mu^*-1}}{|x-y|^\mu} dy dx.
\end{aligned}$$

We remark that such an m exists because of Proposition 4.8. From Theorem 4.6 we know that (u_1, v_1) is a weak solution of $(P_{\lambda,\delta})$. Therefore, we have

$$\begin{aligned}
& I_{\lambda,\delta}((u_1, v_1) + t(u_\epsilon, v_\epsilon)) - I_{\lambda,\delta}(u_1, v_1) \\
&= I_{\lambda,\delta}((u_1, v_1) + t(u_\epsilon, v_\epsilon)) - I_{\lambda,\delta}(u_1, v_1) - t \left(\langle u_1, u_\epsilon \rangle + \langle v_1, v_\epsilon \rangle \right. \\
& \quad \left. - \int_{\Omega} (\lambda |u_1|^{q-2} u_1 u_\epsilon + \delta |v_1|^{q-2} v_1 v_\epsilon) dx - \int_{\Omega} (|x|^{-\mu} * |u_1|^{2_\mu^*}) |v_1|^{2_\mu^*-2} v_1 u_\epsilon dx \right. \\
& \quad \left. - \int_{\Omega} (|x|^{-\mu} * |v_1|^{2_\mu^*}) |u_1|^{2_\mu^*-2} u_1 v_\epsilon dx \right) \\
&= \frac{t^2}{2} C_s^n \|(u_\epsilon, v_\epsilon)\|^2 - \lambda \int_{\Omega} \left(\frac{|u_1 + tu_\epsilon|^q - |u_1|^q}{q} - t |u_1|^{q-2} u_1 u_\epsilon \right) dx
\end{aligned}$$

$$\begin{aligned}
& -\delta \int_{\Omega} \left(\frac{|v_1 + tu_{\epsilon}|^q - |v_1|^q}{q} - t|v_1|^{q-2}v_1u_{\epsilon} \right) dx - \left(\frac{B(u_1 + tu_{\epsilon}, v_1 + tu_{\epsilon}) - B(u_1, v_1)}{2_{\mu}^*} \right. \\
& \left. - \int_{\Omega} (|x|^{-\mu} * |u_1|^{2_{\mu}^*}) |v_1|^{2_{\mu}^*-2} v_1 tu_{\epsilon} dx - \int_{\Omega} (|x|^{-\mu} * |v_1|^{2_{\mu}^*}) |u_1|^{2_{\mu}^*-2} u_1 tu_{\epsilon} dx \right)
\end{aligned}$$

which on using inequality (2) with (4.28)–(4.31) gives

$$\begin{aligned}
& I_{\lambda, \delta}((u_1, v_1) + t(u_{\epsilon}, u_{\epsilon})) - I_{\lambda, \delta}(u_1, v_1) \\
& \leq t^2 C_s^n \left(C(n, \mu)^{\frac{n(n-2s)}{2s(2n-\mu)}} (S_s^H)^{\frac{n}{2s}} + r_1 \epsilon^{n-2s} \right) + (\rho_{1, \lambda} + \rho_{1, \delta}) t^{\rho} \epsilon^{\frac{(n-2s)\rho}{2}} \\
& \quad - \frac{t^{22_{\mu}^*}}{2_{\mu}^*} \left(C(n, \mu)^{\frac{n}{2s}} (S_s^H)^{\frac{2n-\mu}{2s}} - r_2 \epsilon^n \right) - \frac{t^{22_{\mu}^*-1} \rho_2}{(22_{\mu}^* - 1)} r_4 \epsilon^{\frac{n-2s}{2}}.
\end{aligned}$$

Now we define the function $h_{\epsilon} : [0, \infty) \rightarrow \mathbb{R}$ as

$$\begin{aligned}
h_{\epsilon}(t) &= t^2 C_s^n \left(C(n, \mu)^{\frac{n(n-2s)}{2s(2n-\mu)}} (S_s^H)^{\frac{n}{2s}} + r_1 \epsilon^{n-2s} \right) + (\rho_{1, \lambda} + \rho_{1, \delta}) t^{\rho} \epsilon^{\frac{(n-2s)\rho}{2}} \\
& \quad - \frac{t^{22_{\mu}^*}}{2_{\mu}^*} \left(C(n, \mu)^{\frac{n}{2s}} (S_s^H)^{\frac{2n-\mu}{2s}} - r_2 \epsilon^n \right) - \frac{t^{22_{\mu}^*-1} \rho_2}{(22_{\mu}^* - 1)} r_4 \epsilon^{\frac{n-2s}{2}}.
\end{aligned}$$

Then h_{ϵ} attains its maximum at

$$\begin{aligned}
t_{\epsilon} &= (C_s^n)^{\frac{n-2s}{2(n-\mu+2s)}} C(n, \mu)^{-\frac{n(n-2s)}{4s(2n-\mu)}} (S_s^H)^{-\frac{(n-2s)(n-\mu)}{4s(n-\mu+2s)}} \\
& \quad - \frac{\rho_2 r_4 (n-2s)}{4(n-\mu+2s)} C(n, \mu)^{-\frac{n}{2s}} (S_s^H)^{\frac{\mu-2n}{2s}} \epsilon^{\frac{n-2s}{2}} + o(\epsilon^{\frac{n-2s}{2}}).
\end{aligned}$$

Therefore we get

$$\begin{aligned}
& \sup_{t \geq 0} (I_{\lambda, \delta}((u_1, v_1) + t(u_{\epsilon}, u_{\epsilon})) - I_{\lambda, \delta}(u_1, v_1)) \\
& \leq \frac{n-\mu+2s}{2n-\mu} (C_s^n S_s^H)^{\frac{2n-\mu}{n-\mu+2s}} - \frac{\rho_2 r_4 \epsilon^{\frac{n-2s}{2}}}{22_{\mu}^* - 1} C(n, \mu)^{-\frac{n(3n-2\mu+2s)}{4s(2n-\mu)}} (S_s^H)^{\frac{(\mu-n)(3n-2\mu+2s)}{4s(n-\mu+2s)}} + o(\epsilon^{\frac{n-2s}{2}}) \\
& < \frac{n-\mu+2s}{(2n-\mu)} (C_s^n S_s^H)^{\frac{2n-\mu}{n-\mu+2s}} = \frac{n-\mu+2s}{(2n-\mu)} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2n-\mu}{n-\mu+2s}}.
\end{aligned}$$

Choosing $(w_0, z_0) = (u_{\epsilon}, u_{\epsilon})$, for appropriate choice of ϵ as shown above, we obtain the result. \square

Corollary 4.10. *It holds that $l_{\lambda, \delta}^{-} < c_1$.*

Proof. For each $(u, v) \in Y$, by Lemma 3.3 we know that there exists a $t_2(u, v) > 0$ (notation changed to show that t_2 depends on (u, v)) such that $t_2(u, v)(u, v) \in \mathcal{N}_{\lambda, \delta}^{-}$. We consider two sets

$$\begin{aligned}
U_1 &:= \left\{ (u, v) \in Y : \|(u, v)\| < t_2 \left(\frac{(u, v)}{\|(u, v)\|} \right) \right\} \text{ and} \\
U_2 &:= \left\{ (u, v) \in Y : \|(u, v)\| > t_2 \left(\frac{(u, v)}{\|(u, v)\|} \right) \right\}.
\end{aligned}$$

Claim: $Y \setminus \mathcal{N}_{\lambda, \delta}^{-} = U_1 \cup U_2$.

For any $(u, v) \in Y$ we define $(\hat{u}, \hat{v}) := \frac{(u, v)}{\|(u, v)\|}$. Now let $(u, v) \in \mathcal{N}_{\lambda, \delta}^-$. Then we know that there exists a $t_2(\hat{u}, \hat{v}) > 0$ such that $t_2(\hat{u}, \hat{v})(\hat{u}, \hat{v}) \in \mathcal{N}_{\lambda, \delta}^-$. But $(u, v) \in \mathcal{N}_{\lambda, \delta}^-$ implies that it must be that $\frac{t_2(\hat{u}, \hat{v})}{\|(u, v)\|} = 1$ which means $t_2(\hat{u}, \hat{v}) = \|(u, v)\|$. On the other hand, let $(u, v) \in Y$ be such that $t_2(\hat{u}, \hat{v}) = \|(u, v)\|$. By definition $t_2(\hat{u}, \hat{v})(\hat{u}, \hat{v}) \in \mathcal{N}_{\lambda, \delta}^-$ which implies that $(u, v) \in \mathcal{N}_{\lambda, \delta}^-$. This proves the claim.

Next let $(u, v) \in \mathcal{N}_{\lambda, \delta}^+$ then by Lemma 3.3 we know that there exists a $t_1(\hat{u}, \hat{v}) > 0$ such that $t_1(\hat{u}, \hat{v})(\hat{u}, \hat{v}) \in \mathcal{N}_{\lambda, \delta}^+$. But $(u, v) \in \mathcal{N}_{\lambda, \delta}^+$ implies that $\frac{t_1(\hat{u}, \hat{v})}{\|(u, v)\|} = 1$. This gives $t_2(\hat{u}, \hat{v}) > t_1(\hat{u}, \hat{v}) = \|(u, v)\|$ that is $(u, v) \in U_1$. Therefore $\mathcal{N}_{\lambda, \delta}^+ \subset U_1$ and thus $(u_1, v_1) \in U_1$.

We consider the map $\gamma_M \in C([0, 1], Y)$ defined by $\gamma_M(t) := (u_1, v_1) + tM(w_0, z_0)$ for $M > 0$, where (w_0, z_0) is defined Lemma 4.9. Clearly $\gamma(0) = (u_1, v_1)$ and $\gamma(1) = (u_1, v_1) + M(w_0, z_0)$. There exists a $R > 0$ such that $0 < t_2(u, v) < R$ on the set $\{(u, v) \in Y : \|(u, v)\| = 1\}$. Let us choose $M_0 > 0$ such that

$$M_0 \geq \frac{|R^2 - \|(u_1, v_1)\|^2|}{\|(w_0, z_0)\|^2}.$$

Then

$$\begin{aligned} \|(u_1, v_1) + M_0(w_0, z_0)\|^2 &\geq \|(u_1, v_1)\|^2 + M_0^2\|(w_0, z_0)\|^2 + O(M_0) \\ &\geq R^2 > \left(t_2 \left(\frac{(u_1, v_1) + M_0(w_0, z_0)}{\|(u_1, v_1) + M_0(w_0, z_0)\|} \right) \right)^2 \end{aligned}$$

which implies $(u_1, v_1) + M_0(w_0, z_0) \in U_2$. Now since γ_{M_0} is a continuous path starting from (u_1, v_1) to $(u_1, v_1) + M_0(w_0, z_0)$ and $Y \setminus \mathcal{N}_{\lambda, \delta}^- = U_1 \cup U_2$, there must exists a $\hat{t} > 0$ such that $\|(u_1, v_1) + \hat{t}M_0(w_0, z_0)\| = t_2 \left(\frac{(u_1, v_1) + \hat{t}M_0(w_0, z_0)}{\|(u_1, v_1) + \hat{t}M_0(w_0, z_0)\|} \right)$ that is $\gamma_{M_0}(\hat{t}) \in \mathcal{N}_{\lambda, \delta}^-$. Therefore $(u_1, v_1) + \hat{t}M_0(w_0, z_0) \in \mathcal{N}_{\lambda, \delta}^-$. Finally using Lemma 4.9 we obtain

$$l_{\lambda, \delta}^- \leq I_{\lambda, \delta}((u_1, v_1) + \hat{t}M_0(w_0, z_0)) \leq \sup_{t \geq 0} I_{\lambda, \delta}((u_1, v_1) + t(w_0, z_0)) < c_1.$$

This completes the proof. \square

Lemma 4.11. *If $\mu > 4s$ then there exists a $\Upsilon > 0$ such that whenever $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Upsilon$, we have*

$$l_{\lambda, \delta}^- < c_0 := \frac{n - \mu + 2s}{(2n - \mu)} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2n - \mu}{n - \mu + 2s}} - D_0 \left(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} \right)$$

where D_0 has been defined in Lemma 4.2.

Proof. Let $w_0 = z_0 = u_\epsilon$ and define

$$J_{\lambda, \delta}(u, v) = \frac{C_s^n}{2} \|(u, v)\|^2 - \frac{1}{2_\mu^*} B(u, v) \text{ and } f(t) = J_{\lambda, \delta}(tw_0, tz_0).$$

Then $f(0) = 0$ and $f(t) < 0$ if $t \in (0, T)$, $f(t) > 0$ if $t > T$ where $T = \left(\frac{2_\mu^* C_s^n \|(w_0, z_0)\|^2}{2B(w_0, z_0)} \right)^{\frac{1}{2(2_\mu^* - 1)}}$. It is next easy thing to verify that f attains its maximum at $t_* = \left(\frac{C_s^n \|w_0, z_0\|^2}{2B(w_0, z_0)} \right)^{\frac{1}{2(2_\mu^* - 1)}}$. Therefore using (2.10) and Proposition 2.8 we have

$$\begin{aligned}
& \sup_{t \geq 0} J_{\lambda, \delta}(tw_0, tz_0) \\
&= f(t_*) = \frac{C_s^n t_*^2}{2} \|(w_0, z_0)\|^2 - \frac{t_*^{2\frac{2^*}{\mu}}}{2^*} B(w_0, z_0) = \left(\frac{n - \mu + 2s}{2n - \mu} \right) \left(\frac{C_s^n \|u_\epsilon\|^2}{B(u_\epsilon, u_\epsilon)^{\frac{1}{2^*}}} \right)^{\frac{2^*}{2^* - 1}} \\
&\leq \left(\frac{n - \mu + 2s}{2n - \mu} \right) \left(\frac{C_s^n (C(n, \mu)^{\frac{n-2s}{2n-\mu}} (S_s^H)^{\frac{n}{2s}} + O(\epsilon^{n-2s}))}{\left(C(n, \mu)^{\frac{n}{2s}} (S_s^H)^{\frac{2n-\mu}{2s}} - O(\epsilon^n) \right)^{\frac{n-2s}{2n-\mu}}} \right)^{\frac{2^*}{2^* - 1}} \\
&\leq \left(\frac{n - \mu + 2s}{2n - \mu} \right) (C_s^n)^{\frac{2^*}{2^* - 1}} (S_s^H + O(\epsilon^{n-2s}))^{\frac{2^*}{2^* - 1}} \\
&= \left(\frac{n - \mu + 2s}{2n - \mu} \right) \left(\left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^*}{2^* - 1}} + O(\epsilon^{n-2s}) \right).
\end{aligned} \tag{4.32}$$

Recalling the definition of c_0 , we note that if $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Upsilon_1$ where $\Upsilon_1 > 0$ is chosen such that $c_0 > 0$ for example $\Upsilon_1 = \frac{n - \mu + 2s}{2D_0(2n - \mu)} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^*}{n - \mu + 2s}}$. Since $I_{\lambda, \delta}(tw_0, tz_0) \leq \frac{t^2}{2} \|(w_0, z_0)\|^2$ for $t \geq 0$, we can find $\bar{t} > 0$ such that $\sup_{t \in [0, \bar{t}]} I_{\lambda, \delta}(tw_0, tz_0) < c_0$ whenever $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Upsilon_1$. Let us define function $H_{\lambda, \delta} : Y \rightarrow \mathbb{R}$ as $H_{\lambda, \delta}(u, v) := \frac{1}{q} \int_{\Omega} (\lambda |u|^q + \delta |v|^q) dx$. Now using (4.32) we have

$$\begin{aligned}
\sup_{t \geq \bar{t}} I_{\lambda, \delta}(tw_0, tz_0) &= \sup_{t \geq \bar{t}} (J_{\lambda, \delta}(tw_0, tz_0) - H_{\lambda, \delta}(tw_0, tz_0)) \\
&\leq \left(\frac{n - \mu + 2s}{2n - \mu} \right) \left(\left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^*}{2^* - 1}} + O(\epsilon^{n-2s}) \right) - \frac{\bar{t}^q}{q} (\lambda + \delta) \int_{\mathbb{R}^n} |u_\epsilon|^q dx \\
&\leq \left(\frac{n - \mu + 2s}{2n - \mu} \right) \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^*}{2^* - 1}} + O(\epsilon^{n-2s}) - \frac{\bar{t}^q}{q} (\lambda + \delta) \int_0^{\delta_*} |u_\epsilon|^q dx
\end{aligned}$$

for any $\delta_* > 0$. Fix $\delta_* < \delta$ and letting $0 < \epsilon < \delta_*$ we estimate

$$\begin{aligned}
\int_{B(0, \delta_*)} |u_\epsilon|^q dx &= \int_{B(0, \delta_*)} |U_\epsilon|^q dx \geq C_1 |S_{n-1}| \epsilon^{n - \frac{(n-2s)q}{2}} \int_0^{\frac{\delta_*}{\epsilon}} \frac{r^{n-1}}{(1+r^2)^{\frac{(n-2s)q}{2}}} dr \\
&\geq C_2 |S_{n-1}| \epsilon^{n - \frac{(n-2s)q}{2}} \int_0^{\frac{\delta_*}{\epsilon}} r^{n-1-(n-2s)q} dr \\
&\geq C_2 |S_{n-1}| \epsilon^{n - \frac{(n-2s)q}{2}} \begin{cases} \int_1^{\frac{\delta_*}{\epsilon}} r^{n-1-(n-2s)q} dr & \text{if } n \leq (n-2s)q \\ \int_0^{\frac{\delta_*}{\epsilon}} r^{n-1-(n-2s)q} dr & \text{if } n > (n-2s)q \end{cases} \\
&\simeq C_3 \begin{cases} \epsilon^{n - \frac{(n-2s)q}{2}}, & \text{if } n < (n-2s)q \\ \epsilon^{\frac{n}{2}} |\log \epsilon|, & \text{if } n = (n-2s)q \\ \epsilon^{\frac{(n-2s)q}{2}}, & \text{if } n > (n-2s)q \end{cases}
\end{aligned}$$

where C_1, C_2 and C_3 are appropriate positive constants. Therefore

$$\sup_{t \geq t_0} I_{\lambda, \delta}(tw_0, tz_0) \leq \left(\frac{n - \mu + 2s}{2n - \mu} \right) \left(\frac{\tilde{S}_s^H}{2} \right)^{\frac{2^* \mu}{2^* \mu - 1}} + \begin{cases} O(\epsilon^{n-2s}) - (\lambda + \delta)O(\epsilon^{n - \frac{(n-2s)q}{2}}), & \text{if } n < (n-2s)q \\ O(\epsilon^{n-2s}) - (\lambda + \delta)O(\epsilon^{\frac{n}{2}} |\log \epsilon|), & \text{if } n = (n-2s)q \\ O(\epsilon^{n-2s}) - (\lambda + \delta)O(\epsilon^{\frac{(n-2s)q}{2}}), & \text{if } n > (n-2s)q. \end{cases}$$

Since $q/2 < 1$, this implies that when $n > (n-2s)q$ and $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Upsilon := \min\{\Upsilon_1, \Theta\}$, for small enough $\epsilon > 0$ we can get that

$$\sup_{t \geq t_0} I_{\lambda, \delta}(tw_0, tz_0) < \left(\frac{n - \mu + 2s}{2n - \mu} \right) \left(\frac{\tilde{S}_s^H}{2} \right)^{\frac{2^* \mu}{2^* \mu - 1}} - D_0 \left(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} \right) = c_0.$$

In other case, for $\epsilon = (\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{1}{n-2s}} \leq \delta_*$, we get

$$\sup_{t \geq t_0} I_{\lambda, \delta}(tw_0, tz_0) \leq \left(\frac{n - \mu + 2s}{2n - \mu} \right) \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^* \mu}{2^* \mu - 1}} + \begin{cases} C(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}}) - C(\lambda + \delta)(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{1}{n-2s}(n - \frac{(n-2s)q}{2})}, & \text{if } n < (n-2s)q \\ C(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}}) - C(\lambda + \delta)(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{1}{2(n-2s)}} |\log((\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{1}{n-2s}})|, & \text{if } n = (n-2s)q. \end{cases}$$

Let $n < (n-2s)q$ then $1 + \frac{2}{(2-q)(n-2s)} \left(n - \frac{(n-2s)q}{2} \right) < \frac{2}{2-q}$ which implies that we can choose a $\Upsilon_2 > 0$ small enough such that if $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Upsilon_2$ then

$$C \left(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} \right) - C(\lambda + \delta)(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{1}{n-2s}(n - \frac{(n-2s)q}{2})} < -D_0 \left(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} \right).$$

As $\lambda, \delta \rightarrow 0$, $|\log((\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{1}{n-2s}})| \rightarrow \infty$ so in case $n = (n-2s)q$ we can obtain a $\Upsilon_2 > 0$ small enough such that

$$C(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}}) - C(\lambda + \delta)(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{1}{2(n-2s)}} |\log((\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}})^{\frac{1}{n-2s}})| < -D_0 \left(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} \right).$$

Setting $\Upsilon = \min\{\Upsilon_1, \Upsilon_2, \delta_*^{n-2s}\} > 0$ we finally get that

$$\sup_{t \geq 0} I_{\lambda, \delta}(tw_0, tz_0) < c_0$$

whenever $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Upsilon$. To prove the last part of the Lemma, we note that there exists $t_2 > 0$ such that $(t_2 w_0, t_2 z_0) \in \mathcal{N}_{\lambda, \delta}^-$ and

$$l_{\lambda, \delta}^- \leq I_{\lambda, \delta}(t_2 w_0, t_2 z_0) \leq \sup_{t \geq 0} I_{\lambda, \delta}(tw_0, tz_0) < c_0$$

when $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Upsilon$. This concludes the proof. \square

Before proving the existence of second solution, we make a remark at this stage.

Remark 4.12. Using Lemma 4.2 it is easy to see that $c_1 > c_0$, where c_1 is defined in Lemma 4.9 and c_0 is defined in Lemma 4.11.

Theorem 4.13. *There exists a $(u_2, v_2) \in \mathcal{N}_{\lambda, \delta}^-$ such that $I_{\lambda, \delta}(u_2, v_2) = l_{\lambda, \delta}^-$ in each of the following cases:*

- (i) $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$ when $\mu \leq 4s$ and
- (ii) $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta_0 := \min\{\Theta, \Upsilon\}$ when $\mu > 4s$.

Moreover, (u_2, v_2) is a weak solution of $(P_{\lambda, \delta})$.

Proof. Let $\{(u_k, v_k)\} \subset \mathcal{N}_{\lambda, \delta}^-$ be a minimizing sequence such that $\lim_{k \rightarrow \infty} I_{\lambda, \delta}(u_k, v_k) = l_{\lambda, \delta}^-$. By Lemma 4.3(ii), we know that $\{(u_k, v_k)\}$ is a bounded sequence in X_0 . Hence there exists a $(u_2, v_2) \in Y$ such that, upto a subsequence, $(u_k, v_k) \rightharpoonup (u_2, v_2)$ weakly in X_0 as $k \rightarrow \infty$.

Claim (1): As $k \rightarrow \infty$, $u_k \rightarrow u_2$ and $v_k \rightarrow v_2$ strongly in X_0 .

If not, we define $z_k = u_k - u_2$ and $w_k = v_k - v_2$ and assume that as $k \rightarrow \infty$

$$\|(z_k, w_k)\|^2 \rightarrow c^2 \text{ and } B(z_k, w_k) \rightarrow d^{22^*}_\mu,$$

for some $c, d \neq 0$. Then as $k \rightarrow \infty$ we have

$$\|(u_k, v_k)\|^2 = \|(z_k, w_k)\|^2 + \|(u_2, v_2)\|^2 + o_k(1).$$

Before proving claim (1) we state and prove the following.

Claim (2): As $k \rightarrow \infty$, $B(u_k, v_k) - B(z_k, w_k) \rightarrow B(u_2, v_2)$.

From fractional Sobolev embedding we have that

$$|z_k|^{2^*_\mu} - |u_k|^{2^*_\mu} \rightharpoonup |u_2|^{2^*_\mu} \text{ and } |w_k|^{2^*_\mu} - |v_k|^{2^*_\mu} \rightharpoonup |v_2|^{2^*_\mu} \text{ in } L^{\frac{2n}{2n-\mu}}(\mathbb{R}^n).$$

By Proposition 2.1, we have

$$|x|^{-\mu} * (|z_k|^{2^*_\mu} - |u_k|^{2^*_\mu}) \rightharpoonup |x|^{-\mu} * |u_2|^{2^*_\mu} \text{ and } |x|^{-\mu} * (|w_k|^{2^*_\mu} - |v_k|^{2^*_\mu}) \rightharpoonup |x|^{-\mu} * |v_2|^{2^*_\mu} \text{ in } L^{\frac{2n}{\mu}}(\mathbb{R}^n).$$

Also from boundedness of $\{u_k\}$ and $\{v_k\}$ in $L^{\frac{2n}{n-2s}}(\mathbb{R}^n)$ we know that $|z_k|^{2^*_\mu} \rightharpoonup 0$ and $|w_k|^{2^*_\mu} \rightharpoonup 0$ in $L^{\frac{2n}{2n-\mu}}(\mathbb{R}^n)$. This gives $\int_{\Omega} (|x|^{-\mu} * (|z_k|^{2^*_\mu} - |u_k|^{2^*_\mu})) |w_k|^{2^*_\mu} \rightarrow 0$ and $\int_{\Omega} (|x|^{-\mu} * (|w_k|^{2^*_\mu} - |v_k|^{2^*_\mu})) |z_k|^{2^*_\mu} \rightarrow 0$ as $k \rightarrow \infty$. This altogether proves claim (2) because we can write

$$\begin{aligned} B(u_k, v_k) - B(z_k, w_k) &= B(u_k - z_k, v_k - w_k) + \int_{\Omega} (|x|^{-\mu} * (|v_k|^{2^*_\mu} - |w_k|^{2^*_\mu})) |z_k|^{2^*_\mu} \\ &\quad + \int_{\Omega} (|x|^{-\mu} * (|u_k|^{2^*_\mu} - |z_k|^{2^*_\mu})) |w_k|^{2^*_\mu}. \end{aligned}$$

Since $\{(u_k, v_k)\} \subset \mathcal{N}_{\lambda, \delta}^-$, $\lim_{k \rightarrow \infty} \varphi'_{u_k, v_k}(1) = 0$. This gives

$$\varphi'_{u_2, v_2}(1) + C_s^n c^2 - 2d^{22^*}_\mu = 0. \quad (4.33)$$

Claim (3): (u_2, v_2) is non-trivial.

Suppose not and $u_2 \equiv 0 \equiv v_2$. This implies $c \neq 0$ because of Lemma 4.3(ii). Also using definition of \tilde{S}_s^H and $C_s^n c^2 = 2d^{22^*}_\mu$ (by (4.33)), we get

$$\frac{C_s^n c^2}{2} \geq \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{22^*}{2(2^*_{\mu}-1)}}.$$

Therefore

$$\begin{aligned} l_{\lambda,\delta}^- &= \lim_{k \rightarrow \infty} I_{\lambda,\delta}(u_k, v_k) = I_{\lambda,\delta}(0, 0) + \frac{C_s^n c^2}{2} - \frac{2d^{22^*_{\mu}}}{22^*_{\mu}} \\ &= \frac{C_s^n c^2}{2} \left(1 - \frac{1}{2^*_{\mu}} \right) \geq \left(\frac{n - \mu + 2s}{2n - \mu} \right) \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2n - \mu}{n - \mu + 2s}}. \end{aligned} \quad (4.34)$$

If $\mu \leq 4s$, then using (4.34) with Lemma 4.9, we have that $I_{\lambda,\delta}(u_1, v_1) > 0$ but this is a contradiction to $I_{\lambda,\delta}(u_1, v_1) = l_{\lambda,\delta}^+ < 0$ (by Lemma 4.3(i)). Otherwise if $\mu > 4s$, then using (4.34) with Lemma 4.11, we get $-D_0(\lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}}) \geq 0$ which is again a contradiction. This proves claim (3). Since $(u_2, v_2) \in Y \setminus \{(0, 0)\}$ and $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$ for both the cases $\mu \leq 4s$ as well as $\mu > 4s$, by Lemma 3.3 we know that there exist t_1, t_2 such that $0 < t_1 < t_2$, $t_1(u_2, v_2) \in \mathcal{N}_{\lambda,\delta}^+$ and $t_2(u_2, v_2) \in \mathcal{N}_{\lambda,\delta}^-$. That is $\varphi'_{u_2, v_2}(t_1) = 0 = \varphi'_{u_2, v_2}(t_2)$. Let us define the following two functions

$$f(t) = \frac{C_s^n c^2 t^2}{2} - \frac{d^{22^*_{\mu}} t^{22^*_{\mu}}}{2^*_{\mu}} \text{ and } g(t) = \varphi_{u_2, v_2}(t) + f(t).$$

Then we consider the three cases as below:

- (i) $t_2 < 1$,
- (ii) $t_2 \geq 1$ and $d > 0$,
- (iii) $t_2 \geq 1$ and $d = 0$.

(i) Using (4.33) we get $g'(1) = \varphi'_{u_2, v_2}(1) + C_s^n c^2 - 2d^{22^*_{\mu}} = 0$. Since $\{(u_k, v_k)\} \subset \mathcal{N}_{\lambda,\delta}^-$, for all $t > 0$ we get

$$\varphi_{u_k, v_k}(t) \leq \varphi_{u_k, v_k}(1) \quad (4.35)$$

Since $g(t) = \lim_{k \rightarrow \infty} \varphi_{u_k, v_k}(t)$ and passing to the limit as $k \rightarrow \infty$ in (4.35), we obtain $g(t) \leq g(1)$, for $t > 0$. Therefore

$$l_{\lambda,\delta}^- = \lim_{k \rightarrow \infty} \varphi_{u_k, v_k}(1) = g(1) > g(t_2) \geq I_{\lambda,\delta}(t_2 u_2, t_2 v_2) + \frac{t_2^2}{2} (C_s^n c^2 - 2d^{22^*_{\mu}}) > I_{\lambda,\delta}(t_2 u_2, t_2 v_2) \geq l_{\lambda,\delta}^-$$

which is a contradiction.

(ii) We define $t_* = \left(\frac{C_s^n c^2}{2d^{22^*_{\mu}}} \right)^{\frac{1}{22^*_{\mu}-2}}$ and then it is easy to compute that $f(t)$ attains its maximum at $t = t_*$. Also we compute and find that

$$f(t_*) = \frac{n - \mu + 2s}{2n - \mu} \left(\frac{C_s^n c^2}{2d^{22^*_{\mu}}} \right)^{\frac{2^*_{\mu}}{2^*_{\mu}-1}} \geq \frac{n - \mu + 2s}{2n - \mu} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^*_{\mu}}{2^*_{\mu}-1}}.$$

Moreover $f'(t) = t(C_s^n c^2 - 2d^{22^*_{\mu}} t^{22^*_{\mu}-2}) > 0$ if $t \in (0, t_*)$ and $f'(t) \leq 0$ if $t \geq t_*$. Moreover $g(1) = \max_{t > 0} g(t) \geq g(t_*)$. So if $t_* \leq 1$ then

$$\begin{aligned}
l_{\lambda,\delta}^- &= g(1) \geq g(t_*) = I_{\lambda,\delta}(t_* u_2, t_* v_2) + f(t_*) \geq I_{\lambda,\delta}(t_1 u_2, t_1 v_2) + \frac{n-\mu+2s}{2n-\mu} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^*_\mu}{2^*_\mu-1}} \\
&\geq l_{\lambda,\delta}^+ + \frac{n-\mu+2s}{2n-\mu} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^*_\mu}{2^*_\mu-1}} \geq I_{\lambda,\delta}(u_1, v_1) + \frac{n-\mu+2s}{2n-\mu} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^*_\mu}{2^*_\mu-1}} = c_1
\end{aligned} \tag{4.36}$$

which is a contradiction to Lemma 4.9 in the case $\mu \leq 4s$. Whereas when $\mu > 4s$, using Remark 4.12 and (4.36) we get that $l_{\lambda,\delta}^- \geq c_1 > c_0$ which is a contradiction to Lemma 4.11. Therefore we must have $t_* > 1$. Since $g'(t) \leq 0$ for $t \geq 1$, whenever $t \in [1, t_*]$ we get $\varphi'_{u_2, v_2}(t) \leq -f'(t) \leq 0$. This gives either $t_* \leq t_1$ or $t_2 = 1$. If $t_* \leq t_1$ then (4.36) holds true and we arrive at a contradiction whereas if $t_2 = 1$ then $(u_2, v_2) \in \mathcal{N}_{\lambda,\delta}^-$ which implies $C_s^n c^2 = 2d^{2^*_\mu}$ (by (4.33)). This gives

$$\begin{aligned}
l_{\lambda,\delta}^- &= g(1) = I_{\lambda,\delta}(u_2, v_2) + d^{2^*_\mu} \left(1 - \frac{1}{2^*_\mu} \right) \geq I_{\lambda,\delta}(u_2, v_2) + \frac{n-\mu+2s}{2n-\mu} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^*_\mu}{2^*_\mu-1}} \\
&\geq I_{\lambda,\delta}(t_1 u_2, t_1 v_2) + \frac{n-\mu+2s}{2n-\mu} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^*_\mu}{2^*_\mu-1}} \geq I_{\lambda,\delta}(u_1, v_1) + \frac{n-\mu+2s}{2n-\mu} \left(\frac{C_s^n \tilde{S}_s^H}{2} \right)^{\frac{2^*_\mu}{2^*_\mu-1}}
\end{aligned}$$

which contradicts Lemma 4.9 in the case $\mu \leq 4s$. Whereas when $\mu > 4s$, using Remark 4.12 and (4.36) we get that $l_{\lambda,\delta}^- \geq c_1 > c_0$ which is a contradiction to Lemma 4.11.

Hence, only possibility is that (iii) holds true that is $t_2 \geq 1$ and $d = 0$. If $c \neq 0$ then (4.33) implies $\varphi'_{u_2, v_2}(1) = -c^2 < 0$ and also $\varphi''_{u_2, v_2}(1) < 0$ which is a contradiction since $t_2 \geq 1$. Thus $c = 0$ and this proves claim (1). Therefore $I_{\lambda,\delta}(u_2, v_2) = l_{\lambda,\delta}^-$ and obviously $(u_2, v_2) \in \mathcal{N}_{\lambda,\delta}^-$. Finally, (u_2, v_2) is a weak solution of $(P_{\lambda,\delta})$ follows from Lemma 3.2. \square

4.3. Proof of main theorem

Proof of Theorem 1.1. By Theorem 4.6 and 4.13 we know that $(P_{\lambda,\delta})$ has two solutions $(u_1, v_1) \in \mathcal{N}_{\lambda,\delta}^+$ and $(u_2, v_2) \in \mathcal{N}_{\lambda,\delta}^-$ whenever $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta$ if $\mu \leq 4s$ and whenever $0 < \lambda^{\frac{2}{2-q}} + \delta^{\frac{2}{2-q}} < \Theta_0$ if $\mu > 4s$. Obviously they are distinct solutions because $\mathcal{N}_{\lambda,\delta}^+ \cap \mathcal{N}_{\lambda,\delta}^- = \emptyset$. The proof is then completed using Proposition 4.8. \square

Acknowledgments

The authors were funded by IFCAM (Indo-French Centre for Applied Mathematics) UMI CNRS 3494 under the project “Singular phenomena in reaction diffusion equations and in conservation laws”.

References

- [1] C.O. Alves, M.G. Figueiredo, M. Yang, Existence of solutions for a nonlinear Choquard equation with potential vanishing at infinity, *Adv. Nonlinear Anal.* 5 (4) (2016) 331–345.
- [2] C.O. Alves, F. Gao, M. Squassina, M. Yang, Singularly perturbed critical Choquard equations, *J. Differential Equations* 263 (2017) 3943–3988.
- [3] D. Applebaum, Lévy processes—from probability to finance and quantum groups, *Notices Amer. Math. Soc.* 51 (11) (2004) 1336–1347.
- [4] C. Bonanno, P. D’Avenia, M. Ghimenti, M. Squassina, Soliton dynamics for the generalized Choquard equations, *J. Math. Anal. Appl.* 417 (2014) 180–199.
- [5] B. Buffoni, L. Jeanjean, C.A. Stuart, Existence of a nontrivial solution to a strongly indefinite semilinear equation, *Proc. Amer. Math. Soc.* 119 (1) (1993) 179–186.
- [6] Y.-H. Chen, C. Liu, Ground state solutions for non-autonomous fractional Choquard equations, *Nonlinearity* 29 (2016) 1827–1842.
- [7] W. Chen, M. Squassina, Critical nonlocal systems with concave-convex powers, *Adv. Nonlinear Stud.* 16 (4) (2016) 821–842.

- [8] W. Choi, On strongly indefinite systems involving the fractional Laplacian, *Nonlinear Anal.* 120 (2015) 127–153.
- [9] P. D'Avenia, G. Siciliano, M. Squassina, On fractional Choquard equations, *Math. Models Methods Appl. Sci.* 25 (8) (2015) 1447–1476.
- [10] P. D'Avenia, G. Siciliano, M. Squassina, Existence results for a doubly nonlocal equation, *São Paulo J. Math. Sci.* 9 (2) (2015) 311–324.
- [11] H. Fan, Multiple positive solutions for a fractional elliptic system with critical nonlinearities, *Bound. Value Probl.* 2016 (2016) 196, <https://doi.org/10.1186/s13661-016-0703-7>.
- [12] L.F.O. Faria, O.H. Miyagaki, F.R. Pereira, M. Squassina, C. Zhang, The Brezis–Nirenberg problem for nonlocal systems, *Adv. Nonlinear Anal.* 5 (1) (2015) 85–103.
- [13] F. Gao, M. Yang, On nonlocal Choquard equations with Hardy–Littlewood–Sobolev critical exponents, *J. Math. Anal. Appl.* 448 (2) (2017) 1006–1041.
- [14] F. Gao, M. Yang, On the Brezis–Nirenberg type critical problem for nonlinear Choquard equation, *Sci. China Math.* 61 (7) (2018) 1219–1242, <https://doi.org/10.1007/s11425-016-9067-5>.
- [15] M. Ghimenti, J. Van Schaftingen, Nodal solutions for the Choquard equation, *J. Funct. Anal.* 271 (2016) 107–135.
- [16] M. Ghimenti, V. Moroz, J.V. Schaftingen, Least action nodal solutions for the quadratic Choquard equation, *Proc. Amer. Math. Soc.* 145 (2) (2017) 737–747.
- [17] J. Giacomoni, P.K. Mishra, K. Sreenadh, Critical growth fractional elliptic systems with exponential nonlinearity, *Nonlinear Anal.* 136 (2016) 117–135.
- [18] J. Giacomoni, T. Mukherjee, K. Sreenadh, Positive solutions of fractional elliptic equation with critical and singular nonlinearity, *Adv. Nonlinear Anal.* 6 (3) (2017) 327–354.
- [19] Z. Guo, S. Luo, W. Zou, On critical systems involving fractional Laplacian, *J. Math. Anal. Appl.* 446 (1) (2017) 681–706.
- [20] X. He, M. Squassina, W. Zou, The Nehari manifold for fractional systems involving critical nonlinearities, *Commun. Pure Appl. Anal.* 15 (4) (2016) 1285–1308.
- [21] E. Lieb, M. Loss, *Analysis*, Graduate Studies in Mathematics, AMS, Providence, Rhode Island, 2001.
- [22] D. Lü, G. Xu, On nonlinear fractional Schrödinger equations with Hartree-type nonlinearity, *Appl. Anal.* 97 (2) (2018) 255–273, <https://doi.org/10.1080/00036811.2016.1260708>.
- [23] G. Molica Bisci, V.D. Rădulescu, R. Servadei, *Variational Methods for Nonlocal Fractional Problems*, Encyclopedia of Mathematics and Its Applications, Cambridge University Press, Cambridge, 2016.
- [24] V. Moroz, J.V. Schaftingen, A guide to the Choquard equation, *J. Fixed Point Theory Appl.* 19 (1) (2017) 773–813.
- [25] T. Mukherjee, K. Sreenadh, Critical growth fractional elliptic problem with singular nonlinearities, *Electron. J. Differential Equations* 54 (2016) 1–23.
- [26] X. Ros-Oton, Nonlocal equations in bounded domains: a survey, *Publ. Mat.* 60 (2016) 3–26.
- [27] X. Ros-Oton, J. Serra, The extremal solution for the fractional Laplacian, *Calc. Var.* 50 (2014) 723–750.
- [28] R. Servadei, E. Valdinoci, The Brezis–Nirenberg result for the fractional Laplacian, *Trans. Amer. Math. Soc.* 367 (1) (2015) 67–102.
- [29] Z. Shen, F. Gao, M. Yang, Multiple solutions for nonhomogeneous Choquard equation involving Hardy–Littlewood–Sobolev critical exponent, *Z. Angew. Math. Phys.* (2017), <https://doi.org/10.1007/s00033-017-0806-8>.
- [30] L. Silvestre, Regularity of the obstacle problem for a fractional power of the Laplace operator, *Comm. Pure Appl. Math.* 60 (2007) 67–112.
- [31] G. Tarantello, On nonhomogeneous elliptic equations involving critical Sobolev exponent, *Ann. Inst. H. Poincaré Anal. Non Linéaire* 9 (1992) 281–304.
- [32] K. Wang, J. Wei, On the uniqueness of solutions of a nonlocal elliptic system, *Math. Ann.* 365 (1–2) (2016) 105–153.
- [33] M. Xiang, B. Zhang, V.D. Rădulescu, Existence of solutions for perturbed fractional p -Laplacian equations, *J. Differential Equations* 260 (2) (2016) 1392–1413.
- [34] X. Zhang, B. Zhang, M. Xiang, Ground states for fractional Schrödinger equations involving a critical nonlinearity, *Adv. Nonlinear Anal.* 5 (3) (2016) 293–314.